

Resampling in Sequential Monte Carlo

Suzie Brown

12 November 2019

Outline

1. Sequential Monte Carlo
2. How to resample
3. Properties of resampling schemes
4. Link with genealogies


Sequential Monte Carlo

Motivation

- ▶ Want to approximate a sequence of target measures $(\eta_n)_{n \in \mathbb{N}}$
- ▶ Use a system of N particles with dynamics 'mimicking' the target
- ▶ Approximate the measures η_n by (random) empirical measures η_n^N consisting of atoms at the particle positions

Sequential Monte Carlo

Illustration



Resampling

Motivation

- ▶ Resampling is necessary to prevent *weight degeneracy*
- ▶ But resampling causes *ancestral degeneracy*

Resampling

Motivation

- ▶ Resampling is necessary to prevent *weight degeneracy*
- ▶ But resampling causes *ancestral degeneracy*
- ▶ Strategy: resample in a way that minimises 'unnecessary coalescences'

Resampling

Definition

We will take valid resampling schemes to be those satisfying

- ▶ The total number of particles N remains fixed
- ▶ The particles after resampling are equally weighted
- ▶ The scheme is unbiased: the expected number of offspring of particle i is equal to Nw_i for each i

Multinomial Resampling¹

Definition

Parental indices $a_i \in \{1, \dots, N\}$:

$$(a_i \mid w_{1:N}) \stackrel{iid}{\sim} \text{Categorical}(N, w_{1:N})$$


Offspring numbers $v_i \in \{0, \dots, N\}$ such that $\sum v_i = N$:

$$(v_{1:N} \mid w_{1:N}) \sim \text{Multinomial}(1 : N, w_{1:N})$$

¹Efron & Tibshirani (1994) 'An introduction to the bootstrap'


Multinomial Resampling

Inversion Sampling




Multinomial Resampling

Inversion Sampling



Multinomial Resampling

Inversion Sampling



Residual Resampling^{2,3}

Definition

1. Deterministically assign $\lfloor Nw_i \rfloor$ offspring to particle i ; $i=1, \dots, N$
2. There are $R := N - \sum_{i=1}^N \lfloor Nw_i \rfloor$ offspring still to be assigned
3. Assign these randomly according to the residual weights $r_i := \frac{1}{R}(Nw_i - \lfloor Nw_i \rfloor)$

²Liu & Chen (1998) 'Sequential Monte Carlo methods for dynamic systems'

³Whitley (1994) 'A genetic algorithm tutorial'

Residual Resampling

Definition


If residuals are assigned using multinomial resampling, offspring counts are distributed

$$v_{1:N} \stackrel{d}{=} \lfloor Nw_{1:N} \rfloor + \text{Multinomial}(R, r_{1:N})$$


Residual Resampling

Illustration

$w_1 = 0.28$ 


$w_2 = 0.12$ 

$w_3 = 0.51$ 

$w_4 = 0.09$ 


Residual Resampling

Illustration



Residual Resampling

Illustration



Residual Resampling

Illustration

● $r_1 \propto 0.03$ ■


$r_2 \propto 0.12$ ■■■

●● $r_3 \propto 0.01$ ■


$r_4 \propto 0.09$ ■■■

Residual Resampling

Illustration

● $r_1 = 0.12$ 

$r_2 = 0.48$ 

● ● $r_3 = 0.04$ 

$r_4 = 0.36$ 

Stratified Resampling⁴

Definition

Draw uniformly from each stratified interval

$$U_i \sim \text{Uniform} \left(\frac{i-1}{N}, \frac{i}{N} \right); \quad i = 1, \dots, N$$


and determine the parental indices by inversion

$$a_i = \inf \left\{ k : \sum_{j=1}^k w_j \geq U_i \right\}$$

⁴Kitagawa (1996) 'Monte Carlo filter and smoother for non-Gaussian nonlinear state space models'


Stratified Resampling

Inversion Sampling



Stratified Resampling

Inversion Sampling



Systematic Resampling^{5,6}

Definition

Draw uniformly from $[0, \frac{1}{N}]$, and add multiples of $\frac{1}{N}$

$$U_1 \sim \text{Uniform} \left(0, \frac{1}{N} \right)$$

$$U_i = U_1 + \frac{i-1}{N}; \quad i = 2, \dots, N$$

and determine the parental indices by inversion


$$a_i = \inf \left\{ k : \sum_{j=1}^k w_j \geq U_i \right\}$$

⁵Carpenter, Clifford & Fearnhead (1999) 'Improved particle filter for nonlinear problems'

⁶Whitley (1994) 'A genetic algorithm tutorial'


Systematic Resampling

Inversion Sampling



Systematic Resampling

Inversion Sampling



Coalescence Probability

Definition

The probability that a randomly chosen pair of particles at generation t share a common ancestor at generation $(t - 1)$

$$c_N = \frac{1}{N(N-1)} \sum_{i=1}^N v_i(v_i - 1)$$

Coalescence Probability

Example

Consider the case where we have only two particles ($N = 2$)


$$c_2 = \frac{1}{2} [v_1(v_1 - 1) + v_2(v_2 - 1)]$$

The expectation of c_2 conditional on knowing the weights (w_1, w_2) is

$$\begin{aligned} c_2 &= \frac{1}{2} \mathbb{E}[v_1(v_1 - 1) \mid w_{1:2}] + \frac{1}{2} \mathbb{E}[v_2(v_2 - 1) \mid w_{1:2}] \\ &= \mathbb{P}[v_1 = 2 \mid w_{1:2}] + \mathbb{P}[v_2 = 2 \mid w_{1:2}] \end{aligned}$$

Coalescence Probability

Example



References I