

# Asymptotic Genealogies of Sequential Monte Carlo Algorithms


Suzie Brown

26 April 2019

# State space models

Hidden process  $X_0, \dots, X_T \in \mathcal{X}$


Observable process  $Y_0, \dots, Y_T \in \mathcal{Y}$



# State space models

Hidden process  $X_0, \dots, X_T \in \mathcal{X}$

Observable process  $Y_0, \dots, Y_T \in \mathcal{Y}$




$$X_0 \sim \mu(\cdot)$$

$$X_{t+1} \mid (X_t = x_t) \sim f(\cdot \mid x_t)$$

$$Y_t \mid (X_t = x_t) \sim g(\cdot \mid x_t)$$

# Target tracking



use noisy radar data to infer position/trajectory of aircraft:

- $f$  models how aircraft moves
- $g$  models uncertainty in radar measurements

# Inference problems

**Filtering:** where is it now?  $p(x_t|y_{0:t})$

**Prediction:** where will it go next?  $p(x_{t+1}|y_{0:t})$

**Smoothing:** where has it been?  $p(x_{0:t}|y_{0:t})$

Smoothing is “harder” than filtering/prediction [1].

# Deterministic solutions

## Kalman filter [2]

Under a linear Gaussian model:

$$X_0 \sim \mathcal{N}(0, \Sigma_0)$$

$$X_{t+1} \mid (X_t = x_t) \sim \mathcal{N}(Ax_t, \Sigma_x)$$

$$Y_t \mid (X_t = x_t) \sim \mathcal{N}(Bx_t, \Sigma_y)$$

we can recursively compute filtering distributions. Then a backward pass of RTS smoother [3] provides smoothing distributions.

This class of models is rather restrictive.

# Deterministic solutions

## Kalman filter [2]

Under a linear Gaussian model:

$$X_0 \sim \mathcal{N}(0, \Sigma_0)$$

$$X_{t+1} \mid (X_t = x_t) \sim \mathcal{N}(Ax_t, \Sigma_x)$$

$$Y_t \mid (X_t = x_t) \sim \mathcal{N}(Bx_t, \Sigma_y)$$

we can recursively compute filtering distributions. Then a backward pass of RTS smoother [3] provides smoothing distributions.

This class of models is rather restrictive.

## Extended Kalman filter [4]

In non-linear Gaussian models, use a local linear approximation and apply Kalman filter.

This requires gradients and performs poorly in models that are very non-linear.

# Deterministic solutions

## Kalman filter [2]

Under a linear Gaussian model:

$$X_0 \sim \mathcal{N}(0, \Sigma_0)$$

$$X_{t+1} \mid (X_t = x_t) \sim \mathcal{N}(Ax_t, \Sigma_x)$$

$$Y_t \mid (X_t = x_t) \sim \mathcal{N}(Bx_t, \Sigma_y)$$

we can recursively compute filtering distributions. Then a backward pass of RTS smoother [3] provides smoothing distributions.

This class of models is rather restrictive.

## Extended Kalman filter [4]

In non-linear Gaussian models, use a local linear approximation and apply Kalman filter.

This requires gradients and performs poorly in models that are very non-linear.

## Unscented Kalman filter [5]

For highly non-linear Gaussian models, replace the prediction step of Kalman filter by propagating a representative set of points through  $f$ .



## Deterministic solutions

- Kalman filter is optimal for linear Gaussian models, but extended/unscented Kalman filter are no longer optimal.

## Deterministic solutions

- Kalman filter is optimal for linear Gaussian models, but extended/unscented Kalman filter are no longer optimal.
- All of these methods require a Gaussian model. Deterministic solutions are available for some other conjugate families [6], but these are still restrictive.

# Deterministic solutions

- Kalman filter is optimal for linear Gaussian models, but extended/unscented Kalman filter are no longer optimal.
- All of these methods require a Gaussian model. Deterministic solutions are available for some other conjugate families [6], but these are still restrictive.
- Solutions are also available in the case that  $\mathcal{X}$  is finite (integrals become sums), but we will mostly consider  $\mathcal{X}$  to be a continuous space.

# Deterministic solutions

- Kalman filter is optimal for linear Gaussian models, but extended/unscented Kalman filter are no longer optimal.
- All of these methods require a Gaussian model. Deterministic solutions are available for some other conjugate families [6], but these are still restrictive.
- Solutions are also available in the case that  $\mathcal{X}$  is finite (integrals become sums), but we will mostly consider  $\mathcal{X}$  to be a continuous space.

Sequential Monte Carlo (SMC) provides general-purpose (stochastic) methods that do not require a tractable model.

# Deterministic solutions

- Kalman filter is optimal for linear Gaussian models, but extended/unscented Kalman filter are no longer optimal.
- All of these methods require a Gaussian model. Deterministic solutions are available for some other conjugate families [6], but these are still restrictive.
- Solutions are also available in the case that  $\mathcal{X}$  is finite (integrals become sums), but we will mostly consider  $\mathcal{X}$  to be a continuous space.

Sequential Monte Carlo (SMC) provides general-purpose (stochastic) methods that do not require a tractable model.

Only requires sampling from  $f(\cdot|x)$ , and pointwise evaluation of  $g(y|x)$  up to a normalising constant for each  $y$ .

# Sequential Monte Carlo

**Prior:**  $p(x_{0:t}) = \mu(x_0) \prod_{i=1}^t f(x_i|x_{i-1})$

**Likelihood:**  $p(y_{0:t}|x_{0:t}) = \prod_{i=0}^t g(y_i|x_i)$

**Posterior:**  $p(x_{0:t}|y_{0:t}) \propto \mu(x_0)g(y_0|x_0) \prod_{i=1}^t f(x_i|x_{i-1})g(y_i|x_i)$

# Sequential Monte Carlo

**Prior:**  $p(x_{0:t}) = \mu(x_0) \prod_{i=1}^t f(x_i|x_{i-1})$

**Likelihood:**  $p(y_{0:t}|x_{0:t}) = \prod_{i=0}^t g(y_i|x_i)$

**Posterior:**  $p(x_{0:t}|y_{0:t}) \propto \mu(x_0)g(y_0|x_0) \prod_{i=1}^t f(x_i|x_{i-1})g(y_i|x_i)$

- Represent posterior distribution at time  $t$  with  $N$  particles.
- Posterior factorises sequentially, avoiding increase of dimension with  $T$ .

# Sequential Monte Carlo

## Algorithm

After initialisation, iterate these steps:

- **Propagate:** move the particles through the transition  $f$
- **Calculate weights:** weight each particle according to  $g$
- **Resample:** duplicate high-weight particles and kill off low-weight ones



# Sequential Monte Carlo

## Algorithm

After initialisation, iterate these steps:

- **Propagate:** move the particles through the transition  $f$
- **Calculate weights:** weight each particle according to  $g$
- **Resample:** duplicate high-weight particles and kill off low-weight ones


Approximate posterior distribution  $p(dx_{0:t}|y_{0:t})$  by the empirical measure of the particles:

$$\hat{p}(dx_{0:t}|y_{0:t}) = \frac{1}{N} \sum_{i=1}^N \delta_{X_{0:t}^{(i)}}(dx_{0:t})$$


# Ancestral degeneracy

For smoothing we need a sample of trajectories.

Resampling means that trajectories of time  $T$  particles coalesce backwards in time.




# Kingman's coalescent



Looking backwards in time, each pair of lineages coalesces with unit rate.

# Kingman's coalescent



Looking backwards in time, each pair of lineages coalesces with unit rate.

This is the limiting coalescent process in many population models as  $N \rightarrow \infty$  [7, 8].

# Coalescent for SMC

- Analysing the coalescent can help us to understand the performance of various SMC algorithms.

# Coalescent for SMC

- Analysing the coalescent can help us to understand the performance of various SMC algorithms.
- The limiting coalescent of an SMC algorithm may depend on the resampling mechanism.

# Coalescent for SMC

- Analysing the coalescent can help us to understand the performance of various SMC algorithms.
- The limiting coalescent of an SMC algorithm may depend on the resampling mechanism.
- For multinomial resampling, the limiting coalescent is a scaled Kingman coalescent [9].

# Coalescent for SMC

- Analysing the coalescent can help us to understand the performance of various SMC algorithms.
- The limiting coalescent of an SMC algorithm may depend on the resampling mechanism.
- For multinomial resampling, the limiting coalescent is a scaled Kingman coalescent [9].
- What about other (more used) resampling schemes?



## Requirements:

- The total number of particles  $N$  remains fixed.
- The particles after resampling are equally weighted.
- The resampling scheme is unbiased; that is, the expected number of offspring of each particle  $i$  is equal to  $Nw_t^{(i)}$ .

# Resampling

**Multinomial:**  $v_t^{(1:N)} \stackrel{d}{=} \text{Multinomial}(N, w_t^{(1:N)})$

# Resampling

**Multinomial:**  $v_t^{(1:N)} \stackrel{d}{=} \text{Multinomial}(N, w_t^{(1:N)})$

**Residual [10]:**  $v_t^{(1:N)} \stackrel{d}{=} \lfloor Nw_t^{(1:N)} \rfloor + \text{Multinomial}(R, r_t^{(1:N)} / R)$

$$r_t^{(i)} := (Nw_t^{(i)} - \lfloor Nw_t^{(i)} \rfloor)$$

$$R := \sum r_t^{(i)}$$

# Resampling

**Multinomial:**  $v_t^{(1:N)} \stackrel{d}{=} \text{Multinomial}(N, w_t^{(1:N)})$

**Residual [10]:**  $v_t^{(1:N)} \stackrel{d}{=} \lfloor Nw_t^{(1:N)} \rfloor + \text{Multinomial}(R, r_t^{(1:N)} / R)$

$$r_t^{(i)} := (Nw_t^{(i)} - \lfloor Nw_t^{(i)} \rfloor)$$

$$R := \sum r_t^{(i)}$$

- Residual resampling yields lower Monte Carlo variance than multinomial resampling [11].
- Residual resampling is widely used by practitioners.
- Analysing the coalescent for residual resampling is a work in progress.

## Particle Gibbs [12]

Hidden Markov model where transition depends on a hyperparameter  $\theta$ :

$$\theta \sim \nu(\cdot)$$

$$X_0 \sim \mu(\cdot)$$

$$X_{t+1} \mid (X_t = x_t) \sim f_\theta(\cdot \mid x_t) \quad t = 0, \dots, T-1$$

$$Y_t \mid (X_t = x_t) \sim g(\cdot \mid x_t) \quad t = 0, \dots, T$$

## Particle Gibbs [12]

Hidden Markov model where transition depends on a hyperparameter  $\theta$ :

$$\theta \sim \nu(\cdot)$$

$$X_0 \sim \mu(\cdot)$$

$$X_{t+1} \mid (X_t = x_t) \sim f_\theta(\cdot \mid x_t) \quad t = 0, \dots, T-1$$

$$Y_t \mid (X_t = x_t) \sim g(\cdot \mid x_t) \quad t = 0, \dots, T$$

**Gibbs sampler:** alternately sample from  $p(\theta \mid x_{0:t}, y_{0:t})$  and  $p(x_{0:t} \mid \theta, y_{0:t})$ .

## Particle Gibbs [12]

Hidden Markov model where transition depends on a hyperparameter  $\theta$ :

$$\theta \sim \nu(\cdot)$$

$$X_0 \sim \mu(\cdot)$$

$$X_{t+1} \mid (X_t = x_t) \sim f_\theta(\cdot \mid x_t) \quad t = 0, \dots, T-1$$

$$Y_t \mid (X_t = x_t) \sim g(\cdot \mid x_t) \quad t = 0, \dots, T$$

**Gibbs sampler:** alternately sample from  $p(\theta \mid x_{0:t}, y_{0:t})$  and  $p(x_{0:t} \mid \theta, y_{0:t})$ .

- SMC is an appropriate method for sampling from  $p(x_{0:t} \mid \theta, y_{0:t})$
- To target the correct posterior distribution, need to use *conditional SMC*.

## Conditional SMC

- One “immortal” trajectory is conditioned to survive all of the resampling steps.
- Intuitively, this could have a significant effect on the coalescent.



# Conditional SMC

- One “immortal” trajectory is conditioned to survive all of the resampling steps.
- Intuitively, this could have a significant effect on the coalescent.
- But we have this result:

## Corollary 1

*Under the conditions of [9, Lemma 3] , the genealogy of any  $n$  particles from a conditional SMC algorithm with multinomial resampling converges to Kingman's  $n$ -coalescent in the sense of finite-dimensional distributions, under an appropriate time-scaling.*

# Conditional SMC

- One “immortal” trajectory is conditioned to survive all of the resampling steps.
- Intuitively, this could have a significant effect on the coalescent.
- But we have this result:

## Corollary 1

*Under the conditions of [9, Lemma 3] , the genealogy of any  $n$  particles from a conditional SMC algorithm with multinomial resampling converges to Kingman's  $n$ -coalescent in the sense of finite-dimensional distributions, under an appropriate time-scaling.*

- Intuition: as  $N \rightarrow \infty$ , there is zero probability that an arbitrary sample of fixed size  $n$  contains the immortal particle.

# Summary

- SMC is widely used, but doesn't perform well for smoothing.

# Summary

- SMC is widely used, but doesn't perform well for smoothing.
- Analysis of SMC coalescents allows us to quantify the problem of ancestral degeneracy.

# Summary

- SMC is widely used, but doesn't perform well for smoothing.
- Analysis of SMC coalescents allows us to quantify the problem of ancestral degeneracy.
- Proved: SMC with multinomial resampling  $\Rightarrow$  Kingman coalescent.
- (And simulations suggest it is a good model even for small  $N$ .)

# Summary

- SMC is widely used, but doesn't perform well for smoothing.
- Analysis of SMC coalescents allows us to quantify the problem of ancestral degeneracy.
- Proved: SMC with multinomial resampling  $\Rightarrow$  Kingman coalescent.
- (And simulations suggest it is a good model even for small  $N$ .)
- Proved: conditional SMC (with multinomial resampling)  $\Rightarrow$  Kingman coalescent.
- In progress: investigate by simulations whether this is a good model for finite  $N$ .

# Summary

- SMC is widely used, but doesn't perform well for smoothing.
- Analysis of SMC coalescents allows us to quantify the problem of ancestral degeneracy.
- Proved: SMC with multinomial resampling  $\Rightarrow$  Kingman coalescent.
- (And simulations suggest it is a good model even for small  $N$ .)
- Proved: conditional SMC (with multinomial resampling)  $\Rightarrow$  Kingman coalescent.
- In progress: investigate by simulations whether this is a good model for finite  $N$ .
- In progress: analysis of residual resampling (and other resampling schemes).

# References I

- [1] Neil J Gordon, David J Salmond, and Adrian FM Smith. Novel approach to nonlinear/non-Gaussian Bayesian state estimation. In *IEE Proceedings F (Radar and Signal Processing)*, volume 140, pages 107–113. IET, 1993.
- [2] Rudolph Emil Kalman. A new approach to linear filtering and prediction problems. *Journal of Basic Engineering*, 82(1):35–45, 1960.
- [3] Herbert E Rauch, CT Striebel, and F Tung. Maximum likelihood estimates of linear dynamic systems. *AIAA Journal*, 3(8):1445–1450, 1965.
- [4] Andrew H Jazwinski. *Stochastic Processes and Filtering Theory*. Courier Corporation, 2007.
- [5] Eric A Wan and Rudolph Van Der Merwe. The unscented Kalman filter for nonlinear estimation. In *Proceedings of the IEEE 2000 Adaptive Systems for Signal Processing, Communications, and Control Symposium*, pages 153–158. IEEE, 2000.
- [6] Paolo Vidoni. Exponential family state space models based on a conjugate latent process. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 61(1):213–221, 1999.



## References II

- [7] John Kingman. The coalescent. *Stochastic Processes and Their Applications*, 13(3):235–248, 1982.
- [8] Martin Möhle. Robustness results for the coalescent. *Journal of Applied Probability*, 35(2):438–447, 1998.
- [9] Jere Koskela, Paul A Jenkins, Adam M Johansen, and Dario Spanò. Asymptotic genealogies of interacting particle systems with an application to sequential Monte Carlo. *arXiv preprint arXiv:1804.01811*, 2018.
- [10] Jun S Liu and Rong Chen. Sequential Monte Carlo methods for dynamic systems. *Journal of the American Statistical Association*, 93(443):1032–1044, 1998.
- [11] Randal Douc, Olivier Cappé, and Eric Moulines. Comparison of resampling schemes for particle filtering. In *Image and Signal Processing and Analysis, 2005. ISPA 2005. Proceedings of the 4th International Symposium on*, pages 64–69. IEEE, 2005.
- [12] Christophe Andrieu, Arnaud Doucet, and Roman Holenstein. Particle Markov chain Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 72(3):269–342, 2010.