

# Weak convergence proof: an aside

Suzie Brown

October 2, 2020

NB: this document is now obsolete; the subsequent lemma and proof have been added to the document “weak-convproof.tex”.

**Lemma 1.** *For any  $n \leq N \in \mathbb{N}$ , for all  $t > 0$ , define*

$$E_t := \left\{ c_N(t) < \frac{(N-3)_{n-3}}{N^{n-3}} \left( \alpha \left( 1 + \frac{2}{N-2} \right) + B_n \right)^{-1} \right\} \quad (1)$$

where  $\alpha$  and  $B_n$  are positive constants as in equation whatever. Then, for all  $t > 0$ ,

$$\lim_{N \rightarrow \infty} \mathbb{P} \left[ \bigcap_{r=1}^{\tau_N(t)} E_r \right] = 1. \quad (2)$$

*Proof.*

$$\mathbb{P} \left[ \bigcap_{r=1}^{\tau_N(t)} E_r \right] = 1 - \mathbb{P} \left[ \bigcup_{r=1}^{\tau_N(t)} E_r^c \right] \geq 1 - \mathbb{E} \left[ \sum_{r=1}^{\tau_N(t)} \mathbb{P}[E_r^c] \right]. \quad (3)$$

Using the generalised Markov inequality,

$$\begin{aligned} \mathbb{P}[E_r^c] &= \mathbb{P} \left[ c_N(t) \geq \frac{(N-3)_{n-3}}{N^{n-3}} \left( \alpha \left( 1 + \frac{2}{N-2} \right) + B_n \right)^{-1} \right] \\ &\leq \mathbb{E}[c_N(r)^2] \frac{N^{2(n-3)}}{(N-3)_{n-3}^2} \left( \alpha \left( 1 + \frac{2}{N-2} \right) + B_n \right)^2. \end{aligned} \quad (4)$$

Now

$$\begin{aligned} \mathbb{P} \left[ \bigcap_{r=1}^{\tau_N(t)} E_r \right] &\geq 1 - \mathbb{E} \left[ \sum_{r=1}^{\tau_N(t)} \mathbb{E}[c_N(r)^2] \frac{N^{2(n-3)}}{(N-3)_{n-3}^2} \left( \alpha \left( 1 + \frac{2}{N-2} \right) + B_n \right)^2 \right] \\ &= 1 - \frac{N^{2(n-3)}}{(N-3)_{n-3}^2} \left( \alpha \left( 1 + \frac{2}{N-2} \right) + B_n \right)^2 \mathbb{E} \left[ \sum_{r=1}^{\tau_N(t)} c_N(r)^2 \right] \\ &\xrightarrow{N \rightarrow \infty} 1 - (\alpha + B_n)^2 \times 0 = 1. \end{aligned} \quad (5)$$

□