

a
Discovery, pseudobulk (5 lesions)
IL1R1
LIFR
IL1RAP
OSMR
IL6R
CCR2
TNFRSF1A
TNFRSF1B
IL6ST
CXCR4
−0.5
0.0
0.5
1.0
1.5
Δ log2CPM (inflammatory − tissue-forming)
b
CXCR4
IL6ST
TNFRSF1B
TNFRSF1A
CCR2
IL6R
OSMR
IL1RAP
LIFR
IL1R1
inflammatory
tissue-forming
mean log2CPM
0
1
2
3
4
5
6
7
NF-κB family set / with SMAD3
NF-κB family set / without SMAD3
alternative set (ATF1, MTA1) / with SMAD3
alternative set (ATF1, MTA1) / without SMAD3
c
Per-lesion Spearman ρ
(inflammatory vs tissue-forming TF means)
0.3
0.2
0.1
0.0
−0.1
−0.2
−0.3
Discovery (n=5)
Validation (n=3)
d
TF activity correlation (discovery)
RELB
NFKB2
RELA
NFKB1
REL
ATF1
MTA1
SP7
RUNX2
DLX5
MYRF
SMAD3
Spearman ρ
0.8
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
−0.8