

# THE CATEGORICITY SPECTRUM OF LARGE ABSTRACT ELEMENTARY CLASSES

SEBASTIEN VASEY

ABSTRACT. The categoricity spectrum of a class of structures is the collection of cardinals in which the class has a single model up to isomorphism. Assuming that cardinal exponentiation is injective (a weakening of the generalized continuum hypothesis, GCH), we give a complete list of the possible categoricity spectrums of an abstract elementary class with amalgamation and arbitrarily large models. Specifically, the categoricity spectrum is either empty, an end segment starting below the Hanf number, or a closed interval consisting of finite successors of the Löwenheim-Skolem-Tarski number (there are examples of each type). We also prove (assuming a strengthening of the GCH) that the categoricity spectrum of an abstract elementary class with no maximal models is either bounded or contains an end segment. This answers several longstanding questions around Shelah's categoricity conjecture.

## CONTENTS

|   |    |
|---|----|
| 1. Introduction                             | 2  |
| 2. Preliminaries                            | 6  |
| 3. Nice stability and superlimits           | 12 |
| 4. AECs with arbitrarily large models       | 17 |
| 5. Towers and disjoint amalgamation         | 20 |
| 6. Building weakly successful good frames   | 28 |
| 7. Building successful good frames          | 30 |
| 8. The main categoricity transfer           | 32 |
| 9. Categoricity assuming amalgamation       | 33 |
| 10. Categoricity assuming no maximal models | 36 |
| 11. Some auxiliary results                  | 40 |
| References                                  | 42 |

---

*Date:* May 10, 2018

AMS 2010 Subject Classification: Primary 03C48. Secondary: 03C45, 03C52, 03C55, 03C75, 03E05.

*Key words and phrases.* abstract elementary classes, categoricity, amalgamation, no maximal models, good frames, towers, weak generalized continuum hypothesis.

## 1. INTRODUCTION

**1.1. Motivation.** A recurring question in mathematics is whether a list of properties characterize a certain structure. To make this precise, let us restrict ourselves here to structures in the model-theoretic sense: a universe with operations and relations on it. We say that a class  $K$  of such structures is *categorical* if it contains exactly one member up to isomorphism. The Löwenheim-Skolem theorem, a basic result of model theory, says that if a set  $T$  of sentences in first-order logic<sup>1</sup> has an infinite model, then it has models of all infinite sizes (at least  $|T|$ ). In particular, the class of models of  $T$  cannot be categorical.

The following weakening of the definition of categoricity was proposed as a remedy by Łoś [Ło54]: for  $\lambda$  a cardinal, we say that a class  $K$  of structures is *categorical in  $\lambda$*  if it has a single model of cardinality  $\lambda$  up to isomorphism. We say that a theory (i.e. a set of sentences)  $T$  is *categorical in  $\lambda$*  if its class of models is categorical in  $\lambda$ . The following central result of modern model theory describes the behavior of categoricity for countable first-order theories:

**Fact 1.1** (Morley’s categoricity theorem, [Mor65]). If a countable first-order theory is categorical in *some* uncountable cardinal, then it is categorical in *all* uncountable cardinals.

Two examples of classes that are categorical in all uncountable cardinals are the class of vector spaces (over a fixed countable field) and the class of algebraically closed fields (of a fixed characteristic). The reason for categoricity in both classes is that they have well-understood notions of independence (linear independence in vector spaces, algebraic independence in fields). The proof of Morley’s theorem (as well as further developments) tells us that this is not an accident: *any* class of models of an uncountably categorical countable first-order theory will have a nice notion of independence. Such a notion was baptized *forking* by Shelah [She90] and is now central, both for pure model theory and for applications to other fields of mathematics (on the latter, see e.g. Hrushovski’s proof of the Mordell-Lang conjecture for function fields [Hru96]).

**1.2. Shelah’s categoricity conjecture.** One can see Morley’s theorem (and its earlier conjecture by Łoś [Ło54]) as the catalyst that led to the development of forking for first-order theories. It is natural to ask whether Morley’s theorem (and hence forking) can be generalized to other, not necessarily first-order, classes of structures. For uncountable first order theories, Shelah [She74] proved that a first-order theory  $T$  categorical in *some*  $\mu > |T|$  is categorical in *all*  $\mu' > |T|$ . The next step was to look at infinitary logics such as  $\mathbb{L}_{\omega_1, \omega}$  (where countably infinite conjunctions and disjunctions are allowed). The situation here is much more complicated, since the compactness theorem fails, but nevertheless Shelah [She83, Conjecture 2] conjectured the following version of Morley’s categoricity theorem:

**Conjecture 1.2** (Shelah’s categoricity conjecture for  $\mathbb{L}_{\omega_1, \omega}$ ). If an  $\mathbb{L}_{\omega_1, \omega}$ -sentence is categorical in *some*  $\mu \geq \beth_{\omega_1}$ , then it is categorical in *all*  $\mu' \geq \beth_{\omega_1}$ .

---

<sup>1</sup>That is, sentences like  $\forall x \exists y : x \cdot y = 1 \wedge y \cdot x = 1$ : quantification is over elements and only finite conjunctions and disjunctions are allowed.

Here,  $\omega_1$  is the first uncountable ordinal, and for an ordinal  $\alpha$ ,  $\beth_\alpha$  is the cardinal obtained by iterating cardinal exponentiation  $\alpha$ -many times, starting with  $\aleph_0$  (see Section 2.1 for a precise definition). The spirit of the conjecture is that categoricity *somewhere* high-enough should imply categoricity *everywhere* high-enough (and indeed, even that eventual version is open). A construction of Morley (see Example 9.10 here) shows that the “high-enough” threshold must in this case be at least  $\beth_{\omega_1}$ .

As the compactness theorem fails, and moreover there is a plethora of other “reasonable” logics to work with, it turns out to be convenient to work semantically rather than syntactically. The framework of *abstract elementary classes* (AECs), introduced by Shelah in the late seventies [She87a], encompasses “reasonable” infinitary logics such as  $\mathbb{L}_{\omega_1, \omega}$  as well as natural classes of algebraic examples (such as Zilber’s pseudoexponential fields [Zil05] or certain classes of modules [BET07]). Roughly, an AEC is a partially ordered class  $\mathbf{K} = (K, \leq_{\mathbf{K}})$  of structures satisfying some closure properties. For example, it must be closed under unions of chains and any member of the class must have a substructure which is also in the class and has size at most a fixed cardinal, the Löwenheim-Skolem-Tarski number of the class, written  $\text{LS}(\mathbf{K})$  (this is a replacement for the size of the theory). The reader should see 2.1 here for a precise definition, and the introductions of [She09a], [Bal09], or [Gro02] for more motivation on AECs. Note that AECs can also be characterized as certain kind of accessible categories, see [BR12, LR16].

Shelah has given the following version of Conjecture 1.2 for AECs (see [She00, 6.14], [She09a, N.4.3]):

**Conjecture 1.3** (Shelah’s categoricity conjecture for AECs). If an AEC  $\mathbf{K}$  is categorical in *some*  $\mu \geq \beth_{(2^{\text{LS}(\mathbf{K})})^+}$ , then  $\mathbf{K}$  is categorical in *all*  $\mu' \geq \beth_{(2^{\text{LS}(\mathbf{K})})^+}$ .

**1.3. Categoricity in AECs with amalgamation.** One goal of the present paper is to prove (assuming a set-theoretic hypothesis) the statement of Conjecture 1.3 for AECs which satisfy the amalgamation property (i.e. where every span can be completed to a commuting square).

Before going further, let us explain our justification for assuming the amalgamation property. First, amalgamation holds in case the AEC comes from a first-order theory (by compactness). Second, it is known [MS90, SK96, Bon14] that amalgamation holds (eventually) in any AEC that is categorical above a large cardinal. In fact, Grossberg conjectured [Gro02, 2.3] that eventual amalgamation should follow from high-enough categoricity, even without large cardinals. Finally, Shelah showed (assuming a set-theoretic hypothesis) that in an AEC categorical in  $\lambda$  and  $\lambda^+$ , amalgamation holds for models of cardinality  $\lambda$  [She09a, I.3.8]. Thus in an eventually categorical AEC, eventual amalgamation must hold, so amalgamation is a consequence of the eventual categoricity result we want to prove and hence it seems reasonable to assume it as a starting point.

The first milestone result for general AECs with amalgamation is due to Shelah [She99] who showed that categoricity in *some successor*  $\mu \geq H_2$  implies categoricity in *all*  $\mu' \in [H_2, \mu]$ . We have set  $H_1 := \beth_{(2^{\text{LS}(\mathbf{K})})^+}$  and  $H_2 := \beth_{(2^{H_1})^+}$ . Until very recently, it was open whether the successor assumption could be removed, whether one could also go up (i.e. get categoricity in all  $\mu' > \mu$ ), and whether the threshold  $H_2$  could be lowered to the cardinal  $\beth_{(2^{\text{LS}(\mathbf{K})})^+}$  featuring in Conjecture 1.3.

These questions feature in several prominent lists of open problems, for example in [She00, 6.14], [Gro02, Section 9], [Bal06], [Bal09, Appendix D], or the introduction of [She09a].

In his book on AECs, Shelah [She09a, IV.7.12] asserts that he can remove the successor assumption and also get an upward transfer (this is done at the cost of a set-theoretic hypothesis, see below). The  $H_2$  threshold is also slightly lowered, although not quite all the way down to  $H_1$ . However, Shelah's proof relies on a claim that was not proven at the time the book was written (see also [Vas17a, Section 11] for an exposition of this proof, modulo the claim). In very recent joint work of the author with Shelah [SVb], the missing claim was finally proven. It nevertheless remained open whether  $H_2$  could be lowered all the way down to  $H_1$ . This is one contribution of the present paper.

**1.4. Categoricity in AECs with no maximal models.** We also investigate a much weaker framework than AECs with amalgamation: AECs with no maximal models. This was first studied by Shelah and Villaveces [SV99], with later contributions by VanDieren [Van06, Van13]. In these papers, some superstability-like properties were shown to follow from categoricity (assuming again set-theoretic hypotheses). Still, nothing could be said on the categoricity spectrum (i.e. the class of categoricity cardinals above  $\text{LS}(\mathbf{K})$ , see Definition 9.1) and for a long time no further progress was made. Another contribution of the present paper is that the categoricity spectrum in AECs with no maximal models is either bounded or contains an end segment (Corollary 10.7). This implies the eventual categoricity conjecture in this framework (Corollary 10.8), although we are unable to give an explicit bound on the threshold.

**1.5. Main results.** Let us discuss our results more precisely. Assuming a set-theoretic hypothesis, we list *all* possibilities for the categoricity spectrum of an AEC  $\mathbf{K}$  with amalgamation and arbitrarily large models. There are three possibilities (Corollary 9.7): the categoricity spectrum is either empty, an end segment, or a closed interval. Furthermore in case it is an end segment, the first categoricity cardinal must be below  $H_1$ , and in case it is an interval, there must be finitely-many cardinals between the endpoints and  $\text{LS}(\mathbf{K})$ . It was known that each of those three possibilities could happen (see Example 9.10). In particular, for each  $n < \omega$  there is an example (due to Hart and Shelah [HS90] and further analyzed by Baldwin and Kolesnikov [BK09]) of an AEC  $\mathbf{K}$  with  $\text{LS}(\mathbf{K}) = \aleph_0$  which has amalgamation, arbitrarily large models, and is categorical in  $\aleph_0, \aleph_1, \dots, \aleph_n$  but not anywhere above. The present work shows (again assuming amalgamation and arbitrarily large models) that categoricity at  $\aleph_\omega$  or above must imply eventual categoricity, hence that the Hart-Shelah example is in a sense the only obstruction to transferring categoricity upward.

In passing (and still assuming some set theory), we also show as a consequence of our argument that *tameness*, a locality property introduced by Grossberg and VanDieren in [GV06b] and used to prove an upward categoricity transfer in [GV06c, GV06a] follows from categoricity above  $H_1$  in an AEC with amalgamation (Corollary 11.3). This was conjectured in [GV06a, 1.5].

All of the work described in the previous two paragraphs is done assuming a weakening of the generalized continuum hypothesis (GCH). Recall that GCH is the statement that  $2^\lambda = \lambda^+$  for any infinite cardinal  $\lambda$ . We assume here only the *weak generalized continuum hypothesis* (WGCH):  $2^\lambda < 2^{\lambda^+}$  for any infinite cardinal  $\lambda$ . It says that cardinal exponentiation is strictly monotonic. The use of this hypothesis is quite prominent in Shelah's work on AECs [She09a, She09b] and can be traced back to a combinatorial principle, the *weak diamond*, proven by Devlin and Shelah [DS78] to follow from  $2^{\aleph_0} < 2^{\aleph_1}$ .

Regarding AECs with no maximal models, we prove assuming the GCH *and* a strengthening of the weak diamond (see Section 2.1; this hypothesis is already used in [SV99]) that the categoricity spectrum is either bounded or contains an end segment. More precisely, we give a list of five possibilities for the categoricity spectrum above a countable limit of fixed points of the  $\sqcup$  operation (Corollary 10.7). In this case, we do not know whether all of these possibilities can happen.

**1.6. Methods.** The main technical tool of the present paper is the theory of *good frames*, the core object of study in Shelah's two-volume book on AECs [She09a, She09b]. Good frames give a sense in which an AEC is well-behaved locally (i.e. at a single cardinal  $\lambda$ ). Key questions around them is when they exist and when they can be transferred upward. The main technical theorem of this paper, Corollary 6.5, gives sufficient conditions (involving also WGCH) under which certain good frames exist. These sufficient conditions were known to follow from categoricity and amalgamation, and we show here how to derive them from categoricity and no maximal models. This technical theorem thus gives us good frames in many places below the categoricity cardinal. Using the main result of [Vas], these good frames are then shown to be “connected” in the sense that one can describe what happens in the frames above by looking at what happens in the frames below. Such connected good frames (called *successful* by Shelah), turn out to be powerful enough to transfer categoricity and more generally understand the structure of the class completely, as proven in the recent [SVb] (which we rely on as a black box).

Another key is the theory of *towers*, connected to the problem of uniqueness of limit models (see [GVV16] for an overview). We give a simplification of this theory here (Section 5) incorporating recent developments [Vas17b] as well as some new results that may have independent interest. Ultimately, these results are used to build the good frames alluded to in the previous paragraph.

We emphasize that the methods of this paper are usually very local. While AECs with amalgamation or no maximal models are used as convenient test cases, the theorems we obtain make no full use of these assumptions: it often suffices that a condition called *solvability* (introduced by Shelah in [She09a, Chapter IV]), or really a weakening of it called *semisolvability*, holds near the cardinals where we want to build good frames. Also key and closely related is the notion of a *superlimit model* (due to Shelah [She87a, 3.1]). We prove in fact (assuming the WGCH) that classes with superlimit models at every cardinal satisfy the eventual categoricity conjecture (Corollary 11.5). As discussed further at the end of the paper, this sheds light on several other conjectures of Shelah and paves the way for further work on the local superstability theory of AECs.

## 2. PRELIMINARIES

To read the present paper, the reader should preferably have a solid knowledge of AECs and good frames, including knowing Chapter II of [She09a] and [JS13]. Still, we attempt here to give most of the relevant definitions and background facts. The reader may skip this section at first reading and come back to it as needed.

**2.1. Set-theoretic notation.** We will often consider intervals of cardinals and may write  $[\lambda, \infty)$  for the class of cardinals greater than or equal to  $\lambda$  (i.e.  $\infty$  denotes an object that is greater than all cardinals).

We assume basic familiarity with ordinals and cardinals. We identify each cardinal  $\lambda$  with the least ordinal of cardinality  $\lambda$ . For a cardinal  $\lambda$ ,  $\lambda^+$  denotes the successor of  $\lambda$ : the minimal cardinal strictly greater than  $\lambda$ . The *cofinality*  $\text{cf}(\alpha)$  of an ordinal  $\alpha$  is the least cardinality of a cofinal subset of  $\alpha$ . A cardinal  $\lambda$  is *regular* if  $\text{cf}(\lambda) = \lambda$  and singular otherwise.

For  $\lambda$  an infinite cardinal, define the following three statements:

- $\text{WGCH}(\lambda)$  means that  $2^\lambda < 2^{\lambda^+}$  (WGCH stands for “weak generalized continuum hypothesis”).
- $\text{GCH}(\lambda)$  means that  $2^\lambda = \lambda^+$ .
- $\text{GCHWD}(\lambda)$  means that  $2^\lambda = \lambda^+$  and for all regular  $\theta < \lambda^+$ ,  $\Phi_{\lambda^+}(\{\delta < \lambda^+ \mid \text{cf}(\delta) = \theta\})$  holds, where for a set  $S \subseteq \lambda^+$ ,  $\Phi_{\lambda^+}(S)$  holds if and only if for all  $F : {}^{<\lambda^+}2$  there exists  $g : \lambda^+ \rightarrow 2$  such that for every  $f : \lambda^+ \rightarrow 2$  the set  $\{\delta \in S \mid F(f \restriction \delta) = g(\delta)\}$  is stationary (i.e. intersects every closed unbounded subset of  $\lambda^+$ ). This was first studied by Devlin and Shelah [DS78], who proved that  $2^\lambda < 2^{\lambda^+}$  implies  $\Phi_{\lambda^+}(\lambda^+)$  (WD stands for “weak diamond”).

Note that  $\text{GCHWD}(\lambda)$  implies  $\text{GCH}(\lambda)$  which implies  $\text{WGCH}(\lambda)$ .

For  $\Theta$  a class of cardinals, we write  $\text{WGCH}(\Theta)$  if  $\text{WGCH}(\lambda)$  holds for all  $\lambda \in \Theta$ . We write  $\text{WGCH}$  for  $\text{WGCH}(\text{CARD})$ , where  $\text{CARD}$  is the class of all infinite cardinals. Similarly define  $\text{GCH}(\Theta)$ ,  $\text{GCH}$ ,  $\text{GCHWD}(\Theta)$ , and  $\text{GCHWD}$ . It is well known that  $\text{GCHWD}$  is consistent (it holds for example in Gödel’s constructible universe, see [Jec03, p. 550] and [Van06, I.3.4]).

For a cardinal  $\lambda$  and an ordinal  $\alpha$ ,  $\beth_\alpha(\lambda)$  is defined recursively as follows:  $\beth_0(\lambda) = \lambda$ ,  $\beth_{\beta+1}(\lambda) = 2^{\beth_\beta(\lambda)}$ , and  $\beth_\delta(\lambda) = \sup_{\beta < \delta} \beth_\beta(\lambda)$  for  $\delta$  a limit ordinal. We write  $\beth_\alpha$  for  $\beth_\alpha(\aleph_0)$ . Also recursively define  $\lambda^{+\alpha}$ , the  $\alpha$ th successor of  $\lambda$ , as follows:  $\lambda^{+0} = \lambda$ ,  $\lambda^{+(\alpha+1)} = (\lambda^{+\alpha})^+$ , and  $\lambda^{+\delta} = \sup_{\beta < \delta} \lambda^{+\beta}$  for  $\delta$  limit. For an infinite cardinal  $\lambda$ , it will be convenient to write  $h(\lambda)$  instead of  $\beth_{(2^\lambda)^+}$  (this notation is used already in [Bal09, 4.24]; the relevance of this cardinal is given by Fact 4.1).

**2.2. Model-theoretic notation.** Given a structure  $M$ , write  $|M|$  for its universe and  $\|M\|$  for the cardinality of its universe. We often do not distinguish between  $M$  and  $|M|$ , writing e.g.  $a \in M$  instead of  $a \in |M|$ . We write  $M \subseteq N$  to mean that  $M$  is a substructure of  $N$ .

**2.3. Abstract elementary classes.** An *abstract class* is a pair  $\mathbf{K} = (K, \leq_{\mathbf{K}})$ , where  $K$  is a class of structures in a fixed vocabulary  $\tau = \tau(\mathbf{K})$  and  $\leq_{\mathbf{K}}$  is a partial order,  $M \leq_{\mathbf{K}} N$  implies  $M \subseteq N$ , and both  $K$  and  $\leq_{\mathbf{K}}$  respect isomorphisms (the definition is due to Grossberg). We often do not distinguish between  $K$  (the class of structures) and  $\mathbf{K}$  (the *ordered* class of structures). Any abstract class admits a notion of  *$\mathbf{K}$ -embedding*: these are functions  $f : M \rightarrow N$  such that  $f : M \cong f[M]$  and  $f[M] \leq_{\mathbf{K}} N$ . Thus one can naturally see  $\mathbf{K}$  as a category. Unless explicitly stated, any map  $f : M \rightarrow N$  in this paper will be a  $\mathbf{K}$ -embedding. We write  $f : M \xrightarrow{A} N$  to mean that  $f$  is a  $\mathbf{K}$ -embedding from  $M$  into  $N$  which fixes the set  $A$  pointwise (so  $A \subseteq |M|$ ). We similarly write  $f : M \cong_A N$  for isomorphisms from  $M$  onto  $N$  fixing  $A$ .

For  $\lambda$  a cardinal, we will write  $\mathbf{K}_\lambda$  for the restriction of  $\mathbf{K}$  to models of cardinality  $\lambda$ . Similarly define  $\mathbf{K}_{\geq \lambda}$ ,  $\mathbf{K}_{< \lambda}$ , or more generally  $\mathbf{K}_\Theta$ , where  $\Theta$  is a class of cardinals.

For an abstract class  $\mathbf{K}$ , we denote by  $\mathbb{I}(\mathbf{K})$  the number of models in  $\mathbf{K}$  up to isomorphism (i.e. the cardinality of  $\mathbf{K}/\cong$ ). We write  $\mathbb{I}(\mathbf{K}, \lambda)$  instead of  $\mathbb{I}(\mathbf{K}_\lambda)$ . When  $\mathbb{I}(\mathbf{K}) = 1$ , we say that  $\mathbf{K}$  is *categorical*. We say that  $\mathbf{K}$  is *categorical in  $\lambda$*  if  $\mathbf{K}_\lambda$  is categorical, i.e.  $\mathbb{I}(\mathbf{K}, \lambda) = 1$ .

We say that  $\mathbf{K}$  has *amalgamation* if for any  $M_0 \leq_{\mathbf{K}} M_\ell$ ,  $\ell = 1, 2$ , there is  $M_3 \in \mathbf{K}$  and  $\mathbf{K}$ -embeddings  $f_\ell : M_\ell \xrightarrow{M_0} M_3$ ,  $\ell = 1, 2$ .  $\mathbf{K}$  has *joint embedding* if any two models can be  $\mathbf{K}$ -embedded in a common model.  $\mathbf{K}$  has *no maximal models* if for any  $M \in \mathbf{K}$  there exists  $N \in \mathbf{K}$  with  $M \leq_{\mathbf{K}} N$  and  $M \neq N$  (we write  $M <_{\mathbf{K}} N$ ). Localized concepts such as *amalgamation in  $\lambda$*  mean that  $\mathbf{K}_\lambda$  has amalgamation.

The definition of an abstract elementary class is due to Shelah [She87a]:

**Definition 2.1.** An *abstract elementary class (AEC)* is an abstract class  $\mathbf{K}$  in a finitary vocabulary satisfying:

- (1) Coherence: if  $M_0, M_1, M_2 \in \mathbf{K}$ ,  $M_0 \subseteq M_1 \leq_{\mathbf{K}} M_2$  and  $M_0 \leq_{\mathbf{K}} M_2$ , then  $M_0 \leq_{\mathbf{K}} M_1$ .
- (2) Tarski-Vaught chain axioms: if  $\langle M_i : i \in I \rangle$  is a  $\leq_{\mathbf{K}}$ -directed system and  $M := \bigcup_{i \in I} M_i$ , then:
  - (a)  $M \in \mathbf{K}$ .
  - (b)  $M_i \leq_{\mathbf{K}} M$  for all  $i \in I$ .
  - (c) If  $N \in \mathbf{K}$  is such that  $M_i \leq_{\mathbf{K}} N$  for all  $i \in I$ , then  $M \leq_{\mathbf{K}} N$ .
- (3) Löwenheim-Skolem-Tarski axiom: there exists a cardinal  $\lambda \geq |\tau(\mathbf{K})| + \aleph_0$  such that for any  $N \in \mathbf{K}$  and any  $A \subseteq |N|$ , there exists  $M \in \mathbf{K}$  with  $M \leq_{\mathbf{K}} N$ ,  $A \subseteq |M|$ , and  $\|M\| \leq |A| + \lambda$ . We write  $\text{LS}(\mathbf{K})$  for the least such  $\lambda$ .

**2.4. Types.** In any abstract class  $\mathbf{K}$ , we can define a semantic notion of type, called Galois or orbital types in the literature (such types were introduced by Shelah in [She87b]). For  $M \in \mathbf{K}$ ,  $A \subseteq |M|$ , and  $b \in M$ , we write  $\text{tp}_{\mathbf{K}}(b/A; M)$  for the orbital type of  $b$  over  $A$  as computed in  $M$  (usually  $\mathbf{K}$  will be clear from context and we will omit it from the notation). It is the finest notion of type respecting  $\mathbf{K}$ -embeddings, see [Vas16c, 2.16] for a formal definition. For  $M \in \mathbf{K}$ , we write  $\mathbf{S}_{\mathbf{K}}(M) = \mathbf{S}(M)$  for  $\{\text{tp}(b/M; N) \mid M \leq_{\mathbf{K}} N\}$ , the class<sup>2</sup> of all types over  $M$ . We define naturally

<sup>2</sup>If  $\mathbf{K}$  is an AEC,  $\mathbf{S}(M)$  will of course be a set.

what it means for a type to be realized inside a model, to extend another type, and to take the image of a type by a  $\mathbf{K}$ -embedding.

When  $\mathbf{K}$  is an elementary class,  $\mathbf{tp}(b/A; M)$  contains the same information as the usual notion of  $\mathbb{L}_{\omega, \omega}$ -syntactic type. In particular, types in an elementary class are determined by their restrictions to finite sets. This idea was abstracted in [GV06b] and made into the following definition: for  $\chi$  an infinite cardinal, an abstract class  $\mathbf{K}$  is  $(< \chi)$ -*tame* if for any  $M \in \mathbf{K}$  and any distinct  $p, q \in \mathbf{S}(M)$ , there exists  $A \subseteq |M|$  such that  $|A| < \chi$  and  $p \restriction A \neq q \restriction A$ . We say that  $\mathbf{K}$  is  $\chi$ -*tame* if it is  $(< \chi^+)$ -tame. Thus elementary classes are  $(< \aleph_0)$ -tame, but there are examples of non-tame AECs, see e.g. [BV17b, 3.2.2].

**2.5. Stability and saturation.** We say that an abstract class  $\mathbf{K}$ , is *stable in  $\lambda$*  (for  $\lambda$  an infinite cardinal) if  $|\mathbf{S}(M)| \leq \lambda$  for any  $M \in \mathbf{K}_\lambda$ . If  $\mathbf{K}$  is an AEC,  $\lambda \geq \text{LS}(\mathbf{K})$ ,  $\mathbf{K}$  is stable in  $\lambda$  and  $\mathbf{K}$  has amalgamation in  $\lambda$ , then we will often use without comments the *existence of universal extension* [She09a, II.1.16]: for any  $M \in \mathbf{K}_\lambda$ , there exists  $N \in \mathbf{K}_\lambda$  universal over  $M$ . This means that  $M \leq_{\mathbf{K}} N$  and any extension of  $M$  of cardinality  $\lambda$   $\mathbf{K}$ -embeds into  $N$  over  $M$ .

For  $\mathbf{K}$  an AEC and  $\lambda > \text{LS}(\mathbf{K})$ , a model  $N \in \mathbf{K}$  is called  $\lambda$ -*saturated* if for any  $M \in \mathbf{K}_{<\lambda}$  with  $M \leq_{\mathbf{K}} N$ , any  $p \in \mathbf{S}(M)$  is realized in  $N$ .  $N$  is called *saturated* if it is  $\|N\|$ -saturated.

We will also often use without mention the *model-homogeneous = saturated lemma* [She09a, II.1.14]: it says that when  $\mathbf{K}_{<\lambda}$  has amalgamation, a model  $N \in \mathbf{K}$  is  $\lambda$ -saturated if and only if it is  $\lambda$ -model-homogeneous. The latter means that for any  $M \in \mathbf{K}$  with  $\|M\| < \lambda$ ,  $M \leq_{\mathbf{K}} N$ , any  $M' \in \mathbf{K}_{<\lambda} \leq_{\mathbf{K}}$ -extending  $M$  can be  $\mathbf{K}$ -embedded into  $N$  over  $M$ . In particular, assuming amalgamation and joint embedding, there is at most one saturated model of a given cardinality. We write  $\mathbf{K}^{\lambda\text{-sat}}$  for the abstract class of  $\lambda$ -saturated models in  $\mathbf{K}$  (ordered by the appropriate restriction of  $\leq_{\mathbf{K}}$ ).

A local notion of saturation is given by the definition of a *limit model*. For an AEC  $\mathbf{K}$ ,  $\lambda \geq \text{LS}(\mathbf{K})$ , and  $\delta < \lambda^+$  a limit ordinal,  $N$  is  $(\lambda, \delta)$ -*limit over  $M$*  if  $M, N \in \mathbf{K}_\lambda$  and there exists an increasing continuous chain  $\langle M_i : i \leq \delta \rangle$  in  $\mathbf{K}_\lambda$  with  $M_{i+1}$  universal over  $M_i$  for all  $i < \delta$  such that  $M_0 = M$  and  $M_\delta = N$ . We say that  $N$  is *limit over  $M$*  if it is  $(\lambda, \delta)$ -limit over  $M$  for some  $\lambda$  and  $\delta$ . We say that  $N$  is *limit* if it is limit over  $M$  for some  $M$ . By a back and forth argument [SV99, 1.3.6], whenever  $M_0, M_1, M_2 \in \mathbf{K}_\lambda$ ,  $M_\ell$  is  $(\lambda, \delta_\ell)$ -limit over  $M_0$  for  $\ell = 1, 2$ , and  $\text{cf}(\delta_1) = \text{cf}(\delta_2)$ , we have that  $M_1 \cong_{M_0} M_2$ . Similarly, assuming joint embedding any two limit models of the same length (i.e. with the same limit ordinal  $\delta$ ) are isomorphic. The question of *uniqueness of limit models* asks whether the previous two results hold even when the lengths do not have the same cofinality (Fact 2.7 gives a positive answer to this question under some superstability-like assumptions).

**2.6. Superstability, forking, and symmetry.** Let  $\mathbf{K}$  be an AEC. For  $M \leq_{\mathbf{K}} N$  and  $p \in \mathbf{S}(N)$ , we say that  $p$  *does not  $\lambda$ -split over  $M$*  if whenever  $N_1, N_2 \in \mathbf{K}_\lambda$  are such that  $M \leq_{\mathbf{K}} N_\ell \leq_{\mathbf{K}} N$  for  $\ell = 1, 2$  and  $f : N_1 \cong_M N_2$  is given, we have that  $f(p \restriction N_1) = p \restriction N_2$ . This gives a notion of independence with which we will be able, under the definition of superstability given below (already implicit in for example [SV99]), to define a forking-like notion.



**Definition 2.2** ([Vas16a, 10.1]). An AEC  $\mathbf{K}$  is  $\lambda$ -*superstable* if:

- (1)  $\lambda \geq \text{LS}(\mathbf{K})$  and  $\mathbf{K}_\lambda \neq \emptyset$ .
- (2)  $\mathbf{K}_\lambda$  has amalgamation, joint embeddings, and no maximal models.
- (3)  $\mathbf{K}$  is stable in  $\lambda$ .
- (4) Splitting has universal local character: whenever  $\delta < \lambda^+$  is a limit ordinal and  $\langle M_i : i \leq \delta \rangle$  is increasing continuous in  $\mathbf{K}_\lambda$  with  $M_{i+1}$  universal over  $M_i$  for all  $i < \delta$ , then for any  $p \in \mathbf{S}(\bigcup_{i < \delta} M_i)$ , there exists  $i < \delta$  such that  $p$  does not  $\lambda$ -split over  $M_i$ .

Note that the definition is completely local: it only discusses models of cardinality  $\lambda$ . In a  $\lambda$ -superstable AEC, we define nonforking by “shifting” nonsplitting by a universal extension:

**Definition 2.3** ([Vas16b, 3.8]). Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC. Let  $M \leq_{\mathbf{K}} N$  both be limit models and let  $p \in \mathbf{S}(N)$ . We say that  $p$  *does not  $\lambda$ -fork over  $M$*  if there exists  $M_0 \in \mathbf{K}_\lambda$  such that  $M$  is universal over  $M_0$  and  $p$  does not  $\lambda$ -split over  $M_0$ . Usually,  $\lambda$  will be clear from context so we will just say that  $p$  *does not fork over  $M$* .

The definition of forking may seem technical, so the reader can immediately forget it and remember instead the next two facts:

**Fact 2.4.** If  $\mathbf{K}$  is a  $\lambda$ -superstable AEC, then forking has the following properties:

- (1) Invariance: if  $M \leq_{\mathbf{K}} N$  are both limit models in  $\mathbf{K}_\lambda$  and  $p \in \mathbf{S}(N)$  does not fork over  $M$ , then if  $f : N \cong N'$ ,  $f(p)$  does not fork over  $f[M]$ .
- (2) Monotonicity: if  $M \leq_{\mathbf{K}} M' \leq_{\mathbf{K}} N' \leq_{\mathbf{K}} N$  are all limit models in  $\mathbf{K}_\lambda$ ,  $p \in \mathbf{S}(N)$  does not fork over  $M$ , then  $p \upharpoonright N'$  does not fork over  $M'$ .
- (3) Universal local character: if  $\delta < \lambda^+$ ,  $\langle M_i : i \leq \delta \rangle$  is an increasing continuous chain of limit models in  $\mathbf{K}_\lambda$ , with  $M_{i+1}$  universal over  $M_i$  for all  $i < \delta$ , then for any  $p \in \mathbf{S}(M_\delta)$ , there exists  $i < \delta$  such that  $p$  does not fork over  $M_i$ .
- (4) Uniqueness: if  $M \leq_{\mathbf{K}} N$  are both limit models in  $\mathbf{K}_\lambda$  and  $p, q \in \mathbf{S}(N)$  do not fork over  $M$ , then  $p \upharpoonright M = q \upharpoonright M$  implies  $p = q$ .
- (5) Extension: Let  $M \leq_{\mathbf{K}} N$  both be limit models in  $\mathbf{K}_\lambda$ . If  $p \in \mathbf{S}(M)$ , then there exists  $q \in \mathbf{S}(N)$  such that  $q$  does not fork over  $M$  and  $q$  extends  $p$ .
- (6) Transitivity: Let  $M_0 \leq_{\mathbf{K}} M_1 \leq_{\mathbf{K}} M_2$  all be limit models in  $\mathbf{K}_\lambda$ . Let  $p \in \mathbf{S}(M_2)$  and assume that  $p$  does not fork over  $M_1$  and  $p \upharpoonright M_1$  does not fork over  $M_0$ . Then  $p$  does not fork over  $M_0$ .
- (7) Universal continuity: Let  $\delta < \lambda^+$  be a limit ordinal and let  $\langle M_i : i \leq \delta \rangle$  be an increasing continuous chain of limit models with  $M_{i+1}$  universal over  $M_i$ . Let  $\langle p_i : i < \delta \rangle$  be given such that for all  $i < \delta$ ,  $p_i \in \mathbf{S}(M_i)$ , and  $p_i$  is a nonforking extension of  $p_0$ . Then there exists a unique  $p_\delta \in \mathbf{S}(M_\delta)$  such that  $p_\delta$  does not fork over  $M_0$ . In particular,  $p_\delta$  extends each  $p_i$ .
- (8) Disjointness: If  $M \leq_{\mathbf{K}} N$  are both limit models in  $\mathbf{K}_\lambda$  and  $p \in \mathbf{S}(N)$  does not fork over  $M$ , then  $p$  is algebraic if and only if  $p \upharpoonright M$  is algebraic.

*Proof.* Invariance, monotonicity, and universal local character are straightforward to check from the definition. Extension is by [Van06, I.4.10] and universal local character. Uniqueness is [Vas17b, 2.16], and transitivity follows on general grounds (see e.g. the proof of [She09a, II.2.18]). To prove universal continuity, take  $p_\delta$  to

be the nonforking extension of  $p_0$  and use uniqueness and universal local character. Finally, to see disjointness, first use universal local character and transitivity to find  $M_0$  and a limit ordinal  $\delta < \lambda^+$  so that  $M$  is  $(\lambda, \delta)$ -limit over  $M_0$  and  $p$  does not fork over  $M_0$ . Let  $N'$  be  $(\lambda, \delta)$ -limit over  $N$ , hence over  $M_0$ . Let  $q$  be the nonforking extension of  $p$  to  $N'$ . If  $p \upharpoonright M$  is algebraic, then clearly  $p$  is algebraic. Now if  $p$  is algebraic, then  $q$  is algebraic. By [Vas17b, 2.7],  $q$  and  $p \upharpoonright M$  are conjugates, so  $p \upharpoonright M$  is also algebraic, as desired.  $\square$

**Fact 2.5** (The canonicity theorem). Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC. Assume we have a relation “ $p$  is free over  $M$ ” for a type  $p \in \mathbf{S}(N)$  and  $M, N \in \mathbf{K}_\lambda$  limit models. If this relation satisfies invariance, monotonicity, universal local character, uniqueness, and extension (in the sense given in the statement of Fact 2.4), then  $p$  is free over  $M$  if and only if  $p$  does not  $\lambda$ -fork over  $M$ .

*Proof.* As in the proof of [Vas16a, 9.6].  $\square$

The following property of forking is crucial. It is open whether it follows from superstability. The definition we give is not the same as in [Van16a] but is equivalent by [Vas17b, 2.18].

**Definition 2.6.** Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC. We say that  $\mathbf{K}$  has  $\lambda$ -symmetry if for any two limit model  $M \leq_{\mathbf{K}} N$  in  $\mathbf{K}_\lambda$  and any  $a, b \in N$ , the following are equivalent:

- (1) There exists  $M_b \leq_{\mathbf{K}} N_b$  in  $\mathbf{K}_\lambda$  both limits such that  $N \leq_{\mathbf{K}} N_b$ ,  $M \leq_{\mathbf{K}} M_b$ ,  $b \in M_b$ , and  $\text{tp}(a/M_b; N_b)$  does not fork over  $M$ .
- (2) There exists  $M_a \leq_{\mathbf{K}} N_a$  in  $\mathbf{K}_\lambda$  both limits such that  $N \leq_{\mathbf{K}} N_a$ ,  $M \leq_{\mathbf{K}} M_a$ ,  $a \in M_a$ , and  $\text{tp}(b/M_a; N_a)$  does not fork over  $M$ .

Symmetry implies the uniqueness of limit models:

**Fact 2.7** ([Van16a]). Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC with  $\lambda$ -symmetry. Let  $M_0, M_1, M_2 \in \mathbf{K}_\lambda$  be given. If both  $M_1$  and  $M_2$  are limit over  $M_0$ , then  $M_1 \cong_{M_0} M_2$ . In particular, any two limit models in  $\mathbf{K}_\lambda$  are isomorphic.

This leads to yet another key property of forking:

**Fact 2.8** (The conjugation property). Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC with  $\lambda$ -symmetry. Let  $M \leq_{\mathbf{K}} N$  be limit models in  $\mathbf{K}_\lambda$  and let  $p \in \mathbf{S}(N)$ . If  $p$  does not fork over  $M$ , then there is an isomorphism  $f : N \cong M$  such that  $f(p) = p \upharpoonright M$ .

*Proof.* As in [She09a, III.1.21].  $\square$

It is worth noting that both superstability and symmetry follows from categoricity, in an AEC with amalgamation and no maximal models. We will state stronger results in Section 4, but it will be easier to quote from:

**Fact 2.9** (Structure of categorical AECs with amalgamation). Let  $\mathbf{K}$  be an AEC with arbitrarily large models. Let  $\mu > \text{LS}(\mathbf{K})$  be such that  $\mathbf{K}_{<\mu}$  has amalgamation and no maximal models. If  $\mathbf{K}$  is categorical in  $\mu$ , then:

- (1) For any  $\lambda \in [\text{LS}(\mathbf{K}), \mu)$ ,  $\mathbf{K}$  is  $\lambda$ -superstable and has  $\lambda$ -symmetry.

- (2) For any  $\lambda \in (\text{LS}(\mathbf{K}), \mu]$ ,  $\mathbf{K}^{\lambda\text{-sat}}$  is an AEC with  $\text{LS}(\mathbf{K}^{\lambda\text{-sat}}) = \lambda$ . In particular, the model of cardinality  $\mu$  is saturated.

*Proof.* That superstability holds below the categoricity cardinal essentially appears in [SV99, 2.2.1], but a full proof is in [BGVV17]. That symmetry similarly follows from categoricity is in [Vas17c, 5.7], and the statement on saturated models is also proven there (it is a direct consequence of [Van16b]).  $\square$

**2.7. Good frames.** Good  $\lambda$ -frames were introduced by Shelah in [She09a, II] as a bare-bone axiomatization of superstability. We give a simplified definition here.

**Definition 2.10** ([She09a, II.2.1]). A *good  $\lambda$ -frame* is a triple  $\mathfrak{s} = (\mathbf{K}, \perp, \mathbf{S}^{\text{bs}})$  where:

- (1)  $\mathbf{K}$  is an AEC such that:
  - (a)  $\lambda \geq \text{LS}(\mathbf{K})$ .
  - (b)  $\mathbf{K}_\lambda \neq \emptyset$ .
  - (c)  $\mathbf{K}_\lambda$  has amalgamation, joint embedding, and no maximal models.
  - (d)  $\mathbf{K}$  is stable<sup>3</sup> in  $\lambda$ .
- (2) For each  $M \in \mathbf{K}_\lambda$ ,  $\mathbf{S}^{\text{bs}}(M)$  (called the set of *basic types* over  $M$ ) is a set of nonalgebraic types over  $M$  satisfying the *density property*: if  $M <_{\mathbf{K}} N$  are both in  $\mathbf{K}_\lambda$ , there exists  $a \in |N| \setminus |M|$  such that  $\text{tp}(a/M; N) \in \mathbf{S}^{\text{bs}}(M)$ .
- (3)  $\perp$  is an (abstract) independence relation on the basic types satisfying invariance, monotonicity, extension existence, uniqueness, continuity, local character, and symmetry (see [She09a, II.2.1] for the full definition of these properties).

We say that  $\mathfrak{s}$  is *type-full* [She09a, III.9.2(1)] if for any  $M \in \mathbf{K}_\lambda$ ,  $\mathbf{S}^{\text{bs}}(M) = \mathbf{S}^{\text{na}}(M)$ , the set of *all* nonalgebraic types over  $M$ . Rather than explicitly using the relation  $\perp$ , we will say that  $\text{tp}(a/M; N)$  *does not  $\mathfrak{s}$ -fork* over  $M_0$  if  $a \perp_{M_0}^N M$  (this is well-defined by the invariance and monotonicity properties). When  $\mathfrak{s}$  is clear from context, we omit it (this does not conflict with previous terminology by Facts 2.5 and 2.12). We say that a good  $\lambda$ -frame  $\mathfrak{s}$  is *on  $\mathbf{K}$*  if the underlying AEC of  $\mathfrak{s}$  is  $\mathbf{K}$ . We say that  $\mathfrak{s}$  is *categorical* if  $\mathbf{K}$  is categorical in  $\lambda$ .

**Remark 2.11.** We will *not* use the axiom (B) [She09a, II.2.1] requiring the existence of a superlimit model of size  $\lambda$ . In fact many papers (e.g. [JS13]) define good frames without this assumption. Further, we gave a shorter list of properties that in Shelah's original definition, but the other properties follow, see [She09a, II.2].

The reader can forget about the class of basic types: in this paper, we will work exclusively with type-full frames. In this case, the existence of a good frame is stronger than superstability:

**Fact 2.12.** If the AEC  $\mathbf{K}$  has a type-full good  $\lambda$ -frame, then  $\mathbf{K}$  is  $\lambda$ -superstable and has  $\lambda$ -symmetry.

<sup>3</sup>In Shelah's original definition, only the set of basic types is required to be stable. However full stability follows, see [She09a, II.4.2].

*Proof.* By [BGKV16, 4.2],  $\mathfrak{s}$ -nonforking implies  $\lambda$ -nonsplitting. This immediately gives that  $\mathbf{K}$  is  $\lambda$ -superstable. Using canonicity (Fact 2.5),  $\lambda$ -symmetry then follows from the symmetry axiom of good frames.  $\square$

The converse, getting a good frame from superstability and symmetry, is one of the main focus of this paper. One issue is that in a  $\lambda$ -superstable AEC,  $\lambda$ -nonforking is only well-behaved over limit models, and in general the class of limit models may not be closed under unions, hence may not form an AEC. If  $\lambda > \text{LS}(\mathbf{K})$ , the class of limit models is the same as the class of saturated models in  $\mathbf{K}_\lambda$  and we often *do* get that it is closed under unions (see e.g. Fact 2.9). Restricting to the class of limit models in  $\lambda$ , we then get that the AEC is categorical in  $\lambda$ , so we might as well assume this to begin with. A more serious issue is that the local character and continuity properties given by Fact 2.4 are weaker than the corresponding ones for good frames (because in the definition of local character for good frames we do not require that the models in the chain are universal over the previous ones). This will be circumvented by using the theory of towers and previous result on what Jarden and Shelah call *almost good frames*: good frames that still have continuity but have only the weak version of local character of superstable AECs. For now we state what we can get from Fact 2.4:

**Fact 2.13.** If  $\mathbf{K}$  be an AEC which is  $\lambda$ -superstable, has  $\lambda$ -symmetry, and is categorical in  $\lambda$ , then  $\lambda$ -nonforking induces a triple  $\mathfrak{s} = (\mathbf{K}, \perp, \mathbf{S}^{\text{na}})$  which satisfies all the axioms of a type-full good  $\lambda$ -frame, except perhaps for local character and continuity. Moreover, it has the conjugation property (in the sense of Fact 2.8) and satisfies the universal versions of local character and continuity stated in Fact 2.4.

*Proof.* Immediate from Fact 2.4 and the definition of superstability.  $\square$

**Remark 2.14.** Such weak good frames are called (modulo very minor variations in the definition) *H-almost good frames* by Shelah [She09b, VII.5.9].

In [She09a, §III.1], Shelah defines the following positive properties of good frames: weakly successful, successful,  $n$ -successful, and  $\text{good}^+$ . He also defines what it means to take the successor  $\mathfrak{s}^+$  of a successful  $\text{good}^+$  frame and even to take the  $n$ th successor,  $\mathfrak{s}^{+n}$  of an  $n$ -successful  $\text{good}^+$  frame. We do not repeat the definitions here. The reader may look at [BV, §2] for an overview. We will mostly just use these terms as “black boxes”: they will appear in certain facts and theorems, but we will never need their exact definitions. The only fact that the reader must know is that if  $\mathfrak{s}$  is an  $n$ -successful  $\text{good}^+$  frame on the AEC  $\mathbf{K}$ ,  $n \geq 1$ , then  $\mathfrak{s}$  can be extended to be type-full and one can define its  $n$ th successor  $\mathfrak{s}^{+n}$  and this is a type-full good  $\lambda^{+n}$ -frame on  $\mathbf{K}^{\lambda^{+n}\text{-sat}}$ . If  $\mathfrak{s}$  is  $\omega$ -successful (i.e.  $n$ -successful for all  $n < \omega$ ), powerful results from [SVb] imply (assuming WGCH) that the good frame can be lifted to any cardinal above  $\lambda$  and, in a sense, the structure of the AEC above  $\lambda$  is completely understood. For an overview of what we will use, see Fact 8.2.

### 3. NICE STABILITY AND SUPERLIMITS

In this section, we prove several technical results in the setup of nice stability, implicit already in [SV99]. This turns out to be the right context to study stability

in AECs that may not have amalgamation. The main result is Theorem 3.19, showing roughly speaking that restricting to a superlimit (defined below) preserves nice stability and gives a class that is very close to the original one.

The notion of a *superlimit model* is another attempt at defining a local notion of saturation. It was introduced by Shelah [She09a, I.3.3]. We give a definition that makes sense in any abstract class, but we will apply it to  $\mathbf{K}_\lambda$ , for  $\mathbf{K}$  an AEC and  $\lambda \geq \text{LS}(\mathbf{K})$ .

**Definition 3.1.** Let  $\mathbf{K}$  be an abstract class and let  $M \in \mathbf{K}$ .

- (1)  $M$  is *universal* if for every  $M_0 \in \mathbf{K}$  there exists a  $\mathbf{K}$ -embedding  $f : M_0 \rightarrow M$ .
- (2)  $M$  is *superlimit* if:
  - (a)  $M$  has a proper extension.
  - (b)  $M$  is universal.
  - (c) Whenever  $\delta$  is a limit ordinal and  $\langle M_i : i \leq \delta \rangle$  is increasing continuous, if  $M \cong M_i$  for all  $i < \delta$ , then  $M \cong \bigcup_{i < \delta} M_i$ .

**Remark 3.2.** If an abstract class  $\mathbf{K}$  is categorical, then any model in  $\mathbf{K}$  is universal. If in addition  $\mathbf{K}$  has no maximal models, any model in  $\mathbf{K}$  is superlimit.

Given a superlimit, the class of models isomorphic to it generates an AEC:

**Definition 3.3** ([She09a, II.1.25]). Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ . For  $M$  a superlimit in  $\mathbf{K}_\lambda$ , let  $\mathbf{K}^{[M]}$ , the *AEC generated by  $M$* , be defined as follows:  $N \in \mathbf{K}^{[M]}$  if and only if for all  $A \subseteq |N|$  with  $|A| \leq \lambda$ , there exists  $N_0 \in \mathbf{K}$  such that  $N_0 \cong M$ ,  $A \subseteq |N_0|$ , and  $N_0 \leq_{\mathbf{K}} N$ . Order  $\mathbf{K}^{[M]}$  with the restriction of the ordering on  $\mathbf{K}$ .

We will use the following straightforward results without comments:

**Fact 3.4** ([She09a, II.1.26]). Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . If  $M$  is a superlimit in  $\mathbf{K}_\lambda$ , then:

- (1)  $\mathbf{K}^{[M]}$  is an AEC with  $\text{LS}(\mathbf{K}^{[M]}) = \lambda$ .
- (2)  $\mathbf{K}_{\leq \lambda}^{[M]} = \{N_0 \in \mathbf{K}_\lambda \mid N_0 \cong M\}$ . In particular,  $\mathbf{K}_{< \lambda}^{[M]} = \emptyset$  and  $\mathbf{K}^{[M]}$  is categorical in  $\lambda$ .
- (3)  $\mathbf{K}_\lambda^{[M]}$  has no maximal models.
- (4) For any  $N_0 \in \mathbf{K}_\lambda$ , there exists  $N \in \mathbf{K}_\lambda^{[M]}$  such that  $N_0 \leq_{\mathbf{K}} N$ .

Note that a superlimit is unique if it exists:

**Fact 3.5.** Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . If  $M$  and  $N$  are superlimits in  $\mathbf{K}_\lambda$ , then  $M \cong N$ . In particular,  $\mathbf{K}^{[M]} = \mathbf{K}^{[N]}$ .

*Proof.* Straightforward: build a chain with interleaved copies of  $M$  and  $N$ . See for example [She09a, I.3.7(1)].  $\square$

This justifies the following definition:

**Definition 3.6.** Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$  be such that  $\mathbf{K}_\lambda$  has a superlimit. We write  $\mathbf{K}^{\lambda\text{-sl}}$  for the class  $\mathbf{K}^{[M]}$ , where  $M \in \mathbf{K}_\lambda$  is superlimit (the choice of  $M$  does not matter by Fact 3.5).

The class  $\mathbf{K}_\lambda^{\lambda\text{-sl}}$  is dense in  $\mathbf{K}_\lambda$ , in the following sense<sup>4</sup>:

**Definition 3.7.** A class  $K^*$  of structures is *dense* in an abstract class  $\mathbf{K} = (K, \leq_{\mathbf{K}})$  if  $K^* \subseteq K$  and for any  $M \in \mathbf{K}$  there exists  $N \in K^*$  such that  $M \leq_{\mathbf{K}} N$ . We say that an abstract class  $\mathbf{K}^* = (K^*, \leq_{\mathbf{K}^*})$  is *dense in  $\mathbf{K}$*  if  $K^*$  is dense in  $\mathbf{K}$  and for  $M, N \in K^*$ ,  $M \leq_{\mathbf{K}^*} N$  if and only if  $M \leq_{\mathbf{K}} N$ .

**Remark 3.8.** By Fact 3.4, whenever  $\mathbf{K}$  is an AEC,  $\lambda \geq \text{LS}(\mathbf{K})$ , and  $\mathbf{K}_\lambda$  has a superlimit, then  $\mathbf{K}_\lambda^{\lambda\text{-sl}}$  is dense in  $\mathbf{K}_\lambda$ .

We have the following obvious transitivity property:

**Remark 3.9.** If  $\mathbf{K}^{**}$  is dense in  $\mathbf{K}^*$  and  $\mathbf{K}^*$  is dense in  $\mathbf{K}$ , then  $\mathbf{K}^{**}$  is dense in  $\mathbf{K}$ .

Many properties of an abstract class are preserved when passing to a dense subclass. To state the next remark, we need the following definition (which is already used in [SV99]):

**Definition 3.10.** Let  $\mathbf{K}$  be an abstract class. An *amalgamation base* is a model  $M_0 \in \mathbf{K}$  such that for any  $M_1, M_2 \in \mathbf{K}$  with  $M_0 \leq_{\mathbf{K}} M_\ell$ ,  $\ell = 1, 2$ , there exists  $M_3 \in \mathbf{K}$  and  $\mathbf{K}$ -embeddings  $f_\ell : M_\ell \xrightarrow{M_0} M_3$ .

The following combinatorial result of Shelah [She09a, I.3.8] gives us a way to find amalgamation bases:

**Fact 3.11.** Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . Assume  $\text{WGCH}(\lambda)$ . If  $M$  is superlimit in  $\mathbf{K}_\lambda$  and  $\mathbf{K}_{\lambda+}$  has a universal model, then  $M$  is an amalgamation base in  $\mathbf{K}_\lambda$ .

Note that an abstract class has amalgamation precisely when all its models are amalgamation bases. The following are all easy consequences of the definitions (many appear already in [Vas16a, §5]). They will be used without further comments:

**Remark 3.12.** Let  $\mathbf{K}$  be an abstract class and let  $\mathbf{K}^*$  be dense in  $\mathbf{K}$ .

- (1)  $\mathbf{K} \neq \emptyset$  if and only if  $\mathbf{K}^* \neq \emptyset$ .
- (2)  $\mathbf{K}$  has no maximal models if and only if  $\mathbf{K}^*$  has no maximal models.
- (3)  $\mathbf{K}$  has joint embedding if and only if  $\mathbf{K}^*$  has joint embedding.
- (4) For any  $M \in \mathbf{K}^*$ ,  $M$  is an amalgamation base in  $\mathbf{K}^*$  if and only if  $M$  is an amalgamation base in  $\mathbf{K}$ .
- (5) For any  $M \in \mathbf{K}^*$ ,  $M$  is superlimit in  $\mathbf{K}^*$  if and only if  $M$  is superlimit in  $\mathbf{K}$ .
- (6) If  $\mathbf{K}^*$  is closed under unions of  $\omega$ -chains, then any superlimit (in  $\mathbf{K}$ ) is in  $\mathbf{K}^*$ .

Roughly, an AEC is nicely stable if its class of amalgamation bases is dense and behaves like a stable first-order theory. As usual, this is localized to a fixed cardinal  $\lambda$ . The definition appears for the first time in [SVa, 2.3] but is studied already in [SV99].

---

<sup>4</sup>This is a special case of the definition of a skeleton, see [Vas16a, 5.3] but since we have no use for skeletons in this paper, we chose to only study the simpler case.

**Definition 3.13.** An AEC  $\mathbf{K}$  is *nicely  $\lambda$ -stable* (or *nicely stable in  $\lambda$* ) if:

- (1)  $\lambda \geq \text{LS}(\mathbf{K})$  and  $\mathbf{K}_\lambda \neq \emptyset$ .
- (2)  $\mathbf{K}_\lambda$  has joint embedding and no maximal models.
- (3) Density of amalgamation bases: for any  $M \in \mathbf{K}_\lambda$  there exists  $N \in \mathbf{K}_\lambda$  which is an amalgamation base in  $\mathbf{K}_\lambda$  and so that  $M \leq_{\mathbf{K}} N$ .
- (4) For any amalgamation base (in  $\mathbf{K}_\lambda$ )  $M$ , there exists an amalgamation base  $N \in \mathbf{K}_\lambda$  with  $N$  universal over  $M$ .
- (5) Any limit model in  $\mathbf{K}_\lambda$  is an amalgamation base (in  $\mathbf{K}_\lambda$ ).

For  $\Theta$  a class of cardinals, we say that  $\mathbf{K}$  is *nicely  $\Theta$ -stable* (or *nicely stable in  $\Theta$* ) if  $\mathbf{K}$  is nicely  $\lambda$ -stable for every  $\lambda \in \Theta$ .

**Remark 3.14.** Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . If  $\mathbf{K}_\lambda \neq \emptyset$ ,  $\mathbf{K}_\lambda$  has amalgamation, joint embedding, no maximal models, and  $\mathbf{K}$  is stable in  $\lambda$ , then  $\mathbf{K}$  is nicely  $\lambda$ -stable (universal extensions exist by [She09a, II.1.16]; all the other properties are easy to check). In particular, if  $\mathbf{K}$  is  $\lambda$ -superstable then  $\mathbf{K}$  is nicely  $\lambda$ -stable. In setups without amalgamation, it is known [SV99] that nice stability follows from categoricity, no maximal models, and GCHWD. See Fact 4.7 here.

We will often assume in addition to nice stability that there is a superlimit. In this case, the superlimit will be limit, for all possible lengths. For the convenience of the reader, we sketch a proof.

**Lemma 3.15.** Let  $\mathbf{K}$  be a nicely  $\lambda$ -stable AEC. If  $M$  is superlimit in  $\mathbf{K}_\lambda$ , then  $M$  is  $(\lambda, \delta)$ -limit for all limit ordinals  $\delta < \lambda^+$ .

*Proof.* Fix a limit ordinal  $\delta < \lambda^+$ . We build  $\langle M_i : i \leq \delta \rangle$ ,  $\langle N_i : i \leq \delta \rangle$  increasing continuous in  $\mathbf{K}_\lambda$  such that for all  $i < \delta$ :

- (1)  $M_i \cong M$ .
- (2)  $N_i$  is an amalgamation base.
- (3)  $N_{i+1}$  is universal over  $N_i$ .
- (4)  $M_i \leq_{\mathbf{K}} N_i \leq_{\mathbf{K}} M_{i+1}$ .

This is possible: take  $M_0 := M$  and  $N_0$  an amalgamation base containing  $M_0$ . For  $i$  limits, take unions ( $N_i$  is an amalgamation base since it is a limit model, and  $M_i \cong M$  by definition of a superlimit). Given  $M_i$  and  $N_i$ , use universality of the superlimit to pick  $M_{i+1}$  containing  $N_i$  which is isomorphic to  $M$ . Then pick  $N'_{i+1}$  an amalgamation base containing  $M_{i+1}$  and let  $N_{i+1}$  be universal over  $N'_{i+1}$ .

This is enough: the chains are interleaved so  $M_\delta = N_\delta$ . By definition of a superlimit,  $M_\delta \cong M$  and by construction  $N_\delta$  is  $(\lambda, \delta)$ -limit.  $\square$

The following technical result will be used in Section 9:

**Lemma 3.16.** Let  $\mathbf{K}$  be a nicely  $\lambda$ -stable AEC. Let  $M, N \in \mathbf{K}_\lambda$  be superlimits. If  $M \leq_{\mathbf{K}} N$ , then  $M \preceq_{\mathbb{L}_{\infty, \lambda}} N$ .

*Proof.* Let  $\phi(\bar{x})$  be an  $\mathbb{L}_{\infty, \lambda}$  formula and let  $\bar{a}$  be a sequence of length  $\alpha$  in  $M$ , with  $\alpha < \lambda$ . We want to see that  $M \models \phi[\bar{a}]$  if and only if  $N \models \phi[\bar{a}]$ . We proceed by induction on the structure of  $\phi$ . If  $\phi$  is quantifier-free, then this holds because by the definition of an abstract class,  $M \subseteq N$ . If  $\phi$  is a conjunction, disjunction,

or negation, the result follows from the induction hypothesis. Assume now that  $\phi = \exists \bar{y} \psi(\bar{x}; \bar{y})$ , where  $\bar{y}$  has length  $\beta < \lambda$ . If  $M \models \phi[\bar{a}]$ , then by the induction hypothesis  $N \models \phi[\bar{a}]$ . Assume now that  $N \models \phi[\bar{a}]$  and pick  $\bar{b}$  in  $N$  such that  $N \models \psi[\bar{a}; \bar{b}]$ . Let  $\theta := |\alpha|^+ + |\beta|^+ + \aleph_0$ . Note that  $\theta \leq \lambda$  and  $\theta$  is a regular cardinal. By Lemma 3.15,  $M$  and  $N$  are limit models, hence are amalgamation bases. Thus we can pick  $N' \in \mathbf{K}_\lambda$  which is  $(\lambda, \theta)$ -limit over  $N$ , hence also limit over  $M$ . By Lemma 3.15 and uniqueness of limit models of the same length,  $N'$  is superlimit. By what has just been said,  $N' \models \psi[\bar{a}; \bar{b}]$ . Now by Lemma 3.15 again,  $M$  is  $(\lambda, \theta)$ -limit over some  $M_0$ . By making  $M_0$  bigger if necessary (noting that  $\alpha < \theta$ ), we can assume that  $\bar{a}$  is contained entirely in  $M_0$ . Note that  $N'$  is also  $(\lambda, \theta)$ -limit over  $M_0$ , so by uniqueness of limit models of the same length, there is an isomorphism  $f : N' \cong_{M_0} M$ . Thus  $M \models \psi[\bar{a}; f(\bar{b})]$ , hence  $M \models \phi[\bar{a}]$ , as desired.  $\square$

The next result says that nice stability plays very well with taking dense subclasses:

**Theorem 3.17.** Let  $\mathbf{K}$  and  $\mathbf{K}^*$  be AECs and let  $\lambda \geq \text{LS}(\mathbf{K}) + \text{LS}(\mathbf{K}^*)$ . If  $\mathbf{K}_\lambda^*$  is dense in  $\mathbf{K}_\lambda$ , then  $\mathbf{K}$  is nicely  $\lambda$ -stable if and only if  $\mathbf{K}^*$  is nicely  $\lambda$ -stable.

*Proof.* First, check that for any  $M_0 \leq_{\mathbf{K}} M$  both in  $\mathbf{K}_\lambda$ , if  $M$  is limit over  $M_0$  (in  $\mathbf{K}_\lambda$ ), then  $M \in \mathbf{K}_\lambda^*$  (write  $M$  as an increasing union of models in  $\mathbf{K}^*$ ). It follows that limit models in  $\mathbf{K}$  and  $\mathbf{K}^*$  coincide. Notice also that if  $M_0 \leq_{\mathbf{K}} M_1 \leq_{\mathbf{K}} M_2$ ,  $M_2$  is universal over  $M_1$ , and  $M_0$  is an amalgamation base, then  $M_2$  is universal over  $M_0$ . Thus it is enough to check existence of universal extensions on a dense class of amalgamation bases. The rest of the proof is straightforward (see also Remark 3.12).  $\square$

We now aim to show that if  $\mathbf{K}$  is nicely  $[\lambda, \lambda^{+n}]$ -stable and  $\mathbf{K}_\lambda$  is a superlimit, then  $\mathbf{K}^{\lambda\text{-sl}}$  (the class generated by the superlimit) is also nicely  $[\lambda, \lambda^{+n}]$ -stable. The  $n = 0$  case is given by what has just been proven, but for  $n > 0$  we will need to work a little bit more.

For  $\mathbf{K}$  a nicely  $\lambda$ -stable AEC, we call  $M \in \mathbf{K}$   $(\lambda, \lambda^+)$ -limit if there exists an increasing continuous chain  $\langle M_i : i \leq \lambda^+ \rangle$  such that  $M = M_{\lambda^+}$  and for all  $i < \lambda^+$ ,  $M_i \in \mathbf{K}_\lambda$  and  $M_{i+1}$  is universal over  $M_i$ . By the proof of [Van16b, Proposition 14], we have:

**Fact 3.18.** In a nicely  $[\lambda, \lambda^+]$ -stable AEC, a model is  $(\lambda, \lambda^+)$ -limit if and only if it is  $(\lambda^+, \lambda^+)$ -limit.

**Theorem 3.19.** Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ , and let  $n < \omega$ . If  $\mathbf{K}$  is nicely  $[\lambda, \lambda^{+n}]$ -stable and  $\mathbf{K}_\lambda$  has a superlimit, then:

- (1) Any  $(\lambda^{+n}, \lambda^{+(n+1)})$ -limit is in  $\mathbf{K}^{\lambda\text{-sl}}$ . In particular for every  $m \leq n$ ,  $\mathbf{K}_{\lambda^{+m}}^{\lambda\text{-sl}}$  is dense in  $\mathbf{K}_{\lambda^{+m}}$ .
- (2)  $\mathbf{K}^{\lambda\text{-sl}}$  is nicely  $[\lambda, \lambda^{+n}]$ -stable.

*Proof.*

- (1) By induction on  $n$ . If  $n = 0$ , note that by Lemma 3.15, the superlimit in  $\mathbf{K}_\lambda$  is  $(\lambda, \lambda)$ -limit. Moreover, it is easy to check that the  $(\lambda, \lambda^+)$ -limit is a union of an increasing chains of  $(\lambda, \lambda)$ -limits, hence of members of  $\mathbf{K}^{\lambda\text{-sl}}$ . Thus the  $(\lambda, \lambda^+)$ -limit is in  $\mathbf{K}^{\lambda\text{-sl}}$ . If  $n > 0$ , then similarly the



$(\lambda^{+n}, \lambda^{+(n+1)})$ -limit is a union of  $(\lambda^{+n}, \lambda^{+n})$ -limits. By Fact 3.18, these are  $(\lambda^{+(n-1)}, \lambda^{+n})$ -limits. By the induction hypothesis, these are all in  $\mathbf{K}^{\lambda\text{-sl}}$ , hence the  $(\lambda^{+n}, \lambda^{+(n+1)})$ -limit is in  $\mathbf{K}^{\lambda\text{-sl}}$ . The “in particular” part follows: if  $m = 0$ , this is Remark 3.8, and if  $m > 0$  then by definition of nice stability, any model in  $\mathbf{K}_{\lambda^{+m}}$  is contained in an  $(\lambda^{+m}, \lambda^{+m})$ -limit, hence (by Fact 3.18) in an  $(\lambda^{+(m-1)}, \lambda^{+m})$ -limit, which is in  $\mathbf{K}^{\lambda\text{-sl}}$ .

(2) By the previous part and Theorem 3.17.

□

We will often be interested in situations where the AEC is not just nicely stable but nicely superstable, in the sense that it also has a superlimit and the class generated by the superlimit is superstable.

**Definition 3.20.** An AEC  $\mathbf{K}$  is *nicely  $\lambda$ -superstable* (or *nicely superstable in  $\lambda$* ) if:

- (1)  $\mathbf{K}$  is nicely  $\lambda$ -stable.
- (2)  $\mathbf{K}_\lambda$  has a superlimit.
- (3)  $\mathbf{K}^{\lambda\text{-sl}}$  is  $\lambda$ -superstable and has  $\lambda$ -symmetry.

For  $\Theta$  a class of cardinals, we say that  $\mathbf{K}$  is *nicely  $\Theta$ -superstable* (or *nicely superstable in  $\Theta$* ) if  $\mathbf{K}$  is nicely  $\lambda$ -superstable for every  $\lambda \in \Theta$ .

We caution the reader: nice superstability does not immediately imply superstability, since amalgamation is not assumed. However we have:

**Fact 3.21.** Let  $\mathbf{K}$  be an AEC.

- (1) If  $\mathbf{K}$  is  $\lambda$ -superstable, has  $\lambda$ -symmetry, and  $\mathbf{K}_\lambda$  has a superlimit, then  $\mathbf{K}$  is nicely  $\lambda$ -superstable.
- (2) If  $\mathbf{K}$  is  $\lambda$ -superstable,  $\lambda^+$ -superstable, and has  $\lambda^+$ -symmetry, then  $\mathbf{K}$  is nicely  $\lambda^+$ -superstable.
- (3) If  $\mathbf{K}$  is nicely  $\lambda$ -superstable, then  $\mathbf{K}^{\lambda\text{-sl}}$  is  $\lambda$ -superstable, has  $\lambda$ -symmetry, and is categorical in  $\lambda$ .

*Proof.* The first and the second are immediate from the definition. To see the second, apply [Van16b, Theorem 1] and the first. □

#### 4. AECS WITH ARBITRARILY LARGE MODELS

An AEC  $\mathbf{K}$  has *arbitrarily large models* if  $\mathbf{K}_{\geq \chi} \neq \emptyset$  for all cardinals  $\chi$ . Equivalently,  $\mathbf{K}$  is a large category. In this section, we recall some basic facts about AECS with arbitrarily large models. We study in particular a weakening of categoricity called *semisolvability*, equivalent in the first-order case to superstability.

The following is a sufficient condition for an AEC to have arbitrarily large models:

**Fact 4.1** ([She09a, I.1.11]). Let  $\mathbf{K}$  be an AEC. If  $\mathbf{K}_{\geq \chi} \neq \emptyset$  for all  $\chi < h(\text{LS}(\mathbf{K}))$ , then  $\mathbf{K}$  has arbitrarily large models.

Clearly, if an AEC has no maximal models (and is not empty), then it has arbitrarily large models. We will make use of the following weakening of having no maximal models. Such notions are implicit already in both [She09a, Chapter IV] and [She15].

**Definition 4.2.** Let  $\mathbf{K}$  be an AEC and let  $\lambda < \mu$  be infinite cardinals. We call  $\mathbf{K}$   $(\lambda, \mu)$ -*extendible* if for any  $M \in \mathbf{K}_\lambda$  there exists  $N \in \mathbf{K}_\mu$  such that  $M \leq_{\mathbf{K}} N$ . When  $\lambda \geq \mu$ , we assume by convention that  $\mathbf{K}$  is always  $(\lambda, \mu)$ -extendible. We say that  $\mathbf{K}$  is  $\lambda$ -*extendible* if  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible for all  $\mu > \lambda$  and we say that  $\mathbf{K}$  is *extendible* if  $\mathbf{K}$  is  $\lambda$ -extendible for all  $\lambda$ . More generally, for  $\Theta$  a class of cardinals, we say that  $\mathbf{K}$  is  $(\Theta, \mu)$ -*extendible* if  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible for all  $\lambda \in \Theta$ , and we say that  $\mathbf{K}$  is  $\Theta$ -*extendible* if  $\mathbf{K}$  is  $\lambda$ -extendible for all  $\lambda \in \Theta$ .

The following basic properties of extendibility may sometimes be used without mention.

**Remark 4.3.** Let  $\mathbf{K}$  be an AEC.

- (1) Let  $\lambda < \mu_0 \leq \mu_1$ , with  $\mu_0 \geq \text{LS}(\mathbf{K})$ . If  $\mathbf{K}$  is  $(\lambda, \mu_1)$ -extendible, then  $\mathbf{K}$  is  $(\lambda, \mu_0)$ -extendible.
- (2) If  $\mathbf{K}$  is  $(\lambda_0, \lambda_1)$ -extendible and  $(\lambda_1, \lambda_2)$ -extendible, then  $\mathbf{K}$  is  $(\lambda_0, \lambda_2)$ -extendible.
- (3) If  $\mathbf{K}$  is  $\lambda$ -extendible for some  $\lambda$ , then  $\mathbf{K}$  has arbitrarily large models.
- (4) For  $\lambda < \mu$ ,  $\mathbf{K}$  is  $([\lambda, \mu], \mu)$ -extendible if and only if  $\mathbf{K}_{[\lambda, \mu]}$  has no maximal models. In particular,  $\mathbf{K}$  is extendible if and only if  $\mathbf{K}$  has no maximal models.

We recall Shelah's definition of solvability [She09a, Definition IV.1.4], using a more convenient notation for it with only one cardinal parameter, introduced in [Vas17c]. We also introduce a weakening, semisolvability in  $\lambda$ , which only asks for the EM model generated by  $\lambda$  to be universal. This has the same name, but is weaker than, the notion introduced in [Vas17c, 3.1]. Nevertheless, all the proofs there go through with the weaker notion. Both solvability and semisolvability are equivalent to superstability in the first-order case (see [BGVV17] and [GV17, 5.3]). Shelah writes that solvability is perhaps the true analog of superstability in abstract elementary classes [She09a, N§4(B)].

**Definition 4.4.** Let  $\mathbf{K}$  be an AEC and let  $\mu \geq \text{LS}(\mathbf{K})$ .

- (1) Let  $\Upsilon[\mathbf{K}]$  denote the set of Ehrenfeucht-Mostowski (EM) blueprints  $\Phi$  with  $|\tau(\Phi)| \leq \text{LS}(\mathbf{K})$ . See [She09a, IV.0.8] for the full definition. As is standard, for a linear order  $I$  we write  $\text{EM}(I, \Phi)$  for the EM model generated by  $I$  and  $\Phi$ , and  $\text{EM}_\tau(I, \Phi)$  for its reduct to  $\tau$ .
- (2) [She09a, IV.1.4.(1)] We say that  $\Phi$  *witnesses  $\mu$ -solvability* if:
  - (a)  $\Phi \in \Upsilon[\mathbf{K}]$ .
  - (b) If  $I$  is a linear order of size  $\mu$ , then  $\text{EM}_{\tau(\mathbf{K})}(I, \Phi)$  is superlimit in  $\mathbf{K}_\mu$  (recall Definition 3.1).
- (3)  $\Phi$  *witnesses  $\mu$ -semisolvability* if:
  - (a)  $\Phi \in \Upsilon[\mathbf{K}]$
  - (b)  $\text{EM}_{\tau(\mathbf{K})}(\mu, \Phi)$  is universal in  $\mathbf{K}_\mu$ .
- (4)  $\mathbf{K}$  is  $\mu$ -[semi]solvable if there exists  $\Phi$  witnessing  $\mu$ -[semi]solvability.
- (5) For a class  $\Theta$  of cardinals,  $\mathbf{K}$  is  $\Theta$ -[semi]solvable if  $\mathbf{K}$  is  $\mu$ -[semi]solvable for every  $\mu \in \Theta$ .

Directly from the definitions, we have:

**Remark 4.5.** Let  $\mathbf{K}$  be an AEC and let  $\mu \geq \text{LS}(\mathbf{K})$ .

- (1) If  $\mathbf{K}$  has arbitrarily large models and is categorical in  $\mu$ , then  $\mathbf{K}$  is  $\mu$ -solvable.
- (2) If  $\mathbf{K}$  is  $\mu$ -solvable, then  $\mathbf{K}$  is  $\mu$ -semisolvable.
- (3) If  $\mathbf{K}$  is  $\mu$ -semisolvable, then  $\mathbf{K}$  is  $\mu$ -extendible. In particular,  $\mathbf{K}$  has arbitrarily large models.
- (4) If  $\mathbf{K}$  is  $\mu$ -[semi]solvable, then for any  $\lambda \in [\text{LS}(\mathbf{K}), \mu]$ ,  $\mathbf{K}_{\geq \lambda}$  is  $\mu$ -[semi]solvable.

In [Vas17c, 5.1], it was shown that solvability transfers down in AECs with amalgamation and no maximal models (this is of interest, since we do not know whether *categoricity* itself transfers down). The reason for considering semisolvability here is that we can transfer it down using just no maximal models (and in fact extendibility in the relevant cardinals suffices). A very similar argument appears already in [SV99].

**Fact 4.6.** Let  $\mathbf{K}$  be an AEC and let  $\mu > \lambda \geq \text{LS}(\mathbf{K})$ . Assume that  $\mathbf{K}$  is  $\mu$ -semisolvable. The following are equivalent:

- (1)  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible.
- (2)  $\mathbf{K}$  is  $\lambda$ -semisolvable.

*Proof.* If  $\mathbf{K}$  is semisolvable in  $\lambda$ , then  $\mathbf{K}$  is  $\lambda$ -extendible, hence  $(\lambda, \mu)$ -extendible. Conversely, assume that  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible and suppose that  $\Phi$  is an EM blueprint witnessing  $\mu$ -semisolvability. Since  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible, any  $M \in \mathbf{K}_\lambda$  embeds inside  $\text{EM}_{\tau(\mathbf{K})}(\lambda, \Phi)$ , hence inside  $\text{EM}_{\tau(\mathbf{K})}(A, \Phi)$ , for some  $A \subseteq \lambda$  with  $|A| = \lambda$ . In particular,  $\lambda \leq \text{otp}(A) < \lambda^+$ , so by renaming,  $M$  also embeds inside  $\text{EM}_{\tau(\mathbf{K})}(\text{otp}(A), \Phi)$ . We have shown that any model in  $\mathbf{K}_\lambda$  embeds inside  $\text{EM}_{\tau(\mathbf{K})}(\alpha, \Phi)$  for some  $\alpha \in [\lambda, \lambda^+)$ . Now by (for example) [Bal09, 15.5], the linear order  $I := {}^{<\omega}\lambda$  (ordered lexicographically) embeds any ordinal  $\alpha < \lambda^+$ , thus  $\text{EM}_{\tau(\mathbf{K})}(I, \Phi)$  is universal in  $\mathbf{K}_\lambda$ . Now by expanding the vocabulary, build a new blueprint  $\Psi$  such that for any linear order  $J$ ,  $\text{EM}_{\tau(\mathbf{K})}(J, \Psi) = \text{EM}_{\tau(\mathbf{K})}({}^{<\omega}J, \Phi)$ . Then by construction  $\Psi$  witnesses  $\lambda$ -semisolvability.  $\square$

We end this section by stating some known consequences of semisolvability that we will use.

**Fact 4.7.** Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . Assume that  $\mathbf{K}$  is  $[\lambda, \lambda^+]$ -semisolvable.

- (1) If  $\text{GCHWD}(\lambda)$ , then  $\mathbf{K}$  is nicely  $\lambda$ -stable.
- (2) If  $\mathbf{K}_\lambda$  has amalgamation, then  $\mathbf{K}$  is  $\lambda$ -superstable and has  $\lambda$ -symmetry.
- (3) If  $\mathbf{K}_\lambda$  has a superlimit which is an amalgamation base, then  $\mathbf{K}$  is nicely  $\lambda$ -superstable (recall Definition 3.20).
- (4) If  $\mathbf{K}_\lambda$  has a superlimit and  $\text{WGCH}(\lambda)$ , then  $\mathbf{K}$  is nicely  $\lambda$ -superstable.

*Proof.* Note that  $\mathbf{K}_\lambda$  has joint embedding and no maximal models by the semisolvability hypothesis. Of course, it is also not empty. Now:

- (1) By [SV99, §1]. See also [Van06, §3].
- (2) By [BGVV17],  $\mathbf{K}$  is  $\lambda$ -superstable. By [Vas17c, 4.8],  $\mathbf{K}$  has  $\lambda$ -symmetry.
- (3) Note that the usual argument of Morley (see e.g. [Bal09, 8.20] or Claim 1 in the proof of [Vas17c, 3.4]) shows that  $\mathbf{K}^{\lambda\text{-sl}}$  is stable in  $\lambda$ : if  $M$  is

- superlimit in  $\mathbf{K}_\lambda$ , then  $|\mathbf{S}(M)| \leq \lambda$ . Since  $\mathbf{K}_\lambda^{\lambda\text{-sl}}$  has amalgamation, it is nicely  $\lambda$ -stable (see Remark 3.14). By (for example) Theorem 3.17, This implies that  $\mathbf{K}$  is nicely  $\lambda$ -stable. The result now follows from [BGVV17].
- (4) By Fact 3.11 and the previous part.

□

## 5. TOWERS AND DISJOINT AMALGAMATION

In this section, we prove some technical lemmas relative to how much models can be amalgamated to be “as nonforking as possible”. This relies on the theory of towers. Towers were introduced in [SV99], and further studied in several papers since then (e.g. [Van06, Van13, GVV16, Van16a, VV17]). Essentially, everything before Lemma 5.28 is known and appears in some form in either [GVV16] or [Van16a]. We give some proofs and definitions here both because the statements are slightly different and the arguments have been simplified ([GVV16] and [Van16a] had to work without the uniqueness property of nonforking, proven in [Vas17b]; thus the definition of tower there is more complicated).

Throughout this section, we assume:

**Hypothesis 5.1.**  $\mathbf{K}$  is a  $\lambda$ -superstable AEC with  $\lambda$ -symmetry.

We work inside  $\mathbf{K}_\lambda$ : except if said otherwise, all models come from there. The following consequence of symmetry will be crucial. The idea is that we can make sure that two elements are independent “in a uniform way”.

**Fact 5.2** (Nonforking amalgamation). Let  $M_0 \leq_{\mathbf{K}} M_\ell$ ,  $\ell = 1, 2$ , be limit models. Let  $a_\ell \in M_\ell$ . There exists  $f_1, f_2, M_3$  such that  $M_3$  is limit and  $f_\ell : M_\ell \xrightarrow{M_0} M_3$  is such that  $\text{tp}(f_\ell(a_\ell)/f_{3-\ell}(M_{3-\ell}); M_3)$  does not fork over  $M_0$  for  $\ell = 1, 2$ .

*Proof.* As in [She09a, II.2.16].

□

To define towers, we first introduce some notation:

**Notation 5.3.**

- (1) The letter  $I$  will denote a well-ordering  $(|I|, <_I)$ . We usually write  $<$  instead of  $<_I$ .
- (2) For  $I$  a well-ordering, let  $I^-$  be the initial segment of  $I$  which is isomorphic to  $I$  if  $I$  is isomorphic to a limit ordinal or zero, or isomorphic to  $\alpha$  if  $I$  is isomorphic to  $\alpha + 1$ .
- (3) For  $I$  a well-ordering,  $i \in I$ , and  $\alpha$  an ordinal, let  $i +_I \alpha$  denote the unique element  $j \in I$  (if it exists) such that  $\text{otp}(j) = \text{otp}(i) + \alpha$ . We write  $i + \alpha$  instead of  $i +_I \alpha$  when  $I$  is clear from context.

**Definition 5.4.** A *tower*  $\mathcal{T}$  consists of  $\langle M_i : i \in I \rangle \frown \langle a_i : i \in I^- \rangle$ , where:

- (1)  $I$  is a well-ordering of cardinality at most  $\lambda$ .
- (2)  $\langle M_i : i \in I \rangle$  is an increasing chain of limit models, not necessarily continuous.
- (3)  $a_i \in M_{i+1} \setminus M_i$  for each  $i \in I^-$ .

We call  $I$  the *length* (or *index set*) of the tower. We call  $\mathcal{T}$  *continuous* if  $\langle M_i : i \in I \rangle$  is continuous. We say that  $\mathcal{T}$  is *limit* if  $M_{i+1}$  is limit over  $M_i$  for each  $i \in I^-$ . We may often identify a tower  $\mathcal{T}$  indexed by  $I$  with the tower indexed by the ordinal  $\text{otp}(I)$ .

**Definition 5.5.** For  $\mathcal{T} = \langle M_i : i \in I \rangle \frown \langle a_i : i \in I^- \rangle$  and  $I_0 \subseteq I$ , we let  $\mathcal{T} \upharpoonright I_0$  be the sequences  $\langle M_i : i \in I_0 \rangle \frown \langle a_i : i \in I_0 \rangle$ .

**Remark 5.6.** If  $\mathcal{T}$  is a tower indexed by  $I$  and  $I_0 \subseteq I$ , then  $\mathcal{T} \upharpoonright I_0$  is a tower indexed by  $I_0$ .

The reason for indexing towers by a well-ordering instead of just an ordinal is that we will allow towers to be grown by inserting elements not only at the end, but also in the middle. What it means to “grow” a tower is given by the following definition:

**Definition 5.7** (Orderings on towers). For  $\mathcal{T}^\ell = \langle M_i^\ell : i \in I^\ell \rangle \frown \langle a_i^\ell : i \in (I^\ell)^- \rangle$ ,  $\ell = 1, 2$ , two towers, we write  $\mathcal{T}^1 \triangleleft \mathcal{T}^2$  if:

- (1)  $I^1 \subseteq I^2$ .
- (2)  $M_i^2$  is limit over  $M_i^1$  for all  $i \in I^1$ .
- (3)  $a_i^1 = a_i^2$  for all  $i \in (I^1)^-$ .
- (4)  $\text{tp}(a_i^1/M_i^2; M_{i+1}^2)$  does not fork over  $M_i^1$  for all  $i \in (I^1)^-$ .

We write  $\mathcal{T}^1 \leq \mathcal{T}^2$  if  $\mathcal{T}^1 = \mathcal{T}^2$  or  $\mathcal{T}^1 \triangleleft \mathcal{T}^2$ .

**Remark 5.8.**  $\leq$  is a partial order on the class of all towers. Moreover, if  $\mathcal{T}^1$  and  $\mathcal{T}^2$  have index sets  $I^1 \subseteq I^2$  respectively, then  $\mathcal{T}^1 \leq \mathcal{T}^2$  if and only if  $\mathcal{T}^1 \leq \mathcal{T}^2 \upharpoonright I^1$ .

**Definition 5.9.** Let  $\delta < \lambda^+$  be a limit ordinal and let  $\langle \mathcal{T}^j : j < \delta \rangle$  be a  $\leq$ -increasing chain of towers. Assume that  $\mathcal{T}^j = \langle M_i^j : i \in I^j \rangle \frown \langle a_i^j : i \in (I^j)^- \rangle$ . We define  $\mathcal{T}^\delta := \langle M_i^\delta : i \in I^\delta \rangle \frown \langle a_i^\delta : i \in I^\delta \rangle$  as follows:

- (1)  $I^\delta = \bigcup_{j < \delta} I^j$ .
- (2)  $a_i^\delta = a_i^j$  for some (any)  $j < \delta$  such that  $i \in I^j$ .
- (3)  $M_i^\delta = \bigcup_{j < \delta} M_i^j$ .

We write  $\bigcup_{j < \delta} \mathcal{T}^j$  for  $\mathcal{T}^\delta$ .

**Remark 5.10.** If  $\langle \mathcal{T}^j : j < \delta \rangle$  is a  $\leq$ -increasing chain of towers, where  $\mathcal{T}^j$  is indexed by  $I^j$ , and  $\bigcup_{j < \delta} I^j$  is a well-ordering, then  $\bigcup_{j < \delta} \mathcal{T}^j$  is a tower and  $\mathcal{T}^k \leq \bigcup_{j < \delta} \mathcal{T}^j$  for every  $k < \delta$ .

**Definition 5.11.** If a  $\leq$ -increasing chain of towers  $\langle \mathcal{T}^j : j < \gamma \rangle$  is such that for every limit  $j < \gamma$ ,  $\mathcal{T}^j = \bigcup_{k < j} \mathcal{T}^k$ , we call the chain *continuous*.

The following is an interesting property of towers. It says that any extension must be “as disjoint as possible”.

**Definition 5.12** ([SV99, 3.1.11(1)]). A tower  $\mathcal{T} = \langle M_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  is called *reduced* if whenever  $\mathcal{T}' = \langle M'_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  is such that  $\mathcal{T} \leq \mathcal{T}'$ , we have that  $M'_i \cap M_j = M_i$  for any  $i \leq j < \alpha$ .

**Remark 5.13.** If  $\langle \mathcal{T}^j : j < \delta \rangle$  is a chain of towers and  $\mathcal{T}^j$  is reduced for all  $j < \delta$ , then  $\bigcup_{j < \delta} \mathcal{T}^j$  is reduced (provided that its index is a well-ordering).

The following related definition appears (stated differently) in (for example) [JS13, 3.3.2]:

**Definition 5.14.** We say a triple  $(a, M, N)$  is *reduced* if  $(M, N) \frown (a)$  is a reduced tower.

To check that a tower is reduced, it is enough to check its restrictions of length two:

**Lemma 5.15.** Let  $\mathcal{T} = \langle M_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be a continuous tower. If  $(a_i, M_i, M_{i+1})$  is a reduced triple for any  $i + 1 < \alpha$ , then  $\mathcal{T}$  is reduced.

*Proof.* Let  $\mathcal{T}' = \langle M'_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be such that  $\mathcal{T} \trianglelefteq \mathcal{T}'$ . Let  $i \leq j < \alpha$ . We have to see that  $M'_i \cap M_j = M_i$ . We proceed by induction on  $j$ . If  $j = i$ , this is obvious. If  $j$  is limit, this is immediate from the induction hypothesis and (since  $\mathcal{T}$  is assumed to be continuous)  $M_j = \bigcup_{k < j} M_k$ . Assume now that  $j = k + 1$ . By assumption,  $(a_k, M_k, M_{k+1})$  is a reduced triple. Thus  $M'_i \cap M_j \subseteq M'_k \cap M_j = M_k$ , and so  $M'_i \cap M_j = M'_i \cap M_k$  which by the induction hypothesis is just  $M_i$ , as desired.  $\square$

Reduced towers exist: any tower has a reduced extension.

**Fact 5.16** (Density of reduced towers). For any tower  $\mathcal{T}$  of length  $\alpha$ , there exists a reduced tower  $\mathcal{T}'$  of length  $\alpha$  such that  $\mathcal{T} \trianglelefteq \mathcal{T}'$ .

*Proof.* As in (for example) [GVV16, 5.5].  $\square$

So far, we haven't shown that towers have any nontrivial  $\trianglelefteq$ -extensions. In fact, nonforking amalgamation gives a crucially stronger statement.

**Fact 5.17** (Existence of extensions of towers). Let  $\mathcal{T} = \langle M_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be a tower.

- (1) There exists a limit tower  $\mathcal{T}'$  of length  $\alpha$  such that  $\mathcal{T} \triangleleft \mathcal{T}'$ .
- (2) Assume in addition that  $\mathcal{T}$  is limit and continuous. If  $q \in \mathbf{S}(M_0)$ . Then there exists a limit tower  $\mathcal{T}' = \langle M'_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  and  $b \in M'_0$  such that:
  - (a)  $\mathcal{T} \triangleleft \mathcal{T}'$ .
  - (b)  $\mathbf{tp}(b/M_0; M'_0) = q$ .
  - (c)  $\mathbf{tp}(b/\bigcup_{i < \alpha} M_i; \bigcup_{i < \alpha} M'_i)$  does not fork over  $M_0$ .

*Proof.* As in (for example) [She09a, II.4.9].  $\square$

The following technical consequence will be used in the proof of the next theorem.

**Lemma 5.18.** Let  $\delta < \lambda^+$  be a limit ordinal. Let  $\mathcal{T} = \langle M_i : i \leq \delta \rangle \frown \langle a_i : i < \delta \rangle$  be a tower such that  $\mathcal{T} \restriction \delta$  is limit and continuous. Let  $b \in M_\delta$ . If  $\mathbf{tp}(b/\bigcup_{i < \delta} M_i; M_\delta)$  does not fork over  $M_0$ , then there exists a tower  $\mathcal{T}' = \langle M'_i : i \leq \delta \rangle \frown \langle a_i : i < \delta \rangle$  such that  $\mathcal{T} \triangleleft \mathcal{T}'$  and  $b \in M'_0$ .

*Proof.* Write  $M_\delta^0 := \bigcup_{i < \delta} M_i$ . Let  $q := \mathbf{tp}(b/M_0; M_\delta)$ . By Fact 5.17 applied to the tower  $\mathcal{T} \restriction \delta$ , there exists  $\mathcal{T}^* = \langle M_i^* : i < \delta \rangle \frown \langle a_i : i < \delta \rangle$  and  $b^* \in M_0^*$  such that  $\mathcal{T} \restriction \delta \triangleleft \mathcal{T}^*$ ,  $\mathbf{tp}(b^*/M_0; M_0^*) = q$ , and  $\mathbf{tp}(b^*/M_\delta^0; M_\delta^*)$  does not fork over  $M_0$  (we have

set  $M_\delta^* := \bigcup_{i < \delta} M_i^*$ ). Note that we have used that  $\delta$  is a limit ordinal to make sure that all the  $a_i$ 's are still in  $\mathcal{T} \upharpoonright \delta$ . By uniqueness,  $\mathbf{tp}(b^*/M_\delta^0; M_\delta^*) = \mathbf{tp}(b/M_\delta^0; M_\delta)$ . Pick  $M'_\delta$  limit over  $M_\delta$  and  $f : M_\delta^* \xrightarrow{M_\delta^0} M'_\delta$  such that  $f(b^*) = b$ . Let  $M'_i := f[M_i^*]$  for  $i < \delta$ .  $\square$

We obtain the following powerful tool to build continuous towers. This was first proven by VanDieren (using a slightly different notion of tower) [Van16a]. We give a simplification of VanDieren's original proof here.

**Fact 5.19.** Any reduced tower is continuous.

*Proof.* Suppose not. Let  $\alpha$  be the least length of a reduced non-continuous tower. Then it is easy to see that  $\alpha = \delta + 1$ , where  $\delta$  is a limit ordinal. Let  $\mathcal{T} = \langle M_i : i \leq \delta \rangle \frown \langle a_i : i < \delta \rangle$  be such a reduced non-continuous tower. Thus  $\bigcup_{i < \delta} M_i \neq M_\delta$ . Pick  $b \in M_\delta \setminus \bigcup_{i < \delta} M_i$ .

Claim: There is no  $k < \delta$  and no tower  $\mathcal{T}' = \langle M'_i : i \in [k, \delta] \rangle \frown \langle a_i : i \in [k, \delta] \rangle$  such that  $\mathcal{T} \upharpoonright [k, \delta] \triangleleft \mathcal{T}'$  and  $b \in \bigcup_{i \in [k, \delta]} M'_i$ .

Proof of Claim: Suppose  $\mathcal{T}'$  is such a tower and fix  $i < \delta$  such that  $b \in M'_i$ . Then  $b \in M'_i \cap M_\delta$  but  $b \notin M_i$ , so  $M'_i \cap M_\delta \neq M_i$ . Moreover, one can extend  $\mathcal{T}'$  to a tower  $\mathcal{T}''$  of length  $\delta + 1$  so that  $\mathcal{T}'' \upharpoonright [k, \delta] = \mathcal{T}'$  and  $\mathcal{T} \trianglelefteq \mathcal{T}''$  (use universality of  $M'_k$  over  $M_k$ ). This implies that  $\mathcal{T}$  is not reduced, contradiction.  $\uparrow_{\text{Claim}}$

We aim to build a tower as in the claim to get a contradiction. Build  $\langle \mathcal{T}^j : j \leq \delta \rangle$  an  $\triangleleft$ -increasing continuous chain of reduced towers such that  $\mathcal{T}^0 = \mathcal{T}$ . Write  $\mathcal{T}^j = \langle M_i^j : i \leq \delta \rangle \frown \langle a_i : i < \delta \rangle$ . Now consider the diagonal tower  $\mathcal{T}^* := \langle M_i^i : i \leq \delta \rangle \frown \langle a_i : i < \delta \rangle$ . It is easy to check that  $\mathcal{T}^*$  is indeed a tower, and further it is limit. Since  $\delta$  was minimal,  $\mathcal{T}^j \upharpoonright \delta$  is continuous for all  $j \leq \delta$ , and hence  $\mathcal{T}^* \upharpoonright \delta$  is also continuous. Further, it is easy to check that  $\mathcal{T}^* \upharpoonright \delta$  is limit. By local character, there exists  $i < \delta$  such that  $\mathbf{tp}(b/\bigcup_{k < \delta} M_k^k; M_\delta^\delta)$  does not fork over  $M_i^i$ . Let  $\mathcal{T}^{**} := \langle M_i^i : i \in [k+1, \delta] \rangle \frown \langle a_i : i \in [k+1, \delta] \rangle$ . By Lemma 5.18, where  $\mathcal{T}$  there stands for  $\mathcal{T}^{**}$  here, there exists a tower  $\mathcal{T}'$  such that  $\mathcal{T}^{**} \triangleleft \mathcal{T}'$  and  $\mathcal{T}' \upharpoonright \delta$  contains  $b$ . Since  $\mathcal{T}'$  also extends  $\mathcal{T} \upharpoonright [k, \delta]$ , this contradicts the claim.  $\square$

We now want to give conditions under which a tower  $\langle M_i : i < \alpha \rangle \frown \langle a_i : i+1 < \alpha \rangle$  is such that  $\bigcup_{i < \alpha} M_i$  is limit over  $M_0$ . Of course, being a limit tower suffices but it is not clear whether this property is closed under unions. Instead, we will rely on the following weakening (a variation appears in [GVV16, 4.3]):

**Definition 5.20.** A tower  $\mathcal{T} = \langle M_i : i \in I \rangle \frown \langle a_i : i \in I^- \rangle$  is  $I_0$ -full if:

- (1)  $I_0 \subseteq I$ .
- (2) For any  $i \in I_0^-$  and any  $p \in \mathbf{S}^{\text{na}}(M_i)$ , there exists  $k \in [i, i + I_0 \cdot 1)_I$  such that  $\mathbf{tp}(a_k/M_k; M_{k+1})$  is the nonforking extension of  $p$ .

We have the following monotonicity properties:

**Remark 5.21.** Let  $I_0 \subseteq I_1 \subseteq I_2$  be well-orderings and let  $\mathcal{T}$  be a tower indexed by  $I_2$ .

- (1) If  $\mathcal{T} \upharpoonright I_1$  is  $I_0$ -full, then  $\mathcal{T}$  is  $I_0$ -full.

(2) If  $\mathcal{T}$  is  $I_1$ -full, then  $\mathcal{T}$  is  $I_0$ -full.

Intuitively, full towers are those for which the  $a_i$ 's realize all the types many times. To see that full towers generate limit models, we will use:

**Fact 5.22** ([She09a, II.1.16(4)]). If  $\langle M_i : i \leq \lambda \rangle$  is increasing continuous in  $\mathbf{K}_\lambda$  such that  $M_{i+1}$  realizes all types over  $M_i$  for every  $i < \lambda$ , then  $M_\lambda$  is universal over  $M_0$ .

**Lemma 5.23.** If  $\mathcal{T} = \langle M_i : i \in I \rangle \frown \langle a_i : i + 1 < \delta \rangle$  is an  $I_0$ -full tower and  $(1 + \text{otp}(I_0)) \cdot \lambda = \text{otp}(I_0)$ , then  $\bigcup_{i \in I_0} M_i$  is  $(\lambda, \text{cf}(I_0))$ -limit over  $M_0$ .

*Proof.* Fix  $i \in I_0$ . By assumption,  $M_{i+I_0+1}$  realizes all types over  $M_i$ . Thus by Fact 5.22  $M_{i+I_0+\lambda}$  is universal over  $M_i$ . Since  $i$  was arbitrary, we obtain that  $M_0 \frown \langle M_i : i \in I_0 \rangle$  is the desired witness that  $\bigcup_{i \in I_0} M_i$  is  $(\lambda, \text{cf}(I_0))$ -limit over  $M_0$ .  $\square$

Full towers are also preserved by unions:

**Lemma 5.24.** Let  $\delta < \lambda^+$  be a limit ordinal. Let  $\langle \mathcal{T}^j : j < \delta \rangle$  be an increasing chain of towers. Assume that  $\mathcal{T}^j$  is indexed by  $I^j$ , and  $I^\delta := \bigcup_{j < \delta} I^j$  is a well-ordering. Suppose that  $I_0 \subseteq I^0$  is such that  $\mathcal{T}^j$  is  $I_0$ -full for each  $j < \delta$ . Then  $\bigcup_{j < \delta} \mathcal{T}^j$  is  $I_0$ -full.

*Proof.* Let  $\mathcal{T}^\delta := \bigcup_{j < \delta} \mathcal{T}^j$ . For  $j \leq \delta$ , write  $\mathcal{T}^j = \langle M_i^j : i \in I^j \rangle \frown \langle a_i : i \in (I^j)^- \rangle$ . Let  $i \in I_0^-$  and let  $p \in \mathbf{S}^{\text{na}}(M_i^\delta)$ . Pick  $j < \delta$  such that  $p$  does not fork over  $M_i^j$ . Since  $\mathcal{T}^j$  is  $I_0$ -full, there exists  $k \in [i, i +_{I_0} 1)_{I^j}$  such that  $\text{tp}(a_k/M_k^j)$  is the nonforking extension of  $p \upharpoonright M_i^j$ . Clearly,  $k \in [i, i +_{I_0} 1)_{I^\delta}$ . Moreover, by definition of extension of tower,  $\text{tp}(a_k/M_k^\delta; M_{k+1}^\delta)$  does not fork over  $M_k^j$ . By transitivity of forking and uniqueness,  $\text{tp}(a_k/M_k^\delta; M_{k+1}^\delta)$  is the nonforking extension of  $p \upharpoonright M_i^j$ , hence of  $p$ .  $\square$

**Definition 5.25.** For  $I$  and  $J$  linear orders, let  $I \times J$  be the usual cartesian product, ordered lexicographically:  $(i_1, j_1) < (i_2, j_2)$  if and only if either  $i_1 < i_2$ , or  $i_1 = i_2$  and  $j_1 < j_2$ .

We now prove two construction lemmas about full towers. The idea is also described on p. 373 of [GVV16].

**Lemma 5.26.** Let  $I$  be a well-ordering and let  $\alpha < \gamma$  be ordinals in  $[1, \lambda^+)$  with  $\gamma \cdot \lambda = \gamma$ . Let  $\mathcal{T}$  be a limit tower indexed by  $I \times \alpha$ . Then there is a limit tower  $\mathcal{T}'$  indexed by  $I \times \gamma$  such that  $\mathcal{T}'$  is  $I$ -full and  $\mathcal{T}' \upharpoonright (I \times \alpha) = \mathcal{T}$ .

*Proof.* Straightforward: realize all the relevant types.  $\square$

**Lemma 5.27.** Let  $I$  be a well-ordering and let  $\alpha \in [1, \lambda^+)$  be an ordinal. Let  $\mathcal{T}$  be a tower indexed by  $I \times \alpha$ . Then there is an ordinal  $\beta \in [\alpha, \lambda^+)$  and a tower  $\mathcal{T}'$  indexed by  $I \times \beta$  such that  $\mathcal{T} \triangleleft \mathcal{T}'$ ,  $\mathcal{T}'$  is reduced, and  $\mathcal{T}'$  is  $I$ -full.

*Proof.* We build a  $\leq$ -increasing continuous chain of towers  $\langle \mathcal{T}^i : i \leq \omega \rangle$  and an increasing continuous chain of ordinals  $\langle \alpha_i : i \leq \omega \rangle$  such that:

(1)  $\mathcal{T}^i$  is indexed by  $I \times \alpha_i$ .



- (2)  $\mathcal{T} \triangleleft \mathcal{T}^0$ .
- (3) For  $i < \omega$  nonzero and even,  $\mathcal{T}^i$  is reduced.
- (4) For  $i < \omega$  odd,  $\mathcal{T}^i$  is  $I$ -full.

This is enough:  $\mathcal{T}^\omega$  is reduced by Remark 5.13 and  $I$ -full by Lemma 5.24. Thus one can set  $\mathcal{T}' := \mathcal{T}^\omega$ ,  $\beta := \alpha^\omega$ .

This is possible: take  $\alpha_0 := \alpha$  and any  $\mathcal{T}^0$  such that  $\mathcal{T} \triangleleft \mathcal{T}^0$  (exists by Fact 5.17). Now let  $i > 0$  and assume that  $\mathcal{T}^{i-1}$  and  $\alpha_{i-1}$  are given. If  $i$  is even, take  $\alpha_i := \alpha_{i-1}$  and let  $\mathcal{T}^i$  be a reduced tower  $\leq$ -extending  $\mathcal{T}^{i-1}$  which is indexed by  $I \times \alpha_i$  (exists by Fact 5.16). If  $i$  is odd, use Lemma 5.26.  $\square$

In passing, we can now give a proof of Fact 2.7:

*Proof of Fact 2.7.* It is enough to show that for any  $M \in \mathbf{K}_\lambda$  and any limit ordinals  $\delta_1, \delta_2 < \lambda^+$  there exists  $N$  which is both  $(\lambda, \delta_1)$ -limit and  $(\lambda, \delta_2)$ -limit over  $M$ . Extending  $M$  if necessary, we can assume without loss of generality that  $M$  is a limit model. We build a  $\triangleleft$ -increasing continuous chain of towers  $\langle \mathcal{T}^i : i \leq \delta_2 \rangle$  such that:

- (1) The first model in  $\mathcal{T}^0$  is  $M$ .
- (2) For each  $i \leq \delta_2$ ,  $\mathcal{T}^i$  is indexed by  $(\delta_1 + 1) \times \alpha_i$ , for some nonzero ordinal  $\alpha_i$ .
- (3) For each nonzero  $i \leq \delta_2$ ,  $\mathcal{T}^i$  is  $(\delta_1 + 1)$ -full and reduced.

This is possible: use Lemma 5.27 at successor steps, recalling that being full and reduced is preserved at limits (Remark 5.13 and Lemma 5.24).

This is enough: write  $\mathcal{T}^i = \langle M_j^i : j \in (\delta_1 + 1) \times \alpha_i \rangle$ . By Remark 5.21,  $\mathcal{T}^{\delta_2}$  is  $\delta_1$ -full. By Fact 5.19,  $\mathcal{T}^{\delta_2}$  is continuous. Thus by Lemma 5.23,  $M_{\delta_1}^{\delta_2}$  is  $(\lambda, \delta_1)$ -limit over  $M_0^{\delta_2}$ , hence over  $M$ . Moreover by definition of  $\triangleleft$ ,  $M_{\delta_1}^{\delta_2}$  is also  $(\lambda, \delta_1)$ -limit over  $M_{\delta_1}^0$ , hence over  $M$ , as desired.  $\square$

We are now ready to prove that any tower is extended by a continuous limit tower.

**Lemma 5.28.** For any tower  $\mathcal{T}$  there exists a continuous limit tower  $\mathcal{T}'$  such that  $\mathcal{T} \trianglelefteq \mathcal{T}'$ .

*Proof.* By Fact 5.17, we can assume without loss of generality (taking an extension of  $\mathcal{T}$  if necessary) that  $\mathcal{T}$  is a limit tower. Say  $\mathcal{T}$  is indexed by  $I$ . Let  $\gamma < \lambda^+$  be nonzero such that  $\gamma \cdot \lambda = \gamma$ . Let  $\mathcal{T}^*$  be any limit tower indexed by  $I \times \gamma$  such that  $\mathcal{T}^* \upharpoonright I = \mathcal{T}$  (it is easy to see that such towers exist). By Lemma 5.27 (where  $I, \mathcal{T}$  there stands for  $I \times \gamma, \mathcal{T}^*$  here), there is a reduced and  $(I \times \gamma)$ -full tower  $\mathcal{T}^{**}$  indexed by  $I' := I \times \gamma \times \gamma'$  (for some  $\gamma'$ ) such that  $\mathcal{T}^* \leq \mathcal{T}^{**}$ . By Fact 5.19,  $\mathcal{T}^{**}$  is continuous. Let  $\mathcal{T}' := \mathcal{T}^{**} \upharpoonright I$ . Clearly,  $\mathcal{T}'$  is also continuous. To see that  $\mathcal{T}'$  is limit, let  $i \in I^-$ . Observe that  $\mathcal{T}^{**} \upharpoonright [i, i +_I 1]_{I'}$  is  $(\{i\} \times \gamma)$ -full, hence by Lemma 5.23,  $M_{i+1}$  is limit over  $M_i$ , and so  $\mathcal{T}'$  is indeed a limit tower.  $\square$

We can now prove a strengthening of Fact 5.17, where the starting tower is no longer assumed to be continuous and limit.

**Lemma 5.29.** Let  $\mathcal{T} = \langle M_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be a tower and let  $N_0$  be limit over  $M_0$ . Let  $b \in N_0$ . There exists a limit continuous tower  $\mathcal{T}' = \langle M'_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  and an isomorphism  $f : N_0 \cong_{M_0} M'_0$  such that  $\mathcal{T} \leq \mathcal{T}'$  and  $\text{tp}(f(b)/\bigcup_{i < \alpha} M_i; \bigcup_{i < \alpha} M'_i)$  does not fork over  $M_0$ .

*Proof.* First, let  $\mathcal{T}^1 = \langle M_i^1 : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be a limit continuous tower extending  $\mathcal{T}$ , as given by Lemma 5.28. Find  $N'_0$  limit over  $M_0^1$  and  $f_0 : N_0 \cong_{M_0} N'_0$  such that  $q := \text{tp}(f_0(b)/M_0^1; N'_0)$  does not fork over  $M_0$ . By Fact 5.17(2), there exists a limit continuous tower  $\mathcal{T}^2 = \langle M_i^2 : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  and a  $b' \in M_0^2$  such that  $\mathcal{T}^1 \triangleleft \mathcal{T}^2$ ,  $b'$  realizes  $q$ , and  $\text{tp}(b'/\bigcup_{i < \alpha} M_i^1; \bigcup_{i < \alpha} M_i^2)$  does not fork over  $M_0^1$ . Since  $b'$  realizes  $q$ , transitivity and monotonicity imply that  $\text{tp}(b'/\bigcup_{i < \alpha} M_i; \bigcup_{i < \alpha} M_i^2)$  does not fork over  $M_0$ . Since  $M_0^2$  is limit over  $M_0^1$ , there exists  $g : N'_0 \cong_{M_0^1} M_0^2$  such that  $g(f_0(b)) = b'$ . Let  $\mathcal{T}' := \mathcal{T}^2$ ,  $f := gf_0$ .  $\square$

We have arrived to one of the key results of this section. Roughly, it says that any two towers can be amalgamated into a continuous rectangle of models. The statement is similar to [JS13, 3.1.10]

**Lemma 5.30** (Amalgamation of towers). Let  $\mathcal{T} = \langle N_j : j < \beta \rangle \frown \langle b_j : j + 1 < \beta \rangle$  and  $\mathcal{T}' = \langle M_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be towers. Assume that  $N_0 = M_0$ . Then there exists  $\langle f_i : i < \alpha \rangle$  and  $\langle M_{i,j} : i < \alpha, j < \beta \rangle$  such that:

- (1) For all  $i < \alpha$  and  $j < \beta$ ,  $M_{i,j}$  is a limit model.
- (2) For all  $j < \beta$ ,  $M_{0,j} = M_j$ .
- (3) For each  $j < \beta$ ,  $\langle M_{i,j} : i < \alpha \rangle$  is increasing continuous.
- (4) For each  $i < \alpha$ ,  $\langle M_{i,j} : j < \beta \rangle$  is increasing continuous.
- (5)  $\langle f_i : i < \alpha \rangle$  is increasing.
- (6) For all  $i < \alpha$ ,  $f_i : M_i \cong_{M_{0,0}} M_{i,0}$ .
- (7) For all  $i + 1 < \alpha$  and  $j < \beta$ ,  $\text{tp}(f_{i+1}(a_i)/M_{i,j}; M_{i+1,j})$  does not fork over  $M_{i,0}$ .
- (8) For all  $j + 1 < \beta$  and  $i < \alpha$ ,  $\text{tp}(b_j/M_{i,j}; M_{i,j+1})$  does not fork over  $M_{0,j}$ .
- (9) If  $\mathcal{T}$  is reduced, then for any  $i < \alpha$  and  $j < \beta$ ,  $M_{0,j} \cap M_{i,0} = M_{0,0}$ .

*Proof.* First, we claim that we can assume without loss of generality that  $\mathcal{T}'$  is limit and continuous. Indeed, suppose that we could prove the lemma for limit and continuous  $\mathcal{T}'$ . By Lemma 5.28 there exists  $\mathcal{T}^* = \langle M_i^* : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle \leq$ -extending  $\mathcal{T}'$  which is limit and continuous. Let  $\mathcal{T}'' = \langle M'_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  be defined by  $M'_0 = M_0$ ,  $M'_i := M_i^*$  for  $i > 0$ . Then  $\mathcal{T}''$  is also a limit continuous tower. By what we are assuming, there exists  $\langle M'_{i,j} : i < \alpha, j < \beta \rangle$ ,  $\langle g_i : i < \alpha \rangle$  satisfying the conclusion of the lemma, where  $\mathcal{T}'$ ,  $\langle M_{i,j} : i < \alpha, j < \beta \rangle$ ,  $\langle f_i : i < \alpha \rangle$  there is  $\mathcal{T}''$ ,  $\langle M'_{i,j} : i < \alpha, j < \beta \rangle$ ,  $\langle g_i : i < \alpha \rangle$  here. Now define  $f_i := g_i \upharpoonright M_i$  and let  $M_{i,0} := f_i[M_i]$ ,  $M_{i,j} := M'_{i,j}$  for  $j > 0$ . This works. The main point is that, for  $i + 1 < \alpha$ ,  $j \in (0, \beta)$ ,  $\text{tp}(f_{i+1}(a_i)/M_{i,j}; M_{i+1,j})$  does not fork over  $M'_{i,0}$  (by construction), and since  $\mathcal{T}' \leq \mathcal{T}^*$ , using invariance and the definition of  $\leq$ ,  $\text{tp}(f_{i+1}(a_i)/M'_{i,0}; M'_{i+1,0})$  does not fork over  $M_{i,0}$ . Thus by transitivity  $\text{tp}(f_{i+1}(a_i)/M_{i,j}; M_{i+1,j})$  does not fork over  $M_{i,0}$ .

We have shown that we can assume without loss of generality that  $\mathcal{T}'$  is limit and continuous. So assume that it is. We build a  $\leq$ -increasing continuous chain of

towers  $\langle \mathcal{T}^i : i < \alpha \rangle$  and an increasing continuous chain  $\langle f_i : i < \alpha \rangle$  such that, writing  $\mathcal{T}^j = \langle M_{i,j} : j < \beta \rangle \frown \langle b_j : j + 1 < \beta \rangle$ , we have:

- (1)  $\mathcal{T}^0 = \mathcal{T}$ .
- (2) For all  $i < \alpha$ ,  $f_i : N_i \cong_{N_0} M_{i,0}$ .
- (3) For all  $i + 1 < \alpha$ ,  $\mathbf{tp}(f_{i+1}(a_i) / \bigcup_{j < \beta} M_{i,j}; M_{i+1,j})$  does not fork over  $M_{i,0}$ .

This is possible by Lemma 5.29 and some renaming. This is enough: check the requirements and the definition of  $\trianglelefteq$ .  $\square$

We end this section by deducing some results around disjoint amalgamation. The following appears in [JS13, 3.2.3(3)] and the proof carries through to our setup.

**Fact 5.31.** Assume that  $\mathbf{K}$  is categorical in  $\lambda$ . Let  $\mathcal{R}$  be a class of triples  $(a, M, N)$  with  $M \leq_{\mathbf{K}} N$  limit and  $a \in N \setminus M$  such that:

- (1)  $\mathcal{R}$  is closed under isomorphisms:  $(a, M, N) \in \mathcal{R}$  and  $f : N \cong N'$  implies  $(f(a), f[M], N') \in \mathcal{R}$ .
- (2)  $\mathcal{R}$  has the existence property: for any  $M$  and any  $p \in \mathbf{S}^{\text{na}}(M)$ , there exists  $(a, M, N) \in \mathcal{R}$  such that  $p = \mathbf{tp}(a/M; N)$ .

Let  $(a, M, N) \in \mathcal{R}$  and let  $N'$  be limit such that  $N \leq_{\mathbf{K}} N'$ . Then there exists a limit ordinal  $\delta < \lambda^+$  and a continuous tower  $\mathcal{T} = \langle M_i : i \leq \delta \rangle \frown \langle a_i : i < \delta \rangle$  that  $a_0 = a$ ,  $M_0 = M$ ,  $N' \leq_{\mathbf{K}} M_\delta$ , and  $(a_i, M_i, M_{i+1}) \in \mathcal{R}$  for all  $i < \delta$ .

Reduced triples have the existence property in the sense just given:

**Fact 5.32** (Existence property for reduced triples). Assume that  $\mathbf{K}$  is categorical in  $\lambda$ . For any  $M \in \mathbf{K}_\lambda$  and any  $p \in \mathbf{S}^{\text{na}}(M)$ , there exists a reduced triple  $(a, M, N)$  such that  $p = \mathbf{tp}(a/M; N)$ .

*Proof.* Let  $p \in \mathbf{S}(M)$  and write  $p = \mathbf{tp}(b/M; N_0)$ . By Fact 5.16, there exists limit  $M', N'$  such that  $M \leq_{\mathbf{K}} M'$ ,  $N_0 \leq_{\mathbf{K}} N'$ ,  $q := \mathbf{tp}(b/M'; N')$  does not fork over  $M$ , and  $(b, M', N')$  is reduced. By the conjugation property, there exists  $f : M' \cong M$  such that  $f(q) = p$ . Extend  $f$  to some  $g : N' \cong N$  and let  $a := g(b)$ . Then  $(a, M, N)$  is reduced and  $\mathbf{tp}(a/M; N) = f(q) = p$ , as desired.  $\square$

We can now improve Fact 5.2 to make also the amalgams of  $M_1$  and  $M_2$  disjoint over  $M_0$ . Note that this gives in particular disjoint amalgamation.

**Lemma 5.33** (Disjoint nonforking amalgamation). Assume that  $\mathbf{K}$  is categorical in  $\lambda$ . Let  $M_0 \leq_{\mathbf{K}} M_\ell$ ,  $\ell = 1, 2$ , be in  $\mathbf{K}_\lambda$ . Let  $a_\ell \in M_\ell$ . There exists  $f_1, f_2, M_3$  such that  $M_3 \in \mathbf{K}_\lambda$ , and  $f_\ell : M_\ell \xrightarrow{M_0} M_3$  is such that  $\mathbf{tp}(f_\ell(a_\ell) / f_{3-\ell}(M_{3-\ell}); M_3)$  does not fork over  $M_0$  for  $\ell = 1, 2$ . Moreover,  $f_1[M_1] \cap f_2[M_2] = M_0$ .

*Proof.* Extending  $M_1$  if necessary, we may assume without loss of generality that there exists  $M'_1 \leq_{\mathbf{K}} M_1$  containing  $a_1$  such that  $(a_1, M_0, M'_1)$  is reduced. By Facts 5.31 and 5.32, there exists a continuous tower  $\mathcal{T} = \langle N_j : j \leq \delta \rangle \frown \langle b_j : j < \delta \rangle$  such that  $b_0 = a_1$ ,  $M_1 \leq_{\mathbf{K}} N_\delta$ , and  $(b_j, N_j, N_{j+1})$  is reduced for all  $j < \delta$ . This implies in particular that  $\mathcal{T}$  is reduced (Lemma 5.15). Similarly (and after perhaps growing  $M_2$ ), there exists a continuous tower  $\mathcal{T}' = \langle M_i^* : i \leq \delta^* \rangle \frown \langle a_i^* : i < \delta^* \rangle$  such that  $a_2 = a_0^*$ ,  $M_2 \leq_{\mathbf{K}} M_{\delta^*}^*$ , and  $(a_i^*, M_i^*, M_{i+1}^*)$  is reduced for all  $i < \delta^*$ . Let  $\alpha := \delta^* + 1$ ,

$\beta := \delta + 1$ . Let  $\langle M_{i,j} : i < \alpha, j < \beta \rangle$ ,  $\langle g_i : i < \alpha \rangle$  be as given by Lemma 5.30, where  $\langle M_i : i < \alpha \rangle \smallfrown \langle a_i : i + 1 < \alpha \rangle$  there stand for  $\langle M_i^* : i < \alpha \rangle \smallfrown \langle a_i^* : i < \alpha \rangle$  here. Then  $M_3 := M_{\delta^*, \delta}$ ,  $f_1 := \text{id}_{M_1}$ , and  $f_2 := g_{\delta^*} \upharpoonright M_2$  are as desired.  $\square$

## 6. BUILDING WEAKLY SUCCESSFUL GOOD FRAMES

The goal of this section is to build a good  $\lambda$ -frame from  $\lambda$ -superstability,  $\lambda$ -symmetry, and categoricity in  $\lambda$ . We will use uniqueness triples, a key object in (for example) [She09a, §II.6]. We quote the definition from [JS13, §4].

**Definition 6.1.** Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC.

- (1) For  $M_0 \leq_{\mathbf{K}} M_\ell$  all in  $\mathbf{K}_\lambda$ ,  $\ell = 1, 2$ , an *amalgam of  $M_1$  and  $M_2$  over  $M_0$*  is a triple  $(f_1, f_2, N)$  such that  $N \in \mathbf{K}_\lambda$  and  $f_\ell : M_\ell \xrightarrow{M_0} N$ .
- (2) Let  $(f_1^x, f_2^x, N^x)$ ,  $x = a, b$  be amalgams of  $M_1$  and  $M_2$  over  $M_0$ . We say  $(f_1^a, f_2^a, N^a)$  and  $(f_1^b, f_2^b, N^b)$  are *equivalent over  $M_0$*  if there exists  $N_* \in \mathbf{K}_\lambda$  and  $f^x : N^x \rightarrow N_*$  such that  $f^b f_1^b = f^a f_1^a$  and  $f^b f_2^b = f^a f_2^a$ , namely, the following commutes:

$$\begin{array}{ccccc}
 & & N^a & \xrightarrow{f^a} & N_* \\
 & \nearrow f_1^a & \uparrow f_2^a & & \uparrow f^b \\
 M_1 & \xrightarrow{\quad} & & \xrightarrow{f_1^b} & N^b \\
 \uparrow & & \uparrow & \nearrow f_2^b & \\
 M_0 & \longrightarrow & M_2 & & 
 \end{array}$$

Note that being “equivalent over  $M_0$ ” is an equivalence relation ([JS13, 4.3]).

- (3) A triple  $(a, M, N)$  is a *uniqueness triple* if  $M \leq_{\mathbf{K}} N$  are both in  $\mathbf{K}_\lambda$ ,  $a \in |N| \setminus |M|$ , and for any  $M_1 \geq_{\mathbf{K}} M$  in  $\mathbf{K}_\lambda$ , there exists a *unique* (up to equivalence over  $M$ ) amalgam  $(f_1, f_2, N_1)$  of  $N$  and  $M_1$  over  $M$  such that  $\text{tp}(f_1(a)/f_2[M_1]; N_1)$  does not fork over  $M$ .
- (4)  $\mathbf{K}$  has the  *$\lambda$ -existence property for uniqueness triples* if for any  $M \in \mathbf{K}_\lambda$  and any  $p \in \mathbf{S}^{\text{na}}(M)$ , one can write  $p = \text{tp}(a/M; N)$  with  $(a, M, N)$  a uniqueness triple.

Under mild conditions, uniqueness triples are reduced:

**Fact 6.2.** If  $\mathbf{K}$  is a  $\lambda$ -superstable AEC with  $\lambda$ -symmetry which is categorical in  $\lambda$ , then any uniqueness triple is reduced.

*Proof.* By the proof of [JS13, 4.1.11(2)]. Note that conjugation (which is assumed there) holds by Fact 2.8. Further, the disjoint amalgamation statement used in the proof follows from Lemma 5.33.  $\square$

Once we have the existence property for uniqueness triples, we can build the desired good frame:

**Theorem 6.3.** Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC with  $\lambda$ -symmetry which is categorical in  $\lambda$ . If  $\mathbf{K}$  has the  $\lambda$ -existence property for uniqueness triples, then there is a weakly successful good  $\lambda$ -frame on  $\mathbf{K}$ .

*Proof.* We first follow the proof of [JS13, 5.5.4] to show that there exists a 4-ary relation NF on  $\mathbf{K}_\lambda$  satisfying several of the properties of nonforking amalgamation in a stable first-order theory: monotonicity, existence, uniqueness, symmetry and long transitivity (see [JS13, 5.2.1]).

The hypotheses of [JS13, 5.5.4] are that we work with a semi-good  $\lambda$ -frame with the conjugation property which has the existence property for uniqueness triples ([JS13, 5.1.1]). In the context here, we have the existence property for uniqueness triples by assumption and also have the conjugation property (Fact 2.8). Moreover, nonforking induces a good  $\lambda$ -frame, except that it only satisfies weak versions of continuity and local character (Fact 2.13). Now it is pointed out explicitly by Jarden and Shelah [JS13, 5.1.2] that local character is never used in their proof of [JS13, 5.5.4]. As for continuity, it is only used in one place: the proof of [JS13, 5.4.6] (the “opposite uniqueness proposition”). There an appeal to [JS13, 3.1.10] is made and its proof relies on continuity. However [JS13, 3.1.10] is nothing but Lemma 5.30, except that here we have the additional assumption (in clause (9)) that  $\mathcal{T}$  is reduced. This is not an issue, since the appeal to Lemma 5.30 in Jarden-Shelah is with towers  $\langle M_i : i < \alpha \rangle \frown \langle a_i : i + 1 < \alpha \rangle$  where for each  $i + 1 < \alpha$ ,  $(a_i, M_i, M_{i+1})$  is a uniqueness triple, hence in particular (Fact 6.2) is reduced. This implies by Lemma 5.15 that the entire tower is reduced.

Therefore the proof of [JS13, 5.5.4] goes through, and we have the desired relation NF. We now apply [JS, 4.3], which says that given a weak kind of good frame (called an almost good  $\lambda$ -frame) and a nonforking relation NF as here, we get a good  $\lambda$ -frame. In our setup, the frame induced by nonforking satisfies all the axioms of almost good  $\lambda$ -frame except perhaps for continuity. However the proof of [JS, 4.3] does not rely on continuity except to prove stability (which we already have), so we are done.  $\square$

To get the existence property for uniqueness triples, we will use that their existence follows from the weak diamond and some amount of stability in  $\lambda^+$  (the argument is essentially due to Shelah):

**Fact 6.4.** Let  $\mathbf{K}$  be a  $\lambda$ -superstable AEC with  $\lambda$ -symmetry which is categorical in  $\lambda$ . If  $\text{WGCH}(\lambda)$  and for any saturated model  $M$  in  $\mathbf{K}_{\lambda^+}$  there is  $N$  universal over  $M$ , then  $\mathbf{K}$  has the  $\lambda$ -existence property for uniqueness triples.

*Proof.* The proof of [Vas17a, E.8] goes through.  $\square$

**Corollary 6.5** (The local good frame construction theorem). Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . If:

- (1)  $\text{WGCH}(\lambda)$ .
- (2)  $\mathbf{K}$  is  $\lambda$ -superstable, has  $\lambda$ -symmetry, and is categorical in  $\lambda$ .
- (3) For every saturated model  $M$  in  $\mathbf{K}_{\lambda^+}$ , there is  $N$  universal over  $M$ .

Then there is a weakly successful good  $\lambda$ -frame with underlying class  $\mathbf{K}_\lambda$ .

*Proof.* Combine Theorem 6.3 and Fact 6.4.  $\square$

**Remark 6.6.** There are examples of categorical good frames that are not weakly successful [BV]. Thus we cannot expect to remove condition (3) from Corollary 6.5.

We can state a slightly more precise version of Corollary 6.5 using superlimits and the terminology from Definition 3.20. First, one more definition:

**Definition 6.7.** An AEC  $\mathbf{K}$  is *very nicely  $\lambda$ -superstable* (or *very nicely superstable in  $\lambda$* ) if  $\mathbf{K}$  is nicely  $\lambda$ -superstable and  $\mathbf{K}^{\lambda\text{-sl}}$  has the  $\lambda$ -existence property for uniqueness triples. For  $\Theta$  a class of cardinals,  $\mathbf{K}$  is *very nicely  $\Theta$ -superstable* (or *very nicely superstable in  $\Theta$* ) if  $\mathbf{K}$  is very nicely  $\lambda$ -superstable for every  $\lambda \in \Theta$ .

**Corollary 6.8.** If  $\mathbf{K}$  is very nicely  $\lambda$ -superstable, then there is a categorical weakly successful good  $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ .

*Proof.* Apply Theorem 6.3 to  $\mathbf{K}^{\lambda\text{-sl}}$ .  $\square$

**Corollary 6.9.** Let  $\mathbf{K}$  be an AEC and let  $\lambda \geq \text{LS}(\mathbf{K})$ . If:

- (1)  $\text{WGCH}(\lambda)$ .
- (2)  $\mathbf{K}$  is nicely  $\lambda$ -superstable.
- (3)  $\mathbf{K}$  is nicely  $\lambda^+$ -stable.

Then  $\mathbf{K}$  is very nicely  $\lambda$ -superstable.

*Proof.* By definition, it suffices to check that  $\mathbf{K}^{\lambda\text{-sl}}$  has the  $\lambda$ -existence property for uniqueness triples. By Fact 6.4, it suffices to see that for any saturated  $M \in \mathbf{K}_{\lambda^+}^{\lambda\text{-sl}}$ , there is  $N \in \mathbf{K}_{\lambda^+}^{\lambda\text{-sl}}$  universal over  $M$ . We know that  $\mathbf{K}$  is nicely  $[\lambda, \lambda^+]$ -stable, so by Theorem 3.19,  $\mathbf{K}^{\lambda\text{-sl}}$  is nicely  $\lambda^+$ -stable. Now note that a saturated model in  $\mathbf{K}_{\lambda^+}^{\lambda\text{-sl}}$  is nothing but a  $(\lambda, \lambda^+)$ -limit, which by Fact 3.18 is a  $(\lambda^+, \lambda^+)$ -limit. By definition of nice stability, this means that  $M$  is an amalgamation base, and hence has a universal extension, as desired.  $\square$

## 7. BUILDING SUCCESSFUL GOOD FRAMES

Suppose that  $\mathbf{K}$  is an AEC with some good structural properties,  $\lambda \geq \text{LS}(\mathbf{K})$ . In this section, we want to apply Corollary 6.5 to all successors of  $\lambda$ . What do we get if we succeed? In a nutshell, we get a sequence  $\langle \mathfrak{s}^n : n < \omega \rangle$ , where each  $\mathfrak{s}^n$  is a good  $\lambda^{+n}$ -frame. The following crucial result tells us that in fact the frames will be “connected”, in the sense that  $\mathfrak{s}^{n+1}$  is the successor of  $\mathfrak{s}^n$  (so forking in  $\mathfrak{s}^{n+1}$  is described in terms of forking in  $\mathfrak{s}^n$ ).

**Fact 7.1** ([Vas, 5.15]). Let  $\mathfrak{s}$  be a weakly successful categorical good  $\lambda$ -frame on an AEC  $\mathbf{K}$ . If there is a good  $\lambda^+$ -frame on  $\mathbf{K}^{\lambda^+\text{-sat}}$ , then  $\mathfrak{s}$  is successful good $^+$ .

We deduce a close relationship between nice superstability in  $[\lambda, \lambda^{+n}]$  and the existence of an  $n$ -successful good $^+$   $\lambda$ -frame:

**Theorem 7.2.** Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ , and let  $n \in [1, \omega]$ .

- (1) If  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+n}]$ -superstable, then there is an  $n$ -successful good<sup>+</sup>  $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ .
- (2) If  $\mathbf{K}_\lambda$  has a superlimit and there is an  $n$ -successful good<sup>+</sup>  $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ , then  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+(n-1)}]$ -superstable and nicely  $\lambda^{+n}$ -superstable.

*Proof.*

- (1) By Corollary 6.8, there is a weakly successful good  $\lambda$ -frame  $\mathfrak{s}$  on  $\mathbf{K}^{\lambda\text{-sl}}$ . We prove by induction on  $n \in [1, \omega)$  that  $\mathfrak{s}^{+(n-1)}$  is defined and is a successful good<sup>+</sup> frame on  $\mathbf{K}^{\lambda^{+(n-1)}\text{-sl}}$ . When  $n = 1$ ,  $\mathfrak{s}^{+0} = \mathfrak{s}$ . Moreover by Corollary 6.8 applied to  $\mathbf{K}_{\lambda^+}$ , there is a weakly successful good  $\lambda^+$ -frame on  $\mathbf{K}^{\lambda^+\text{-sl}}$ . Now,  $\mathbf{K}^{\lambda\text{-sl}}$  is dense in  $\mathbf{K}_{[\lambda, \lambda^+]}$  (Theorem 3.19), so the superlimit in  $\mathbf{K}_{\lambda^+}$  must be in  $\mathbf{K}^{\lambda\text{-sl}}$  and must be superlimit there too. Also, the superlimit in  $\mathbf{K}_{\lambda^+}$  is  $(\lambda^+, \lambda^+)$ -limit by Lemma 3.15, so it is saturated, i.e.  $\mathbf{K}^{\lambda^+\text{-sl}} = \mathbf{K}^{\lambda^+\text{-sat}}$ . By Fact 7.1,  $\mathfrak{s}$  is successful good<sup>+</sup>. For the inductive case, apply the base case to  $\mathbf{K}^{\lambda^{+(n-1)}\text{-sl}}$  (recalling the canonicity theorem, Fact 2.5).
- (2) Similarly to before (using Fact 2.12 to derive superstability from a good frame and [Vas, 5.12] to get the existence property for uniqueness triples), we get that  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+(n-1)}]$ -superstable. At the last step, we do not know that we get a weakly successful good frames, but we still have enough to derive that  $\mathbf{K}$  is nicely  $\lambda^{+n}$ -superstable.

□

**Remark 7.3.** As in Remark 6.6, the Hart-Shelah example shows [BV] that the second conclusion of Theorem 7.2 *cannot* be strengthened to  $\mathbf{K}$  being very nicely  $[\lambda, \lambda^{+n}]$ -superstable.

We now iterate Corollary 6.5. To get the best possible results, it will be handy to know we get superlimits automatically after one iteration:

**Fact 7.4.** If  $\mathfrak{s}$  is a successful categorical good<sup>+</sup>  $\lambda$ -frame on an AEC  $\mathbf{K}$ , then  $\mathfrak{s}^+$  is a good<sup>+</sup>  $\lambda$ -frame. In particular,  $\mathbf{K}_{\lambda^{++}}^{\lambda^+\text{-sat}}$  has a superlimit.

*Proof.* The frame  $\mathfrak{s}^+$  is good<sup>+</sup> by [She09a, III.1.9]. The “in particular” part then follows from [SVa, 3.14]. □

**Theorem 7.5.** Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ , and let  $n \in [1, \omega)$ . If:

- (1)  $\text{WGCH}([\lambda, \lambda^{+n}])$ .
- (2)  $\mathbf{K}$  is nicely  $[\lambda^{+2}, \lambda^{+(n+1)}]$ -stable.
- (3)  $\mathbf{K}_\lambda$  has a superlimit.
- (4)  $\mathbf{K}_{\lambda^+}$  has a superlimit.
- (5)  $\mathbf{K}$  is  $\lambda^{+(n+1)}$ -semisolvable (recall Definition 4.4).

Then  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+n}]$ -superstable.

*Proof.* Since  $\mathbf{K}_\lambda$  and  $\mathbf{K}_{\lambda^+}$  have superlimits,  $\mathbf{K}_{[\lambda, \lambda^+]}$  has no maximal models. Since  $\mathbf{K}$  is nicely  $[\lambda^{+2}, \lambda^{+n}]$ -stable,  $\mathbf{K}_{[\lambda, \lambda^{+n}]}$  has no maximal models. Thus  $\mathbf{K}$  is  $(\lambda^{+m}, \lambda^{+(n+1)})$ -extendible for any  $m \leq n+1$ . By Fact 4.6,  $\mathbf{K}$  is  $[\lambda, \lambda^{+(n+1)}]$ -semisolvable. By Fact 4.7,  $\mathbf{K}$  is also nicely  $[\lambda, \lambda^+]$ -stable, hence  $\mathbf{K}$  is nicely  $[\lambda, \lambda^{+(n+1)}]$ -stable. By Fact

4.7 again (used twice),  $\mathbf{K}$  is nicely  $[\lambda, \lambda^+]$ -superstable and by Corollary 6.9 (used twice),  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^+]$ -superstable. In particular (Theorem 7.2),  $\mathbf{K}^{\lambda\text{-sl}}$  has a successful good<sup>+</sup>  $\lambda$ -frame. By Fact 7.4,  $\mathbf{K}^{\lambda\text{-sl}}$  has a superlimit in  $\lambda^{++}$ . By Remark 3.12, this implies that  $\mathbf{K}_{\lambda^{++}}$  has a superlimit. Now iterate what has just been proven, replacing  $\lambda$  by  $\lambda^+$ .  $\square$

The assumption of nice  $\lambda^{+(n+1)}$ -stability in Theorem 7.5 cannot be removed because of the Hart-Shelah example (see Remark 7.3). If we are only interested in building an  $n$ -successful good<sup>+</sup>-frame, we can replace it by categoricity in  $\lambda^{+(n+1)}$ , or even by few models in  $\lambda^{+(n+1)}$ . To see this, we will use two more facts:

**Fact 7.6** ([JS13, 7.1.3]). Let  $\mathfrak{s}$  be a weakly successful categorical good  $\lambda$ -frame on an AEC  $\mathbf{K}$ . If  $\mathbb{I}(\mathbf{K}, \lambda^{++}) < 2^{\lambda^{++}}$ , then  $\mathfrak{s}$  is successful.

**Fact 7.7.** Let  $\mathfrak{s}$  be a successful categorical good  $\lambda$ -frame on an AEC  $\mathbf{K}$ . If  $\mathbf{K}_{\lambda^+}$  has a superlimit, then  $\mathfrak{s}$  is good<sup>+</sup>.

*Proof.* Combine [Vas, 3.3] and [Vas, 5.12].  $\square$

**Theorem 7.8.** Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ , and let  $n \in [1, \omega)$ . If:

- (1)  $\text{WGCH}([\lambda, \lambda^{+n}])$ .
- (2)  $\mathbf{K}$  is nicely  $[\lambda^{+2}, \lambda^{+n}]$ -stable.
- (3)  $\mathbf{K}_{\lambda}$  has a superlimit.
- (4)  $\mathbf{K}_{\lambda^+}$  has a superlimit.
- (5)  $\mathbf{K}$  is  $\lambda^{+(n+1)}$ -semisolvable.
- (6)  $\mathbb{I}(\mathbf{K}, \lambda^{+(n+1)}) < 2^{\lambda^{+(n+1)}}$ .

Then there is an  $n$ -successful good<sup>+</sup>  $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ .

*Proof.* As in the proof of Theorem 7.5, we can replace  $\lambda$  by  $\lambda^{+(n-1)}$  if necessary to assume without loss of generality that  $n = 1$ . Also as in the proof of Theorem 7.5,  $\mathbf{K}$  is very nicely  $\lambda$ -superstable. By Corollary 6.8, there is a weakly successful good  $\lambda$ -frame  $\mathfrak{s}$  on  $\mathbf{K}^{\lambda\text{-sl}}$ . By Fact 7.6,  $\mathfrak{s}$  is successful. By Fact 7.7,  $\mathfrak{s}$  is good<sup>+</sup>.  $\square$

**Remark 7.9.** In Theorem 7.8, if the superlimit in  $\mathbf{K}_{\lambda^+}$  is an amalgamation base, the proof shows that we can replace  $\text{WGCH}([\lambda, \lambda^{+n}])$  by  $\text{WGCH}([\lambda, \lambda^{+n}])$ .

## 8. THE MAIN CATEGORICITY TRANSFER

Having successful good frames implies that the categoricity spectrum is quite well understood, as the next two facts show:

**Fact 8.1.** Let  $\mathfrak{s}$  be a categorical  $n$ -successful good<sup>+</sup>  $\lambda$ -frame on an AEC  $\mathbf{K}$ . If  $\mathbf{K}$  is categorical in *some*  $\mu \in (\lambda, \lambda^{+(n+1)}]$ , then  $\mathbf{K}$  is categorical in *all*  $\mu' \in (\lambda, \lambda^{+(n+1)}]$ . Moreover,  $\mathbf{K}_{[\lambda, \lambda^{+n}]}$  has amalgamation, no maximal models, and is  $\lambda$ -tame.

*Proof.* Combining [She09a, III.2.3(4)], [She09a, III.2.9], and [She09a, III.2.12], we get that  $\mathbf{K}$  is categorical in every  $\mu' \in (\lambda, \lambda^{+(n+1)}]$ . Tameness then follows from [She09a, III.1.10]. As for amalgamation and no maximal models, they are immediate from categoricity and the definition of a good frame.  $\square$



**Fact 8.2** ([SVb]). Let  $\mathfrak{s}$  be a categorical  $\omega$ -successful  $\text{good}^+$   $\lambda$ -frame on an AEC  $\mathbf{K}$ . If  $\text{WGCH}([\lambda^{+n}, \lambda^{+\omega}))$  holds for some  $n < \omega$ , then  $\mathbf{K}$  has arbitrarily large models, and if  $\mathbf{K}$  is categorical in *some*  $\mu > \lambda$ , then  $\mathbf{K}$  is categorical in *all*  $\mu' > \lambda$  and moreover  $\mathbf{K}_{\geq \lambda}$  has amalgamation, no maximal models, and is  $\lambda$ -tame.

We can now prove the main technical categoricity transfer theorem of this paper. The assumption are essentially the following: no maximal models, some superlimits, and some instances of nice stability. They all readily follow in categorical AECS with amalgamation (see e.g. Fact 2.9), and we will see (Theorem 10.2) that they can also be derived from only no maximal models. The price to paid is a set-theoretic hypothesis: the WGCH.

**Theorem 8.3** (Main theorem). Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ ,  $\mu > \lambda^+$ , and set  $\mu_0 := \min(\mu, \lambda^{+\omega})$ . If:

- (1)  $\text{WGCH}([\lambda, \mu_0))$ .
- (2)  $\mathbf{K}$  has arbitrarily large models.
- (3)  $\mathbf{K}_\lambda$  has a superlimit.
- (4)  $\mathbf{K}_{\lambda^+}$  has a superlimit.
- (5)  $\mathbf{K}$  is nicely  $[\lambda^{+2}, \mu_0)$ -stable.
- (6)  $\mathbf{K}$  is  $(\mu_0, \mu)$ -extendible.
- (7)  $\mathbf{K}$  is categorical in  $\mu$ .

Then  $\mathbf{K}_{[\lambda, \mu]}^{\lambda\text{-sl}}$  is categorical in every  $\mu' \in [\lambda, \mu]$ , has amalgamation, no maximal models, and is  $\lambda$ -tame. Moreover if  $\mu \geq \lambda^{+\omega}$ , then  $\mathbf{K}^{\lambda\text{-sl}}$  is categorical in every  $\mu' \geq \lambda$ , has amalgamation, no maximal models, and is  $\lambda$ -tame.

*Proof.* Either  $\mu < \lambda^{+\omega}$  or  $\mu \geq \lambda^{+\omega}$ . We consider these cases separately.

- If  $\mu < \lambda^{+\omega}$ , fix  $n \in [1, \omega)$  such that  $\mu = \lambda^{+(n+1)}$  (recall that we are assuming that  $\mu > \lambda^+$ ). By Theorem 7.8, there is an  $n$ -successful  $\text{good}^+$   $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ . Now apply Fact 8.1.
- If  $\mu \geq \lambda^{+\omega}$ , first fix  $n \in [1, \omega)$ . Note that the hypotheses imply that  $\mathbf{K}_{[\lambda, \lambda^{+\omega})}$  has no maximal models, so  $\mathbf{K}$  must be  $(\lambda^{+(n+1)}, \mu)$ -extendible. By Fact 4.6,  $\mathbf{K}$  is  $\lambda^{+(n+1)}$ -semisolvable. By Theorem 7.5,  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+n}]$ -superstable. Since this is valid for all  $n < \omega$ ,  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+\omega})$ -superstable. By Theorem 7.2 (and canonicity, see Fact 2.5), there is an  $\omega$ -successful  $\text{good}^+$   $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ . Now apply Fact 8.2.

□

**Remark 8.4.** For any AEC  $\mathbf{K}$  and any  $\mu > \lambda \geq \text{LS}(\mathbf{K})$  such that  $\mathbf{K}_\lambda$  has a superlimit and  $\mathbf{K}$  is categorical in  $\mu$ ,  $\mathbf{K}_{\geq \mu}^{\lambda\text{-sl}} \subseteq \mathbf{K}_{\geq \mu}$ . In the conclusion of the above theorem, equality will hold, because  $\mathbf{K}^{\lambda\text{-sl}}$  has a model of cardinality  $\mu$ .

## 9. CATEGORICITY ASSUMING AMALGAMATION

In this section, we use Theorem 8.3 to fully describe the categoricity spectrum of AECS with amalgamation. Recall the definition of the categoricity spectrum:

**Definition 9.1.** The *categoricity spectrum* of an AEC  $\mathbf{K}$  is the class of cardinals  $\mu \geq \text{LS}(\mathbf{K})$  such that  $\mathbf{K}$  is categorical in  $\mu$ . We write  $\text{Cat}(\mathbf{K})$  for the categoricity spectrum of  $\mathbf{K}$ .

**Remark 9.2.** For any AEC  $\mathbf{K}$  and any cardinal  $\lambda$ ,  $\text{Cat}(\mathbf{K}_{\geq \lambda}) = \text{Cat}(\mathbf{K}) \cap [\lambda, \infty)$ .

We will use the following known results about the categoricity spectrum of AECs with amalgamation and arbitrarily large models. First, it is not too difficult to see that it is a closed class (see for example the end of the proof of [GV06c, 4.3]):

**Fact 9.3.** If  $\mathbf{K}$  is an AEC with amalgamation and arbitrarily large models, then  $\text{Cat}(\mathbf{K})$  is a closed class. More precisely, let  $\mathbf{K}$  be an AEC with arbitrarily large models and let  $\langle \mu_i : i < \delta \rangle$  be an increasing sequence in  $\text{Cat}(\mathbf{K})$ . If  $\mathbf{K}_{\leq \mu_i}$  has amalgamation for each  $i < \delta$ , then  $\sup_{i < \delta} \mu_i \in \text{Cat}(\mathbf{K})$ .

We will also use that a version of Morley's omitting type holds in AECs. This is essentially due to Shelah. See [Vas17a, 9.2] for a full proof.

**Fact 9.4** (Morley's omitting type theorem for AECs). Let  $\mathbf{K}$  be an AEC with amalgamation and let  $\mu > \text{LS}(\mathbf{K})$ . If every model in  $\mathbf{K}_\mu$  is  $\text{LS}(\mathbf{K})^+$ -saturated, then there exists  $\chi < h(\text{LS}(\mathbf{K}))$  such that every model in  $\mathbf{K}_{\geq \chi}$  is  $\text{LS}(\mathbf{K})^+$ -saturated.

Specializing Theorem 8.3 to AECs with amalgamation, we get that the categoricity spectrum has no gaps:

**Lemma 9.5.** Assume WGCH. Let  $\mathbf{K}$  be an AEC with amalgamation and arbitrarily large models. If  $\mu_1 < \mu_2$  are both in  $\text{Cat}(\mathbf{K})$ , then  $[\mu_1, \mu_2] \subseteq \text{Cat}(\mathbf{K})$ .

*Proof.* If  $\mu_2 = \mu_1^+$ , there is nothing to prove, so assume that  $\mu_2 > \mu_1^+$ . We apply Theorem 8.3, where  $\lambda, \mu$  there stand for  $\mu_1, \mu_2$  here. We have to check that its hypotheses are satisfied. By assumption WGCH holds and  $\mathbf{K}$  has arbitrarily large models. By categoricity in  $\mu_1$ ,  $\mathbf{K}_{\mu_1}$  has a superlimit. Since  $\mathbf{K}$  is categorical in  $\mu_1$ ,  $\mathbf{K}$  has joint embedding in  $\mu_1$ , and so a diagram chase (using amalgamation) shows that  $\mathbf{K}_{\geq \mu_1}$  also has joint embedding. In particular,  $\mathbf{K}_{\geq \mu_1}$  has no maximal models. In particular,  $\mathbf{K}$  is  $(\mu_0, \mu_2)$ -extendible, where  $\mu_0 := \min(\mu_2, \mu_1^{+\omega})$ . By Fact 4.6,  $\mathbf{K}$  is  $[\mu_1, \mu_2]$ -semisolvable. By Fact 4.7,  $\mathbf{K}$  is  $\mu$ -superstable and has  $\mu$ -symmetry, for every  $\mu \in [\mu_1, \mu_2)$ . In particular,  $\mathbf{K}$  is nicely  $[\mu_2^{+2}, \mu_0)$ -stable. By Fact 3.21, it follows that  $\mathbf{K}$  is nicely  $\mu_1^+$ -superstable. In particular,  $\mathbf{K}_{\mu_1^+}$  has a superlimit. This completes the verification that the hypotheses of Theorem 8.3 hold.  $\square$

Further, if we are only categorical in one cardinal we can apply Theorem 8.3 to a subclass of saturated models:

**Lemma 9.6.** Assume WGCH. Let  $\mathbf{K}$  be an AEC with amalgamation and arbitrarily large models. If  $\text{Cat}(\mathbf{K}) \cap [\text{LS}(\mathbf{K})^{+\omega}, \infty) \neq \emptyset$ , then there exists  $\chi < h(\text{LS}(\mathbf{K}))$  such that  $[\chi, \infty) \subseteq \text{Cat}(\mathbf{K})$ .

*Proof.* Let  $\mu \in \text{Cat}(\mathbf{K}) \cap [\text{LS}(\mathbf{K})^{+\omega}, \infty)$ . Partitioning the AEC into disjoint classes, each of which has joint embedding, we can assume without loss of generality that  $\mathbf{K}_{< \mu}$  has joint embedding (see [Vas16a, 10.13]). In particular,  $\mathbf{K}_{< \mu}$  has no maximal models. By Fact 2.9,  $\mathbf{K}^{\text{LS}(\mathbf{K})^+ \text{-sat}}$  is an AEC and  $\mathbf{K}^{\text{LS}(\mathbf{K})^+ \text{-sat}}$  is categorical in both

$\text{LS}(\mathbf{K})^+$  and  $\mu$ . Proceeding as in the proof of Lemma 9.5 (where  $\mathbf{K}, \mu_1, \mu_2$  there stand for  $\mathbf{K}^{\text{LS}(\mathbf{K})^+ \text{-sat}}, \text{LS}(\mathbf{K})^+, \mu$  here), we see that Theorem 8.3 applies, and hence  $\mathbf{K}^{\text{LS}(\mathbf{K})^+ \text{-sat}}$  is categorical in every  $\mu' \geq \text{LS}(\mathbf{K})^+$ . Now apply Fact 9.4.  $\square$

The two lemmas just proven are enough to give the full description of the categoricity spectrum:

**Corollary 9.7.** Assume WGCH. If  $\mathbf{K}$  is an AEC with amalgamation and arbitrarily large models, then exactly one of the following holds:

- (1)  $\text{Cat}(\mathbf{K}) = \emptyset$ .
- (2)  $\text{Cat}(\mathbf{K}) = [\text{LS}(\mathbf{K})^+{}^m, \text{LS}(\mathbf{K})^+{}^n]$  for some  $m \leq n < \omega$ .
- (3)  $\text{Cat}(\mathbf{K}) = [\chi, \infty)$  for some  $\chi \in [\text{LS}(\mathbf{K}), h(\text{LS}(\mathbf{K}))]$ .

*Proof.* Assume first that  $\text{Cat}(\mathbf{K}) \cap [\text{LS}(\mathbf{K})^+{}^\omega, \infty) \neq \emptyset$ . By Lemma 9.6, there is  $\chi < h(\text{LS}(\mathbf{K}))$  such that  $[\chi, \infty) \subseteq \text{Cat}(\mathbf{K})$ . Take the minimal such  $\chi$  with  $\chi \geq \text{LS}(\mathbf{K})$ . We have that  $\text{Cat}(\mathbf{K}) = [\chi, \infty)$ . Indeed, if  $\chi_0 \in \text{Cat}(\mathbf{K})$  and  $\chi_0 < \chi$ , then by Lemma 9.5,  $[\chi_0, \chi] \subseteq \text{Cat}(\mathbf{K})$ , and hence  $[\chi_0, \infty) \subseteq \text{Cat}(\mathbf{K})$ , contradicting the minimality of  $\chi$ .

Assume now that  $\text{Cat}(\mathbf{K}) \cap [\text{LS}(\mathbf{K})^+{}^\omega, \infty) = \emptyset$  but  $\text{Cat}(\mathbf{K}) \neq \emptyset$ . Let  $m < \omega$  be least such that  $\text{LS}(\mathbf{K})^+{}^m \in \text{Cat}(\mathbf{K})$  and let  $n < \omega$  be maximal such that  $\text{LS}(\mathbf{K})^+{}^n \in \text{Cat}(\mathbf{K})$ . Note that such an  $n$  exists: otherwise by Fact 9.3,  $\text{LS}(\mathbf{K})^+{}^\omega \in \text{Cat}(\mathbf{K})$ , which we have assumed does not happen. By Lemma 9.5,  $[\text{LS}(\mathbf{K})^+{}^m, \text{LS}(\mathbf{K})^+{}^n] \subseteq \text{Cat}(\mathbf{K})$ , and by minimality of  $m$  and maximality of  $n$ , this must be an equality:  $\text{Cat}(\mathbf{K}) = [\text{LS}(\mathbf{K})^+{}^m, \text{LS}(\mathbf{K})^+{}^n]$ , as desired.  $\square$

**9.1. Examples.** We end with examples showing that each of the cases of Corollary 9.7 may happen. We will need:

**Fact 9.8.** For each infinite cardinal  $\lambda$  and each ordinal  $\alpha < (2^\lambda)^+$ , there exists AECs  $\mathbf{K}^\alpha, \mathbf{K}^{*,\alpha}$  such that:

- (1)  $\text{LS}(\mathbf{K}^\alpha) = \text{LS}(\mathbf{K}^{*,\alpha}) = \lambda$ .
- (2)  $\mathbf{K}^\alpha, \mathbf{K}^{*,\alpha}$  have amalgamation.
- (3)  $\mathbf{K}^\alpha$  has a model of every cardinality  $\mu \in [\lambda, \beth_\alpha(\lambda)]$ , but no model of cardinality  $(\beth_\alpha(\lambda))^+$ .
- (4)  $\mathbf{K}^{*,\alpha}$  has a model of every cardinality  $\mu \in [\lambda, \lambda^{+\alpha})$  but no model of cardinality  $\lambda^{+\alpha}$ .

*Proof sketch.* By [She90, VII.5.5(6)],  $\delta(\lambda, 2^\lambda) = (2^\lambda)^+$ , where  $\delta(\lambda, 2^\lambda)$  is the least order type of a well-ordering not definable using the class of models of a first order theory in a vocabulary of cardinality  $\lambda$  omitting at most  $2^\lambda$ -many types. Thus by (the proof of [She90, VII.5.4]), for any  $\alpha < (2^\lambda)^+$ , we can code the cumulative hierarchy: there is an AEC  $\mathbf{K}^\alpha$  such that  $\text{LS}(\mathbf{K}^\alpha) = \lambda$ ,  $\mathbf{K}^\alpha$  has a model of cardinality  $\beth_\alpha(\lambda)$  but not models of cardinality  $(\beth_\alpha(\lambda))^+$ .

Coding successors is done similarly to obtain  $\mathbf{K}^{*,\alpha}$ : the basic template to code the successor operation is to work in a vocabulary with a linear order  $<$ , a predicate  $P$ , and a binary function  $F$ . We then consider structures where for any  $x$  not in  $P$ ,  $F(x, \cdot)$  codes an injection from the predecessors of  $x$  into  $P$ . The size of the

universe of such a structure is at most the successor of the size of  $P$  (the idea is probably folklore; it is described in [Gro]). Using this idea, one can define  $\mathbf{K}^{*,\alpha}$  when  $\alpha$  is a successor. When  $\alpha$  is zero, we can take  $\mathbf{K}^{*,0} = \mathbf{K}^0$ , and when  $\alpha$  is limit we can take a disjoint union of  $\mathbf{K}^{*,\beta+1}$  for  $\beta < \alpha$ .  $\square$

**Fact 9.9** ([HS90, BK09]). For each  $n < \omega$ , there is an AEC  $\mathbf{K}^{\text{hs},n}$  such that:

- (1)  $\text{LS}(\mathbf{K}^{\text{hs},n}) = \aleph_0$ .
- (2)  $\mathbf{K}^{\text{hs},n}$  has amalgamation and arbitrarily large models.
- (3)  $\text{Cat}(\mathbf{K}^{\text{hs},n}) = [\aleph_0, \aleph_n]$ .

**Example 9.10.**

- (1) The AEC of all fields (ordered by subfield) has empty categoricity spectrum, amalgamation, and arbitrarily large models. There are of course many other such examples.
- (2) Fix an infinite cardinal  $\lambda$  and an ordinal  $\alpha < (2^\lambda)^+$ . Take the disjoint union of the AEC  $\mathbf{K}^\alpha$  given by Fact 9.8 with a totally categorical AEC to get an AEC  $\mathbf{K}$  with amalgamation and arbitrarily large models such that  $\text{LS}(\mathbf{K}) = \lambda$  and  $\text{Cat}(\mathbf{K}) = [\beth_\alpha(\lambda)^+, \infty)$ . Similarly, using  $\mathbf{K}^{*,\alpha}$  we can get an AEC  $\mathbf{K}'$  with amalgamation and arbitrarily large models such that  $\text{LS}(\mathbf{K}') = \lambda$  and  $\text{Cat}(\mathbf{K}') = [\lambda^{+\alpha}, \infty)$ . If GCH holds, any  $\chi \in [\lambda, h(\lambda))$  is of the form  $\lambda^{+\beta}$  for some  $\beta < (2^\lambda)^+$ , so this gives a complete list of examples for the third case of Corollary 9.7.
- (3) Fix  $m \leq n < \omega$ . Take a disjoint union of the Hart-Shelah example ( $\mathbf{K}^{\text{hs},n}$  given by Fact 9.9) and the AEC  $\mathbf{K}^{*,m}$  given by Fact 9.8. We obtain an AEC  $\mathbf{K}$  with amalgamation and arbitrarily large models such that  $\text{LS}(\mathbf{K}) = \aleph_0$  and  $\text{Cat}(\mathbf{K}) = [\text{LS}(\mathbf{K})^{+m}, \text{LS}(\mathbf{K})^{+n}]$ . This gives a complete list of examples for the second case of Corollary 9.7 in case  $\text{LS}(\mathbf{K}) = \aleph_0$ . In case  $\text{LS}(\mathbf{K}) > \aleph_0$ , work in preparation of Shelah and Villaveces [SVc] constructs for each  $n < \omega$  (assuming GCH) an AEC with Löwenheim-Skolem-number  $\lambda$  which has categoricity spectrum  $[\lambda, \lambda^{+n}]$ . Using this, one could then also get a complete list of example for the second case of Corollary 9.7 in case  $\text{LS}(\mathbf{K}) = \lambda$ .

**Remark 9.11.** The last two examples given above all fail joint embedding. It would be interesting to know whether there are examples with joint embedding, or whether (assuming WGCH) for AECs with amalgamation, joint embedding, and arbitrarily large models, categoricity above  $\text{LS}(\mathbf{K})^{+\omega}$  implies categoricity everywhere strictly above  $\text{LS}(\mathbf{K})$ .

## 10. CATEGORICITY ASSUMING NO MAXIMAL MODELS

In this section, we study the categoricity spectrum of AECs with no maximal models. Take an AEC with no maximal models categorical in a high-enough cardinal. By Fact 4.7, nice stability holds in this setup (assuming GCHWD), so to apply Theorem 8.3, it remains to check that there are superlimits in  $\lambda$  and  $\lambda^+$ , for some suitable  $\lambda$ . The existence of superlimits may well follow from the uniqueness of limit models, but the latter is not known in this context (see the discussion in [Van06, Van13]). Instead, we will take  $\lambda$  a suitable fixed point of the  $\beth$  function

and rely heavily on [She09a, Chapter IV] (we do *not* rely on [She09a, §IV.2], which contains a gap – see the discussion at the beginning of [Vas17d, §4]).

We give a name to the kind of fixed points we will consider:

**Definition 10.1.** A cardinal  $\lambda$  is a *nice fixed point* if there exists a strictly increasing sequence  $\langle \lambda_n : n < \omega \rangle$  such that  $\lambda = \sup_{n < \omega} \lambda_n$  and  $\lambda_n = \beth_{\lambda_n}$  for all  $n < \omega$ .

Note that such fixed points can easily be found by iterating the  $\beth$  functions enough times. The following result is where we will use [She09a, Chapter IV]. Essentially, everything in the conclusion below except the existence of the superlimit in  $\mathbf{K}_{\lambda^+}$  is proven by Shelah there.

**Theorem 10.2.** Let  $\mathbf{K}$  be an AEC and let  $\mu > \lambda > \text{LS}(\mathbf{K})$ . If:

- (1)  $\lambda$  is a nice fixed point.
- (2)  $\mathbf{K}$  is  $\mu$ -solvable.
- (3) For  $M, N$  in  $\mathbf{K}_{\mu}^{\mu\text{-sl}}$ ,  $M \leq_{\mathbf{K}} N$  implies  $M \preceq_{\mathbb{L}_{\infty, \lambda}} N$ .
- (4)  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible.

Then  $\mathbf{K}$  is  $\lambda$ -solvable,  $\mathbf{K}_{\lambda^+}$  has a superlimit, and there is a type-full good  $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ .

*Proof.* By [She09a, IV.1.41],  $\mathbf{K}$  is pseudo  $\lambda$ -solvable, which means [She09a, IV.1.4(3)] that in the definition of solvability,  $\text{EM}_{\tau(\mathbf{K})}(I, \Phi)$  is not required to be universal (but must satisfy the other properties of a superlimit). In the present setup, we are assuming that  $\mathbf{K}$  is  $(\lambda, \mu)$ -extendible, and so in particular any  $M \in \mathbf{K}_{\lambda}$  embeds inside an EM model. This implies that the EM model must be universal, hence that  $\mathbf{K}$  is  $\lambda$ -solvable. In particular,  $\mathbf{K}_{\lambda}$  has a superlimit.

By [She09a, IV.4.10], there is a type-full good  $\lambda$ -frame on the class generated by the EM models witnessing pseudo-solvability. As argued in the previous paragraph, this class must be  $\mathbf{K}^{\lambda\text{-sl}}$ .

It remains to see that  $\mathbf{K}_{\lambda^+}$  has a superlimit. Since  $\mathbf{K}$  has a superlimit in  $\lambda$  and  $\mathbf{K}^{\lambda\text{-sl}}$  has a good  $\lambda$ -frame, we have by Fact 2.12 and Theorem 3.17 that  $\mathbf{K}$  is nicely  $\lambda$ -superstable. It is easy to check that the  $(\lambda, \lambda^+)$ -limit model is the saturated model in  $\mathbf{K}_{\lambda^+}^{\lambda\text{-sl}}$ . Thus (using Theorem 3.19 in the background), proving that  $\mathbf{K}_{\lambda^+}$  has a superlimit is equivalent to showing that for any limit  $\delta < \lambda^{++}$ , if  $\langle M_i : i < \delta \rangle$  is an increasing chain of saturated models (in  $\mathbf{K}_{\lambda^+}^{\lambda\text{-sl}}$ ), then  $\bigcup_{i < \delta} M_i$  is saturated. Work inside some saturated  $M$  with  $M_i \leq_{\mathbf{K}} M$  for all  $i < \delta$ .

The proof of [She09a, IV.4.4] gives that  $\mathbf{K}_{\lambda}$  is  $(< \kappa)$ -tame, for some  $\kappa < \lambda$ . We claim that  $\mathbf{K}$  (and hence  $M$ ) does not have the  $(\kappa, 0)$ -order property of length  $h(\kappa)$  (see [She99, 4.3] for the definition). Otherwise, by [She99, 4.7.2] (see [BGKV16, 5.13] for a proof), the EM model of cardinality  $\lambda$  would have such an order property, and this is impossible by [Vas17c, 3.4]. Note that  $h(\kappa) < \lambda$ , so we can apply [BV17a, 4.27] to prove using stability theory inside a model and averages that  $\bigcup_{i < \delta} M_i$  is saturated. Note that hypothesis (3) of [BV17a, 4.27] holds because of the definition of the good frame in [She09a, IV.4.10] (see the proof of the local character property there).  $\square$

To obtain hypothesis (3) in Theorem 10.2, we will use:

**Fact 10.3** ([She09a, IV.1.12]). Let  $\mathbf{K}$  be an AEC and let  $\mu \geq \lambda > \text{LS}(\mathbf{K})$ . If  $\mathbf{K}$  is categorical in  $\mu$  and  $\mu = \mu^{<\lambda}$ , then whenever  $M \leq_{\mathbf{K}} N$  are both in  $\mathbf{K}_\mu$ , we have that  $M \preceq_{\mathbb{L}_{\infty, \lambda}} N$ .

We can now prove an upward categoricity transfer from categoricity in one cardinal in AECs with no maximal models. The categoricity cardinal is assumed to satisfy some cardinal arithmetic – this will be remedied in the next result, at the cost of assuming categoricity in two cardinals.

**Theorem 10.4.** Let  $\mathbf{K}$  be an AEC and let  $\mu > \lambda > \text{LS}(\mathbf{K})$ . If:

- (1)  $\text{GCHWD}([\lambda, \lambda^{+\omega}])$ .
- (2)  $\mu = \mu^{<\lambda}$  (or just:  $M \leq_{\mathbf{K}} N$  both in  $\mathbf{K}_\mu$  implies  $M \preceq_{\mathbb{L}_{\infty, \lambda}} N$ ).
- (3)  $\lambda$  is a nice fixed point.
- (4)  $\lambda^{+\omega} \leq \mu$ .
- (5)  $\mathbf{K}$  is  $([\lambda, \lambda^{+\omega}], \mu)$ -extendible.
- (6)  $\mathbf{K}$  is categorical in  $\mu$ .

Then  $\mathbf{K}$  is categorical in every  $\mu' \geq \mu$ .

*Proof.* By Fact 10.3,  $M \leq_{\mathbf{K}} N$  both in  $\mathbf{K}_\mu$  implies  $M \preceq_{\mathbb{L}_{\infty, \lambda}} N$ . We want to apply Theorem 8.3. We check its hypotheses. Since  $\text{GCHWD}([\lambda, \lambda^{+\omega}])$ , we have that  $\text{WGCH}([\lambda, \lambda^{+\omega}])$ . Since  $\lambda$  is a nice fixed point, we have that  $h(\text{LS}(\mathbf{K})) < \lambda$ , so  $\mathbf{K}$  has arbitrarily large models (Fact 4.1). By Theorem 10.2,  $\mathbf{K}_\lambda$  has a superlimit and  $\mathbf{K}_{\lambda^+}$  has a superlimit. By Fact 4.6 and the extendibility hypothesis,  $\mathbf{K}$  is  $[\lambda, \lambda^{+\omega}]$ -semisolvable. In particular by Fact 4.7,  $\mathbf{K}$  is nicely  $[\lambda^{+2}, \lambda^{+\omega}]$ -stable. This completes the verification that the hypotheses of Theorem 8.3 hold.  $\square$

**Remark 10.5.** The proof shows that  $\text{GCHWD}([\lambda, \lambda^{+\omega}])$  can be replaced with the conjunction of  $\text{WGCH}([\lambda, \lambda^+])$  and  $\text{GCHWD}([\lambda^{+2}, \lambda^{+\omega}])$ .

The next result removes the assumption that  $\mu = \mu^{<\lambda}$  in Theorem 10.4 at the cost of assuming categoricity in two cardinals.

**Theorem 10.6.** Let  $\mathbf{K}$  be an AEC and let  $\mu_2 > \mu_1 \geq \lambda > \text{LS}(\mathbf{K})$ . If:

- (1)  $\text{GCHWD}([\lambda, \lambda^{+\omega}])$ .
- (2)  $\text{WGCH}(\mu_1)$ .
- (3)  $\lambda$  is a nice fixed point.
- (4)  $\mathbf{K}$  is  $([\lambda, \lambda^{+\omega}], \mu_2)$ -extendible.
- (5)  $\mathbf{K}$  is categorical in  $\mu_1$ .
- (6)  $\mathbf{K}$  is  $(\mu_1^+, \mu_2)$ -extendible.
- (7)  $\mathbf{K}$  is categorical in  $\mu_2$ .
- (8) If  $\mu_1 = \lambda$ , then  $\mu_2 = \mu_2^{<\lambda}$ .

Then  $\mathbf{K}$  is categorical in every  $\mu \in [\mu_1, \mu_2]$ . Moreover if  $\mu_2 \geq \lambda^{+\omega}$ ,  $\mathbf{K}$  is categorical in every  $\mu' \geq \mu_1$ .

*Proof.* If  $\mu_2 = \lambda^+$ , the conclusion is vacuous, so assume that  $\mu_2 > \lambda^+$ . As in the proof of Theorem 10.4,  $\mathbf{K}$  has arbitrarily large models. In particular (Remark 4.5),

$\mathbf{K}$  is  $\mu_1$ -solvable and  $\mu_2$ -solvable. Since  $\mathbf{K}$  is  $(\mu_1^+, \mu_2)$ -extendible, Fact 4.6 says that  $\mathbf{K}$  is  $\mu_1^+$ -semisolvable. By Fact 4.7,  $\mathbf{K}$  is nicely  $\mu_1$ -stable. By Lemma 3.16, for  $M \leq_{\mathbf{K}} N$  both in  $\mathbf{K}_{\mu_1}$ ,  $M \preceq_{\mathbb{L}_{\infty, \mu_1}} N$ . In particular,  $M \preceq_{\mathbb{L}_{\infty, \lambda}} N$ . By Theorem 10.2 (using Fact 10.3 if  $\mu_1 = \lambda$ ),  $\mathbf{K}_{\lambda}$  has a superlimit and  $\mathbf{K}_{\lambda^+}$  has a superlimit. Let  $\mu'_2 := \min(\mu_2, \lambda^{+\omega})$ . We apply Theorem 8.3, where  $\mu_0, \mu$  there stand for  $\mu'_2, \mu_2$  here. We first have to check its hypotheses.  $\text{WGCH}([\lambda, \mu'_2])$  holds because we are assuming more:  $\text{GCHWD}([\lambda, \lambda^{+\omega}])$ . We have already established that  $\mathbf{K}$  has arbitrarily large models,  $\mathbf{K}_{\lambda}$  has a superlimit, and  $\mathbf{K}_{\lambda^+}$  has a superlimit. That  $\mathbf{K}$  is  $(\mu'_2, \mu_2)$ -extendible follows from hypothesis (4) here, and that  $\mathbf{K}$  is categorical in  $\mu_2$  is also assumed. It remains to see that  $\mathbf{K}$  is nicely  $[\lambda^{+2}, \mu'_2)$ -stable: use Facts 4.6 and 4.7 again.  $\square$

We are now ready to give a short list of possibilities for the categoricity spectrum above a nice fixed point. This shows in particular that the categoricity spectrum of an AEC with no maximal models is either bounded or contains an end segment (assuming GCHWD). Here, we do not know whether the three intermediate possibilities listed below can happen (Conjecture 1.3 would imply they cannot happen).

**Corollary 10.7.** Assume GCHWD. Let  $\mathbf{K}$  be an AEC and let  $\lambda > \text{LS}(\mathbf{K})$  be a nice fixed point. If  $\mathbf{K}_{\geq \lambda}$  has no maximal models, then exactly one of the following possibilities holds:

- (1)  $\text{Cat}(\mathbf{K}_{\geq \lambda}) = \emptyset$ .
- (2)  $\text{Cat}(\mathbf{K}_{\geq \lambda}) = [\lambda^{+m}, \lambda^{+n}]$  for some  $m \leq n < \omega$ .
- (3)  $\text{Cat}(\mathbf{K}_{\geq \lambda}) = \{\mu\}$ , for some  $\mu \geq \lambda^{+\omega}$  with  $\mu < \mu^{<\lambda}$ .
- (4)  $\text{Cat}(\mathbf{K}_{\geq \lambda}) = \{\lambda, \mu\}$ , for some  $\mu \geq \lambda^{+\omega}$  with  $\mu < \mu^{<\lambda}$ .
- (5)  $\text{Cat}(\mathbf{K}_{\geq \lambda}) = [\chi, \infty)$  for some  $\chi \geq \lambda$ .

*Proof.* First, we give a couple of consequences of the two theorems proven above:

Claim 1: If  $\lambda < \mu_1 < \mu_2$  are such that  $\mu_1, \mu_2 \in \text{Cat}(\mathbf{K})$ , then  $[\mu_1, \mu_2] \subseteq \text{Cat}(\mathbf{K})$ . Moreover if in addition  $\mu_2 \geq \lambda^{+\omega}$  then  $[\mu_2, \infty) \subseteq \text{Cat}(\mathbf{K})$ .

Proof of Claim 1: By Theorem 10.6.  $\dagger_{\text{Claim 1}}$

Claim 2: If there exists  $n < \omega$  such that  $\lambda, \lambda^{+n} \in \text{Cat}(\mathbf{K})$ , then  $[\lambda, \lambda^{+n}] \subseteq \text{Cat}(\mathbf{K})$ .

Proof of Claim 2: By Theorem 10.6, using that by GCH,  $(\lambda^{+n})^{<\lambda} = \lambda^{+n}$ .  $\dagger_{\text{Claim 2}}$

Claim 3: If  $\mu \geq \lambda^{+\omega}$  is such that  $\mu = \mu^{<\lambda}$  and  $\mu \in \text{Cat}(\mathbf{K})$ , then  $[\mu, \infty) \subseteq \text{Cat}(\mathbf{K})$ .

Proof of Claim 3: By Theorem 10.4 and Fact 10.3.  $\dagger_{\text{Claim 3}}$

Now assume first that  $\text{Cat}(\mathbf{K})$  is unbounded. By Claim 1, there must exist a cardinal  $\chi \geq \lambda$  such that  $[\chi, \infty) \subseteq \text{Cat}(\mathbf{K}_{\geq \lambda})$ . Take the least such  $\chi$ . By Claim 1 and Claim 2, we must have that  $[\chi, \infty) = \text{Cat}(\mathbf{K}_{\geq \lambda})$ . Thus (5) holds.

Now assume that  $\text{Cat}(\mathbf{K})$  is bounded but (1) fails, i.e.  $\text{Cat}(\mathbf{K}_{\geq \lambda}) \neq \emptyset$ . If  $\text{Cat}(\mathbf{K}_{\geq \lambda^{+\omega}}) \neq \emptyset$ , then by the moreover part of Claim 1,  $\text{Cat}(\mathbf{K}_{> \lambda}) = \{\mu\}$  for some  $\mu \geq \lambda^{+\omega}$ , and by Claim 3  $\mu < \mu^{<\lambda}$ . In this case either (3) or (4) must happen. Finally, if  $\text{Cat}(\mathbf{K}_{\geq \lambda^{+\omega}}) = \emptyset$  but  $\text{Cat}(\mathbf{K}_{\geq \lambda}) \neq \emptyset$ , pick  $m < \omega$  minimal such that  $\lambda^{+m} \in \text{Cat}(\mathbf{K})$  and  $n < \omega$  maximal such that  $\lambda^{+n} \in \text{Cat}(\mathbf{K})$  (as in the proof of Corollary 9.7, such an  $n$  must exist). By Claim 1 and Claim 2,  $[\lambda^{+m}, \lambda^{+n}] \subseteq \text{Cat}(\mathbf{K})$ . By minimality of  $m$  and maximality of  $n$ , this must be an equality, so (2) holds.  $\square$

We conclude with the eventual categoricity conjecture for AECs with no maximal models. Note that the proof of existence for the map  $\lambda \mapsto \mu_\lambda$  below is not constructive: it does not give an explicit bound on  $\mu_\lambda$ .

**Corollary 10.8** (Eventual categoricity for AECs with no maximal models). Assuming GCHWD, there exists a map  $\lambda \mapsto \mu_\lambda$  such that for any AEC  $\mathbf{K}$  with no maximal models, if  $\mathbf{K}$  is categorical in *some*  $\mu \geq \mu_{\text{LS}(\mathbf{K})}$ , then  $\mathbf{K}$  is categorical in *all*  $\mu' \geq \mu_{\text{LS}(\mathbf{K})}$ .

*Proof.* A consequence of Corollary 10.7 is that in any AEC with no maximal models, the categoricity spectrum is either bounded or contains an end segment. As shown in (for example) [Bal09, 15.13], this suffices: for a fixed  $\lambda$ , there is only a set of AECs with Löwenheim-Skolem-Tarski number  $\lambda$ . By the axiom of replacement, there will exist  $\mu_\lambda^0$  bounding the categoricity spectrum of all those where it is bounded. Similarly, there will exist  $\mu_\lambda^1$  bounding the start of the end-segment for all those where it is unbounded. Take  $\mu_\lambda := \mu_\lambda^0 + \mu_\lambda^1$ .  $\square$

## 11. SOME AUXILIARY RESULTS

We give here some easy corollaries of our methods that were buried in our notation. Specifically, we study tameness and existence of good frames in AECs with amalgamation, and also start the investigation of AECs with a lot of superlimits.

First, we show that good frames, and in fact successful ones, exist as low as possible in categorical AECs with amalgamation. This implies tameness and (if the categoricity cardinal is high-enough) a transfer of the structure of the class upward.

**Corollary 11.1.** Assume WGCH. Let  $\mathbf{K}$  be an AEC with arbitrarily large models and let  $\mu > \text{LS}(\mathbf{K})^+$  be such that  $\mathbf{K}_{<\mu}$  has amalgamation. If  $\mathbf{K}$  is categorical in both  $\text{LS}(\mathbf{K})$  and  $\mu$ , then:

- (1) There is a good<sup>+</sup>  $\text{LS}(\mathbf{K})$ -frame on  $\mathbf{K}$ , which is  $n$ -successful whenever  $\text{LS}(\mathbf{K})^{+(n+1)} \leq \mu$ .
- (2)  $\mathbf{K}_{<\mu}$  is  $\text{LS}(\mathbf{K})$ -tame and categorical in all  $\mu' \in [\text{LS}(\mathbf{K}), \mu]$ .
- (3) If  $\mu \geq \text{LS}(\mathbf{K})^{+\omega}$ , then  $\mathbf{K}$  has amalgamation, no maximal models, is  $\text{LS}(\mathbf{K})$ -tame, and is categorical in all  $\mu' \geq \text{LS}(\mathbf{K})$ .

*Proof.* The proof of Lemma 9.5 (where  $\mu_1, \mu_2$  there stand for  $\text{LS}(\mathbf{K}), \mu$  here) shows that the hypotheses of Theorem 7.8 (with  $\text{LS}(\mathbf{K})$  here standing for  $\lambda$  there) are satisfied. Now apply Facts 8.1 and 8.2.  $\square$

If we assume just categoricity in one cardinal, we obtain the previous result on a subclass of saturated models:

**Corollary 11.2.** Assume WGCH. Let  $\mathbf{K}$  be an AEC with arbitrarily large models and let  $\mu > \text{LS}(\mathbf{K})^{+2}$  be such that  $\mathbf{K}_{<\mu}$  has amalgamation and no maximal models. If  $\mathbf{K}$  is categorical in  $\mu$ , then:

- (1) There is a good<sup>+</sup>  $\text{LS}(\mathbf{K})^+$ -frame on  $\mathbf{K}^{\text{LS}(\mathbf{K})^+ \text{-sat}}$ , which is  $n$ -successful whenever  $\text{LS}(\mathbf{K})^{+(n+2)} \leq \mu$ .



- (2)  $\mathbf{K}_{<\mu}^{\text{LS}(\mathbf{K})^+-\text{sat}}$  is  $\text{LS}(\mathbf{K})^+$ -tame. In particular (using the notation of [Bal09, 11.6]),  $\mathbf{K}$  is  $(\text{LS}(\mathbf{K})^+, <\mu)$ -weakly tame.
- (3) If  $\mu \geq \text{LS}(\mathbf{K})^{+\omega}$ , then  $\mathbf{K}$  has amalgamation, no maximal models, and is  $\text{LS}(\mathbf{K})^+$ -weakly tame.

*Proof.* Apply Corollary 11.1 to  $\mathbf{K}^{\text{LS}(\mathbf{K})^+-\text{sat}}$  (it is an AEC by Fact 2.9).  $\square$

Grossberg and VanDieren have conjectured [GV06a, 1.5] that tameness should follow from high-enough categoricity. We are now in a position to prove that this holds, assuming WGCH and amalgamation. This also proves Conjecture 1.3 for AECs with amalgamation.

**Corollary 11.3.** Assume WGCH. Let  $\mathbf{K}$  be an AEC with amalgamation. If  $\mathbf{K}$  is categorical in *some*  $\mu \geq h(\text{LS}(\mathbf{K}))$ , then there is  $\chi < h(\text{LS}(\mathbf{K}))$  such that  $\mathbf{K}$  is categorical in *all*  $\mu' \geq \chi$  and  $\mathbf{K}$  is  $\chi$ -tame.

*Proof.* By Fact 4.1,  $\mathbf{K}$  has arbitrarily large models. By Corollary 9.7, there is  $\chi < h(\text{LS}(\mathbf{K}))$  such that  $\mathbf{K}$  is categorical in all  $\mu' \geq \chi$ . By Corollary 11.1,  $\mathbf{K}$  is  $\chi$ -tame.  $\square$

In [She09a, II.3.7], Shelah gives a construction of a good frame using WGCH, categoricity in  $\lambda$ ,  $\lambda^+$ , and few models in  $\lambda^{++}$ . This can be iterated to get  $n$ -successful frames (see [She09a, II.9.1]). One can see Theorems 7.5 and 7.8 as variations where less structural assumptions are made in the  $\lambda^{+m}$ 's, but some global assumptions (semisolvability, which implies arbitrarily large models) are also made. We give here a consequence of Theorem 7.5 which shows that it is enough to assume that the class has superlimits in the  $\lambda^{+m}$ 's. Thus in a sense we replace categoricity in Shelah's theorem by superlimits, and we replace few models by semisolvability:

**Corollary 11.4.** Let  $\mathbf{K}$  be an AEC, let  $\lambda \geq \text{LS}(\mathbf{K})$ , and let  $n < \omega$ . If:

- (1)  $\text{WGCH}([\lambda, \lambda^{+(n+1)}])$ .
- (2)  $\mathbf{K}_{\lambda^{+m}}$  has a superlimit for every  $m \leq n+1$ .
- (3)  $\mathbf{K}$  is  $\lambda^{+(n+2)}$ -semisolvable.

Then  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+n}]$ -superstable. In particular, there is a weakly successful good  $\lambda$ -frame on  $\mathbf{K}^{\lambda\text{-sl}}$ , which will be  $n$ -successful good $^+$  when  $n \geq 1$ .

*Proof.* The “in particular” part follows from Corollary 6.8 and Theorem 7.2. To see that  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+n}]$ -superstable, first observe that by the existence of superlimits,  $\mathbf{K}_{[\lambda, \lambda^{+(n+1)}]}$  has no maximal models, so  $\mathbf{K}$  is  $([\lambda, \lambda^{+(n+1)}], \lambda^{+(n+2)})$ -extendible. By Fact 4.6, this implies that  $\mathbf{K}$  is  $[\lambda, \lambda^{+(n+2)}]$ -semisolvable. By Fact 4.7, we get that  $\mathbf{K}$  is nicely  $[\lambda, \lambda^{+(n+1)}]$ -superstable. Now apply Corollary 6.9 to get that  $\mathbf{K}$  is very nicely  $[\lambda, \lambda^{+n}]$ -superstable.  $\square$

Similarly, we get that in an AEC with a superlimit at every cardinal, the categoricity spectrum has no gaps. More precisely:

**Corollary 11.5.** Let  $\mathbf{K}$  be an AEC and let  $\mu > \lambda \geq \text{LS}(\mathbf{K})$ . If:

- (1)  $\text{WGCH}([\lambda, \min(\mu, \lambda^{+\omega})])$ .

- (2)  $\mathbf{K}$  has arbitrarily large models.
- (3)  $\mathbf{K}$  is categorical in  $\lambda$ .
- (4) For every  $\lambda' \in (\lambda, \mu)$ ,  $\mathbf{K}_{\lambda'}$  has a superlimit.
- (5)  $\mathbf{K}$  is categorical in  $\mu$ .

Then  $\mathbf{K}$  is categorical in every  $\mu' \in [\lambda, \mu]$ . Moreover if  $\mu \geq \lambda^{+\omega}$ ,  $\mathbf{K}$  is categorical in every  $\mu' \geq \lambda$ .

*Proof.* By definition of superlimits,  $\mathbf{K}_{[\lambda, \mu]}$  has no maximal models. By Fact 4.6,  $\mathbf{K}$  is  $[\lambda, \mu]$ -semisolvability. By Fact 4.7,  $\mathbf{K}$  is nicely  $[\lambda, \mu]$ -superstable. Thus the hypotheses of Theorem 8.3 hold, hence its conclusion.  $\square$

As in the proof of Corollary 10.8, we deduce that AECs with a superlimit everywhere satisfy the eventual categoricity conjecture. Note that having superlimits imply no maximal models, but as opposed to Corollary 10.8, Corollary 11.5 assumes only WGCH instead of GCHWD.

In the opinion of the author, Corollaries 11.4 and 11.5 suggest that AECs with superlimits are an interesting framework to study. This was already foreshadowed by Shelah in the introduction to [She09a]. Specifically, Shelah conjectures [She09a, N.4.6] that a variation of the eventual categoricity conjecture holds, but with solvability replacing categoricity. Corollary 11.5 shows that (a weakening of) this conjecture implies the eventual categoricity conjecture, assuming WGCH. Thus the superlimit and (semi)solvability spectrums seem to be the key to an understanding of superstability and categoricity in abstract elementary classes.

## REFERENCES

- [Bal06] John T. Baldwin, *Abstract elementary classes: some answers, more questions*, URL: <http://homepages.math.uic.edu/~jbaldwin/pub/turino2.pdf>, 2006.
- [Bal09] ———, *Categoricity*, University Lecture Series, vol. 50, American Mathematical Society, 2009.
- [BET07] John T. Baldwin, Paul C. Eklof, and Jan Trlifaj,  ${}^{\perp}N$  as an abstract elementary class, *Annals of Pure and Applied Logic* **149** (2007), 25–39.
- [BGKV16] Will Boney, Rami Grossberg, Alexei Kolesnikov, and Sebastien Vasey, *Canonical forking in AECs*, *Annals of Pure and Applied Logic* **167** (2016), no. 7, 590–613.
- [BGVV17] Will Boney, Rami Grossberg, Monica VanDieren, and Sebastien Vasey, *Superstability from categoricity in abstract elementary classes*, *Annals of Pure and Applied Logic* **168** (2017), no. 7, 1383–1395.
- [BK09] John T. Baldwin and Alexei Kolesnikov, *Categoricity, amalgamation, and tameness*, *Israel Journal of Mathematics* **170** (2009), 411–443.
- [Bon14] Will Boney, *Tameness from large cardinal axioms*, *The Journal of Symbolic Logic* **79** (2014), no. 4, 1092–1119.
- [BR12] Tibor Beke and Jiří Rosický, *Abstract elementary classes and accessible categories*, *Annals of Pure and Applied Logic* **163** (2012), 2008–2017.
- [BV] Will Boney and Sebastien Vasey, *Good frames in the Hart-Shelah example*, *Archive for Mathematical Logic*, to appear. URL: <http://arxiv.org/abs/1607.03885v4>. DOI: 10.1007/s00153-017-0599-7.
- [BV17a] ———, *Chains of saturated models in AECs*, *Archive for Mathematical Logic* **56** (2017), no. 3, 187–213.
- [BV17b] ———, *A survey on tame abstract elementary classes*, *Beyond first order model theory* (José Iovino, ed.), CRC Press, 2017, pp. 353–427.
- [DS78] Keith J. Devlin and Saharon Shelah, *A weak version of  $\diamond$  which follows from  $2^{\aleph_0} < 2^{\aleph_1}$* , *Israel Journal of Mathematics* **29** (1978), no. 2, 239–247.

- [Gro] Rami Grossberg, *A course in model theory I*, A book in preparation.
- [Gro02] ———, *Classification theory for abstract elementary classes*, Contemporary Mathematics **302** (2002), 165–204.
- [GV06a] Rami Grossberg and Monica VanDieren, *Categoricity from one successor cardinal in tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 2, 181–201.
- [GV06b] ———, *Galois-stability for tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 1, 25–49.
- [GV06c] ———, *Shelah’s categoricity conjecture from a successor for tame abstract elementary classes*, The Journal of Symbolic Logic **71** (2006), no. 2, 553–568.
- [GV17] Rami Grossberg and Sebastien Vasey, *Equivalent definitions of superstability in tame abstract elementary classes*, The Journal of Symbolic Logic **82** (2017), no. 4, 1387–1408.
- [GVV16] Rami Grossberg, Monica VanDieren, and Andrés Villaveces, *Uniqueness of limit models in classes with amalgamation*, Mathematical Logic Quarterly **62** (2016), 367–382.
- [Hru96] Ehud Hrushovski, *The Mordell-Lang conjecture for function fields*, Journal of the American Mathematical Society **9** (1996), no. 3, 667–690.
- [HS90] Bradd Hart and Saharon Shelah, *Categoricity over  $P$  for first order  $T$  or categoricity for  $\phi \in L_{\omega_1, \omega}$  can stop at  $\aleph_k$  while holding for  $\aleph_0, \dots, \aleph_{k-1}$* , Israel Journal of Mathematics **70** (1990), 219–235.
- [Jec03] Thomas Jech, *Set theory*, 3rd ed., Springer-Verlag, 2003.
- [JS] Adi Jarden and Saharon Shelah, *Non forking good frames without local character*, Preprint. URL: <http://arxiv.org/abs/1105.3674v1>.
- [JS13] ———, *Non-forking frames in abstract elementary classes*, Annals of Pure and Applied Logic **164** (2013), 135–191.
- [LR16] Michael J. Lieberman and Jiří Rosický, *Classification theory for accessible categories*, The Journal of Symbolic Logic **81** (2016), no. 1, 151–165.
- [Mor65] Michael Morley, *Categoricity in power*, Transactions of the American Mathematical Society **114** (1965), 514–538.
- [MS90] Michael Makkai and Saharon Shelah, *Categoricity of theories in  $L_{\kappa, \omega}$ , with  $\kappa$  a compact cardinal*, Annals of Pure and Applied Logic **47** (1990), 41–97.
- [She74] Saharon Shelah, *Categoricity of uncountable theories*, Proceedings of the Tarski symposium (Leon Henkin, John Addison, C. C. Chang, William Craig, Dana Scott, and Robert Vaught, eds.), American Mathematical Society, 1974, pp. 187–203.
- [She83] ———, *Classification theory for non-elementary classes I: The number of uncountable models of  $\psi \in L_{\omega_1, \omega}$ . Part A*, Israel Journal of Mathematics **46** (1983), no. 3, 214–240.
- [She87a] ———, *Classification of non elementary classes II. Abstract elementary classes*, Classification Theory (Chicago, IL, 1985) (John T. Baldwin, ed.), Lecture Notes in Mathematics, vol. 1292, Springer-Verlag, 1987, pp. 419–497.
- [She87b] ———, *Universal classes*, Classification theory (Chicago, IL, 1985) (John T. Baldwin, ed.), Lecture Notes in Mathematics, vol. 1292, Springer-Verlag, 1987, pp. 264–418.
- [She90] ———, *Classification theory and the number of non-isomorphic models*, 2nd ed., Studies in logic and the foundations of mathematics, vol. 92, North-Holland, 1990.
- [She99] ———, *Categoricity for abstract classes with amalgamation*, Annals of Pure and Applied Logic **98** (1999), no. 1, 261–294.
- [She00] ———, *On what I do not understand (and have something to say), model theory*, Math. Japonica **51** (2000), 329–377.
- [She09a] ———, *Classification theory for abstract elementary classes*, Studies in Logic: Mathematical logic and foundations, vol. 18, College Publications, 2009.
- [She09b] ———, *Classification theory for abstract elementary classes 2*, Studies in Logic: Mathematical logic and foundations, vol. 20, College Publications, 2009.
- [She15] ———, *A.E.C. with not too many models*, pp. 367–402, De Gruyter, 2015.
- [SK96] Saharon Shelah and Oren Kolman, *Categoricity of theories in  $L_{\kappa, \omega}$ , when  $\kappa$  is a measurable cardinal. Part I*, Fundamentae Mathematica **151** (1996), 209–240.
- [SVa] Saharon Shelah and Sebastien Vasey, *Abstract elementary classes stable in  $\aleph_0$* , Annals of Pure and Applied Logic, To appear. URL: <http://arxiv.org/abs/1702.08281v5>. DOI: 10.1016/j.apal.2018.02.004.
- [SVb] ———, *Categoricity and multidimensional diagrams*, In preparation.

- [SVc] Saharon Shelah and Andrés Villaveces, *Categoricity may fail late*, Preprint. URL: <http://arxiv.org/abs/math/0404258v1>.
- [SV99] ———, *Toward categoricity for classes with no maximal models*, Annals of Pure and Applied Logic **97** (1999), 1–25.
- [Van06] Monica VanDieren, *Categoricity in abstract elementary classes with no maximal models*, Annals of Pure and Applied Logic **141** (2006), 108–147.
- [Van13] ———, *Erratum to "Categoricity in abstract elementary classes with no maximal models" [Ann. Pure Appl. Logic 141 (2006) 108–147]*, Annals of Pure and Applied Logic **164** (2013), no. 2, 131–133.
- [Van16a] ———, *Superstability and symmetry*, Annals of Pure and Applied Logic **167** (2016), no. 12, 1171–1183.
- [Van16b] ———, *Symmetry and the union of saturated models in superstable abstract elementary classes*, Annals of Pure and Applied Logic **167** (2016), no. 4, 395–407.
- [Vas] Sebastien Vasey, *Tameness from two successive good frames*, Preprint. URL: <http://arxiv.org/abs/1707.09008v3>.
- [Vas16a] ———, *Building independence relations in abstract elementary classes*, Annals of Pure and Applied Logic **167** (2016), no. 11, 1029–1092.
- [Vas16b] ———, *Forking and superstability in tame AECs*, The Journal of Symbolic Logic **81** (2016), no. 1, 357–383.
- [Vas16c] ———, *Infinitary stability theory*, Archive for Mathematical Logic **55** (2016), 567–592.
- [Vas17a] ———, *Downward categoricity from a successor inside a good frame*, Annals of Pure and Applied Logic **168** (2017), no. 3, 651–692.
- [Vas17b] ———, *On the uniqueness property of forking in abstract elementary classes*, Mathematical Logic Quarterly **63** (2017), no. 6, 598–604.
- [Vas17c] ———, *Saturation and solvability in abstract elementary classes with amalgamation*, Archive for Mathematical Logic **56** (2017), 671–690.
- [Vas17d] ———, *Shelah's eventual categoricity conjecture in universal classes: part I*, Annals of Pure and Applied Logic **168** (2017), no. 9, 1609–1642.
- [VV17] Monica VanDieren and Sebastien Vasey, *Symmetry in abstract elementary classes with amalgamation*, Archive for Mathematical Logic **56** (2017), no. 3, 423–452.
- [Zil05] Boris Zilber, *Pseudo-exponentiation on algebraically closed fields of characteristic zero*, Annals of Pure and Applied Logic **132** (2005), 67–95.
- [Ło54] Jerzy Łoś, *On the categoricity in power of elementary deductive systems and some related problems*, Colloquium Mathematicae **3** (1954), no. 1, 58–62.

*E-mail address:* `sebv@math.harvard.edu`

*URL:* <http://math.harvard.edu/~sebv/>

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MASSACHUSETTS, USA