

SYMMETRY IN ABSTRACT ELEMENTARY CLASSES WITH AMALGAMATION

MONICA M. VANDIEREN AND SEBASTIEN VASEY

ABSTRACT. This paper is part of a program initiated by Saharon Shelah to extend the model theory of first order logic to the non-elementary setting of abstract elementary classes (AECs). An abstract elementary class is a semantic generalization of the class of models of a complete first order theory with the elementary substructure relation. We examine the symmetry property of splitting (previously isolated by the first author) in AECs with amalgamation that satisfy a local definition of superstability.

The key results are a downward transfer of symmetry and a deduction of symmetry from failure of the order property. These results are then used to prove several structural properties in categorical AECs, improving classical results of Shelah who focused on the special case of categoricity in a successor cardinal.

We also study the interaction of symmetry with tameness, a locality property for Galois (orbital) types. We show that superstability and tameness together imply symmetry. This sharpens previous work of Boney and the second author.

CONTENTS

1. Introduction	2
2. Background	6
3. Transferring symmetry	12
4. A hierarchy of symmetry properties	19
5. Symmetry from no order property	25
6. Symmetry and tameness	29

Date: February 16, 2016

AMS 2010 Subject Classification: Primary 03C48. Secondary: 03C45, 03C52, 03C55, 03C75, 03E55.

Key words and phrases. Abstract elementary classes; Categoricity; Superstability; Tameness; Symmetry; Splitting; Good frame; Limit models; Saturated models.

This material is based upon work done while the second author was supported by the Swiss National Science Foundation under Grant No. 155136.

7. Symmetry and categoricity	33
References	36

1. INTRODUCTION

The guiding conjecture for the classification of abstract elementary classes (AECs) is Shelah’s categoricity conjecture. For an introduction to AECs and Shelah’s categoricity conjecture, see [Bal09].

Although most progress towards Shelah’s categoricity conjecture has been made under the assumption that the categoricity cardinal is a successor, e.g. [She99, GV06a, Bon14b], recently, the second author has proved a categoricity transfer theorem without assuming that the categoricity cardinal is a successor, but assuming that the class is universal [Vasc, Vasd] (other partial results not assuming categoricity in a successor cardinal are in [Vasb] and [She09, Chapter IV]). In this paper, we work in a more general framework than [Vasc, Vasd]. We assume the amalgamation property and no maximal models and deduce new structural results without having to assume that the categoricity cardinal is a successor, or even has “high-enough” cofinality.

Beyond Shelah’s categoricity conjecture, a major focus in developing a classification theory for AECs has been to find an appropriate generalization of first-order superstability. Approximations isolated in [She99] and [SV99] have provided a mechanism for proving categoricity transfer results (see also [GV06a], [Vasc, Vasd]). In Chapter IV of [She09], Shelah introduced *solvability* and claims it should be the true definition of superstability in AECs (see Discussion 2.9 in the introduction to [She09]). It seems, however, that under the assumption that the class has amalgamation, a more natural definition is a version of “ $\kappa(T) = \aleph_0$ ”, first considered by Shelah and Villaveces in [SV99]. In [GV], it is shown that this definition is equivalent to many others (including solvability and the existence of a *good frame*, a local notion of independence), provided that the AEC satisfies a locality property for types called *tameness* [GV06b].

Without tameness, progress has been made in the study of structural consequences of the Shelah-Villaveces definition of superstability such as the uniqueness of limit models (e.g. [GVV]) or the property that the union of saturated models is saturated ([BVb, Vanb]). Recently in [Vana], the first author isolated a symmetry property for splitting that turns out to be closely related to the uniqueness of limit models.

1.1. Transferring symmetry. In this paper we prove a downward transfer theorem for this symmetry property. This allows us to gain insight into all of the aspects of superstability mentioned above.

Theorem 1.1. *Let \mathcal{K} be an AEC. Suppose λ and μ are cardinals so that $\lambda > \mu \geq \text{LS}(\mathcal{K})$ and \mathcal{K} is superstable in every $\chi \in [\mu, \lambda]$. Then λ -symmetry implies μ -symmetry.*

Theorem 1.1 (proven at the end of Section 3) improves Theorem 2 of [Vanb] which transfers symmetry from μ^+ to μ . We also clarify the relationship between μ -symmetry (as a property of μ -splitting) and the symmetry property in good frames (see Section 4). The latter is older and has been studied in the literature: the work of Shelah in [She01] led to [She09, Theorem 3.7], which gives conditions under which a good frame (satisfying a version of symmetry) exists (but uses set-theoretic axioms beyond ZFC and categoricity in two successive cardinals). One should also mention [She09, Theorem IV.4.10] which builds a good frame (in ZFC) from categoricity in a high-enough cardinal. Note, however, the cardinal is very high and the underlying class of the frame is a smaller class of Ehrenfeucht-Mostowski models, although this can be fixed by taking an even larger cardinal. Moreover Shelah's work contains a gap, see [BVa] for a more thorough discussion.

It was observed in [BGKV, Theorem 5.14] that Shelah's proof of symmetry of first-order forking generalizes naturally to give that the symmetry property of any reasonable global independence notion follows from the assumption of no order property. This is used in [Vasa] to build a good frame from tameness and categoricity (the results there are improved in [Vasb, BVb]). As for symmetry *transfers*, Boney [Bon14a] has shown how to transfer symmetry of a good frame *upward* using tameness for types of length two. This was later improved to tameness for types of length one with a more conceptual proof in [BVc].

Theorem 1.1 differs from these works in a few ways. First, we do not assume tameness nor set-theoretic assumptions, and we do not work within the full strength of a frame or with categoricity (only with superstability). Also, we obtain a *downward* and not an upward transfer. The arguments of this paper use *towers* whereas the aforementioned result of Boney and the second author use independent sequences to transfer symmetry upward.

1.2. Symmetry and superstability. Another consequence of our work is a better understanding of the relationship between superstability and symmetry. It was claimed in an early version of [GVV] that

μ -superstability directly implies the uniqueness of limit models of size μ but an error was later found in the proof. Here we show that this *is* true provided we have enough instances of superstability:

Theorem 5.8. *Let \mathcal{K} be an AEC and let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is superstable in all $\mu' \in [\mu, \beth_{(2^\mu)^+})$, then \mathcal{K} has μ -symmetry.*

The main idea is to imitate the proof of the aforementioned [BGKV, Theorem 5.14] to get the order property from failure of symmetry. However we do not have as much global independence as there so the proof here is quite technical.

1.3. Implications in categorical AECs. As a corollary of Theorem 5.8, we obtain several applications to categorical AECs. A notable contribution of this paper is an improvement on a 1999 result of Shelah (see [She99, Theorem 6.5]):

Fact 1.2. *Let \mathcal{K} be an AEC with amalgamation and no maximal models. Let λ and μ be cardinals such that $\text{cf}(\lambda) > \mu > \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in λ , then any limit model of size μ is saturated.*

Shelah claims in a remark immediately following his result that this can be generalized to show that for $M_0, M_1, M_2 \in \mathcal{K}_\mu$, if M_1 and M_2 are limit over M_0 , then $M_1 \cong_{M_0} M_2$ (that is, the isomorphism also fixes M_0). This is however not what his proof gives (see the discussion after Theorem 10.17 in [Bal09]). Here we finally prove this stronger statement. Moreover, we can replace the hypothesis that $\text{cf}(\lambda) > \mu$ by $\lambda \geq \beth_{(2^\mu)^+}$. That is, it is enough to ask for λ to be high-enough (but of arbitrary cofinality):

Corollary 7.3. *Let \mathcal{K} be an AEC with amalgamation and no maximal models. Let λ and μ be cardinals so that $\lambda > \mu \geq \text{LS}(\mathcal{K})$ and assume that \mathcal{K} is categorical in λ . If either $\text{cf}(\lambda) > \mu$ or $\lambda \geq \beth_{(2^\mu)^+}$, then whenever $M_0, M_1, M_2 \in \mathcal{K}_\mu$ are such that both M_1 and M_2 are limit models over M_0 , we have that $M_1 \cong_{M_0} M_2$.*

This gives a proof (assuming amalgamation, no maximal models, and a high-enough categoricity cardinal) of the (in)famous [SV99, Theorem 3.3.7], where a gap was identified in the first author's Ph.D. thesis. The gap was fixed assuming categoricity in μ^+ in [Van06, Van13] (see also the exposition in [GVV]). In [BG, Corollary 6.10], this was improved to categoricity in an arbitrary $\lambda > \mu$ provided that μ is big-enough and the class satisfies strong locality assumptions (full tameness and shortness and the extension property for coheir). In [Vas, Theorem 7.11], only tameness was required but the categoricity had to be in a λ

with $\text{cf}(\lambda) > \mu$. Still assuming tameness, this is shown for categoricity in any $\lambda \geq \beth_{(2^\mu)^+}$ in [BVb, Theorem 7.1]. Here assuming tameness we will improve this to categoricity in any $\lambda > \text{LS}(\mathcal{K})$ (see Corollary 7.10).

In general, we obtain that an AEC with amalgamation categorical in a high-enough cardinal has several structural properties that were previously only known for AECs categorical in a cardinal of high-enough *cofinality*, or even just in a successor.

Corollary 1.3. *Let \mathcal{K} be an AEC with amalgamation. Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$ and assume that \mathcal{K} is categorical in λ . Let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in a $\lambda > \mu$, then:*

- (1) (see Corollary 7.4) *If $\lambda \geq \beth_{(2^\mu)^+}$ and $\mu > \text{LS}(\mathcal{K})$, then the model of size λ is μ -saturated.*
- (2) (see Corollary 7.9) *If $\mu \geq \beth_{(2^{\text{LS}(\mathcal{K})})^+}$ and the model of size λ is μ^+ -saturated, then there exists a type-full good μ -frame with underlying class the saturated models in \mathcal{K}_μ .*

This improves several classical results from Shelah's milestone study of categorical AECs with amalgamation [She99]:

- Corollary 1.3.(1) partially answers Baldwin [Bal09, Problem D.1.(2)] which asked if in any AEC with amalgamation categorical in a high-enough cardinal, then model in the categoricity cardinal is saturated.
- Corollary 1.3.(2) partially answers the question in [She99, Remark 4.9.(1)] of whether there is a parallel to forking in categorical AECs with amalgamation. It also improves on [Vasb, Theorem 7.4], which assumed categoricity in a successor (and a higher Hanf number bound).
- As part of the proof of Corollary 1.3.(2), we derive *weak tameness* (i.e. tameness over saturated models) from categoricity in a big-enough cardinal (this is Corollary 7.5). It was previously only known how to do so assuming that the categoricity cardinal has high-enough cofinality [She99, Main Claim II.2.3].

We deduce a downward categoricity transfer in AECs with amalgamation (see also Corollary 7.7):

Corollary 7.8. *Let \mathcal{K} be an AEC with amalgamation. Let $\text{LS}(\mathcal{K}) < \mu = \beth_\mu < \lambda$. If \mathcal{K} is categorical in λ , then \mathcal{K} is categorical in μ .*

This improves on [Vasb, Theorem 10.16] where the result is stated with the additional assumption of $(< \mu)$ -tameness.

1.4. Implications in tame AECs. This paper also combines our results with tameness: in Section 6, we improve Hanf number bounds for several consequences of superstability. With Will Boney, the second author has shown [BVb, Theorem 7.1] that μ -superstability and μ -tameness imply that for all high-enough λ , limit models of size λ are unique (in the strong sense discussed above), unions of chains of λ -saturated models are saturated, and there exists a type-full good λ -frame. We transfer this behavior downward using our symmetry transfer theorem to get that the latter result is actually true starting from $\lambda = \mu^+$, and the former starting from $\lambda = \mu$:

Corollary 1.4. *Let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is μ -superstable and μ -tame, then:*

- (1) (see Corollary 6.9) *If $M_0, M_1, M_2 \in \mathcal{K}_\mu$ are such that both M_1 and M_2 are limit models over M_0 , then $M_1 \cong_{M_0} M_2$.*
- (2) (see Corollary 6.10) *For any $\lambda > \mu$, the union of an increasing chain of λ -saturated models is λ -saturated.*
- (3) (see Corollary 6.14) *There exists a type-full good μ^+ -frame with underlying class the saturated models in \mathcal{K}_{μ^+} .*

In fact, μ -tameness along with μ -superstability already implies μ -symmetry. Many assumptions weaker than tameness (such as the existence of a good μ^+ -frame, see Theorem 4.15) suffice to obtain such a conclusion.

1.5. Notes. A word on the background needed to read this paper: It is assumed that the reader has a solid knowledge of AECs (including the material in [Bal09]). Some familiarity with good frames, in particular the material of [Vasa] would be very helpful. In addition to classical results, e.g. in [She99], the paper uses heavily the results of [Vana, Vanb] on limit models and the symmetry property of splitting. It also relies on the construction of a good frame in [Vasa]. At one point we also use the canonicity theorem for good frames [Vasb, Theorem 9.7].

This paper was written while the second author was working on a Ph.D. thesis under the direction of Rami Grossberg at Carnegie Mellon University and he would like to thank Professor Grossberg for his guidance and assistance in his research in general and in this work specifically. We also thank a referee for a report which helped improve the presentation of this paper.

2. BACKGROUND

All throughout this paper, we assume the amalgamation property:

Hypothesis 2.1. \mathcal{K} is an AEC with amalgamation.

For convenience, we fix a big-enough monster model \mathfrak{C} and work inside \mathfrak{C} . This is possible since by Remark 2.9, we will have the joint embedding property in addition to the amalgamation property for models of the relevant cardinalities. At some point, we will also use the following fact whose proof is folklore (see e.g. [Vasb, Proposition 10.13])

Fact 2.2. *Assume that \mathcal{K} has joint embedding in some $\lambda \geq \text{LS}(\mathcal{K})$. Then there exists $\chi < \beth_{(2^{\text{LS}(\mathcal{K})})^+}$ and an AEC \mathcal{K}^* such that:*

- (1) $\mathcal{K}^* \subseteq \mathcal{K}$ and \mathcal{K}^* has the same strong substructure relation as \mathcal{K} .
- (2) $\text{LS}(\mathcal{K}^*) = \text{LS}(\mathcal{K})$.
- (3) \mathcal{K}^* has amalgamation, joint embedding, and no maximal models.
- (4) $\mathcal{K}_{\geq \min(\lambda, \chi)} = (\mathcal{K}^*)_{\geq \min(\lambda, \chi)}$.

Many of the pre-requisite definitions and notations used in this paper can be found in [GVV]. Here we recall the more specialized concepts that we will be using explicitly.

We write $\text{ga-tp}(\bar{a}/M)$ for the Galois type of the sequence \bar{a} over M (and we write $\text{ga-S}(M)$ for the set of all Galois types over M). While the reader can think of $\text{ga-tp}(\bar{a}/M)$ as the orbit of \bar{a} under the action of $\text{Aut}_M(\mathfrak{C})$, $\text{ga-tp}(\bar{a}/M)$ is really defined as the equivalence class of the triple $(\bar{a}, M, \mathfrak{C})$ under a certain equivalence relation (see for example [Gro02, Definition 6.4]). This allows us to define the restriction of a Galois type to any strong substructure of its domain, as well as its image under any automorphism of \mathfrak{C} (and by extension any \mathcal{K} -embedding whose domain contains the domain of the type).

With that remark in mind, we can state the definition of non-splitting, a notion of independence from [She99, Definition 3.2].

Definition 2.3. *A type $p \in \text{ga-S}(N)$ does not μ -split over M if and only if for any $N_1, N_2 \in \mathcal{K}_\mu$ such that $M \leq N_\ell \leq N$ for $\ell = 1, 2$, and any $f : N_1 \cong_M N_2$, we have $f(p \upharpoonright N_1) = p \upharpoonright N_2$*

We will use the definition of universality from [Van06, Definition I.2.1]:

Definition 2.4. *Let $M, N \in \mathcal{K}$ be such that $M \leq N$. We say that N is μ -universal over M if for any $M' \geq M$ with $\|M'\| \leq \mu$, there exists $f : M' \xrightarrow[M]{\rightarrow} N$. We say that N is universal over M if N is $\|M\|$ -universal over M .*

A fundamental concept in the study of superstable AECs is the notion of a *limit model*, first introduced in [She99]. We only give the definition here and refer the reader to [GVV] for more history and motivation.

Definition 2.5. *Let $\mu \geq \text{LS}(\mathcal{K})$ and let $\alpha < \mu^+$ be a limit ordinal. Let $M \in \mathcal{K}_\mu$. We say that N is (μ, α) -limit over M (or a (μ, α) -limit model over M) if there exists a strictly increasing continuous chain $\langle M_i : i \leq \alpha \rangle$ in \mathcal{K}_μ such that $M_0 = M$, $M_\alpha = N$, and M_{i+1} is universal over M_i for all $i < \alpha$. We say that N is limit over M (or a limit model over M) if it is (μ, β) -limit over M for some $\beta < \mu^+$. Finally, we say that N is limit if it is limit over N_0 for some $N_0 \in \mathcal{K}_{\|N\|}$.*

Towers were introduced in Shelah and Villaveces [SV99] as a tool to prove the uniqueness of limit models. A tower is an increasing sequence of length α of limit models, denoted by $\bar{M} = \langle M_i \in \mathcal{K}_\mu \mid i < \alpha \rangle$, along with a sequence of designated elements $\bar{a} = \langle a_i \in M_{i+1} \setminus M_i \mid i + 1 < \alpha \rangle$ and a sequence of designated submodels $\bar{N} = \langle N_i \mid i + 1 < \alpha \rangle$ for which $N_i \leq M_i$, $\text{ga-tp}(a_i/M_i)$ does not μ -split over N_i , and M_i is universal over N_i (see Definition I.5.1 of [Van06]).

Now we recall a bit of terminology regarding towers. The collection of all towers $(\bar{M}, \bar{a}, \bar{N})$ made up of models of cardinality μ and sequences indexed by α is denoted by $\mathcal{K}_{\mu, \alpha}^*$. For $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \alpha}^*$, if $\beta < \alpha$ then we write $(\bar{M}, \bar{a}, \bar{N}) \upharpoonright \beta$ for the tower made of the subsequences $\bar{M} \upharpoonright \beta = \langle M_i \mid i < \beta \rangle$, $\bar{a} \upharpoonright \beta = \langle a_i \mid i + 1 < \beta \rangle$, and $\bar{N} \upharpoonright \beta = \langle N_i \mid i + 1 < \beta \rangle$. We sometimes abbreviate the tower $(\bar{M}, \bar{a}, \bar{N})$ by \mathcal{T} .

Definition 2.6. *For towers $(\bar{M}, \bar{a}, \bar{N})$ and $(\bar{M}', \bar{a}', \bar{N}')$ in $\mathcal{K}_{\mu, \alpha}^*$, we say*

$$(\bar{M}, \bar{a}, \bar{N}) \leq (\bar{M}', \bar{a}', \bar{N}')$$

if for all $i < \alpha$, $M_i \leq M'_i$, $\bar{a} = \bar{a}'$, $\bar{N} = \bar{N}'$ and whenever M'_i is a proper extension of M_i , then M'_i is universal over M_i . If for each $i < \alpha$, M'_i is universal over M_i we will write $(\bar{M}, \bar{a}, \bar{N}) < (\bar{M}', \bar{a}', \bar{N}')$.

In order to transfer symmetry from λ to μ we will need to consider a generalization of these towers where the models M_i and N_i may have different cardinalities. Fix $\lambda \geq \mu \geq \text{LS}(\mathcal{K})$ and α a limit ordinal $< \mu^+$. We will write $\mathcal{K}_{\lambda, \alpha, \mu}^*$ for the collection of towers of the form $(\bar{M}, \bar{a}, \bar{N})$ where $\bar{M} = \langle M_i \mid i < \alpha \rangle$ is a sequence of models each of cardinality λ and $\bar{N} = \langle N_i \mid i + 1 < \alpha \rangle$ is a sequence of models of cardinality μ . We require that for $i < \alpha$, M_i is μ -universal over N_i and $\text{ga-tp}(a_i/M_i)$ does not μ -split over N_i .

In a natural way we order these towers by the following adaptation of Definition 2.6.

Definition 2.7. Let $\lambda \geq \chi \geq \mu \geq \text{LS}(\mathcal{K})$ be cardinals and fix $\alpha < \mu^+$ an ordinal. For towers $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda, \alpha, \mu}^*$ and $(\bar{M}', \bar{a}', \bar{N}') \in \mathcal{K}_{\chi, \alpha, \mu}^*$, we say

$$(\bar{M}, \bar{a}, \bar{N}) <_\mu (\bar{M}', \bar{a}', \bar{N}')$$

if for all $i < \alpha$, $M_i \leq M'_i$, $\bar{a} = \bar{a}'$, $\bar{N} = \bar{N}'$, and there is $\theta < \lambda^+$ so that M'_i is a (λ, θ) -limit model witnessed by a sequence $\langle M_i^{\gamma'} \mid \gamma' < \theta \rangle$ with $M_i < M_0^{\gamma'}$.

Note that Definition 2.6 is defined only on towers in $\mathcal{K}_{\mu, \alpha}^*$ and is slightly weaker from the ordering $<_\mu$ when restricted to $\mathcal{K}_{\mu, \alpha}^*$. In particular, the models M'_i in the tower $(\bar{M}', \bar{a}, \bar{N}) <_\mu (\bar{M}, \bar{a}, \bar{N})$ are only required to be universal over M_i and limit. It is not necessary that M'_i is limit over M_i as we require if $(\bar{M}', \bar{a}, \bar{N}) <_\mu (\bar{M}, \bar{a}, \bar{N})$.

Towers are particularly suited for superstable abstract elementary classes, in which they are known to exist and in which the union of an increasing chain of towers will be a tower. The definition below is already implicit in [SV99] and has since then been studied in many papers, e.g. [Van06, GVV, Vasb, BVb, GV]. We will use the definition from [Vasb, Definition 10.1]:

Definition 2.8. \mathcal{K} is μ -superstable (or superstable in μ) if:

- (1) $\mu \geq \text{LS}(\mathcal{K})$.
- (2) \mathcal{K}_μ is nonempty, has joint embedding, and no maximal models.
- (3) \mathcal{K} is stable in μ . That is, $|\text{ga-S}(M)| \leq \mu$ for all $M \in \mathcal{K}_\mu$. Some authors call this “Galois-stable,” and:
- (4) μ -splitting in \mathcal{K} satisfies the “no long splitting chains” property:
For any limit ordinal $\alpha < \mu^+$, for every sequence $\langle M_i \mid i < \alpha \rangle$ of models of cardinality μ with M_{i+1} universal over M_i and for every $p \in \text{ga-S}(\bigcup_{i < \alpha} M_i)$, there exists $i < \alpha$ such that p does not μ -split over M_i .

Remark 2.9. By our global hypothesis of amalgamation (Hypothesis 2.1), if \mathcal{K} is μ -superstable, then $\mathcal{K}_{\geq \mu}$ has joint embedding.

Remark 2.10. By the weak transitivity property of μ -splitting [Vasb, Proposition 3.7], μ -superstability implies the following continuity property (which is sometimes also stated as part of the definition): For any limit ordinal $\alpha < \mu^+$, for every sequence $\langle M_i \mid i < \alpha \rangle$ of models of cardinality μ with M_{i+1} universal over M_i and for every $p \in \text{ga-S}(\bigcup_{i < \alpha} M_i)$, if $p \restriction M_i$ does not μ -split over M_0 for all $i < \alpha$, then p does not μ -split over M_0 . We will use this freely.

Proposition 2.11. *Let $\mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K}_μ has joint embedding and \mathcal{K} is stable in μ . The following are equivalent:*

- (1) \mathcal{K} has a model of size μ^+ .
- (2) \mathcal{K}_μ is nonempty and has no maximal models.
- (3) \mathcal{K} has a limit model of size μ .
- (4) There exists $M_0, M_1, M_2 \in \mathcal{K}_\mu$ such that $M_0 < M_1 < M_2$ and M_1 is universal over M_0 .

Proof. (1) implies (2) is by joint embedding. (2) implies (3) implies (4) is straightforward. We show (4) implies (1). Assume $M_0 < M_1 < M_2$ and M_1 is universal over M_0 . It is enough to show that M_2 has a proper extension. By universality, there exists $f : M_2 \xrightarrow{M_0} M_1$. Now extend f to $g : M'_2 \cong_{M_0} M_2$. Since $M_1 < M_2$, $M_2 < M'_2$, as desired. \square

The main results of this paper involve the concept of symmetry over limit models and its equivalent formulation involving towers which was identified in [Vana]:

Definition 2.12. *We say that an abstract elementary class exhibits symmetry for non- μ -splitting if whenever models $M, M_0, N \in \mathcal{K}_\mu$ and elements a and b satisfy the conditions 1-4 below, then there exists M^b a limit model over M_0 , containing b , so that $\text{ga-tp}(a/M^b)$ does not μ -split over N . See Figure 1.*

- (1) M is universal over M_0 and M_0 is a limit model over N .
- (2) $a \in M \setminus M_0$.
- (3) $\text{ga-tp}(a/M_0)$ is non-algebraic and does not μ -split over N .
- (4) $\text{ga-tp}(b/M)$ is non-algebraic and does not μ -split over M_0 .

We end by recalling a few results of the first author showing the importance of the symmetry property:

Fact 2.13 (Theorem 4 in [Vana]). *If \mathcal{K} is μ -superstable and the union of any chain (of length less than μ^{++}) of saturated models of size μ^+ is saturated, then \mathcal{K} has μ -symmetry.*

Many of the results on symmetry rely on the equivalent formulation of μ -symmetry in terms of reduced towers.

Definition 2.14. *A tower $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \alpha}^*$ is reduced if it satisfies the condition that for every $<$ -extension $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \alpha}^*$ of $(\bar{M}, \bar{a}, \bar{N})$ and for every $i < \alpha$, $M'_i \cap (\bigcup_{j < \alpha} M_j) = M_i$.*




FIGURE 1. A diagram of the models and elements in the definition of symmetry. We assume the type $\text{ga-tp}(b/M)$ does not μ -split over M_0 and $\text{ga-tp}(a/M_0)$ does not μ -split over N . Symmetry implies the existence of M^b a limit model over M_0 containing b , so that $\text{ga-tp}(a/M^b)$ does not μ -split over N .

Definition 2.15. A tower $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \alpha}^*$ is continuous if for any limit $i < \alpha$, $M_i = \bigcup_{j < i} M_j$.

Fact 2.16 (Theorem 5 in [Vana]). Assume \mathcal{K} is μ -superstable. The following are equivalent:

- (1) \mathcal{K} has μ -symmetry.
- (2) Any reduced tower in $\mathcal{K}_{\mu, \alpha}^*$ is continuous.

It was previously established (in [SV99] or more explicitly in [GVV]) that the continuity of reduced towers gives uniqueness of limit models:

Fact 2.17. Assume \mathcal{K} is μ -superstable. If any reduced tower in $\mathcal{K}_{\mu, \alpha}^*$ is continuous (or equivalently by Fact 2.16 if \mathcal{K} has μ -symmetry), then for any $M_0, M_1, M_2 \in \mathcal{K}_\mu$, if M_1 and M_2 are limit over M_0 , then $M_1 \cong_{M_0} M_2$.

Symmetry also has implications to chains of saturated models. For $\lambda > \text{LS}(\mathcal{K})$, write $\mathcal{K}^{\lambda\text{-sat}}$ for the class of λ -saturated models in $\mathcal{K}_{\geq \lambda}$. We also define $\mathcal{K}^{0\text{-sat}} := \mathcal{K}$. Using this notation, we have:

Fact 2.18 (Theorem 22 in [Vanb]). Assume \mathcal{K} is μ -superstable, μ^+ -superstable, and every limit model in \mathcal{K}_{μ^+} is saturated. Then $\mathcal{K}^{\mu^+\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\mu^+\text{-sat}}) = \mu^+$.

Remark 2.19. By Fact 2.17, the hypotheses of Fact 2.18 hold if \mathcal{K} is μ -superstable, μ^+ -superstable, and has μ^+ -symmetry.

We will also use the following easy lemma:

Lemma 2.20. *Let λ be a limit cardinal and let $\lambda_0 < \lambda$. Assume that for all $\mu \in [\lambda_0, \lambda)$, $\mathcal{K}^{\mu\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\mu\text{-sat}}) = \mu$. Then $\mathcal{K}^{\lambda\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\lambda\text{-sat}}) = \lambda$.*

Proof. That $\mathcal{K}^{\lambda\text{-sat}}$ is closed under chains is easy to check. To see $\text{LS}(\mathcal{K}^{\lambda\text{-sat}}) = \lambda$, let $M \in \mathcal{K}^{\lambda\text{-sat}}$ and let $A \subseteq |M|$. Without loss of generality, $\chi := |A| \geq \lambda$. Let $\delta := \text{cf}(\lambda)$ and let $\langle \lambda_i : i < \delta \rangle$ be an increasing sequence of cardinals with limit λ . Build $\langle M_i : i \leq \delta \rangle$ increasing continuous in \mathcal{K}_χ such that for all $i < \delta$, M_{i+1} is λ_i^+ -saturated and $A \subseteq |M_0|$. This is possible by assumption. Then M_δ is λ_i^+ -saturated for all $i < \delta$, hence is λ -saturated. Thus it is as needed. \square

3. TRANSFERRING SYMMETRY

In this section we prove Theorem 1.1 which is key to the results in the following sections. We start with a few observations which will allow us to extend the tower machinery from [GVV] and [Vanb] to include towers composed of models of different cardinalities. In particular, we derive an extension property for towers of different cardinalities, Lemma 3.7. This will allow us to adapt the arguments from [Vanb] to prove Theorem 1.1.

We start with a study of chains where each model indexed by a successor is universal over its predecessor:

Proposition 3.1. *Suppose that $\lambda \geq \text{LS}(\mathcal{K})$ is a cardinal. Assume that \mathcal{K} is stable in λ with no maximal models of cardinality λ . Let θ be a limit ordinal. Assume $\langle M_i \in \mathcal{K}_{\geq \text{LS}(\mathcal{K})} \mid i < \theta \rangle$ is a strictly increasing and continuous sequence of models so that for all $i < \theta$, M_{i+1} is universal over M_i . If $M := \bigcup_{i < \theta} M_i$ has size λ , then M is a (λ, θ) -limit model over some model containing M_0 .*

Proof. By cardinality considerations, $\theta < \lambda^+$. Replacing θ by $\text{cf}(\theta)$ if necessary, we can assume without loss of generality that θ is regular. By λ -stability and the assumption that \mathcal{K} has no maximal models of cardinality λ , we can fix a (λ, θ) -limit model M^* witnessed by $\langle M_i^* \mid i < \theta \rangle$ with $M_0 \leq M_0^*$. If there exists $i < \theta$ such that $M_i \in \mathcal{K}_\lambda$, then the sequence $\langle M_j \mid j \in [i, \theta) \rangle$ witnesses that M is (λ, θ) -limit and $M_0 \leq M_i$; so assume that $\lambda > \text{LS}(\mathcal{K})$ and $M_i \in \mathcal{K}_{< \lambda}$ for all $i < \theta$. Then we must have that $\theta = \text{cf}(\lambda)$. If λ is a successor, we must have that $\theta = \lambda$ and we obtain the result from [Vanb, Proposition 14]; so assume λ is limit. For $i < \theta$, let $\lambda_i := \|M_i\|$.

Fix $\langle a_\alpha \mid \alpha < \lambda \rangle$ an enumeration of M^* . Using the facts that M_{i+1} is universal over M_i and that M_{i+1}^* is universal over M_i^* , we can build an isomorphism $f : M \cong M^*$ inductively by defining an increasing and continuous sequence of \mathcal{K} -embeddings f_i so that $f_i : M_i \rightarrow M_i^*$, $f_0 = \text{id}_{M_0}$, and $\{a_\alpha \mid \alpha < \lambda_i\} \subseteq \text{rg}(f_{i+1})$. \square

We will use the following generalization of the weak transitivity property of μ -splitting proven in [Vas, Proposition 3.7]. The difference here is that the models are allowed to be of size bigger than μ .

Proposition 3.2. *Let $\mu \geq \text{LS}(\mathcal{K})$ be such that \mathcal{K} is stable in μ . Let $M_0 \leq M_1 < M'_1 \leq M_2$ all be in $\mathcal{K}_{\geq \mu}$. Assume that M'_1 is universal over M_1 . Let $p \in \text{ga-S}(M_2)$. If $p \upharpoonright M'_1$ does not μ -split over M_0 and p does not μ -split over some $N \in \mathcal{K}_\mu$ with $N \leq M_1$, then p does not μ -split over M_0 .*

Proof. Note that by definition of μ -splitting, $M_0 \in \mathcal{K}_\mu$. Thus by making N larger if necessary we can assume that $M_0 \leq N$. By basic properties of universality we have that M'_1 is universal over N , hence without loss of generality $M_1 = N$. In particular, $M_1 \in \mathcal{K}_\mu$. By stability, build $M''_1 \in \mathcal{K}_\mu$ universal over M_1 such that $M_1 < M''_1 \leq M'_1$. By monotonicity, $p \upharpoonright M''_1$ does not μ -split over M_0 . Thus without loss of generality also $M'_1 \in \mathcal{K}_\mu$. By definition of μ -splitting, it is enough to check that $p \upharpoonright M'_2$ does not μ -split over M_0 for all $M'_2 \in \mathcal{K}_\mu$ with $M'_2 \leq M_2$. Thus without loss of generality again $M_2 \in \mathcal{K}_\mu$. Now use the weak transitivity property of μ -splitting [Vas, Proposition 3.7]. \square

We use the previous proposition to extend the continuity property of μ -splitting to models of size bigger than μ . This is very similar to the argument in [She09, Claim II.2.11].

Proposition 3.3. *Let $\mu \geq \text{LS}(\mathcal{K})$ and assume that \mathcal{K} is μ -superstable.*

Suppose $\langle M_i \in \mathcal{K}_{\geq \mu} \mid i < \delta \rangle$ is an increasing sequence of models so that, for all $i < \delta$, M_{i+1} is universal over M_i . Let $p \in \text{ga-S}(\bigcup_{i < \delta} M_i)$. If $p \upharpoonright M_i$ does not μ -split over M_0 for each $i < \delta$, then p does not μ -split over M_0 .

Proof. Without loss of generality, $\delta = \text{cf}(\delta)$. Let $M_\delta := \bigcup_{i < \delta} M_i$. There are two cases to check. If $\delta > \mu$, then by [She99, Claim 3.3], there exists $N \in \mathcal{K}_\mu$ with $N \leq M_\delta$ such that p does not μ -split over N . Pick $i < \delta$ such that $N \leq M_i$. Then p does not μ -split over M_i . By Proposition 3.2 (where $(M_0, M_1 M'_1, M_2, N)$ there stand for $(M_0, M_i, M_{i+1}, M_\delta, N)$ here), p does not μ -split over M_0 .

Suppose then that $\delta \leq \mu$ and for sake of contradiction that M^* of cardinality μ witnesses the splitting of p over M_0 , i.e. $p \restriction M^*$ μ -splits over M_0 . We can find $\langle M_i^* \in \mathcal{K}_\mu \mid i < \delta \rangle$ an increasing resolution of M^* so that $M_i^* \leq M_i$ for all $i < \delta$. By monotonicity of splitting, stability in μ , and the fact that each M_{i+1} is universal over M_i , we can increase M^* , if necessary, to arrange that M_{i+1}^* is universal over M_i^* . Since $p \restriction M_i$ does not μ -split over M_0 , monotonicity of non-splitting implies that $p \restriction M_i^*$ does not μ -split over M_0 . Then, by μ -superstability $p \restriction M^*$ does not μ -split over M_0 . This contradicts our choice of M^* . \square

We adapt the proof of the extension property for non- μ -splitting ([Van06, Theorem I.4.10]) to handle models of different sizes under the additional assumption of superstability in the size of the bigger model. The conclusion can also be achieved using the assumption of tameness instead of superstability (since μ -splitting and λ -splitting coincide if \mathcal{K} is μ -tame and $\mu \leq \lambda$, see [BGKV, Proposition 3.12]).

Proposition 3.4. *Fix cardinals $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Suppose that \mathcal{K} is μ -stable and λ -superstable.*

Let $M \in \mathcal{K}_\mu$ and $M^\lambda, M' \in \mathcal{K}_\lambda$ be such that $M \leq M^\lambda \leq M'$ and M^λ is limit over some model containing M . Let $p \in \text{ga-S}(M^\lambda)$ be such that p does not μ -split over M . Then there exists $q \in \text{ga-S}(M')$ extending p so that q does not μ -split over M . Moreover q is algebraic if and only if p is.

Proof. Let $\theta < \lambda^+$ and $\langle M_i^\lambda \mid i < \theta \rangle$ witness that M^λ is (λ, θ) -limit with $M \leq M_0^\lambda$. Write $p := \text{ga-tp}(a/M^\lambda)$. By λ -superstability there exists $i < \theta$ so that p does not λ -split over M_i^λ . Since M_{i+2}^λ is universal over M_{i+1}^λ there exists $f : M' \xrightarrow{M_{i+1}^\lambda} M_{i+2}^\lambda$. Extend f to $g \in \text{Aut}_{M_{i+1}^\lambda}(\mathfrak{C})$.

Let $q := g^{-1}(p) \restriction M' = \text{ga-tp}(g^{-1}(a)/M')$. Note that q is nonalgebraic if p is nonalgebraic (the converse will follow once we have shown that q extends p). By monotonicity, invariance, and our assumption that p does not μ -split over M , we can conclude that q does not μ -split over N . By similar reasoning also q does not λ -split over M_i^λ . In particular $\text{ga-tp}(g^{-1}(a)/M^\lambda) = q \restriction M^\lambda$ does not λ -split over M_i^λ . Since g fixes M_{i+1}^λ , we know that $g^{-1}(a)$ realizes $p \restriction M_{i+1}^\lambda$. Therefore, we get by the uniqueness of non- λ -splitting extensions that $q \restriction M^\lambda = \text{ga-tp}(f^{-1}(a)/M^\lambda) = \text{ga-tp}(a/M^\lambda) = p$. This shows that q extends p , as desired. \square

We can now prove an extension property for towers in $\mathcal{K}_{\lambda,\alpha,\mu}^*$.

Lemma 3.5. *Let λ and μ be cardinals satisfying $\lambda \geq \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in μ and in λ . For any $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda,\alpha,\mu}^*$, there exists $(\bar{M}', \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda,\alpha,\mu}^*$ so that:*

$$(\bar{M}, \bar{a}, \bar{N}) <_{\mu} (\bar{M}', \bar{a}, \bar{N})$$

Proof. If $\lambda = \mu$, the result follows from infinitely many (for example $\text{cf}(\lambda)$ many) applications of [GVV, Lemma 5.3] which is the extension property for towers. If $\lambda > \mu$, the result follows similarly from the proof of the extension property for towers using Proposition 3.4. \square

We also have a continuity property:

Lemma 3.6. *Let $\mu \geq \text{LS}(\mathcal{K})$ be such that \mathcal{K} is μ -superstable. Let $\langle \lambda_i : i < \delta \rangle$ be an increasing sequence of cardinals with $\lambda_0 \geq \mu$. Let $\langle (\bar{M}^i, \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda_i,\alpha,\mu}^* \mid i < \delta \rangle$ be a sequence of towers such that $(\bar{M}^i, \bar{a}, \bar{N}) <_{\mu} (\bar{M}^{i+1}, \bar{a}, \bar{N})$ for all $i < \delta$.*

Let \bar{M}^{δ} be the sequence composed of models of the form $M_{\beta}^{\delta} := \bigcup_{i < \delta} M_{\beta}^i$ for $\beta < \alpha$. Let $\lambda := \sum_{i < \delta} \lambda_i$.

Then $(\bar{M}^{\delta}, \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda,\alpha,\mu}^$ and $(\bar{M}^i, \bar{a}, \bar{N}) <_{\mu} (\bar{M}^{\delta}, \bar{a}, \bar{N})$ for all $i < \delta$.*

Proof. Working by induction on δ , we can assume without loss of generality that the sequence of tower is continuous. That is, for each $\beta < \alpha$ and limit $i < \delta$, $M_{\beta}^i = \bigcup_{j < i} M_{\beta}^j$. Of course, it is enough to show that $(\bar{M}^0, \bar{a}, \bar{N}) <_{\mu} (\bar{M}^{\delta}, \bar{a}, \bar{N})$. Let $\beta < \alpha$. There are two things to check: M_{β}^{λ} is a limit model over a model that contains M_{β}^0 , and $\text{ga-tp}(a_{\beta}/M_{\beta}^{\lambda})$ does not μ -split over N_{β} . Proposition 3.1 confirms that M_{β}^{λ} is a (λ, δ) -limit model over some model containing M_{β}^0 . Because each $(\bar{M}^i, \bar{a}, \bar{N})$ is a tower, we know that $\text{ga-tp}(a_{\beta}/M_{\beta}^i)$ does not μ -split over N_{β} . This allows us to apply Proposition 3.3 to conclude that $\text{ga-tp}(a_{\beta}/M_{\beta}^{\lambda})$ does not μ -split over N_{β} . \square

We conclude an extension property for towers of different sizes

Lemma 3.7. *Let κ , λ and μ be cardinals satisfying $\lambda \geq \kappa \geq \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in μ and in every $\chi \in [\kappa, \max(\kappa^+, \lambda))$*

Let $(\bar{M}^{\kappa}, \bar{a}, \bar{N}) \in \mathcal{K}_{\kappa,\alpha,\mu}^$.*

(1) *There exists $(\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda,\alpha,\mu}^*$ so that*

$$(\bar{M}^{\kappa}, \bar{a}, \bar{N}) <_{\mu} (\bar{M}, \bar{a}, \bar{N}).$$

- (2) If in addition \mathcal{K} is λ -superstable, then there exists a sequence $\langle N_\beta^\lambda \mid \beta < \alpha \rangle$ so that $N_i \leq N_\beta^\lambda$ for all $\beta < \alpha$ and $(\bar{M}, \bar{a}, \bar{N}^\lambda) \in \mathcal{K}_{\lambda, \alpha}^*$.

Proof. We prove the first statement in the lemma by induction on λ . If $\lambda = \kappa$, this is given by Lemma 3.5. Now assume that $\lambda > \kappa$. Fix an increasing continuous sequence $\langle \lambda_i \mid i < \text{cf}(\lambda) \rangle$ which is cofinal in λ and so that $\lambda_0 = \kappa$ (if $\lambda = \chi^+$ is a successor we can take $\lambda_i = \chi$ for all $i < \lambda$). We build a sequence $\langle (\bar{M}^{\lambda_i}, \bar{a}, \bar{N}) \in \mathcal{K}_{\lambda_i, \alpha, \mu}^* \mid i < \text{cf}(\lambda) \rangle$ which is increasing (that is, $(\bar{M}^{\lambda_i}, \bar{a}, \bar{N}) <_\mu (\bar{M}^{\lambda_{i+1}}, \bar{a}, \bar{N})$ for all $i < \text{cf}(\lambda)$) and continuous (in the obvious sense, see Lemma 3.6). This is possible by the induction hypothesis. Now by Lemma 3.6, the union of the chain of towers (defined there) is as desired.

For part (2), recall from Definition 2.7 that for each $\beta < \alpha$, M_β is a limit model over some model containing M_β^κ . Let $\langle M_{\beta, i}^* \in \mathcal{K}_\lambda \mid i < \theta_\beta \rangle$ witness this. By λ -superstability, for each $\beta < \alpha$, there exists $i_\beta < \theta_\beta$ so that $\text{ga-tp}(a_\beta/M_\beta^\lambda)$ does not λ -split over M_{β, i_β}^* . By our choice of $M_{\beta, 0}^*$ containing M_β^κ , and consequently N_β^κ , we can take $N_\beta^\lambda := M_{\beta, i_\beta}^*$. \square

We now begin the proof of Theorem 1.1. The structure of the proof is similar to the proof of Theorem 2 of [Vanb]; only here we work with towers in $\mathcal{K}_{\lambda, \alpha, \mu}^*$ as opposed to only towers in $\mathcal{K}_{\mu, \alpha}^*$.

Proof of Theorem 1.1. Suppose for the sake of contradiction that \mathcal{K} does not have symmetry for μ -non-splitting. By Fact 2.16 and our μ -superstability assumption, \mathcal{K} has a reduced discontinuous tower in $\mathcal{K}_{\mu, \alpha}^*$ for some $\alpha < \mu^+$. Let α be the minimal ordinal for which there is a reduced, discontinuous tower in $\mathcal{K}_{\mu, \alpha}^*$. By Lemma 5.7 of [GVV], we may assume that $\alpha = \delta + 1$ for some limit ordinal δ . Fix $\mathcal{T} = (\bar{M}, \bar{a}, \bar{N}) \in \mathcal{K}_{\mu, \alpha}^*$ a reduced discontinuous tower with $b \in M_\delta \setminus \bigcup_{\beta < \alpha} M_\beta$.

Let $I := \text{cf}(\lambda)$. By Lemma 3.7, we can build an increasing and continuous chain of towers $\langle \mathcal{T}^i \mid i \in I \rangle$ extending $\mathcal{T} \upharpoonright \delta$. If $\lambda = \kappa^+$ for some κ , then select each $\mathcal{T}^i \in \mathcal{K}_{\kappa, \delta, \mu}^*$. If λ is a limit cardinal, fix $\langle \lambda_i \mid i < \text{cf}(\lambda) \rangle$ to be an increasing and continuous sequence of cardinals cofinal in λ , with $\lambda_0 > \mu$ and choose $\mathcal{T}^i \in \mathcal{K}_{\lambda_i, \delta, \mu}^*$. Let $\mathcal{T}^\lambda := \bigcup_{i \in I} \mathcal{T}^i$.

Notice that by Lemma 3.6, and our assumptions on the towers \mathcal{T}^i , we can conclude that $\mathcal{T}^\lambda \in \mathcal{K}_{\lambda, \delta, \mu}^*$ and \mathcal{T}^λ extends $\mathcal{T} \upharpoonright \delta$. In particular, for each $\beta < \alpha$,

- (1) $\text{ga-tp}(a_\beta/M_\beta^\lambda)$ does not μ -split over N_β .

Furthermore by the second part of Lemma 3.7 we can find N_β^λ so that the tower defined by $(\bar{M}^\lambda, \bar{a}, \bar{N}^\lambda)$ is in $\mathcal{K}_{\lambda,\delta}^*$ and each M_β^λ is a limit over N_β^λ . We can extend this to a tower of length $\delta + 1$ by appending to $(\bar{M}^\lambda, \bar{a}, \bar{N}^\lambda)$ a model M_δ^λ of cardinality λ containing $\bigcup_{\beta < \delta} M_\beta^\lambda$ and M^δ . Call this tower \mathcal{T}^b , since it contains b .

By λ -symmetry and Fact 2.16, we know that all reduced towers in $\mathcal{K}_{\lambda,\alpha}^*$ are continuous. Therefore \mathcal{T}^b is not reduced. However, by the density of reduced towers [GVV, Theorem 5.6], we can find a reduced, continuous extension of \mathcal{T}^b in $\mathcal{K}_{\lambda,\delta+1}^*$. By λ -applications of this theorem, we may assume that for each $\beta < \alpha$, the model indexed by β in this reduced tower is a $(\lambda, \text{cf}(\lambda))$ -limit over M_β^λ . Refer to this tower as \mathcal{T}^* . See Fig. 2.




FIGURE 2. The towers in the proof of Theorem 1.1

Claim 3.8. *For every $\beta < \alpha$, $\text{ga-tp}(a_\beta/M_\beta^*)$ does not μ -split over N_β .*

Proof. Since M_β^λ and M_β^* are both limit models over N_β^λ , by λ -symmetry and Fact 2.17, there exists $f : M_\beta^\lambda \cong_{N_\beta^\lambda} M_\beta^*$. Since \mathcal{T}^* is a tower extending \mathcal{T}^b , we know $\text{ga-tp}(a_\beta/M_\beta^*)$ does not λ -split over N_β^λ . Therefore by

the definition of non-splitting, it must be the case that $\text{ga-tp}(f(a_\beta)/M_\beta^*) = \text{ga-tp}(a_\beta/M_\beta^*)$. From this equality of types we can fix $g \in \text{Aut}_{M_\beta^*}(\mathfrak{C})$ with $g(f(a_\beta)) = a_\beta$. An application of $g \circ f$ to (1) yields the statement of the claim. \square

We can now complete the proof of Theorem 1.1. By the continuity of \mathcal{T}^* there exists $\beta < \delta$ so that $b \in M_\beta^*$. We can then use \mathcal{T}^* to construct a tower $\dot{\mathcal{T}}$ in $\mathcal{K}_{\mu, \delta+1}^*$ extending \mathcal{T} so that $b \in \dot{M}_\beta$ contradicting our assumption that \mathcal{T} was reduced. This is possible by the downward Löwenheim property of abstract elementary classes, μ -stability, universality of the models in \mathcal{T}^* , monotonicity of non- μ -splitting, and Claim 3.8. \square

Similar to the proof of [Vana, Theorem 4] we can use Lemma 3.7 to derive symmetry from categoricity. More precisely, it is enough to assume that all the models in the top cardinal have enough saturation.

Theorem 3.9. *Suppose λ and μ are cardinals so that $\lambda > \mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is superstable in every $\chi \in [\mu, \lambda)$, and all the models of size λ are μ^+ -saturated, then \mathcal{K} has μ -symmetry.*

Proof. Suppose that \mathcal{K} does not satisfy μ -symmetry. Then by Fact 2.16 there is a reduced discontinuous tower in $\mathcal{K}_{\mu, \alpha}^*$. As in the proof of Theorem 1.1, we can find a discontinuous reduced tower $\mathcal{T} \in \mathcal{K}_{\mu, \alpha}^*$ with $\alpha = \delta + 1$ with the witness of discontinuity $b \in M_\delta \setminus \bigcup_{\beta < \delta} M_\beta$. As in the proof of Theorem 1.1, we can use Lemma 3.7 (note that we only use the first part so not assuming λ -superstability is okay) to find a tower $\mathcal{T}^\lambda \in \mathcal{K}_{\lambda, \mu, \delta}^*$ extending $\mathcal{T} \upharpoonright \delta$.

By our assumption that all the models of size λ are μ^+ -saturated, $\text{ga-tp}(b/\bigcup_{\beta < \delta} M_\beta)$ is realized in $\bigcup_{\beta < \delta} M_\beta^\lambda$. Let b' and $\beta' < \delta$ be such that $b' \models \text{ga-tp}(b/\bigcup_{\beta < \delta} M_\beta)$ and $b' \in M_{\beta'}^\lambda$. Fix $f \in \text{Aut}_{\bigcup_{\beta < \delta} M_\beta}(\mathfrak{C})$ so that $f(b') = b$. Notice that $\mathcal{T}^b := f(\mathcal{T}^\lambda)$ is a tower in $\mathcal{K}_{\lambda, \delta, \mu}^*$ extending $\mathcal{T} \upharpoonright \delta$ with $b \in M_{\beta'}^b$.

We can now use the downward Löwenheim-Skolem property of abstract elementary classes, stability in μ , μ^+ -saturation of models of cardinality λ , and monotonicity of non- μ -splitting to construct from \mathcal{T}^b a discontinuous tower in $\mathcal{K}_{\mu, \alpha}^*$ extending \mathcal{T} so that b appears in the model indexed by β' in the tower. This will contradict our choice of \mathcal{T} being reduced. \square

Remark 3.10. *Instead of assuming that all the models of size λ are μ^+ -saturated, it is enough to assume the following weaker property. For any $\delta < \mu^+$ and any increasing chain $\langle M_i : i < \delta \rangle$ in \mathcal{K}_λ of $(< \lambda, \text{cf}(\lambda))$ -limit models (i.e. for each $i < \delta$, there exists a resolution of M_i $\langle M_i^j \in \mathcal{K}_{<\lambda} : j < \text{cf}(\lambda) \rangle$ such that M_i^{j+1} is universal over M_i^j for each $j < \text{cf}(\lambda)$), $\bigcup_{i < \delta} M_i$ is μ^+ -saturated.*

4. A HIERARCHY OF SYMMETRY PROPERTIES

We discuss the relationship between the symmetry property of Definition 2.12 and other symmetry properties previously defined in the literature, especially the symmetry property in the definition of a good μ -frame. This expands on the short remark after Definition 3 of [Vana] and on Corollary 2 there. It will be convenient to use the following terminology. A minor variation (where “limit over” is replaced by “universal over”) appears in [Vasa, Definition 3.8].

Definition 4.1. *Let $M_0 \leq M \leq N$ be models in \mathcal{K}_μ . We say a type $p \in \text{ga-S}(N)$ explicitly does not μ -fork over (M_0, M) if:*

- (1) *M is limit over M_0 .*
- (2) *p does not μ -split over M_0 .*

We say that p does not μ -fork over M if there exists M_0 so that p explicitly does not μ -fork over (M_0, M) .

Remark 4.2. *Assuming μ -superstability, the relation “ p does not μ -fork over M ” is very close to defining an independence notion with the properties of forking in a first-order superstable theory (i.e. a good μ -frame, see below). In fact using tameness it can be used to do precisely that, see [Vasa] or Theorem 6.4. Moreover forking in any categorical good μ -frame has to be μ -forking, see Fact 4.10.*

We now give several variations on μ -symmetry. We will show that variation (1) is equivalent to (2) which implies (3) which implies (4). Moreover variation (1) is equivalent to the μ -symmetry of Definition 2.12 and variation (4) is equivalent to the symmetry property of good frames. We do not know if any of the implications can be reversed, or even if all the variations already follow from superstability (see Question 4.14).

For clarity, we have highlighted the differences between each property.

Definition 4.3. *Let $\mu \geq \text{LS}(\mathcal{K})$.*

- (1) \mathcal{K} has uniform μ -symmetry if for any limit models N, M_0, M in \mathcal{K}_μ where M is limit over M_0 and M_0 is limit over N , if $\text{ga-tp}(b/M)$ does not μ -split over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ explicitly does not μ -fork over (N, M_0) , there exists $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ **explicitly does not μ -fork over** (N, M_0) .
- (2) \mathcal{K} has weak uniform μ -symmetry if for any limit models N, M_0, M in \mathcal{K}_μ where M is limit over M_0 and M_0 is limit over N , if $\text{ga-tp}(b/M)$ does not μ -fork over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ explicitly does not μ -fork over (N, M_0) , there exists $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ **explicitly does not μ -fork over** (N, M_0) . See Figure 3.
- (3) \mathcal{K} has non-uniform μ -symmetry if for any limit models M_0, M in \mathcal{K}_μ where M is limit over M_0 , if $\text{ga-tp}(b/M)$ does not μ -split over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ does not μ -fork over M_0 , there exists $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ **does not μ -fork over** M_0 .
- (4) \mathcal{K} has weak non-uniform μ -symmetry if for any limit models M_0, M in \mathcal{K}_μ where M is limit over M_0 , if $\text{ga-tp}(b/M)$ does not μ -fork over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ does not μ -fork over M_0 , there exists $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ **does not μ -fork over** M_0 .




FIGURE 3. A diagram of the models and elements in the definition of weak uniform μ -symmetry. We require that $\text{ga-tp}(b/M)$ does not μ -fork over M_0 in the weak version, so there exists M'_0 such that M_0 is limit over M'_0 and $\text{ga-tp}(b/M)$ does not μ -split over M'_0

The difference between the uniform and non-uniform variations is in the conclusion: in the uniform case, we start with $\text{ga-tp}(a/M_0)$ which explicitly does not μ -fork over (N, M_0) and get $\text{ga-tp}(a/M_b)$ explicitly does not μ -fork over (N, M_0) . Thus both types do not μ -split over N . In the non-uniform case, we start with $\text{ga-tp}(a/M_0)$ which does not μ -fork over M_0 , hence explicitly does not μ -fork over (N, M_0) for some N , but we only get that $\text{ga-tp}(a/M_b)$ does not μ -fork over M_0 , so it explicitly does not μ -fork over (N', M_0) , for some N' potentially different from N .

The difference between weak and non-weak is in the starting assumption: in the weak case, we assume that $\text{ga-tp}(b/M)$ does not μ -fork over M_0 , hence there exists M'_0 so that M_0 is limit over M'_0 and $\text{ga-tp}(b/M)$ does not μ -split over M'_0 . In the non-weak case, we assume only that $\text{ga-tp}(b/M)$ does not μ -split over M_0 . Even under μ -superstability, it is open whether this implies that there must exist a smaller M'_0 so that $\text{ga-tp}(b/M)$ does not μ -split over M'_0 . The problem is that μ -splitting need not satisfy the transitivity property, see the discussion after Definition 3.8 in [Vas].

Using the monotonicity property of μ -splitting, we get the easy implications:

Proposition 4.4. *Let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} has uniform μ -symmetry, then it has non-uniform μ -symmetry and weak uniform μ -symmetry. If \mathcal{K} has non-uniform μ -symmetry, then it has weak non-uniform μ -symmetry.*

Playing with the definitions and monotonicity of μ -splitting (noting that cases ruled out by Definition 2.12 such as $a \in |M_0|$ are easy to handle), we also have:

Proposition 4.5. *\mathcal{K} has uniform μ -symmetry if and only if it has μ -symmetry (in the sense of Definition 2.12).*

Surprisingly, uniform symmetry and weak uniform symmetry are also equivalent assuming superstability. We will use the characterization of symmetry in terms of reduced towers provided by Fact 2.16.

Lemma 4.6. *If \mathcal{K} is μ -superstable, then weak uniform μ -symmetry is equivalent to uniform μ -symmetry.*

Proof. By Proposition 4.4, uniform symmetry implies weak uniform symmetry. Now assuming weak uniform symmetry, the proof of (1) \Rightarrow (2) of Fact 2.16 still goes through. The point is that whenever we

consider $\text{ga-tp}(b/M)$ in the proof, $M = \bigcup_{i < \delta} M_i$ for some increasing continuous $\langle M_i : i < \delta \rangle$ with M_{i+1} universal over M_i for all $i < \delta$, and we simply use that by superstability $\text{ga-tp}(b/M)$ does not μ -split over M_i for some $i < \delta$. However we also have that $\text{ga-tp}(b/M)$ explicitly does not μ -fork over (M_i, M_{i+1}) .

Therefore reduced towers are continuous, and hence by Fact 2.16 \mathcal{K} has μ -symmetry (and so by Proposition 4.5 uniform μ -symmetry). \square

How do these definitions compare to the symmetry property in good μ -frames? Recall [She09, Definition II.2.1] that a good μ -frame is a triple $\mathfrak{s} = (\mathcal{K}_\mu, \perp, \text{ga-S}^{\text{bs}})$ where:

- (1) \mathcal{K} is an AEC.
- (2) For each $M \in \mathcal{K}_\mu$, $\text{ga-S}^{\text{bs}}(M)$ (called the set of *basic types* over M) is a set of nonalgebraic Galois types over M satisfying (among others) the *density property*: if $M < N$ are in \mathcal{K}_μ , there exists $a \in |N| \setminus |M|$ such that $\text{ga-tp}(a/M; N) \in \text{ga-S}^{\text{bs}}(M)$.
- (3) \perp is an (abstract) independence relation on types of length one over models in \mathcal{K}_μ satisfying several basic properties (that we will not list here) of first-order forking in a superstable theory.

Remark 4.7. *We will not use the axiom (B) [She09, Definition II.2.1] requiring the existence of a superlimit model of size μ . In fact many papers (e.g. [JS13]) define good frames without this assumption.*

As in [She09, Definition II.6.35], we say that a good μ -frame \mathfrak{s} is *type-full* if for each $M \in \mathcal{K}_\mu$, $\text{ga-S}^{\text{bs}}(M)$ consists of *all* the nonalgebraic types over M . For simplicity, we focus on type-full good frames in this paper. Given a type-full good μ -frame $\mathfrak{s} = (\mathcal{K}_\mu, \perp, \text{ga-S}^{\text{bs}})$ and $M_0 \leq M$ both in \mathcal{K}_μ , we say that a nonalgebraic type $p \in \text{ga-S}(M)$ *does not \mathfrak{s} -fork over M_0* if it does not fork over M_0 according to the abstract independence relation \perp of \mathfrak{s} . We say that a good μ -frame \mathfrak{s} is *on \mathcal{K}_μ* if its underlying class is \mathcal{K}_μ .

The existence of a good μ -frame gives quite a lot of information about the class.

Fact 4.8. *Assume there is a good μ -frame on $\mathcal{K}_\mu^{\lambda\text{-sat}}$, for $\lambda \leq \mu$ (so in particular, unions of chains of λ -saturated models are λ -saturated). Then:*

- (1) *For any $M_0, M_1, M_2 \in \mathcal{K}_\mu$ such that M_1 and M_2 are limit over M_0 , $M_1 \cong_{M_0} M_2$.*
- (2) *\mathcal{K} is μ -superstable.*

Proof. The first part is [She09, Lemma II.4.8] (or see [Bon14a, Theorem 9.2]). Note that by the usual back and forth argument (as made explicit in the proof of [BVb, Theorem 7.1]), any limit model is isomorphic to a limit model where the models witnessing it are λ -saturated. The second part is because:

- By definition of a good μ -frame, $\mu \geq \text{LS}(\mathcal{K})$, \mathcal{K}_μ is nonempty, has amalgamation, joint embedding, and no maximal models.
- By [She09, Claim II.4.2.(1)], \mathcal{K} is stable in μ .
- By the uniqueness property of \mathfrak{s} -forking, if a type does not \mathfrak{s} -fork over M_0 (where \mathfrak{s} is a good μ -frame on \mathcal{K}_μ), then it does not μ -split over M_0 (see [BGKV, Lemma 4.2]). Thus we obtain the “no long splitting chains” condition in Definition 2.8 when the M_i s are λ -saturated, and as noted above (or in [Vasb, Proposition 10.6]) we can do a back and forth argument to get that no long splitting chains holds even when the members of the chain are not saturated.

□

Among the axioms a good μ -frame must satisfy is the symmetry axiom:

Definition 4.9. *The symmetry axiom for a good μ -frame $\mathfrak{s} = (\mathcal{K}_\mu, \perp, \text{ga-S}^{bs})$ is the following statement: For any $M_0 \leq M$ in \mathcal{K}_μ , if $\text{ga-tp}(b/M)$ does not \mathfrak{s} -fork over M_0 and $a \in |M| \setminus |M_0|$ is so that $\text{ga-tp}(a/M_0) \in \text{ga-S}^{bs}(M_0)$, there exists $M_b \in \mathcal{K}_\mu$ containing b and extending M_0 so that $\text{ga-tp}(a/M_b)$ does not \mathfrak{s} -fork over M_0 .*

Note that the good frame axioms imply that \mathcal{K} has amalgamation in μ , so for this definition (and for simplicity only) we work inside a saturated model \mathfrak{C} of size μ^+ .

Since the symmetry properties of Definition 4.3 are all over limit models only, we will discuss only frames whose models are the limit models. By Fact 4.8, such frames are categorical (that is, their underlying class has a single model up to isomorphism). This is not a big loss since most known general constructions of a good μ -frame (e.g. [She09, Theorem II.3.7], [Vasb, Theorem 1.3]) assume categoricity in μ . In the known constructions when categoricity in μ is not assumed (such as in [Vasb, Corollary 10.19]), it holds that the union of a chain of μ -saturated model is μ -saturated, so we can simply restrict the frame to the saturated models of size μ .

We will use [Vasb, Theorem 9.7] that categorical good μ -frames are canonical:

Fact 4.10 (The canonicity theorem for categorical good frames). *Let $\mathfrak{s} = (\mathcal{K}_\mu, \perp, \text{ga-S}^{bs})$ be a categorical good μ -frame. Let $p \in \text{ga-S}^{bs}(M)$ and let $M_0 \leq M$ be in \mathcal{K}_μ . Then p does not \mathfrak{s} -fork over M_0 if and only if p does not μ -fork over M_0 (recall Definition 4.1).*

Remark 4.11. *The statement of Theorem 9.7 in [Vasb] uses the definition of μ -forking with “universal over” instead of “limit over,” but the proof goes through also for the “limit over” definition.*

Remark 4.12. *The proof of the second part of Fact 4.8 and Fact 4.10 do not use the symmetry axiom (but the first part of Fact 4.8 does).*

Using the canonicity theorem, we obtain:

Theorem 4.13. *Let \mathfrak{s} be a type-full categorical good μ -frame on \mathcal{K}_μ , except that we do not assume that it satisfies the symmetry axiom. The following are equivalent:*

- (1) *\mathfrak{s} satisfies the symmetry axiom (Definition 4.9).*
- (2) *\mathcal{K} has weak non-uniform μ -symmetry (Definition 4.3.(4)).*

Proof. By Fact 4.10 (and Remark 4.12), μ -forking and \mathfrak{s} -forking coincide. Now replace \mathfrak{s} -forking by μ -forking in the symmetry axiom and expand the definition. \square

One can ask whether weak non-uniform symmetry can be replaced by the uniform version:

Question 4.14. *Assume there is a type-full categorical good μ -frame on \mathcal{K}_μ . Does \mathcal{K} have μ -symmetry? More generally, if \mathcal{K} is μ -superstable, does \mathcal{K} have μ -symmetry?*

We will show (Corollary 6.9) that the answer is positive if \mathcal{K} is μ -tame. Still much less suffices:

Theorem 4.15. *If \mathcal{K} is μ -superstable and has a good μ^+ -frame on $\mathcal{K}_{\mu^+}^{\lambda\text{-sat}}$ for some $\lambda \leq \mu^+$, then \mathcal{K} has μ -symmetry.*

Proof. By Fact 4.8, all limit models in \mathcal{K}_{μ^+} are saturated and \mathcal{K} is μ^+ -superstable. By Fact 2.18, the remark following it, and Fact 2.13, \mathcal{K} has μ -symmetry. \square

We end this section with a partial answer to Question 4.14 assuming that the good frame satisfies several additional technical properties of frames introduced by Shelah (see [She09, Definitions III.1.1, III.1.3]). For this result amalgamation (Hypothesis 2.1) is not necessary.

Corollary 4.16. *Assume there is a successful good^+ μ -frame with underlying class \mathcal{K}_μ . Then \mathcal{K} has μ -symmetry.*

Proof. Let \mathfrak{s} be a successful good^+ μ -frame with underlying class \mathcal{K}_μ . By [She09, Claim II.6.36], we can assume without loss of generality that \mathfrak{s} is type-full (note that by [BGKV, Theorem 6.13] there can be only one such type-full frame). By Fact 4.8, \mathcal{K} is μ -superstable. By [She09, III.1.6, III.1.7, III.1.8], there is a good μ^+ -frame \mathfrak{s}^+ on $\mathcal{K}_{\mu^+}^{\mu^+\text{-sat}}$. By Theorem 4.15, \mathcal{K} has μ -symmetry. \square

5. SYMMETRY FROM NO ORDER PROPERTY

In this section, we give another way to derive symmetry. The idea is to imitate the argument from [BGKV, Theorem 5.14], but we first have to obtain enough properties of independence. We will work with μ -forking (Definition 4.1). We start by improving Proposition 3.4.

Proposition 5.1 (Extension property of forking). *Let $\text{LS}(\mathcal{K}) \leq \mu \leq \lambda$. Let $M \leq N$ be in $K_{[\mu, \lambda]}$. Let $p \in \text{ga-S}(M)$ be such that p explicitly does not μ -fork over (M_0, M) . If \mathcal{K} is superstable in every $\chi \in [\mu, \lambda]$, then there exists $q \in \text{ga-S}(N)$ extending p and explicitly not μ -forking over (M_0, M) . Moreover q is algebraic if and only if p is.*

Proof. By induction on $\|N\|$. Let a realize p . If $\|N\| = \|M\|$ this is given by Proposition 3.4 (if $\|M\| = \|N\| = \mu$, this is [Van06, Theorem I.4.10]). If $\|M\| < \|N\|$, build $\langle N_i \in \mathcal{K}_{\|M\|+|i|} : i \leq \|N\| \rangle$ increasing continuous such that $N_0 = M$, N_{i+1} is limit over N_i , and $\text{ga-tp}(a/N_i)$ explicitly does not μ -fork over (M_0, M) . This is possible by the induction hypothesis and the continuity property of splitting (Proposition 3.3). Now N_λ is $\|N\|$ -universal over $N_0 = M$, so let $f : N \xrightarrow[M]{} N_\lambda$. Let $q := f^{-1}(\text{ga-tp}(a/f[N]))$. It is easy to check that q is as desired. \square

Next we recall the definition of the order property in AECs [She99, Definition 4.3].

Definition 5.2. *Let α and λ be cardinals. A model $M \in \mathcal{K}$ has the α -order property of length λ if there exists $\langle \bar{a}_i : i < \lambda \rangle$ inside M with $\ell(\bar{a}_i) = \alpha$ for all $i < \lambda$, such that for any $i_0 < j_0 < \lambda$ and $i_1 < j_1 < \lambda$, $\text{ga-tp}(\bar{a}_{i_0}\bar{a}_{j_0}/\emptyset) \neq \text{ga-tp}(\bar{a}_{i_1}\bar{a}_{j_1}/\emptyset)$.*

We say that \mathcal{K} has the α -order property of length λ if some $M \in \mathcal{K}$ has it. We say that \mathcal{K} has the α -order property if it has the α -order property of length λ for all cardinals λ .

We will use two important facts. The first says that it is enough to look at length up to the Hanf number. The second is that the order property implies instability.

Definition 5.3. For λ an infinite cardinal, $h(\lambda) := \beth_{(2^\lambda)^+}$.

Fact 5.4 (Claim 4.5.3 in [She99]). *Let α be a cardinal. If \mathcal{K} has the α -order property of length λ for all $\lambda < h(\alpha + \text{LS}(\mathcal{K}))$, then \mathcal{K} has the α -order property.*

Fact 5.5. *If \mathcal{K} has the α -order property and $\mu \geq \text{LS}(\mathcal{K})$ is such that $\mu = \mu^\alpha$, then \mathcal{K} is not stable in μ .*

Proof. By [She99, Claim 4.8.2] (see [BGKV, Fact 5.13] for a proof), there exists $M \in \mathcal{K}_\mu$ such that $|\text{ga-S}^\alpha(M)| > \mu$. By [Bon, Theorem 3.1], \mathcal{K} is not stable in μ . \square

The following lemma appears in some more abstract form in [BGKV, Lemma 5.6]. The lemma says that if we assume that p does not μ -fork over M , then in the definition of non-splitting (Definition 2.3) we can replace the N_ℓ by arbitrary sequences in N of length at most μ . In the proof of Lemma 5.7, this will be used for sequences of length one.

Lemma 5.6. *Let $\mu \geq \text{LS}(\mathcal{K})$. Let $M \in \mathcal{K}_\mu$ and $N \in \mathcal{K}_{\geq \mu}$ be such that $M \leq N$. Assume that \mathcal{K} is stable in μ . If $p \in \text{ga-S}(N)$ does not μ -fork over M (Definition 4.1), a realizes p , and $\bar{b}_1, \bar{b}_2 \in {}^{\leq \mu}N$ are such that $\text{ga-tp}(\bar{b}_1/M) = \text{ga-tp}(\bar{b}_2/M)$, then $\text{ga-tp}(a\bar{b}_1/M) = \text{ga-tp}(a\bar{b}_2/M)$.*

Proof. Pick $N_0 \in \mathcal{K}_\mu$ containing $\bar{b}_1\bar{b}_2$ with $M \leq N_0 \leq N$. Then $p \upharpoonright N_0$ does not μ -fork over M . Replacing N by N_0 if necessary, we can assume without loss of generality that $N \in \mathcal{K}_\mu$. By definition of μ -forking, there exists $M_0 \in \mathcal{K}_\mu$ such that $M_0 \leq M$ and p does not μ -split over M_0 . By the extension and uniqueness property for μ -splitting there exists N' extending N of cardinality μ so that N' is universal over both N and M , and $\text{ga-tp}(a/N')$ does not μ -split over M_0 . Since $\text{ga-tp}(\bar{b}_1/M) = \text{ga-tp}(\bar{b}_2/M)$ and since N' is universal over N , we can find $f : N \xrightarrow{M} N'$ so that $f(\bar{b}_1) = \bar{b}_2$. Since $\text{ga-tp}(a/N')$ does not μ -split over M_0 we know $\text{ga-tp}(f(a)/f(N)) = \text{ga-tp}(a/f(N))$. By our choice of f this implies that there exists $g \in \text{Aut}_{f(N)}(\mathfrak{C})$ so that $g(f(a)) = a$, $g \upharpoonright M = \text{id}_M$, and $g(\bar{b}_2) = \bar{b}_1$. In other words $\text{ga-tp}(f(a)\bar{b}_2/M) = \text{ga-tp}(a\bar{b}_1/M)$. Moreover f^{-1} witnesses that $\text{ga-tp}(a\bar{b}_1/M) = \text{ga-tp}(f(a)\bar{b}_2/M)$, which we have seen is equal to $\text{ga-tp}(a\bar{b}_2/M)$. \square

The next lemma shows that failure of symmetry implies the order property. The proof is similar to that of [BGKV, Theorem 5.14], the difference is that we use Lemma 5.6 and the equivalence between symmetry and weak uniform symmetry (Lemma 4.6).

Lemma 5.7. *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, \lambda)$. If \mathcal{K} does not have μ -symmetry, then it has the μ -order property of length λ .*

Proof. By Lemma 4.6, \mathcal{K} does not have weak uniform μ -symmetry. We first pick witnesses to that fact. Pick limit models $N, M_0, M \in \mathcal{K}_\mu$ such that M is limit over M_0 and M_0 is limit over N . Pick b such that $\text{ga-tp}(b/M)$ does not μ -fork over M_0 , $a \in |M|$, and $\text{ga-tp}(a/M_0)$ explicitly does not μ -fork over (N, M_0) , and there does *not* exist $M_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a/M_b)$ explicitly does not μ -fork over (N, M_0) . We will show that \mathfrak{C} has the μ -order property of length λ .

We build increasing continuous $\langle N_\alpha : \alpha < \lambda \rangle$ and $\langle a_\alpha, b_\alpha, N'_\alpha : \alpha < \lambda \rangle$ by induction so that for all $\alpha < \lambda$:

- (1) $N_\alpha, N'_\alpha \in \mathcal{K}_{\mu+|\alpha|}$.
- (2) N_0 is limit over M and $b \in |N_0|$.
- (3) $\text{ga-tp}(a_\alpha/M_0) = \text{ga-tp}(a/M_0)$ and $a_\alpha \in |N'_\alpha|$.
- (4) $\text{ga-tp}(b_\alpha/M) = \text{ga-tp}(b/M)$ and $b_\alpha \in |N_{\alpha+1}|$.
- (5) N'_α is limit over N_α and $N_{\alpha+1}$ is limit over N'_α .
- (6) $\text{ga-tp}(a_\alpha/N_\alpha)$ explicitly does not μ -fork over (N, M_0) and $\text{ga-tp}(b_\alpha/N'_\alpha)$ does not μ -fork over M_0 .

This is possible. Let N_0 be any model in \mathcal{K}_μ containing M and a and limit over M . At α limits, let $N_\alpha := \bigcup_{\beta < \alpha} N_\beta$. Now assume inductively that N_β has been defined for $\beta \leq \alpha$, and $a_\beta, b_\beta, N'_\beta$ have been defined for $\beta < \alpha$. By extension for splitting, find $q \in \text{ga-S}(N_\alpha)$ that explicitly does not μ -fork over (N, M_0) and extends $\text{ga-tp}(a/M_0)$. Let a_α realize q and pick N'_α limit over N_α containing a_α . Now by extension again, find $q' \in \text{ga-S}(N'_\alpha)$ that does not μ -fork over M_0 and extends $\text{ga-tp}(b/M)$. Let b_α realize q' and pick $N_{\alpha+1}$ limit over N'_α containing b_α .

This is enough. We show that for $\alpha, \beta < \lambda$:

- (1) $\text{ga-tp}(a_\alpha b/M_0) \neq \text{ga-tp}(ab/M_0)$
- (2) If $\beta < \alpha$, $\text{ga-tp}(ab/M_0) \neq \text{ga-tp}(a_\alpha b_\beta/M_0)$.
- (3) If $\beta \geq \alpha$, $\text{ga-tp}(ab/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$.

For (1), observe that $b \in |N_0| \subseteq |N_\alpha|$ and $\text{ga-tp}(a_\alpha/N_\alpha)$ explicitly does not μ -fork over (N, M_0) . Therefore by monotonicity N_α witnesses that there exists $N_b \in \mathcal{K}_\mu$ containing b and limit over M_0 so that $\text{ga-tp}(a_\alpha/M_b)$ explicitly does not μ -fork over (N, M_0) . By failure of symmetry and invariance, we must have that $\text{ga-tp}(a_\alpha b/M_0) \neq \text{ga-tp}(ab/M_0)$.

For (2), suppose $\beta < \alpha$. We know that $\text{ga-tp}(a_\alpha/N_\alpha)$ does not μ -fork over M_0 . Since $\beta < \alpha$, $b, b_\beta \in |N_\alpha|$ and $\text{ga-tp}(b/M) = \text{ga-tp}(b_\beta/M)$, we must have by Lemma 5.6 that $\text{ga-tp}(a_\alpha b/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$. Together with (1), this implies $\text{ga-tp}(ab/M_0) \neq \text{ga-tp}(a_\alpha b_\beta/M_0)$. This is really where we use the equivalence between uniform μ -symmetry and weak uniform μ -symmetry: if we only had failure of uniform μ -symmetry, then we would only know that $\text{ga-tp}(b/M)$ does not μ -split over M_0 , so would be unable to use Lemma 5.6.

To see (3), suppose $\beta \geq \alpha$ and recall that (by (1)) $\text{ga-tp}(ab/M_0) = \text{ga-tp}(a_\beta b/M_0)$. We also have that $\text{ga-tp}(b_\beta/N'_\beta)$ does not μ -fork over M_0 . Moreover $\text{ga-tp}(a/M_0) = \text{ga-tp}(a_\alpha/M_0)$, and $a, a_\alpha \in N'_\beta$. By Lemma 5.6 again, $\text{ga-tp}(ab_\beta/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$. This gives us that $\text{ga-tp}(ab/M_0) = \text{ga-tp}(a_\alpha b_\beta/M_0)$.

Now let \bar{d} be an enumeration of M_0 and for $\alpha < \lambda$, let $\bar{c}_\alpha := a_\alpha b_\alpha \bar{d}$. Then (2) and (3) together tell us that the sequence $\langle \bar{c}_\alpha \mid \alpha < \lambda \rangle$ witnesses the μ -order property of length λ . \square

We conclude that symmetry follows from enough instances of superstability.

Theorem 5.8. *Let $\mu \geq \text{LS}(\mathcal{K})$. Then there exists $\lambda < h(\mu)$ such that if \mathcal{K} is superstable in every $\chi \in [\mu, \lambda)$, then \mathcal{K} has μ -symmetry.*

Proof. If \mathcal{K} is unstable in 2^μ , then we can set $\lambda := (2^\mu)^+$ and get a vacuously true statement; so assume that \mathcal{K} is stable in 2^μ . By Fact 5.5, \mathcal{K} does not have the μ -order property. By Fact 5.4, there exists $\lambda < h(\mu)$ such that \mathcal{K} does not have the μ -order property of length λ . By Lemma 5.7, it is as desired. \square

Remark 5.9. *How can one obtain many instances of superstability as in the hypothesis of Theorem 5.8? One way is categoricity, see Fact 7.1. Another way is to start with one instance of superstability and transfer it up using tameness, see Fact 6.7.*

6. SYMMETRY AND TAMENESS

Tameness is a locality property for types introduced by Grossberg and VanDieren in [GV06b] and used to prove Shelah's eventual categoricity conjecture from a successor in [GV06c]. It has also played a key roles in the proof of several other categoricity transfers, for example [Bon14b, Vasc, Vasd].

Definition 6.1 (Tameness). *Let $\mu \geq \text{LS}(\mathcal{K})$. \mathcal{K} is μ -tame if for every $M \in \mathcal{K}$ and every $p, q \in \text{ga-S}(M)$, if $p \neq q$, then there exists $M_0 \in \mathcal{K}_{\leq \mu}$ with $M_0 \leq M$ such that $p \restriction M_0 \neq q \restriction M_0$.*

In this section, we study the combination of tameness (and its relatives, see below) with superstability. In Section 7, we will combine tameness and categoricity.

6.1. Weak tameness. We will start by studying a weaker, more local, variation that appears already in [She99]. We use the notation in [Bal09, Definition 11.6].

Definition 6.2 (Weak tameness). *Let χ, μ be cardinals with $\text{LS}(\mathcal{K}) \leq \chi \leq \mu$. \mathcal{K} is (χ, μ) -weakly tame if for any saturated $M \in \mathcal{K}_\mu$, any $p, q \in \text{ga-S}(M)$, if $p \neq q$, there exists $M_0 \in \mathcal{K}_\chi$ with $M_0 \leq M$ and $p \restriction M_0 \neq q \restriction M_0$.*

Tameness says that type over *any* models are determined by their small restrictions. Weak tameness says that only types over *saturated* models have this property.

While there is no known example of an AEC that is weakly tame but not tame, it is known that weak tameness follows from categoricity in a suitable cardinal (but the corresponding result for non-weak tameness is open, see [GV06a, Conjecture 1.5]): this appears as [She99, Main Claim II.2.3] and a simplified argument is in [Bal09, Theorem 11.15].

Fact 6.3. *Let $\lambda > \mu \geq h(\text{LS}(\mathcal{K}))$. Assume that \mathcal{K} is categorical in λ , and the model of cardinality λ is μ^+ -saturated. Then there exists $\chi < h(\text{LS}(\mathcal{K}))$ such that \mathcal{K} is (χ, μ) -weakly tame.*

It was shown in [Vasa] (and further improvements in [Vasb, Section 10] and [BVb]) that tameness can be combined with superstability to build a good frame at a high-enough cardinal. At a meeting in the winter of 2015 in San Antonio, the first author asked whether weak tameness could be used instead. This is not a generalization for the sake of

generalization because weak tameness (but not tameness) is known to follow from categoricity. We can answer in the affirmative:

Theorem 6.4. *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, \lambda]$ and has λ -symmetry.*

If \mathcal{K} is (μ, λ) -weakly tame, then there exists a type-full good λ -frame with underlying class $\mathcal{K}_\lambda^{\lambda\text{-sat}}$.

Proof. First observe that limit models in \mathcal{K}_λ are unique (by Fact 2.17), hence saturated. By Theorem 1.1, \mathcal{K} has χ -symmetry for every $\chi \in [\mu, \lambda]$. By Fact 2.18, for every $\chi \in [\mu, \lambda)$, $\mathcal{K}^{\chi^+\text{-sat}}$, the class of χ^+ -saturated models in $\mathcal{K}_{\geq \chi^+}$ is an AEC with $\text{LS}(\mathcal{K}^{\chi^+\text{-sat}}) = \chi^+$. Therefore by Lemma 2.20 $\mathcal{K}^{\lambda\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\lambda\text{-sat}}) = \lambda$. By the λ -superstability assumption, $\mathcal{K}_\lambda^{\lambda\text{-sat}}$ is nonempty, has amalgamation, no maximal models, and joint embedding. It is also stable in λ . We want to define a type-full good λ -frame \mathfrak{s} on $\mathcal{K}_\lambda^{\lambda\text{-sat}}$. We define forking in the sense of \mathfrak{s} (\mathfrak{s} -forking) as follows: For $M \leq N$ saturated of size λ , a non-algebraic $p \in \text{ga-S}(N)$ does not \mathfrak{s} -fork over M if and only if it does not μ -fork over M .

Now most of the axioms of good frames are verified in Section 4 of [Vasa], the only properties that remain to be checked are extension, uniqueness, and symmetry. Extension is by Proposition 5.1, and uniqueness is by uniqueness of splitting in μ ([Van06, I.4.12]) and the weak tameness assumption. As for symmetry, we know that λ -symmetry holds, hence by Proposition 4.4, Proposition 4.5, and Theorem 4.13 the symmetry property of good frame follows. \square

Remark 6.5. *If $\lambda = \mu^+$ above, then the hypotheses reduce to “ \mathcal{K} is superstable in μ and μ^+ and \mathcal{K} has μ^+ -symmetry”.*

We can combine this construction with the results of Section 5:

Corollary 6.6. *Let $\lambda > \mu \geq \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is superstable in every $\chi \in [\mu, h(\lambda))$. If \mathcal{K} is (μ, λ) -weakly tame, then there exists a type-full good λ -frame with underlying class $\mathcal{K}_\lambda^{\lambda\text{-sat}}$.*

Proof. Combine Theorem 6.4 and Theorem 5.8. \square

6.2. Global tameness. For the rest of this section, we will work with global non-weak tameness. Superstability has been studied together with amalgamation and tameness in works from the second author [Vasa, Vasb, BVb, GV]. We will use the following upward transfer of superstability:

Fact 6.7 (Proposition 10.10 in [Vasb]). *Assume \mathcal{K} is μ -superstable and μ -tame. Then for all $\mu' \geq \mu$, \mathcal{K} is μ' -superstable. In particular, $\mathcal{K}_{\geq \mu}$ has no maximal models and is stable in all cardinals.*

An application of tameness and superstability is to chains of saturated models. Recall from Section 4 that $\mathcal{K}^{\lambda\text{-sat}}$ denotes the class of λ -saturated models in $\mathcal{K}_{\geq \lambda}$. We would like to give conditions under which $\mathcal{K}^{\lambda\text{-sat}}$ is an AEC – in particular unions of chains of λ -saturated models are λ -saturated. From superstability and tameness, it is known that one eventually obtains this behavior:

Fact 6.8 (Theorem 7.1 in [BVb]). *Assume \mathcal{K} is μ -superstable and μ -tame. Then there exists $\lambda_0 < \beth_{(2^{\mu^+})^+}$ such that for any $\lambda \geq \lambda_0$, $\mathcal{K}^{\lambda\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\lambda\text{-sat}}) = \lambda$.*

We can use this to show that superstability implies symmetry in tame AECs (obtaining another partial answer to Question 4.14). We also give another, more self-contained proof that does not rely on Fact 6.8.

Corollary 6.9. *If \mathcal{K} is μ -superstable and μ -tame, then \mathcal{K} has μ -symmetry.*

First proof. First observe that by Fact 6.7, \mathcal{K} is superstable in every $\mu' \geq \mu$. By Fact 6.8, there exists $\lambda_0 \geq \mu$ such that $\mathcal{K}^{\lambda_0^+\text{-sat}}$ is an AEC. Therefore the hypotheses of Fact 2.13 are satisfied, so \mathcal{K} has λ_0 -symmetry. By Theorem 1.1, \mathcal{K} has μ -symmetry. \square

Second proof. As in the first proof, \mathcal{K} is superstable in every $\mu' \geq \mu$. By Theorem 5.8, \mathcal{K} has μ -symmetry. \square

Thus we obtain an improvement on the Hanf number of Fact 6.8:

Corollary 6.10. *Assume \mathcal{K} is μ -superstable and μ -tame. For every $\lambda > \mu$, $\mathcal{K}^{\lambda\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\lambda\text{-sat}}) = \lambda$.*

Proof. By Fact 6.7 and Corollary 6.9, \mathcal{K} is λ -superstable and has λ -symmetry for any $\lambda > \mu$. By Fact 2.18, $\mathcal{K}^{\mu^+\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\mu^+\text{-sat}}) = \mu^+$. We can replace μ^+ with any successor $\lambda > \mu$. To take care of limit cardinals λ , use Lemma 2.20. \square

Note that Corollary 6.10 is an improvement on Fact 6.8 and its second proof does not rely on Fact 6.8. However beyond Fact 6.8, the arguments of [BVb] (in particular the use of averages) have other applications (see for example the proof of solvability in [GV, Theorem 5.43]).

We can also say more on another result of Boney and the second author: [BVb, Lemma 6.9.(2)] implies that, assuming μ -superstability, there is a $\lambda_0 \geq \mu$ such that if $\langle M_i : i < \delta \rangle$ is a chain of λ_0 -saturated models where $\delta \geq \lambda_0$ and M_{i+1} is universal over M_i , then $\bigcup_{i < \delta} M_i$ is saturated. We can improve this too:

Corollary 6.11. *Assume \mathcal{K} is μ -superstable and μ -tame. Let δ be a limit ordinal and $\langle M_i : i < \delta \rangle$ is increasing in $\mathcal{K}_{\geq \mu}$ and M_{i+1} is universal over M_i for all $i < \delta$. Let $M_\delta := \bigcup_{i < \delta} M_i$. If $\|M_\delta\| > \text{LS}(\mathcal{K})$, then M_δ is saturated.*

Proof. By Proposition 3.1, M_δ is a $(\lambda, \text{cf}(\delta))$ -limit model, where $\lambda = \|M_\delta\|$. By Fact 6.7, \mathcal{K} is λ -superstable. By Corollary 6.9, \mathcal{K} has λ -symmetry. By Fact 2.17, M_δ is saturated. \square

One can ask whether Corollary 6.10 can be improved further by also getting the conclusion for $\lambda = \mu$. If $\mu = \text{LS}(\mathcal{K})$, it is not clear that $\text{LS}(\mathcal{K})$ -saturated models are the right notion so perhaps the right question is to be framed in terms of a superlimit. Recall from [She09, Definition N.2.2.4] that a superlimit model is a universal model M with a proper extension so that if $\langle M_i : i < \delta \rangle$ is an increasing chain with $M \cong M_i$ for all $i < \delta$, then (if $\delta < \|M\|^+$), $M \cong \bigcup_{i < \delta} M_i$. Note that, assuming μ -superstability and uniqueness of limit models of size μ , it is easy to see that the existence of a superlimit of size μ is equivalent to the statement that the union of an increasing chain of limit models in μ (of length less than μ^+) is limit.

Question 6.12. *Assume \mathcal{K} is μ -tame and there is a type-full good μ -frame on \mathcal{K}_μ (or just that \mathcal{K} is μ -superstable). Is there a superlimit model of size μ ?*

We now turn to good frames and show that, assuming tameness, the statement of Theorem 6.4 can be simplified. Recall that previous work of the second author gives a condition under which good frames can be constructed from tameness:

Fact 6.13 (Theorem 10.8 in [Vasb]). *Assume \mathcal{K} is μ -superstable and μ -tame. If for any $\delta < \mu^+$, any chain of length δ of saturated models in \mathcal{K}_{μ^+} has a saturated union, then there is a type-full good μ^+ -frame with underlying class $\mathcal{K}_{\mu^+}^{\mu^+-\text{sat}}$.*

Combining this with Fact 6.8 it was proven in [BVb] that μ -superstability and μ -tameness implies the existence of a good λ -frame on the saturated models of size λ , for some high-enough $\lambda > \mu$. Now we show that

we can take $\lambda = \mu^+$. We again give two proofs: one uses Theorem 6.4 and the other relies on Fact 6.13.

Corollary 6.14. *If \mathcal{K} is μ -superstable and μ -tame, then there is a type-full good μ^+ -frame with underlying class $\mathcal{K}_{\mu^+}^{\mu^+-sat}$.*

First proof. Combine Fact 6.13 and Corollary 6.10. \square

Second proof. By Fact 6.7, \mathcal{K} is superstable in every $\mu' \geq \mu$. Now apply Corollary 6.6 (with λ there standing for μ^+ here). \square

Remark 6.15. *To obtain a type-full good λ -frame for $\lambda > \mu^+$, we can either make a slight change to the second proof of Corollary 6.14, or use the upward frame transfer of Boney and the second author [Bon14a, BVc].*

7. SYMMETRY AND CATEGORICITY

Theorem 1.1 has several applications to categorical AECs. We will use the following result, an adaptation of an argument of Shelah and Villaveces [SV99, Theorem 2.2.1], to settings with amalgamation:

Fact 7.1 (The Shelah-Villaveces theorem, Theorem 6.3 in [GV]). *Assume that \mathcal{K} has no maximal models. Let $\mu \geq \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in a $\lambda > \mu$, then \mathcal{K} is μ -superstable.*

Corollary 7.2. *Assume that \mathcal{K} has no maximal models. Suppose λ and μ are cardinals so that $\lambda > \mu \geq \text{LS}(\mathcal{K})$ and assume that \mathcal{K} is categorical in λ . Then \mathcal{K} is μ -superstable and it has μ -symmetry if at least one of the following conditions hold:*

- (1) *The model of size λ is μ^+ -saturated.*
- (2) *$\lambda \geq h(\mu)$.*

Proof. By Fact 7.1, \mathcal{K} is χ -superstable in every $\chi \in [\mu, \lambda)$. Now:

- (1) If the model of size λ is μ^+ -saturated, then by Theorem 3.9, \mathcal{K} has μ -symmetry.
- (2) If $\lambda \geq h(\mu)$, then by Theorem 5.8, \mathcal{K} has μ -symmetry.

\square

As announced in the introduction, we can combine Corollary 7.2 with Fact 2.17 to improve on [She99, Theorem 6.5]. The following result also improves on Corollary 18 of [Vanb], by removing the successor

assumption in the categoricity cardinal and obtaining uniqueness of limit models in much smaller cardinalities as well.

Corollary 7.3. *Assume that \mathcal{K} has no maximal models. Suppose λ and μ are cardinals so that $\lambda > \mu \geq \text{LS}(\mathcal{K})$ and assume that \mathcal{K} is categorical in λ . If either $\text{cf}(\lambda) > \mu$ or $\lambda \geq h(\mu)$, then \mathcal{K} has uniqueness of limit models of cardinality μ . That is, if $M_0, M_1, M_2 \in \mathcal{K}_\mu$ are such that both M_1 and M_2 are limit models over M_0 , then $M_1 \cong_{M_0} M_2$.*

Proof. Categoricity in λ , the assumption that $\text{cf}(\lambda) > \mu$, and Fact 7.1 imply that the model of cardinality λ is μ^+ -saturated. We can now apply Corollary 7.2, to get that \mathcal{K} is μ -superstable and has μ -symmetry. Then Fact 2.17 finishes the proof. \square

Once we have obtained symmetry from a high-enough categoricity cardinal, we can deduce that the model in the categoricity cardinal has some saturation:

Corollary 7.4. *Let $\mu > \text{LS}(\mathcal{K})$. Assume that \mathcal{K} is categorical in a $\lambda \geq \sup_{\mu_0 < \mu} h(\mu_0^+)$. Then the model of size λ is μ -saturated.*

Proof. By Fact 2.2, we can assume without loss of generality that \mathcal{K} has no maximal models. We check that the model of size λ is μ_0^+ -saturated for every $\mu_0 \in [\text{LS}(\mathcal{K}), \mu)$. Fix such a μ_0 . By Corollary 7.2, \mathcal{K} is μ_0 -superstable, μ_0^+ -superstable, and has μ_0^+ -symmetry. By Fact 2.18 $\mathcal{K}^{\mu_0^+ \text{-sat}}$, the class of μ_0^+ -saturated models in $\mathcal{K}_{\geq \mu_0^+}$, is an AEC with Löwenheim-Skolem number μ_0^+ . Since it has arbitrarily large models, it must have a model of size λ , which is unique by categoricity. \square

We conclude that categoricity in a high-enough cardinal implies some amount of weak tameness:

Corollary 7.5. *Let $\mu \geq \text{LS}(\mathcal{K})$. Let $\lambda \geq h(\mu^+)$. If \mathcal{K} is categorical in λ , then there exists $\chi < H_1$ such that \mathcal{K} is (χ, μ) -weakly tame.*

Proof. By Corollary 7.4, the model of size λ is μ^+ -saturated. Now apply Fact 6.3. \square

We can derive a downward categoricity transfer. We will use the notation from Chapter 14 of [Bal09]: we write H_1 for $h(\text{LS}(\mathcal{K}))$ and H_2 for $h(H_1) = h(h(\text{LS}(\mathcal{K})))$. We will use the following fact, given by the proof of [Bal09, Theorem 14.9] (originally [She99, II.1.6]):

Fact 7.6. *If \mathcal{K} is categorical in a $\lambda > H_2$, \mathcal{K} is (χ, H_2) -weakly tame for some $\chi < H_1$, and the model of size λ is χ -saturated, then \mathcal{K} is categorical in H_2 .*

Corollary 7.7. *If \mathcal{K} is categorical in a $\lambda \geq h(H_2^+)$, then \mathcal{K} is categorical in H_2 .*

Proof. By Corollary 7.5, there exists $\chi < H_1$ such that \mathcal{K} is (χ, H_2) -weakly tame. By Corollary 7.4, the model of size λ is χ -saturated. Now apply Fact 7.6. \square

We obtain in particular:

Corollary 7.8. *Let $\mu = \beth_\mu > \text{LS}(\mathcal{K})$. If \mathcal{K} is categorical in some $\lambda > \mu$, then \mathcal{K} is categorical in μ .*

Proof. Without loss of generality (Fact 2.2), \mathcal{K} has no maximal models. Applying Corollary 7.7 to $\mathcal{K}_{\geq \mu_0}$ for each $\mu_0 < \mu$, we get that \mathcal{K} is categorical in unboundedly many $\mu_0 < \mu$. By (for example) Fact 7.1, \mathcal{K} is stable in every $\mu_0 < \mu$. Thus the models in the categoricity cardinals below μ are saturated, hence every model of size μ is also saturated. \square

We can also build a good frame assuming categoricity in a high-enough cardinal (of arbitrary cofinality).

Corollary 7.9. *Let $\mu \geq H_1$. Assume that \mathcal{K} is categorical in a $\lambda > \mu$. If the model of size λ is μ^+ -saturated (e.g. if $\text{cf}(\lambda) > \mu$ or by Corollary 7.4 if $\lambda \geq h(\mu^+)$), then there exists a type-full good μ -frame with underlying class $\mathcal{K}_\mu^{\mu\text{-sat}}$.*

Proof. By Fact 2.2, we can assume without loss of generality that \mathcal{K} has no maximal models. By Fact 6.3, there exists $\chi < H_1$ such that \mathcal{K} is (χ, μ) -weakly tame. By Corollary 7.2, \mathcal{K} has χ' -symmetry and is χ' -superstable for every $\chi' \in [\chi, \mu]$. Now apply Theorem 6.4 with (μ, λ) there standing for (χ, μ) -here. \square

The Hanf number of Corollary 7.9 can be improved if we assume that the AEC is tame. We state a more general corollary summing up our results in tame categorical AECs:

Corollary 7.10. *Assume that \mathcal{K} has no maximal models and is $\text{LS}(\mathcal{K})$ -tame. If \mathcal{K} is categorical in a $\lambda > \text{LS}(\mathcal{K})$, then:*

- (1) For any $\mu \geq \text{LS}(\mathcal{K})$, \mathcal{K} has uniqueness of limit models in μ : if $M_0, M_1, M_2 \in \mathcal{K}_\mu$ are such that both M_1 and M_2 are limit over models M_0 , then $M_1 \cong_{M_0} M_2$.
- (2) For any $\mu > \text{LS}(\mathcal{K})$, $\mathcal{K}^{\mu\text{-sat}}$ is an AEC with $\text{LS}(\mathcal{K}^{\mu\text{-sat}}) = \mu$ and there exists a type-full good μ -frame with underlying class $\mathcal{K}_\mu^{\mu\text{-sat}}$.

Proof. By Fact 7.1, \mathcal{K} is superstable in $\text{LS}(\mathcal{K})$. By Fact 6.7, \mathcal{K} is superstable in every $\mu \geq \text{LS}(\mathcal{K})$. By Corollary 6.9, \mathcal{K} has symmetry in every $\mu \geq \text{LS}(\mathcal{K})$. The first part now follows from Fact 2.17 and the second from Corollary 6.10 and Corollary 6.14 (together with Remark 6.15). \square

Remark 7.11. By [BVc], we can transfer the type-full good $\text{LS}(\mathcal{K})^+$ -frame on $\mathcal{K}_{\text{LS}(\mathcal{K})^+}^{\text{LS}(\mathcal{K})^+-\text{sat}}$ given by the previous corollary to a type-full good ($\geq \text{LS}(\mathcal{K})^+$)-frame with underlying class $\mathcal{K}^{\text{LS}(\mathcal{K})^+-\text{sat}}$. That is, the non-forking relation of the frame can be extended to types over all $\text{LS}(\mathcal{K})^+$ -saturated models.

REFERENCES

- [Bal09] John T. Baldwin, *Categoricity*, University Lecture Series, vol. 50, American Mathematical Society, 2009.
- [BG] Will Boney and Rami Grossberg, *Forking in short and tame AECs*, Preprint. URL: <http://arxiv.org/abs/1306.6562v9>.
- [BGKV] Will Boney, Rami Grossberg, Alexei Kolesnikov, and Sebastien Vasey, *Canonical forking in AECs*, Preprint. URL: <http://arxiv.org/abs/1404.1494v2>.
- [Bon] Will Boney, *Computing the number of types of infinite length*, Notre Dame Journal of Formal Logic, To appear. URL: <http://arxiv.org/abs/1309.4485v2>.
- [Bon14a] ———, *Tameness and extending frames*, Journal of Mathematical Logic **14** (2014), no. 2, 1450007.
- [Bon14b] ———, *Tameness from large cardinal axioms*, The Journal of Symbolic Logic **79** (2014), no. 4, 1092–1119.
- [BVa] Will Boney and Sebastien Vasey, *Categoricity and infinitary logics*, Preprint. URL: <http://arxiv.org/abs/1508.03316v2>.
- [BVb] ———, *Chains of saturated models in AECs*, Preprint. URL: <http://arxiv.org/abs/1503.08781v3>.
- [BVc] ———, *Tameness and frames revisited*, Preprint. URL: <http://arxiv.org/abs/1406.5980v5>.
- [Gro02] Rami Grossberg, *Classification theory for abstract elementary classes*, Contemporary Mathematics **302** (2002), 165–204.
- [GV] Rami Grossberg and Sebastien Vasey, *Superstability in abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1507.04223v2>.

- [GV06a] Rami Grossberg and Monica VanDieren, *Categoricity from one successor cardinal in tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 2, 181–201.
- [GV06b] ———, *Galois-stability for tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 1, 25–49.
- [GV06c] ———, *Shelah’s categoricity conjecture from a successor for tame abstract elementary classes*, The Journal of Symbolic Logic **71** (2006), no. 2, 553–568.
- [GVV] Rami Grossberg, Monica VanDieren, and Andrés Villaveces, *Uniqueness of limit models in classes with amalgamation*, Mathematical Logic Quarterly, To appear. URL: <http://arxiv.org/abs/1507.02118v1>.
- [JS13] Adi Jarden and Saharon Shelah, *Non-forking frames in abstract elementary classes*, Annals of Pure and Applied Logic **164** (2013), 135–191.
- [She99] Saharon Shelah, *Categoricity for abstract classes with amalgamation*, Annals of Pure and Applied Logic **98** (1999), no. 1, 261–294.
- [She01] ———, *Categoricity of an abstract elementary class in two successive cardinals*, Israel Journal of Mathematics **126** (2001), 29–128.
- [She09] ———, *Classification theory for abstract elementary classes*, Studies in Logic: Mathematical logic and foundations, vol. 18, College Publications, 2009.
- [SV99] Saharon Shelah and Andrés Villaveces, *Toward categoricity for classes with no maximal models*, Annals of Pure and Applied Logic **97** (1999), 1–25.
- [Vana] Monica VanDieren, *Superstability and symmetry*, Preprint. URL: <http://arxiv.org/abs/1507.01990v1>.
- [Vanb] ———, *Symmetry and the union of saturated models in superstable abstract elementary classes*, Annals of Pure and Applied Logic, To appear. URL: <http://arxiv.org/abs/1512.01786v2>.
- [Van06] ———, *Categoricity in abstract elementary classes with no maximal models*, Annals of Pure and Applied Logic **141** (2006), 108–147.
- [Van13] ———, *Erratum to "Categoricity in abstract elementary classes with no maximal models" [Ann. Pure Appl. Logic 141 (2006) 108–147]*, Annals of Pure and Applied Logic **164** (2013), no. 2, 131–133.
- [Vasa] Sebastien Vasey, *Forking and superstability in tame AECs*, The Journal of Symbolic Logic, To appear. URL: <http://arxiv.org/abs/1405.7443v3>.
- [Vasb] ———, *Independence in abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1503.01366v5>.
- [Vasc] ———, *Shelah’s eventual categoricity conjecture in universal classes. Part I*, Preprint. URL: <http://arxiv.org/abs/1506.07024v9>.
- [Vasd] ———, *Shelah’s eventual categoricity conjecture in universal classes. Part II*, Preprint. URL: <http://arxiv.org/abs/1602.02633v1>.

DEPARTMENT OF MATHEMATICS, ROBERT MORRIS UNIVERSITY, MOON TOWNSHIP PENNSYLVANIA USA

E-mail address: vandieren@rmu.edu

URL: <https://sites.google.com/a/rmu.edu/vandieren/home>

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PENNSYLVANIA, USA

E-mail address: sebv@cmu.edu

URL: <http://math.cmu.edu/~svasey/>