

INDISCERNIBLE EXTRACTION AND MORLEY SEQUENCES

SEBASTIEN VASEY

ABSTRACT. We present a new proof of the existence of Morley sequences in simple theories. We avoid using the Erdős-Rado theorem and instead use Ramsey's theorem. The proof shows that the basic theory of forking in simple theories can be developed inside $\langle H((2^{2^{|T|}})^+), \in \rangle$, answering a question of Grossberg, Iovino and Lessmann, as well as a question of Baldwin.

1. INTRODUCTION

Shelah [She80, Lemma 9.3] has shown that, in a simple first-order theory T , Morley sequences exist for every type. The proof proceeds by first building an independent sequence of length $\beth_{(2^{|T|})^+}$ for the given type and then using the Erdős-Rado theorem together with Morley's method to extract the desired indiscernibles.

After slightly improving on the length of the original independent sequence [GIL02, Appendix A], Grossberg, Iovino and Lessmann observed that, in contrast, most of the theory of forking in a stable first-order theory T can be carried out inside $\langle H(\chi), \in \rangle$ for $\chi := (2^{2^{|T|}})^+$. The authors then asked whether the same could be said about simple theories, and so in particular whether there was another way to build Morley sequences there.

Similarly and independently, Baldwin [Bal, Question 3.1.9] has asked whether the equivalence between forking and dividing in simple theories could be proven without using the axiom of replacement.

We answer those questions in the affirmative by showing how to build a Morley sequence from any infinite independent sequence. We avoid using cardinals like $\beth_{(2^{|T|})^+}$, whose existence need powerful set-theoretic principles, and use only axioms from “ordinary mathematics”.

Date: March 14, 2014.

Our construction actually works whenever forking has what we call *dual finite character*, a property that holds in simple theories and has seldom been studied before.

This paper was written while working on a Ph.D. under the direction of Rami Grossberg at Carnegie Mellon University and I would like to thank Professor Grossberg for his guidance and assistance in my research in general and in this work specifically. I also thank John Baldwin, José Iovino, Itay Kaplan, Alexei Kolesnikov, and Anand Pillay for valuable comments on earlier versions of this paper.

2. PRELIMINARIES

For the rest of this paper, fix a complete first-order theory T in a language $L(T)$ and work inside its monster model \mathfrak{C} . We write $|T|$ for $|L(T)| + \aleph_0$. We denote by $\text{Fml}(L(T))$ the set of first-order formulas in the language $L(T)$. For a tuple \bar{a} in \mathfrak{C} and ϕ a formula, we write $\models \phi[\bar{a}]$ instead of $\mathfrak{C} \models \phi[\bar{a}]$.

When I is a linearly ordered set, $(\bar{a}_i)_{i \in I}$ are tuples, and $i \in I$, we write $\bar{a}_{<i}$ for $(\bar{a}_j)_{j < i}$. It is often assumed without comments that all the \bar{a}_i s have the same (finite) arity.

One could, in the spirit of reverse mathematics [Fri74], try to find the exact proof-theoretic strength of some of the facts of simple theories, like Harnik did for stable theories [Har85, Har87]. We do not attempt this here and focus on the model theory: For simplicity, we will state and prove our results in ZFC.

We point out, however, that all the results of this paper (except for Fact 5, Corollary 6, and Proposition 7 which are background facts not used anywhere else in the paper) can be formalized in ZC^0 , a much weaker theory which we now define¹:

Definition 1 (ZC^0). Work in the language of set theory with a constant symbol Θ (intended to denote $|T|$). There is a version of ZFC for that language, which we will also denote by ZFC. Following the terminology of [Kun80], let ZC denote ZFC without replacement (but with full comprehension). Let $\text{ZC} - P$ be ZC without power set. Let ZC^0 be $\text{ZC} - P$, together with the following two axioms:

- (1) Θ is an infinite cardinal.
- (2) For any set X of size $\leq \Theta$, $\mathcal{P}(\mathcal{P}(X))$ exists.

¹Although we have not verified this, we suspect ZC^0 could be replaced by a suitably modified version of Woodin's ZFC* [Woo10].

Note that the universe of “ordinary mathematics” $\langle V(\omega + \omega), \in, \aleph_0 \rangle$ is a model of ZC , hence of ZC^0 . Letting $\chi := \left(2^{2^{|T|}}\right)^+$, $\langle H(\chi), \in, |T| \rangle$ is also a model of ZC^0 . When formalizing our statements in ZC^0 , some obvious changes have to be made: for example, we need to work inside local monster models, assume $|T| \leq \Theta$, and similarly bound the sizes of other sets.

We assume the reader is familiar with forking. As a brief reminder, forking is an independence notion, originally developed by Shelah to solve the stability spectrum problem, which has turned out to be central in classification theory. We will use the following definition:

Definition 2 (Forking). Let \bar{b} be a tuple, and let A be a set. A formula $\phi(\bar{x}, \bar{b})$ *divides over* A if there is a sequence $\langle \bar{b}_i \mid i < \omega \rangle$ of indiscernibles over A such that $\bar{b}_0 = \bar{b}$, and $\{\phi(\bar{x}, \bar{b}_i) \mid i < \omega\}$ is inconsistent.

A formula $\phi(\bar{x}, \bar{b})$ *forks over* A if there are formulas $\phi_0(\bar{x}, \bar{b}_0), \dots, \phi_{m-1}(\bar{x}, \bar{b}_{m-1})$, each dividing over A , such that $\phi(\bar{x}, \bar{b}) \vdash \bigvee_{i < m} \phi_i(\bar{x}, \bar{b}_i)$.

A type p *forks over* A if $p \vdash \phi(\bar{x}, \bar{b})$ for some formula $\phi(\bar{x}, \bar{b})$ that forks over A .

The following concepts are central:

Definition 3 (Morley sequence). Let I be a linearly ordered set. Let $\mathbf{I} := \langle \bar{a}_i \mid i \in I \rangle$ be a sequence of finite tuples of the same arity. Let $A \subseteq B$ be sets, and let $p \in S(B)$ be a type that does not fork over A .

\mathbf{I} is said to be an *independent sequence for* p *over* A if:

- (1) For all $i \in I$, $\bar{a}_i \models p$.
- (2) For all $i \in I$, $\text{tp}(\bar{a}_i / B\bar{a}_{<i})$ does not fork over A .

\mathbf{I} is said to be a *Morley sequence for* p *over* A if:

- (1) \mathbf{I} is an independent sequence for p over A .
- (2) \mathbf{I} is indiscernible over B .

The following fact about forking (which holds in all complete first-order theories) follows from the extension property and the compactness theorem:

Fact 4 (Existence of independent sequences). Let $A \subseteq B$ be sets, and let $p \in S(B)$ be a type that does not fork over A . Let I be a linearly ordered set. Then there is an independent sequence $\mathbf{I} := \langle \bar{a}_i \mid i \in I \rangle$ for p over A .

3. INDISCERNIBLE EXTRACTION

Fact 4 tells us it is easy to build independent sequences. What about Morley sequences? If a sufficiently long sequence always contains an indiscernible subsequence, the existence of Morley sequences follows from Fact 4. This is the case in stable theories, but not in general: There are counterexamples among both simple [She85, p. 209] and dependent [KS] theories. Thus a different approach is needed in the unstable case. Shelah observed [She80, Lemma 9.3] the following:

Fact 5 (The indiscernible extraction theorem). Let A be a set, and let I be a linearly ordered set. Let $\gamma := (2^{|T|+|A|})^+$, $\mu := \beth_\gamma$, and let $\langle \bar{a}_j \mid j < \mu \rangle$ be a sequence of finite tuples of the same arity. Then there exists a sequence $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$, indiscernible over A such that:

For any $i_0 < \dots < i_{n-1}$ in I , there exists $j_0 < \dots < j_{n-1} < \mu$ so that $\text{tp}(\bar{b}_{i_0} \dots \bar{b}_{i_{n-1}}/A) = \text{tp}(\bar{a}_{j_0} \dots \bar{a}_{j_{n-1}}/A)$.

Corollary 6 (Existence of Morley sequences in arbitrary theories). Let $A \subseteq B$. Let $p \in S(B)$ be a type that does not fork over A . Let I be a linearly ordered set. Then there is a Morley sequence $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$ for p over A .

Proof sketch. Build a long-enough independent sequence for p over A using Fact 4, then use the monotonicity, finite character and invariance properties of forking to see that the extracted sequence \mathbf{I} given by Fact 5 is as desired. \square

It is shown in [GIL02, Theorem A.2] that the length μ of the original sequence in Fact 5 can be decreased to $\beth_{\delta(|T|+|A|)}$, where $\delta(\lambda)$ is defined to be the least ordinal not definable in the logic $L_{\lambda^+, \omega}$. The proof is set-theoretic. The main idea is to use the Erdős–Rado theorem infinitely many times inside an ill-founded model of (a large fragment of) set theory.

Observe that in the indiscernible extraction theorem, the bound $\mu := \beth_{\delta(|T|+|A|)}$ is optimal:

Proposition 7. For every infinite cardinal λ , and every $\mu < \beth_{\delta(\lambda)}$, there is a theory T with $|T| = \lambda$ such that the indiscernible extraction theorem (with $A = \emptyset$) fails for sequences of length μ .

Proof. Fix λ . Pick $\mu < \beth_{\delta(\lambda)}$. By building on Morley’s idea for lower bounds of Hanf numbers (see [She90], Theorem VII.5.4), we can get a complete theory T of size λ with built-in Skolem functions, and a type

p such that $\text{EC}(T, p)$ (the class of all models of T omitting the type p) contains a model M of size $\geq \mu$, but no model of size $\beth_{\delta(\lambda)}$. Without loss of generality, $\|M\| = \mu$. Let $\{a_j \mid j < \mu\}$ enumerate M . Work in the monster model for T . By assumption, a_j does not realize p , for all $j < \mu$.

Assume for a contradiction there exists a sequence of indiscernibles $\mathbf{I} := \langle b_i \mid i < \beth_{\delta(\lambda)} \rangle$ satisfying the conclusion of the indiscernible extraction theorem. Then in particular, b_i does not realize p for any $i < \beth_{\delta(\lambda)}$. Let $N := \text{EM}(\mathbf{I})$ be the Skolem hull of \mathbf{I} . Then $\|N\| = \beth_{\delta(\lambda)}$, so by construction, $N \notin \text{EC}(T, p)$. This means there is $i_0 < \dots < i_n < \beth_{\delta(\lambda)}$ and a term τ such that $\tau(b_{i_0}, \dots, b_{i_n}) \models p$, so $b_{i_0} \dots b_{i_n} \models q$, where $q(\bar{x}) := p(\tau(\bar{x}))$. But then for some $j_0 < \dots < j_n < \mu$, $a_{j_0} \dots a_{j_n} \models q$, i.e. $\tau(a_{j_0}, \dots, a_{j_n}) \models p$. But $\tau(a_{j_0}, \dots, a_{j_n}) \in M \in \text{EC}(T, p)$, so $\tau(a_{j_0}, \dots, a_{j_n})$ does not realize p , a contradiction. \square

Proposition 7 does not rule out a smaller upper bound μ for particular classes of theories: As was hinted at earlier, if T is stable $\mu := (2^{|T|+|A|})^+$ is enough (see [She90, Theorem I.2.8]). We do not know if there is also a smaller bound for simple theories. Restricting the initial sequence to be independent may also give additional information.

We emphasize once again that Fact 5, Corollary 6, and Proposition 7 are ZFC results. *They will not be used in the rest of this paper.*

We give a weaker version of the indiscernible extraction theorem which holds for $\mu = \omega$. This is the key to obtain existence of Morley sequences. As the proof makes clear, this is really just a slight variation on the Ehrenfeucht-Mostowski theorem.

Theorem 8 (The weak indiscernible extraction theorem). Let A be a set, and let I be a linearly ordered set. Let $\mathbf{J} := \langle \bar{a}_j \mid j < \omega \rangle$ be a sequence of finite tuples of the same arity. Then there exists a sequence $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$, indiscernible over A such that:

For any $i_0 < \dots < i_{n-1}$ in I , for all *finite* $q \subseteq \text{tp}(\bar{b}_{i_0} \dots \bar{b}_{i_{n-1}}/A)$, there exists $j_0 < \dots < j_{n-1} < \omega$ so that $\bar{a}_{j_0} \dots \bar{a}_{j_{n-1}} \models q$.

Proof. By adding new constant symbols to T , we can without loss of generality assume that $A = \emptyset$.

Using the terminology of [TZ12, Definition 5.1.2], let Γ be the Ehrenfeucht-Mostowski type of \mathbf{J} . That is:

$$\Gamma := \{\phi(\bar{x}_0, \dots, \bar{x}_{n-1}) \in \text{Fml}(L(T)) \mid \models \phi[\bar{a}_{j_0}, \dots, \bar{a}_{j_{n-1}}] \text{ for all } j_0 < \dots < j_{n-1} < \omega\}$$

Using Ramsey's theorem (see [TZ12, Lemma 5.1.3]), there is a sequence of indiscernibles $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$ realizing Γ , i.e. for any $n < \omega$, any $\phi(\bar{x}_0, \dots, \bar{x}_{n-1}) \in \Gamma$, and any $i_0 < \dots < i_{n-1}$ in I , we have $\models \phi[\bar{b}_{i_0}, \dots, \bar{b}_{i_{n-1}}]$.

Now, observe that \mathbf{I} is as desired: Let $i_0 < \dots < i_{n-1}$ in I be given, and let q be a finite subset of $\text{tp}(\bar{b}_{i_0} \dots \bar{b}_{i_{n-1}} / \emptyset)$. Without loss of generality, q consists of a single formula $\phi(\bar{x})$. Assume for a contradiction there is no $j_0 < \dots < j_{n-1} < \omega$ such that $\models \phi[\bar{a}_{j_0}, \dots, \bar{a}_{j_{n-1}}]$. By definition, this implies $\neg\phi(\bar{x}) \in \Gamma$, and hence we must have $\models \neg\phi[\bar{b}_{i_0}, \dots, \bar{b}_{i_{n-1}}]$, a contradiction. \square

The reader should be wary of concluding the existence of Morley sequences directly from Theorem 8 and the finite character of forking. Indeed, Theorem 8 does not give us enough invariance to imitate the proof of Corollary 6. In fact, we suspect that a sequence extracted from an independent sequence using Theorem 8 need not in general be Morley.

4. EXTRACTING MORLEY SEQUENCES IN SIMPLE THEORIES

To continue, we become interested in the following property of forking:

Definition 9 (Dual finite character). Forking is said to have *dual finite character (DFC)* if whenever $p := \text{tp}(\bar{c}/A\bar{b})$ forks over A , there exists $\bar{a} \in A$, and a formula $\phi(\bar{x}, \bar{b}, \bar{a}) \in p$ such that $\models \phi[\bar{c}, \bar{b}', \bar{a}]$ implies $\text{tp}(\bar{c}/A\bar{b}')$ forks over A .

Notice that this immediately implies something stronger:

Lemma 10. Assume DFC. Assume $p := \text{tp}(\bar{c}/A\bar{b})$ forks over A , and $\phi(\bar{x}, \bar{b}, \bar{a}) \in p$ is as given by Definition 9. Then $\text{tp}(\bar{c}'/A) = \text{tp}(\bar{c}/A)$ and $\models \phi[\bar{c}', \bar{b}', \bar{a}]$ imply $\text{tp}(\bar{c}'/A\bar{b}')$ forks over A .

Proof. Assume $\text{tp}(\bar{c}'/A) = \text{tp}(\bar{c}/A)$. Let f be an automorphism of \mathfrak{C} fixing A such that $f(\bar{c}') = \bar{c}$. Assume $\models \phi[\bar{c}', \bar{b}', \bar{a}]$. Applying f , $\models \phi[\bar{c}, f(\bar{b}'), \bar{a}]$. Since ϕ witnesses DFC, $\text{tp}(\bar{c}/A f(\bar{b}'))$ forks over A . Applying f^{-1} and using invariance of forking, $\text{tp}(\bar{c}'/A\bar{b}')$ forks over A . \square

Dual finite character is a sort of weak local definability of forking. It says that forking is witnessed by a formula, in a way that allows us to change the *domain* of the type under consideration. This allows us to complete the proof of existence of Morley sequences:

Theorem 11. Assume forking has DFC. Let $A \subseteq B$ be sets. Let $p \in S(B)$ be a type that does not fork over A . Let I be a linearly ordered set. Then there is a Morley sequence $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$ for p over A .

Proof. Use Fact 4 to build an independent sequence $\mathbf{J} := \langle \bar{a}_j \mid j < \omega \rangle$ for p over A . Let $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$ be indiscernible over B , as described by Theorem 8. We claim \mathbf{I} is as required.

It is indiscernible over B , and for every $i \in I$, every \bar{b}_i realizes p : If $\bar{b}_i \not\models p$, fix a formula $\phi(\bar{x}, \bar{b}) \in p$ so that $\models \neg \phi[\bar{b}_i, \bar{b}]$. By the defining property of \mathbf{I} , there exists $j < \omega$ so that $\models \neg \phi[\bar{a}_j, \bar{b}]$, so $\bar{a}_j \not\models p$, a contradiction.

It remains to see that for every $i \in I$, $p_i := \text{tp}(\bar{b}_i/B\bar{b}_{<i})$ does not fork over A . Assume not, and fix $i \in I$ so that p_i forks over A . Fix $\bar{b} \in B$ and $i_0 < \dots < i_{n-1} < i$ such that $p'_i := \text{tp}(\bar{b}_i/A\bar{b}_{i_0} \dots \bar{b}_{i_{n-1}}\bar{b})$ forks over A . Fix $\bar{a} \in A$ and $\phi(\bar{x}, \bar{b}_{i_0} \dots \bar{b}_{i_{n-1}}\bar{b}, \bar{a}) \in p'_i$ witnessing DFC.

Find $j_0 < \dots < j_n < \omega$ such that $\models \phi[\bar{a}_{j_n}, \bar{a}_{j_0} \dots \bar{a}_{j_{n-1}}\bar{b}, \bar{a}]$. Since it has already been observed that $\text{tp}(\bar{a}_{j_n}/A) = \text{tp}(\bar{b}_i/A) = p \upharpoonright A$, Lemma 10 implies that $\text{tp}(\bar{a}_{j_n}/A\bar{a}_{j_0} \dots \bar{a}_{j_{n-1}}\bar{b})$ forks over A , contradicting the independence of \mathbf{J} . \square

A variation of DFC appears as property A.7' in [Mak84], but we haven't found any other occurrence in the literature. Makkai observed that forking symmetry implies DFC, so this is how we will define simplicity:

Definition 12. A first-order theory T is *simple* if its forking has the symmetry property, i.e. whenever $\text{tp}(\bar{c}/A\bar{b})$ forks over A , $\text{tp}(\bar{b}/A\bar{c})$ forks over A .

This is equivalent to T not having the tree property, or to forking having local character [Kim01, Theorem 2.4]. Moreover, the methods of [Adl09] show that the equivalence can be proven in ZC^0 , without using Morley sequences. As an example, we outline why symmetry follows from local character: The key is [Adl09, Theorem 3.6], which shows (without using Morley sequences) that if the D -rank is bounded, then symmetry holds. One can then use [Adl09, Lemma 4.1], which says that the D -rank is bounded if and only if forking has local character.

Lemma 13. Assume T is simple. Then forking has DFC.

Proof. Assume $p := \text{tp}(\bar{c}/A\bar{b})$ fork over A . By symmetry, $q := \text{tp}(\bar{b}/A\bar{c})$ forks over A . Fix $\bar{a} \in A$ and $\psi(\bar{y}, \bar{x}, \bar{z})$ such that $\psi(\bar{y}, \bar{c}, \bar{a}) \in q$ witnesses forking, i.e. if $\models \psi[\bar{b}', \bar{c}, \bar{a}]$ then $\text{tp}(\bar{b}'/A\bar{c})$ forks over A .

Let $\phi(\bar{x}, \bar{y}, \bar{z}) := \psi(\bar{y}, \bar{x}, \bar{z})$. Then $\phi(\bar{x}, \bar{b}, \bar{a}) \in p$, and if $\models \phi[\bar{c}, \bar{b}', \bar{a}]$, then $\models \psi[\bar{b}', \bar{c}, \bar{a}]$, so $\text{tp}(\bar{b}'/A\bar{c})$ forks over A , so by symmetry, $\text{tp}(\bar{c}/A\bar{b}')$ forks over A . This shows $\phi(\bar{x}, \bar{b}, \bar{a})$ witnesses DFC. \square

Corollary 14 (Existence of Morley sequences in simple theories). Assume T is simple. Let $A \subseteq B$ be sets. Let $p \in S(B)$ be a type that does not fork over A . Let I be a linearly ordered set. Then there is a Morley sequence $\mathbf{I} := \langle \bar{b}_i \mid i \in I \rangle$ for p over A .

Proof. Combine Lemma 13 and Theorem 11. \square

We do not know whether forking having DFC is equivalent to simplicity, or whether forking has DFC in a larger class of theories. The following example shows DFC does not always hold:

Example 15. Let T be the first-order theory of dense linear orderings. Let A be the set of nonzero rational numbers. Let $b_0 < c < b_1$ be positive infinitesimals (i.e. greater than zero but smaller than any positive rational). Then $p := \text{tp}(c/Ab_0b_1)$ divides over A , and for any $\phi(x, b_0, b_1) \in p$, one can find nonzero rationals b'_0, b'_1 such that $\models \phi[c, b'_0, b'_1]$. But then $\text{tp}(c/Ab'_0b'_1) = \text{tp}(c/A)$ does not fork over A . Thus forking does not have DFC in T .

We conclude by answering the questions mentioned in the introduction:

- (1) [GIL02, Question A.1] asked whether the first three sections of [GIL02] could be formalized inside $\langle H(\chi), \in \rangle$. Corollary 14 establishes the existence of Morley sequences (Theorem 1.14 there). The only other problematic results are the characterization of forking using the D -rank (Corollary 3.21 there), and Corollary 3.20 preceding it. To formalize these in ZC^0 , one can use the (equivalent) definition of the D -rank given in [Wag00, Definition 2.3.4], together with the proof of [Wag00, Proposition 2.3.9].
- (2) [Bal, Question 3.1.9] asked whether the equivalence between forking and dividing in simple theories could be formalized in ZC . ZC^0 is a subtheory of ZC , and the equivalence between forking and dividing is proven in [GIL02, Theorem 2.5]. Thus (1) also answers Baldwin's question.

REFERENCES

- [Adl09] Hans Adler, *Thorn-forking as local forking*, Journal of Mathematical Logic **9** (2009), no. 1, 21–38.
- [Bal] John T. Baldwin, *How big should the monster model be?*, Version from July 2, 2013. Available online. URL: <http://www.math.uic.edu/~jbaldwin/pub/monster4.pdf>.
- [Fri74] Harvey M. Friedman, *Some systems of second order arithmetic and their use*, Proceedings of the international congress of mathematicians (Ralph D. James, ed.), vol. 1, 1974, pp. 235–242.
- [GIL02] Rami Grossberg, José Iovino, and Olivier Lessmann, *A primer of simple theories*, Archive for Mathematical Logic **41** (2002), no. 6, 541–580.
- [Har85] Victor Harnik, *Stability theory and set existence axioms*, The Journal of Symbolic Logic **50** (1985), no. 1, 123–137.
- [Har87] ———, *Set existence axioms for general (not necessarily countable) stability theory*, Annals of Pure and Applied Logic **34** (1987), 231–243.
- [Kim01] Byunghan Kim, *Simplicity, and stability in there*, The Journal of Symbolic Logic **66** (2001), no. 2, 822–836.
- [KS] Itay Kaplan and Saharon Shelah, *A dependent theory with few indiscernibles*, Preprint. URL: <http://arxiv.org/abs/1010.0388v4>.
- [Kun80] Kenneth Kunen, *Set theory: an introduction to independence proofs*, Studies in logic and the foundations of mathematics, vol. 102, Elsevier, 1980.
- [Mak84] Michael Makkai, *A survey of basic stability theory, with particular emphasis on orthogonality and regular types*, Israel Journal of Mathematics **49** (1984), no. 1, 181–238.
- [She80] Saharon Shelah, *Simple unstable theories*, Annals of Mathematical Logic **19** (1980), no. 3, 177–203.
- [She85] ———, *Monadic logic: Hanf numbers*, Around Classification Theory of Models, Springer-Verlag, 1985, pp. 203–223.
- [She90] ———, *Classification theory and the number of non-isomorphic models*, 2nd ed., Studies in logic and the foundations of mathematics, vol. 92, North-Holland, 1990.
- [TZ12] Katrin Tent and Martin Ziegler, *A course in model theory*, Lecture Notes in Logic, vol. 40, Cambridge University Press, 2012.
- [Wag00] Frank O. Wagner, *Simple theories*, Mathematics and its Applications, Kluwer Academic Publishers, 2000.
- [Woo10] Hugh Woodin, *The axiom of determinacy, forcing axioms, and the non-stationary ideal*, 2nd ed., De Gruyter, 2010.