

MATH 269X - MODEL THEORY FOR ABSTRACT ELEMENTARY CLASSES, SPRING 2018 LECTURE NOTES

SEBASTIEN VASEY

CONTENTS

1.	Notation	1
2.	Universal classes	1
3.	Abstract elementary classes and the presentation theorem	4
4.	Abstract elementary classes with intersections	7
5.	μ -AECs and accessible categories	9
	References	15

1. NOTATION

We use the letter τ for a vocabulary, K for a class of structures. For M a τ -structure, we write $|M|$ for its universe and $\|M\|$ for the cardinality of its universe. We often abuse notation and write for example $a \in M$ instead of $a \in |M|$. We write $M \subseteq N$ for M is a substructure of N .

For I, A sets, we let ${}^I A$ be the set of functions from I to A (we think of them as I -indexed sequences of elements of A). We write \bar{a} for a sequence of elements. We write ${}^{<\infty} A$ for $\bigcup_{\kappa} {}^\kappa A$, where κ ranges over all ordinals. For $\bar{a} \in {}^I A$ and $I_0 \subseteq I$, we write $\bar{a} \upharpoonright I_0$ for the restriction of \bar{a} to I_0 , $\ell(\bar{a}) = I$ (usually used when I is an ordinal), $\text{dom}(\bar{a}) = I$, and $\text{ran}(\bar{a})$ be the range of \bar{a} : the set of elements in the sequence.

For λ a cardinal, we write $[A]^\lambda$ for the subsets of A of cardinality λ . Similarly, $[A]^{<\lambda}$ denotes the subsets of A of cardinality less than λ .

2. UNIVERSAL CLASSES

We start by studying a simple model-theoretic framework. It was first studied by Tarski under the assumption that the vocabulary is finite [Tar54].

Definition 2.1 (Tarski). A *universal class* is a class K of structures in a fixed vocabulary $\tau = \tau(K)$ that is fixed under isomorphisms, substructures, and unions of chains (according to the substructure relation).

Example 2.2. The class of all fields, of all locally finite groups, of all vector spaces over \mathbb{Q} are universal classes. The class of all algebraically closed fields is not (\mathbb{Q} is a subfield of \mathbb{C} which is not algebraically closed).

In the definition, we could have required closure under directed unions instead of just unions of chains. However it turns out that this follows. This is due to Iwamura [Iwa44]:

Exercise 2.3. Let K be a universal class. Let $\langle M_i : i \in I \rangle$ be a directed (according to substructure) system in K . Then $\bigcup_{i \in I} M_i \in K$.

The following is an important basic result about universal classes. We will see it generalizes to AECs.

Definition 2.4. Call a τ -structure M *finitely-generated* if there exists a *finite* subset $A \subseteq |M|$ such that M is the closure of A under its functions.

Theorem 2.5. Let K be a universal class in a vocabulary τ and let M be a τ -structure. The following are equivalent:

- (1) $M \in K$.
- (2) $M_0 \in K$ for all finitely-generated substructures of M .

Proof. If $M \in K$, then by closure under substructure any substructure of it is in K as well. Conversely, if all finitely-generated substructures of M , then they form a directed system in K whose union is M , hence by Exercise 2.3 $M \in K$. \square

There is a correspondence between universal classes and classes axiomatized by universal sentences in infinitary logics. When the vocabulary is finitary (and relational), this was observed by Tarski [Tar54] (in this case universal classes correspond to classes of models of a universal first-order theory). Tarski's proof generalizes.

Definition 2.6. We call an $\mathbb{L}_{\infty, \omega}$ -sentence *universal* if it is of the form $\forall \bar{x} \psi(\bar{x})$, where ψ is quantifier-free.

Theorem 2.7 (Tarski's presentation theorem). Let K be a class of structures in some vocabulary τ . The following are equivalent:

- (1) There is a set Γ of quantifier-free (first-order) types such that K is the class of all τ -structures omitting Γ .
- (2) K is the class of models of a universal $\mathbb{L}_{\infty, \omega}$ theory.
- (3) K is a universal class.

Proof.

- (1) implies (2): Assume that K is the class of τ -structures omitting Γ . For each $p(\bar{x}) \in \Gamma$, let ϕ_p be the sentence $\forall \bar{x} \bigvee_{\psi \in p} \neg \psi(\bar{x})$. Let $T := \{\phi_p \mid p \in \Gamma\}$. It is easy to check that K is the class of models of T .
- (2) implies (3): This is straightforward to check.

- (3) implies (1): Let K_0 be the class of τ -structures that are generated by a finite set (i.e. the class of τ -structures M_0 such that M_0 is the closure under its functions of some finite subset $A \subseteq |M_0|$) and are *not* substructures of any member of K . For each $M_0 \in K_0$, let $p_{M_0}(\bar{x})$ be a type coding it. That is, for any N , if $N \models p[\bar{a}]$, then N is generated by \bar{a} and $N \cong M_0$. Let $\Gamma := \{p_{M_0} \mid M_0 \in K_0\}$. We claim that K is the set of τ -structures omitting Γ . To see this, first notice that any member of K omits Γ by definition of Γ . Conversely, if M omits Γ , then any finitely-generated substructure of M omits Γ , hence is in K . By Theorem 2.5, $M \in K$.

□

Remark 2.8. The proof of Tarski's presentation theorem shows that any universal class K is axiomatized by a universal $\mathbb{L}_{(2^{|\tau(K)|+\aleph_0})^+, \omega}$ sentence.

The following definition was somewhat implicit in the proof above:

Definition 2.9. Let K be a universal class. For $M \in K$ and $A \subseteq |M|$, let $\text{cl}^M(A)$ be the closure of A under the functions of M . Equivalently, $\text{cl}^M(A)$ is the intersection of all $M_0 \subseteq M$ which contain A . Note that $\text{cl}^M(A)$ is a substructure of M , hence is itself in K .

2.1. Tameness in universal classes. It is natural to ask how much of the compactness theorem is lost in the setup of universal classes. We have seen that locally finite groups are universal classes, so clearly we cannot expect the compactness theorem to hold in full generality. However, consider the following interesting consequence of compactness:

Exercise 2.10. Let T be a first-order theory. Let \mathfrak{C} be a monster model for T (i.e. it is λ -saturated, where λ is much bigger than any of the other objects appearing in the statement). Let α be an ordinal and let $\bar{a}, \bar{b} \in {}^\alpha \mathfrak{C}$. The following are equivalent:

- (1) $\mathfrak{C} \models \phi[\bar{a}] \leftrightarrow \phi[\bar{b}]$ for all first-order formulas ϕ .
- (2) There exists an automorphism of \mathfrak{C} taking \bar{a} to \bar{b} .

In other words, syntactic first-order types contain the same information as “semantic” types (defined in terms of orbit of a monster model). Is there a version of such a statement for universal classes? Note that universal classes may fail the amalgamation property (e.g. locally finite groups do [Neu60]), so it may not be possible to build a monster model in this case. Further, first-order types are not the right notion here, since they are not necessarily preserved by substructure. Quantifier-free types should be used and we then have the following result, due to Will Boney, which appears in [Vas17, 3.7].

Theorem 2.11 (Boney). Let K be a universal class. Let $M_1, M_2 \in K$ and let $\bar{a}_\ell \in {}^\alpha M_\ell$, $\ell = 1, 2$. The following are equivalent:

- (1) For any quantifier-free formula ϕ , $M_1 \models \phi[\bar{a}_1]$ if and only if $M_2 \models \phi[\bar{a}_2]$.
- (2) There exists $f : \text{cl}^{M_1}(\bar{a}_1) \cong \text{cl}^{M_2}(\bar{a}_2)$ such that $f(\bar{a}_1) = \bar{a}_2$.

Proof. (2) implies (1) is obvious: quantifier-free formulas are preserved by taking substructures and isomorphisms. We show (1) implies (2). For each $I \subseteq \alpha$ and

$\ell = 1, 2$, write $M_\ell^I := \text{cl}^{M_\ell}(\bar{a}_\ell \restriction I)$. We will build by induction on $|I|$ maps $f_I : M_1^I \cong M_2^I$ such that $f_I(\bar{a}_1 \restriction I) = \bar{a}_2 \restriction I$. This will clearly be enough: take $I = \alpha$.

This is possible: for I finite, M_ℓ^I is coded by its quantifier-free type, hence such a map exists by equality of the quantifier-free types of $\bar{a}_1 \restriction I$ and $\bar{a}_2 \restriction I$. Now if $|I|$ is infinite, observe that for $I_0 \subseteq J_0 \subseteq I$ with $|I_0| + |J_0| < |I|$, $f_{I_0} \subseteq f_{J_0}$. This is because we know that $f_{J_0}(\bar{a}_1 \restriction I_0) = \bar{a}_2 \restriction I_0 = f_{I_0}(\bar{a}_1 \restriction I_0)$ and for any $b \in M_{I_0}$, $b = \sigma(\bar{a}_1 \restriction I_0)$, for σ a term (*this is the key feature of universal classes used in the proof*). Thus $f_{J_0}(b) = \sigma(f_{J_0}(\bar{a}_1 \restriction I_0)) = \sigma(f_{I_0}(\bar{a}_1 \restriction I_0)) = f_{I_0}(\sigma(\bar{a}_1 \restriction I_0)) = f_{I_0}(b)$. Therefore $f_I := \bigcup_{I_0 \subseteq I, |I_0| < |I|}$ is a directed union of a system of isomorphisms, and therefore an isomorphism itself. By definition, it must take $\bar{a}_1 \restriction I$ to $\bar{a}_2 \restriction I$, as desired. \square

Remark 2.12. We could have added a parameter set A contained in both M_1 and M_2 , but this is not needed: one can take \bar{a}_1 and \bar{a}_2 to include an enumeration of it.

We will later see that this result says in technical terms, than “universal classes are fully ($< \aleph_0$)-tame and short over the empty set”. A little less formally, orbital types in universal classes are determined by their finite restrictions.

3. ABSTRACT ELEMENTARY CLASSES AND THE PRESENTATION THEOREM

Not all elementary classes are universal (algebraically closed fields are one example). Thus the framework of universal classes is limited. Shelah introduced in the late 70s AECs as a semantic framework encompassing in particular classes of models of $\mathbb{L}_{\infty, \omega}(Q)$ (the paper that introduced them was [She87], but Shelah lectured on them many years before 1985¹). We will first give the definition of an abstract class.

Definition 3.1. An *abstract class* is a pair $\mathbf{K} = (K, \leq_{\mathbf{K}})$, where K is a class of structures in a fixed vocabulary $\tau = \tau(\mathbf{K})$ and $\leq_{\mathbf{K}}$ is a partial order, $M \leq_{\mathbf{K}} N$ implies $M \subseteq N$, and both K and $\leq_{\mathbf{K}}$ respect isomorphisms (the definition is due to Grossberg). Any abstract class admits a notion of *\mathbf{K} -embedding*: these are functions $f : M \rightarrow N$ such that $f : M \cong f[M]$ and $f[M] \leq_{\mathbf{K}} N$. We sometimes think of \mathbf{K} as the category whose objects are elements in K and whose morphisms are \mathbf{K} -embeddings.

We often do not distinguish between K and \mathbf{K} . For λ a cardinal, we will write \mathbf{K}_λ for the restriction of \mathbf{K} to models of cardinality λ . Similarly define $\mathbf{K}_{\geq \lambda}$ or more generally \mathbf{K}_S , where S is a class of cardinals. We will also use the following notation:

Notation 3.2. For \mathbf{K} an abstract class and $N \in \mathbf{K}$, write $\mathcal{P}_{\mathbf{K}}(N)$ for the set of $M \in \mathbf{K}$ with $M \leq_{\mathbf{K}} N$. Similarly define $\mathcal{P}_{\mathbf{K}_\lambda}(N)$, $\mathcal{P}_{\mathbf{K}_{<\lambda}}(N)$, etc.

For an abstract class \mathbf{K} , we denote by $\mathbb{I}(\mathbf{K})$ the number of models in \mathbf{K} up to isomorphism (i.e. the cardinality of \mathbf{K}/\cong). We write $\mathbb{I}(\mathbf{K}, \lambda)$ instead of $\mathbb{I}(\mathbf{K}_\lambda)$. When $\mathbb{I}(\mathbf{K}) = 1$, we say that \mathbf{K} is *categorical*. We say that \mathbf{K} is *categorical in λ* if \mathbf{K}_λ is categorical, i.e. $\mathbb{I}(\mathbf{K}, \lambda) = 1$.

¹SV: fix?

We say that \mathbf{K} has *amalgamation* if for any $M_0 \leq_{\mathbf{K}} M_\ell$, $\ell = 1, 2$ there is $M_3 \in \mathbf{K}$ and \mathbf{K} -embeddings $f_\ell : M_\ell \rightarrow M_3$, $\ell = 1, 2$. \mathbf{K} has *joint embedding* if any two models can be \mathbf{K} -embedded in a common model. \mathbf{K} has *no maximal models* if for any $M \in \mathbf{K}$ there exists $N \in \mathbf{K}$ with $M \leq_{\mathbf{K}} N$ and $M \neq N$ (we write $M <_{\mathbf{K}} N$). Localized concepts such as *amalgamation in λ* mean that \mathbf{K}_λ has amalgamation.

Definition 3.3 (Shelah). An *abstract elementary class (AEC)* is an abstract class \mathbf{K} in a finitary vocabulary satisfying:

- (1) Coherence: if $M_0, M_1, M_2 \in \mathbf{K}$, $M_0 \subseteq M_1 \leq_{\mathbf{K}} M_2$ and $M_0 \leq_{\mathbf{K}} M_2$, then $M_0 \leq_{\mathbf{K}} M_1$.
- (2) Tarski-Vaught axioms: if δ is a limit ordinal, $\langle M_i : i < \delta \rangle$ is a $\leq_{\mathbf{K}}$ -increasing chain and $M := \bigcup_{i < \delta} M_i$, then:
 - (a) $M \in \mathbf{K}$.
 - (b) $M_j \leq_{\mathbf{K}} M$ for all $j < \delta$.
 - (c) Smoothness: if $N \in \mathbf{K}$ is such that $M_i \leq_{\mathbf{K}} N$ for all $i < \delta$, then $M \leq_{\mathbf{K}} N$.
- (3) Löwenheim-Skolem-Tarski (LST) axiom: there exists a cardinal $\lambda \geq |\tau(\mathbf{K})| + \aleph_0$ such that for any $N \in \mathbf{K}$ and any $A \subseteq |N|$, there exists $M \in \mathcal{P}_{\mathbf{K}_{\lambda+|A|}}(N)$ such that $A \subseteq |M|$ and $M \leq_{\mathbf{K}} N$. We write $\text{LS}(\mathbf{K})$ for the least such λ .

Similarly to Exercise 2.3, the following holds:

Exercise 3.4. Let \mathbf{K} be an AEC. Then the Tarski-Vaught axioms holds for directed systems. That is, let $\langle M_i : i \in I \rangle$ be a $\leq_{\mathbf{K}}$ -directed system. Let $M := \bigcup_{i \in I} M_i$. Then:

- (1) $M \in \mathbf{K}$.
- (2) $M_i \leq_{\mathbf{K}} M$ for all $i \in I$.
- (3) Smoothness: if $N \in \mathbf{K}$ is such that $M_i \leq_{\mathbf{K}} N$ for all $i \in I$, then $M \leq_{\mathbf{K}} N$.

Example 3.5.

- (1) $\mathbf{K} = (\text{Mod}(T), \preceq)$, where T is any first-order theory, is an AEC with $\text{LS}(\mathbf{K}) = |\tau(T)| + \aleph_0$.
- (2) $\mathbf{K} = (K, \subseteq)$, where K is a universal class, is an AEC with $\text{LS}(\mathbf{K}) = |\tau(K)| + \aleph_0$. We may abuse notation and call also such a \mathbf{K} a universal class (or even a universal AEC).
- (3) $\mathbf{K} = (\text{Mod}(\psi), \preceq_\Phi)$, where $\psi \in \mathbb{L}_{\infty, \omega}$ and Φ is a fragment containing ψ , is an AEC with $\text{LS}(\mathbf{K}) \leq |\Phi| + |\tau(\psi)| + \aleph_0$.
- (4) For a fixed infinite cardinal λ , the class of well-orderings of type at most λ^+ ordered by being an initial segment is an AEC \mathbf{K} with $\text{LS}(\mathbf{K}) = \lambda$.
- (5) The class of well-orderings ordered by being an initial segment is not an AEC (it fails the Löwenheim-Skolem-Tarski axiom).
- (6) The class of well-orderings ordered by being a subordering is not an AEC (it fails to be closed under chains).
- (7) See more examples in [BV17, §3].

How are AECs related to universal classes? The following result of Shelah says that any AEC is the reduct of a universal class [She09, 1.9(1)] (the presentation we give combines [Vas17, §2] and [LRVb, 6.4]):

Theorem 3.6 (Shelah's presentation theorem). Let \mathbf{K} be an AEC with vocabulary $\tau = \tau(\mathbf{K})$. Then there exists a universal class \mathbf{K}^+ in an expansion τ^+ of τ with $|\tau^+| = \text{LS}(\mathbf{K})$ and such that the reduct map is a faithful functor from \mathbf{K}^+ into \mathbf{K} which is surjective on objects. In other words:

- (1) For any $M \in \mathbf{K}$, there exists $M^+ \in \mathbf{K}^+$ such that $M^+ \upharpoonright \tau = M$.
- (2) For any $M^+ \subseteq N^+$ both in \mathbf{K}^+ , letting $M := M^+ \upharpoonright \tau$, $N := N^+ \upharpoonright \tau$, we have that $M, N \in \mathbf{K}$ and $M \leq_{\mathbf{K}} N$.

Corollary 3.7. For any AEC \mathbf{K} , there exists a universal $\mathbb{L}_{(2^{\text{LS}(\mathbf{K})})^+, \omega}$ -sentence ψ in an expansion of $\tau(\mathbf{K})$ such that the models in \mathbf{K} are exactly the $\tau(\mathbf{K})$ -reducts of models of ψ .

Proof. By Theorem 3.6, Tarski's presentation Theorem 2.7, and Remark 2.8. \square

To prove Theorem 3.6, the following notion will be useful [Vas17, 2.9]:

Definition 3.8. Let \mathbf{K} be an abstract class and let $N \in K$. We say \mathcal{F} is a *set of Skolem functions for N* if:

- (1) \mathcal{F} is a non-empty set, and each element f of \mathcal{F} is a function from N^n to N , for some $n < \omega$.
- (2) For all $A \subseteq |N|$, $M := \mathcal{F}[A] := \bigcup \{f[A] \mid f \in \mathcal{F}\}$ is such that $M \leq_{\mathbf{K}} N$ and contains A .

Remark 3.9. Let \mathbf{K} be an AEC, let $N \in \mathbf{K}$, \mathcal{F} be a set of Skolem functions for N , and $A \subseteq |N|$. Then (by the smoothness axiom) the closure of A under the functions in \mathcal{F} is also a \mathbf{K} -substructure of N containing A .

Lemma 3.10. Let \mathbf{K} be an AEC. For any $N \in \mathbf{K}$, there exists a set \mathcal{F} of Skolem functions for N with $|\mathcal{F}| = \text{LS}(\mathbf{K})$.

Proof. We build $\langle N_s \mid s \in [N]^{<\aleph_0} \rangle$ such that for each $s, t \in [N]^{<\aleph_0}$:

- (1) $N_s \in \mathcal{P}_{\mathbf{K} \leq \text{LS}(\mathbf{K})}(N)$.
- (2) $s \subseteq |N_s|$.
- (3) $s \subseteq t$ implies $N_s \leq_{\mathbf{K}} N_t$.

This is possible by inductive applications of the LST and coherence axioms. This is enough: for each $s \in [N]^{<\aleph_0}$, let $\{a_i^s : i < \text{LS}(\mathbf{K})\}$ be an enumeration (possibly with repetitions) of N_s . Now for each $n < \omega$, each $i < \text{LS}(\mathbf{K})$, and each $\bar{a} \in {}^n N$, we let $f_i^n(\bar{a})$ be $a_i^{\text{ran}(\bar{a})}$. Let $\mathcal{F} := \{f_i^n : i < \text{LS}(\mathbf{K}), n < \omega\}$. This is as desired: let $A \subseteq |N|$ and let $M := \mathcal{F}[A]$. Then it is easy to check that $M = \bigcup_{s \in [A]^{<\aleph_0}} M_s$. Note that $\langle M_s : s \in [A]^{<\aleph_0} \rangle$ is a directed system and since $N_s \leq_{\mathbf{K}} N$ for all s , it follows from the smoothness axiom that $M \leq_{\mathbf{K}} N$. \square

Proof of Theorem 3.6. Let τ^+ consist of $\tau \cup \{f_i^n : i < \text{LS}(\mathbf{K}), n < \omega\}$, where f_i^n is a new function symbol of arity n . Let \mathbf{K}^+ be class of τ^+ -structures M^+ such that $M_0^+ \upharpoonright \tau \leq_{\mathbf{K}} M^+ \upharpoonright \tau$ for any $M_0^+ \subseteq M^+$. Let $\mathbf{K}^+ := (\mathbf{K}^+, \subseteq)$. It is easy to check that \mathbf{K}^+ is a universal class and by definition, (2) is satisfied. To see (1), let $M \in \mathbf{K}$. By Lemma 3.10, M has a set of Skolem functions \mathcal{F} . Expand M to $M^+ := (M, g)_{g \in \mathcal{F}}$. Then by definition of Skolem functions, $M^+ \in \mathbf{K}^+$. \square

4. ABSTRACT ELEMENTARY CLASSES WITH INTERSECTIONS

The following generalizes Definition 2.9:

Definition 4.1. For \mathbf{K} an AEC, $N \in \mathbf{K}$ and $A \subseteq |N|$, let $\text{cl}^N(A) := \bigcap \{M \in \mathbf{K} \mid M \leq_{\mathbf{K}} N, A \subseteq |M|\}$. We see it as a $\tau(\mathbf{K})$ -substructure of N .

Exercise 4.2. Let \mathbf{K} be an AEC, $M \leq_{\mathbf{K}} N$ be in \mathbf{K} , and $A, B \subseteq |N|$.

- (1) Invariance: If $f : N \cong N'$, then $f[\text{cl}^N(A)] = \text{cl}^{N'}(f[A])$.
- (2) Monotonicity 1: $A \subseteq \text{cl}^N(A)$.
- (3) Monotonicity 2: $A \subseteq B$ implies $\text{cl}^N(A) \subseteq \text{cl}^N(B)$.
- (4) Monotonicity 3: If $A \subseteq |M|$, then $\text{cl}^N(A) \subseteq \text{cl}^M(A)$.
- (5) Idempotence: $\text{cl}^N(M) = M$ and $\text{cl}^N(\text{cl}^N(A)) = \text{cl}^N(A)$.

The notion of having (or admitting) intersections is introduced for AECs in [BS08, 1.2] and further studied in [Vas17, §2].

Definition 4.3. Let \mathbf{K} be an abstract class, $N \in \mathbf{K}$, and $A \subseteq |N|$.

- (1) We say that N has intersections over A if $\text{cl}^N(A) \leq_{\mathbf{K}} M$.
- (2) We say that N has intersections if it has intersections over all $A \subseteq |N|$.
- (3) We say that \mathbf{K} has intersections if all $N \in \mathbf{K}$ have intersections.

Remark 4.4. Formally, $\text{cl}^N(A)$ also depends on \mathbf{K} but usually \mathbf{K} is clear from context. We may write $\text{cl}_{\mathbf{K}}^N(A)$ to make \mathbf{K} explicit.

Example 4.5. Any universal class has intersections. Algebraically closed fields also have intersections. See more examples in [Vas17, 2.6].

Definition 4.6. Let \mathbf{K} be an AEC. Let $M \in \mathbf{K}$ and let $A \subseteq |M|$ be a set. M is *minimal over A* if whenever $M \leq_{\mathbf{K}} N$ and $M' \leq_{\mathbf{K}} N$ contains A , then $M' = M$. M is *minimal over A in N* if $M \leq_{\mathbf{K}} N$ and this holds whenever $N' \leq_{\mathbf{K}} N$.

The following characterization of having intersections is [Vas17, 2.11]:

Theorem 4.7. Let \mathbf{K} be an AEC and let $N \in \mathbf{K}$. The following are equivalent:

- (1) N admits intersections.
- (2) There is an operator $\text{cl} := \text{cl}^N : \mathcal{P}(|N|) \rightarrow \mathcal{P}(|N|)$ such that for all $A, B \subseteq |N|$ and all $M \leq_{\mathbf{K}} N$:
 - (a) $\text{cl}(A) \leq_{\mathbf{K}} N$.
 - (b) $A \subseteq \text{cl}(A)$.
 - (c) $A \subseteq B$ implies $\text{cl}(A) \subseteq \text{cl}(B)$.
 - (d) $\text{cl}(M) = M$.
- (3) For each $A \subseteq |N|$, there is a unique minimal model over A in N .
- (4) There is a set \mathcal{F} of Skolem functions for N such that:
 - (a) $|\mathcal{F}| \leq \text{LS}(K)$.
 - (b) For all $M \leq_{\mathbf{K}} N$, we have $\mathcal{F}[M] = M$.

Moreover the operator $\text{cl}^N : \mathcal{P}(|N|) \rightarrow \mathcal{P}(|N|)$ with the properties in (2) is unique and if it exists then it has the following characterizations:

- $\text{cl}^N(A) = \bigcap \{M \leq_{\mathbf{K}} N \mid A \subseteq |M|\}$.

- $\text{cl}^N(A) = \mathcal{F}[A]$, for any set of Skolem functions \mathcal{F} for N such that $\mathcal{F}[M] = M$ for all $M \leq_{\mathbf{K}} N$.
- $\text{cl}^N(A)$ is the unique minimal model over A in N .

Proof.

- (1) implies (2): Let $\text{cl}^N(A) := \bigcap \{M \leq_{\mathbf{K}} N \mid A \subseteq |M|\}$. Even without hypotheses on N , (2b), (2c), and (2d) are satisfied. Since N admits intersections, (2a) is also satisfied.
- (2) implies (3): Let $A \subseteq |N|$. Let cl be as given by (2). Let $M := \text{cl}(A)$. By (2a), $M \leq_{\mathbf{K}} N$. By (2b), $A \subseteq |M|$. Moreover if $M' \leq_{\mathbf{K}} N$ contains A , then by (2c), $|M| \subseteq |\text{cl}(M')|$ but by (2d), $\text{cl}(M') = M'$. Thus by coherence and (2a) $M \leq_{\mathbf{K}} M'$. This shows both that M is minimal over A and that it is unique.
- (3) implies (4): We slightly change the proof of Lemma 3.10 as follows: in the construction of the N_s 's, let N_s be the unique minimal model over s in N . Now let \mathcal{F} be as obtained by the rest of the construction there. Let $A \subseteq |N|$. We claim that $\mathcal{F}[A]$ is minimal over A in N . This shows in particular that \mathcal{F} is as required.
Let $M := \mathcal{F}[A]$. Since \mathcal{F} is a set of Skolem functions, $M \leq_{\mathbf{K}} N$ and M contains A . Moreover, $M = \bigcup_{s \in [A]^{<\aleph_0}} N_s$. Now if $M' \leq_{\mathbf{K}} N$ contains A , then for all $s \in [A]^{<\aleph_0}$, $s \in [M']^{<\aleph_0}$, so as N_s is minimal over s in N , $N_s \leq_{\mathbf{K}} M'$. It follows that $M \leq_{\mathbf{K}} M'$, so $M = M'$.
- (4) implies (1): Let \mathcal{F} be as given by (4). Let $A \subseteq |N|$. Let $M := \mathcal{F}[A]$. By definition of Skolem functions, M contains A and $M \leq_{\mathbf{K}} N$. We claim that $M = \bigcap \{M' \leq_{\mathbf{K}} N \mid A \subseteq |M'|\}$. Indeed, if $M' \leq_{\mathbf{K}} N$ contains A , then by the hypothesis on \mathcal{F} , $M = \mathcal{F}[A] \subseteq \mathcal{F}[M'] = M'$.

The moreover part follows from the arguments above. \square

Exercise 4.8 ([Vas, 3.6]). Let \mathbf{K} be an AEC. Show that if N has intersections for all $N \in \mathbf{K}_{\leq \text{LS}(\mathbf{K})}$, then \mathbf{K} has intersections.

We obtain the following properties of the closure operator, which complement Exercise 4.2.

Theorem 4.9. Let \mathbf{K} be an AEC with intersections, let $M \leq_{\mathbf{K}} N$ and let $A \subseteq |M|$.

- (1) Monotonicity 3: $\text{cl}^M(A) = \text{cl}^N(A)$.
- (2) (Finite character) For any $b \in \text{cl}^N(A)$, there exists a finite $A_0 \subseteq A$ such that $b \in \text{cl}^N(A_0)$.

Proof. Finite character follows from the characterization of cl^N in terms of Skolem functions (Theorem 4.7). For monotonicity 3, let $M_0 := \text{cl}^N(A)$. We have $M_0 \leq_{\mathbf{K}} N$ since N admits intersections over A . Since $M \leq_{\mathbf{K}} N$ contains A , we must have $|M_0| \subseteq |M|$. By coherence, $M_0 \leq_{\mathbf{K}} M$. Now M_0 is the unique minimal model over A in N , so it must be minimal in M as well, and hence $M_0 = \text{cl}^M(A)$. \square

Remark 4.10. There is a generalization of Tarski's presentation Theorem 2.7 to AECs with intersections [BV].

5. μ -AECs AND ACCESSIBLE CATEGORIES

The following naturally generalizes the definition of an AEC to classes that are only closed under sufficiently directed unions:

Definition 5.1 ([BGL⁺16, 2.2]). Let μ be a regular cardinal. A μ -abstract elementary class (or μ -AEC for short) is an abstract class \mathbf{K} (where we allow here the vocabulary to be infinitary) satisfying:

- (1) Coherence: if $M_0, M_1, M_2 \in \mathbf{K}$, $M_0 \subseteq M_1 \leq_{\mathbf{K}} M_2$ and $M_0 \leq_{\mathbf{K}} M_2$, then $M_0 \leq_{\mathbf{K}} M_1$.
- (2) Tarski-Vaught axioms: if $\langle M_i : i \in I \rangle$ is a μ -directed system (where I is μ -directed if every subset of I of size strictly less than μ has a least upper bound) and $M := \bigcup_{i \in I} M_i$, then:
 - (a) $M \in \mathbf{K}$.
 - (b) $M_i \leq_{\mathbf{K}} M$ for all $i \in I$.
 - (c) Smoothness: if $N \in \mathbf{K}$ is such that $M_i \leq_{\mathbf{K}} N$ for all $i \in I$, then $M \leq_{\mathbf{K}} N$.
- (3) Löwenheim-Skolem-Tarski (LST) axiom: there exists a cardinal $\lambda \geq |\tau(\mathbf{K})| + \mu$ such that $\lambda = \lambda^{<\mu}$ and for any $N \in \mathbf{K}$ and any $A \subseteq |N|$, there exists $M \in \mathcal{P}_{\mathbf{K}_{\lambda+|A|<\mu}}(N)$ such that $A \subseteq |M|$ and $M \leq_{\mathbf{K}} N$. We write $\text{LS}(\mathbf{K})$ for the least such λ .

Remark 5.2. Technically, $\text{LS}(\mathbf{K})$ depends on μ , but this should not cause any problems, so we remove this from the notation.

Note that, in contrast to Exercise 2.3, asking only that the class be closed under *chains* of cofinality at least μ is a significantly weaker condition:

Exercise 5.3 ([AR94, 1.c.(2)]). For $n < \omega$, let P_n be the ordinal $\omega_n + 1$, ordered as usual. Let $Q := \prod_{1 \leq n < \omega} P_n$ and let P be the subposet of Q consisting of those sequences $(x_n)_{n < \omega}$ with only finitely many $n < \omega$ so that $x_n = \omega_n$.

- (1) Check that Q is a complete lattice.
- (2) Check that P is closed (in Q) under joins of chains of uncountable cofinality.
- (3) Check that P is not closed under joins of \aleph_1 -directed sets. *Hint:* Consider $\prod_{1 \leq n < \omega} \omega_n$.

Example 5.4.

- (1) AECs are exactly the \aleph_0 -AECs.
- (2) The class of well-orderings ordered by being a suborder is an \aleph_1 -AEC.
- (3) The class of well-founded models of ZFC, ordered by elementary substructure, is an \aleph_1 -AEC.
- (4) The class of well-orderings ordered by being an initial segment is *not* a μ -AEC for any μ (the LST axiom fails).
- (5) The class of all Banach spaces (ordered by being a complete subspace) is an \aleph_1 -AEC.
- (6) The class of all μ -complete Boolean algebras (ordered by being a subalgebra) is a μ -AEC. However the class of all complete Boolean algebras is not.

- (7) The class of models of any $\mathbb{L}_{\infty, \mu}$ sentence can be made into a μ -AEC by ordering it with elementarity according to a fragment.
- (8) See more examples in [BGL⁺16, §2].

Accessible categories were introduced by Lair [Lai81] (he called them “catégorie modelable”). The standard textbooks on them are [MP89, AR94] (see also the following basic references on category theory [AHS04, Lan98]). One can see them as axiomatizing the category-theoretic essence of classes of models of $\mathbb{L}_{\infty, \infty}$ sentences:

Definition 5.5. Let \mathcal{K} be a category and let λ be a regular cardinal.

- (1) An object M is λ -*presentable* if its hom-functor $\mathcal{K}(M, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits. Put another way, M is λ -presentable if for any morphism $f : M \rightarrow N$ with N a λ -directed colimit $\langle \phi_\alpha : N_\alpha \rightarrow N \rangle$ with diagram maps $\phi_{\alpha\beta} : N_\alpha \rightarrow N_\beta$, f factors essentially uniquely through one of the N_α . That is, $f = \phi_\alpha f_\alpha$ for some $f_\alpha : M \rightarrow N_\alpha$, and if $f = \phi_\beta f_\beta$ as well, there is $\gamma > \alpha, \beta$ such that $\phi_{\gamma\alpha} f_\alpha = \phi_{\gamma\beta} f_\beta$.
- (2) \mathcal{K} is λ -*accessible* if it has λ -directed colimits and \mathcal{K} contains a set S of λ -presentable objects such that every object of \mathcal{K} is isomorphic to a λ -directed colimit of objects in S .
- (3) \mathcal{K} is *accessible* if it is λ' -accessible for some regular cardinal λ' .

Intuitively, an accessible category is a category with all sufficiently directed colimits and such that every object can be written as a highly directed colimit of “small” objects. Here “small” is interpreted in terms of *presentability*, a notion of size that makes sense in any (possibly non-concrete) category. In the category of sets, of course, a set is λ -presentable if and only if its cardinality is less than λ ; in an AEC \mathbf{K} , the same is true for all regular $\lambda > \text{LS}(\mathbf{K})$. More generally:

Exercise 5.6. Let \mathbf{K} be a μ -AEC, let $\lambda = \lambda^{<\mu} \geq \text{LS}(\mathbf{K})$, and let $M \in \mathbf{K}$. Show that M is λ^+ -presentable if and only if $\|M\| \leq \lambda$.

When $\lambda < \lambda^{<\mu}$, presentability still gives a natural notion of size in several categories. For example, in Banach spaces it corresponds to the *density character* [LR17, 3.1]. From Exercise 5.6, it is easy to see the following:

Exercise 5.7. Prove that if \mathbf{K} is a μ -AEC, then it is an $\text{LS}(\mathbf{K})^+$ -accessible category.

There are examples of accessible categories that are *not* (equivalent to) μ -AECs. The simplest one is the category of sets (where the morphisms are functions). The problem is that the morphisms need not be monomorphisms. If we assume that all morphisms are mono, then we will see (Theorem 5.21) that we do in some sense have a μ -AEC. Before proving this, we take a second look at presentability. First, we prove the following generalization of the fact that a small union of small sets is not too big:

Lemma 5.8. Let \mathcal{K} be a λ -accessible category. Then any λ -directed colimit of at most θ -many λ -presentable objects is $(\theta + \lambda)^+$ -presentable.

Proof. Let M be a λ -directed colimit $\langle \phi_i : M_i \rightarrow M, i \in I \rangle$, where $|I| \leq \theta$ and each M_i is λ -presentable. Let $\mu := (\theta + \lambda)^+$. Let $f : M \rightarrow N$ be a morphism, with N a μ -directed colimit of objects $\langle N_j : j \in J \rangle$. Let $f_i := f \phi_i$. By λ -presentability of

M_i , f_i factors (essentially uniquely) through some N_{j_i} , $j_i \in J$. Now there are at most θ -many j_i 's, so since J is μ -directed, there is $j \in J$ with $j_i \leq j$ for all $i \in I$. It follows that f must factor through N_j , showing that M is μ -presentable. \square

Recall that a *retract* is a map $f : M \rightarrow N$ such that there is $g : N \rightarrow M$ so that fg is the identity on N . We also say that N is a *retract* of M . The following is easy to check:

Exercise 5.9. Prove that if $f_1 : M \rightarrow N_1$ and $f_2 : M \rightarrow N_2$ are retracts, as witnessed by g_1 and g_2 , and $g_1 f_1 = g_2 f_2$, then N_1 and N_2 are isomorphic. Conclude that there is only a set (up to isomorphism) of retracts of any given object M .

The following follows from the definition of λ -presentability and playing with morphisms:

Exercise 5.10. Let \mathcal{K} be a λ -accessible category and let S be a set of λ -presentable objects such that any object in \mathcal{K} is a λ -directed colimit of members of S . Prove that any λ -presentable object is a retract of a member of S . Thus \mathcal{K} has only a set (up to isomorphism) of λ -presentable objects. Conversely, show that a retract of a μ -presentable object is μ -presentable, for any regular $\mu \geq \lambda$.

Toward understanding presentability further, we prove a technical lemma saying when an object resolves into a sufficiently directed colimit. We will use the following definitions:

Definition 5.11. For μ a cardinal, μ^* is μ^+ if μ is successor, and μ if μ is limit.

Definition 5.12. For κ, μ infinite cardinals, we say that μ is κ -closed if $\theta^{<\kappa} < \mu$ for all $\theta < \mu$.

Definition 5.13. For λ an uncountable cardinal, we call an object M in a category \mathcal{K} ($< \lambda$)-presentable if it is λ_0 -presentable for some regular $\lambda_0 < \lambda$.

The following is given by the proof of [MP89, 2.3.10]. It is stated as [LRVa, 3.8].

Lemma 5.14. Let $\kappa < \mu \leq \lambda$ be cardinals with κ and μ regular and $\text{cf}(\lambda) \geq \mu$. Let \mathcal{K} be a category with κ -directed colimits. If $M \in \mathcal{K}$ is a κ -directed colimit of ($< \lambda$)-presentable objects and μ is κ -closed, then M is a μ -directed colimit of ($< \lambda + \mu^*$)-presentable objects.

Proof sketch. Suppose that M is a κ -directed colimit of the ($< \lambda$)-presentable objects $\langle M_i : i \in I \rangle$. Since μ is κ -closed, any subset of I of cardinality strictly less than μ is contained inside a κ -directed subset of I of cardinality strictly less than μ . Thus the set \mathbb{P} of all κ -directed subsets of I of cardinality strictly less than μ is μ -directed. For $s \in \mathbb{P}$, let M_s be the colimit of the M_i 's with $i \in s$. Now the induced system $\langle M_s : s \in \mathbb{P} \rangle$ has M as its colimit and:

- (1) μ -directed, since \mathbb{P} is μ -directed.
- (2) Made of ($< \lambda + \mu^*$)-presentable objects.

\square

We deduce several interesting results:

Theorem 5.15. Let \mathcal{K} be a λ -accessible category. If $\mu > \lambda$ is a λ -closed regular cardinal, then \mathcal{K} is μ -accessible.

Proof. Directly from Lemma 5.14. \square

Remark 5.16. We cannot in general remove the assumption that μ is λ -closed from Theorem 5.15 (see [AR94, 2.11]). In fact, for $\mu > 2^{<\lambda}$ regular, the statements “ μ is λ -closed” and “every λ -accessible category is μ -accessible” are equivalent (see [LR17, 4.11] or [LRVa, 2.6]).

Theorem 5.17. Let \mathcal{K} be an accessible category. Then:

- (1) Any object of \mathcal{K} is λ -presentable, for some λ .
- (2) For any regular cardinal λ , there is only a set (up to isomorphism) of λ -presentable objects.

Proof. Let μ be such that \mathcal{K} is μ -accessible. Let S be a set of μ -presentable objects so that any object is isomorphic to a μ -directed colimit of members of S . It follows from Lemma 5.8 that any object must be λ -presentable, for some λ . This proves the first item. For the second, Exercise 5.10 shows that there is only a set of μ -presentable objects. By Theorem 5.15, \mathcal{K} is moreover λ -accessible for arbitrarily large λ , so the result follows. \square

As mentioned before, in the category of sets, an object is λ -presentable if and only if its cardinality is strictly less than λ . Thus the least cardinal λ such that an object is λ -presentable (we call this the *presentability rank*) is always a successor. This cannot be expected to be true in all cases:

Example 5.18 ([LRVa, 6.1]). Let λ be an infinite cardinal and let \mathbf{K} be the AEC of all well-orderings of type at most λ^+ , ordered by initial segment. An object in that category is μ -presentable if and only if it has cofinality at most μ . In particular, taking λ above a weakly inaccessible, there is an object whose presentability rank is a limit.

The following question of Beke and Rosický [BR12] remains open:

Question 5.19. For a fixed accessible category, is every *high-enough* presentability rank a successor?

We can give the following approximation [LRVa, 3.11]:

Theorem 5.20. Let \mathcal{K} be a λ -accessible category. If $\mu > \lambda$ is weakly inaccessible and λ -closed, then any μ -presentable object is $(< \mu)$ -presentable.

Proof. Let M be μ -presentable. By definition, M can be resolved into a λ -directed colimit of λ -presentable objects, hence of $(< \mu)$ -presentables. By Lemma 5.14, M can be resolved into a μ -directed colimit of $(< \mu)$ -presentable objects. By μ -presentability of M , this means that M is a retract of a $(< \mu)$ -presentable object, hence is itself $(< \mu)$ -presentable, as desired. \square

Note that assuming the singular cardinal hypothesis, every weakly inaccessible above $2^{<\lambda}$ is λ -closed. Since Solovay showed that the singular cardinal hypothesis holds above certain large cardinals (see [Sol74] or [Jec03, 20.8]) it follows that Question 5.19 has a positive answer assuming a large cardinal axiom (a proper class of strongly compact cardinals).

5.1. From accessible category to μ -AEC. We now aim to show²:

Theorem 5.21 ([BGL⁺16, 4.5]). For any μ -accessible category \mathcal{K} whose morphisms are monomorphisms, \mathcal{K} is equivalent to a μ -AEC.

Recall that two categories \mathcal{K}_1 and \mathcal{K}_2 are *equivalent* if there is a functor $F : \mathcal{K}_1 \rightarrow \mathcal{K}_2$ which is:

- (1) Full: its restriction to sets of the form $\text{Hom}(M, N)$ is onto $\text{Hom}(FM, FN)$.
- (2) Faithful: its restriction to sets of the form $\text{Hom}(M, N)$ is injective.
- (3) Essentially surjective: any object N in \mathcal{K}_2 is isomorphic to FM for some object M in \mathcal{K}_1 .

This is weaker than an isomorphism of category, but preserves all reasonable category-theoretic notions. Intuitively, we allow isomorphic objects inside the category to be identified. One example to keep in mind is that the category of a single object with only the identity morphism is equivalent (but not isomorphic) to the category of all singleton sets.

The proof of Theorem 5.21 proceeds in two steps. The first shows that \mathcal{K} is equivalent to a certain accessible category of structures. The second shows that this category must actually be a μ -AEC. Let us implement the first step. For τ a vocabulary, we denote by $\text{Emb}(\tau)$ the category whose objects are τ -structures and whose morphisms are injective homomorphisms.

Lemma 5.22 ([BGL⁺16, 4.8]). Let \mathcal{K} be a λ -accessible category whose morphisms are monomorphisms. Then there is a (finitary) vocabulary τ and a functor $E : \mathcal{K} \rightarrow \text{Emb}(\tau)$ which is full and faithful and preserves λ -directed colimits.

Proof. Let \mathcal{K}_0 be a small full subcategory of \mathcal{K} containing (up to isomorphism) all the λ -presentable objects. For each $M \in \mathcal{K}_0$, let S_M be a unary relation symbol and for each morphism f in \mathcal{K}_0 , let \underline{f} be a binary function symbol. The vocabulary τ will consist of all such S_M and \underline{f} . Now map each $M \in \mathcal{K}$ to the following τ -structure EM :

- (1) Its universe are the morphisms $g : M_0 \rightarrow M$, where $M_0 \in \mathcal{K}_0$.
- (2) For each $M_0 \in \mathcal{K}_0$, $S_{M_0}^{EM}$ is the set of morphisms $g : M_0 \rightarrow M$.
- (3) For each morphism $f : M_0 \rightarrow M_1$ of \mathcal{K}_0 , and each $g : M_1 \rightarrow M$, $\underline{f}^{EM}(g) = gf$. When $g \notin S_{M_1}^{EM}$, just let $\underline{f}^{EM}(g) = g$.

Map each morphism $f : M \rightarrow N$ to the function $\bar{f} : EM \rightarrow EN$ given by $\bar{f}(g) = fg$. That E is full and faithful and preserves λ -directed colimits is a long but crucial exercise in diagram chasing (closely related to the Yoneda lemma). For example,

²It was known since Rosický's thesis [Ros83, Ros81] that accessible categories are classes of models of certain $\mathbb{L}_{\infty, \infty}$ sentence, but seeing them as μ -AEC is more direct.

to see that E is full, assume first that $M \in \mathcal{K}_0$. Then id_M is a morphism in \mathcal{K}_0 so given $g : EM \rightarrow EN$, we can let $f := g(\text{id}_M)$ and it turns out that $E(f) = g$. When M is not λ -presentable, resolve it into a λ -directed colimit of λ -presentable objects. \square

The second step shows that any coherent abstract class which looks like an accessible category is in fact a μ -AEC. First, it is not too hard to show (using resolutions into directed systems again) that only a weak version of the LST axiom suffices:

Exercise 5.23. Let \mathbf{K} be an abstract class satisfying all the axioms of a μ -AEC except possibly the LST axiom. Let $\theta \geq \mu + |\tau(\mathbf{K})|$ be such that:

- (1) θ is μ -closed.
- (2) $\text{cf}(\theta) \geq \mu$.
- (3) For any $M \in \mathbf{K}$ and any $A \subseteq |M|$ with $|A| < \theta$, there exists $M_0 \in \mathcal{P}_{\mathbf{K}_{<\theta}}(M)$ with $A \subseteq |M_0|$.

Then \mathbf{K} is a μ -AEC with $\text{LS}(\mathbf{K}) \leq \theta$.

Lemma 5.24. Let \mathbf{K} be an abstract class satisfying the coherence axiom and let μ be a regular cardinal. Assume that \mathbf{K} is μ -accessible and further the μ -directed colimits are concrete (given by unions, i.e. they are the same as in $\text{Emb}(\tau(\mathbf{K}))$). Let C be the set of cardinals λ such that for any $M \in \mathbf{K}$, $\|M\| < \lambda$ if and only if M is $(< \lambda)$ -presentable. Then C is closed unbounded. In particular, \mathbf{K} is a μ -AEC.

Proof. C is clearly closed. Now given any cardinal λ , there is (up to isomorphism) only a set of λ^+ -presentable objects (Theorem 5.17) and only a set of objects of cardinality λ . Thus there is a cardinal λ' such that any λ^+ -presentable object has cardinality strictly less than λ' and any object of cardinality at most λ is $(< \lambda')$ -presentable. Thus given any cardinal λ_0 , we can build an increasing sequence $\langle \lambda_i : i < \omega \rangle$ such that for any $i < \omega$, any λ_i^+ -presentable object has cardinality strictly less than λ_{i+1} and any object of cardinality λ_i is $(< \lambda_{i+1})$ -presentable. Now by construction $\sup_{i < \omega} \lambda_i$ is in C . Thus C is unbounded.

To see the “in particular” part, we have to prove the LST axiom. Pick $\theta \in C$ a limit cardinal such that θ is μ -closed and $\text{cf}(\theta) \geq \mu + |\tau(\mathbf{K})|$. Now let $M \in \mathbf{K}$ and let $A \subseteq |M|$ with $|A| < \theta$ be given. Let $\theta_0 := ((|A| + \aleph_0)^{<\mu})^+$. Note that θ_0 is μ -closed so by Theorem 5.15, \mathbf{K} is θ_0 -accessible. Thus M is a θ_0 -directed colimit of θ_0 -presentable objects $\langle M_i : i \in I \rangle$. Since θ_0 -directed colimits are concrete, this implies that A is contained inside some M_i . Now by definition of C , M_i has cardinality strictly less than θ . This shows that the hypotheses of Exercise 5.23 are satisfied. \square

Proof of Theorem 5.21. Let \mathcal{K} be a μ -accessible category whose morphisms are monomorphisms. By Lemma 5.22, there is a vocabulary τ such that \mathcal{K} is equivalent to a full subcategory of $\text{Emb}(\tau)$ which is closed under μ -directed colimits inside $\text{Emb}(\tau)$. Equivalently, it is closed under μ -directed unions. Closing such a category under isomorphism, we obtain an abstract class \mathbf{K} (the ordering is just substructure) which satisfies the hypotheses of Lemma 5.24, hence is a μ -AEC. \square

REFERENCES

- [AHS04] Jiří Adámek, Horst Herrlich, and George E. Strecker, *Abstract and concrete categories*, online edition ed., 2004, Available from <http://katmat.math.uni-bremen.de/acc/>.
- [AR94] Jiří Adamek and Jiří Rosický, *Locally presentable and accessible categories*, London Math. Society Lecture Notes, Cambridge University Press, 1994.
- [BGL⁺16] Will Boney, Rami Grossberg, Michael J. Lieberman, Jiří Rosický, and Sebastien Vasey, μ -Abstract elementary classes and other generalizations, *The Journal of Pure and Applied Algebra* **220** (2016), no. 9, 3048–3066.
- [BR12] Tibor Beke and Jiří Rosický, *Abstract elementary classes and accessible categories*, *Annals of Pure and Applied Logic* **163** (2012), 2008–2017.
- [BS08] John T. Baldwin and Saharon Shelah, *Examples of non-locality*, *The Journal of Symbolic Logic* **73** (2008), 765–782.
- [BV] Will Boney and Sebastien Vasey, *Structural logic and abstract elementary classes with intersections*, Preprint. URL: <http://arxiv.org/abs/1801.01908v2>.
- [BV17] ———, *A survey on tame abstract elementary classes*, *Beyond first order model theory* (José Iovino, ed.), CRC Press, 2017, pp. 353–427.
- [Iwa44] Tsurane Iwamura, *A lemma on directed sets*, *Zenkoku Shijo Sugaku Danwakai* **262** (1944), 107–111.
- [Jec03] Thomas Jech, *Set theory*, 3rd ed., Springer-Verlag, 2003.
- [Lai81] Christian Lair, *Catégories modelables et catégories esquissables*, *Diagrammes* **6** (1981), L1–L20.
- [Lan98] Saunders Mac Lane, *Categories for the working mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer, 1998.
- [LR17] Michael J. Lieberman and Jiří Rosický, *Metric abstract elementary classes as accessible categories*, *The Journal of Symbolic Logic* **82** (2017), no. 3, 1022–1040.
- [LRVa] Michael J. Lieberman, Jiří Rosický, and Sebastien Vasey, *Internal sizes in μ -abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1708.06782v2>.
- [LRVb] ———, *Universal abstract elementary classes and locally multipresentable categories*, Preprint. URL: <http://arxiv.org/abs/1707.09005v2>.
- [MP89] Michael Makkai and Robert Paré, *Accessible categories: The foundations of categorical model theory*, *Contemporary Mathematics*, vol. 104, American Mathematical Society, 1989.
- [Neu60] B. H. Neumann, *On amalgams of periodic groups*, *Proceedings of the Royal Society (London)*, Series A **255** (1960), 477–489.
- [Ros81] Jiří Rosický, *Categories of models of languages $L_{\kappa, \lambda}$* , Abstracts of the 9th winter school on abstract analysis, Mathematics institute Prague, 1981, pp. 153–157.
- [Ros83] ———, *Representace konkrétních kategorií*, 1983, Doctoral thesis, University of Brno.
- [She87] Saharon Shelah, *Classification of non elementary classes II. Abstract elementary classes*, *Classification Theory* (Chicago, IL, 1985) (John T. Baldwin, ed.), *Lecture Notes in Mathematics*, vol. 1292, Springer-Verlag, 1987, pp. 419–497.
- [She09] ———, *Classification theory for abstract elementary classes*, *Studies in Logic: Mathematical logic and foundations*, vol. 18, College Publications, 2009.
- [Sol74] Robert M. Solovay, *Strongly compact cardinals and the GCH*, *Proceedings of the Tarski Symposium*, *Proc. Sympos. Pure Math.*, vol. 25, American Mathematical Society, 1974, pp. 365–372.
- [Tar54] Alfred Tarski, *Contributions to the theory of models I*, *Indagationes Mathematicae* **16** (1954), 572–581.
- [Vas] Sebastien Vasey, *Quasiminimal abstract elementary classes*, *Archive for Mathematical Logic*, To appear. DOI: 10.1007/s00153-017-0570-7. URL: <https://arxiv.org/abs/1611.07380v7>.
- [Vas17] ———, *Shelah’s eventual categoricity conjecture in universal classes: part I*, *Annals of Pure and Applied Logic* **168** (2017), no. 9, 1609–1642.

E-mail address: sebv@math.harvard.edu

URL: <http://math.harvard.edu/~sebv/>

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MASSACHUSETTS, USA