

# Adding a third normal to CLUBB

Sven Bergmann

University of Wisconsin Milwaukee

May 3, 2024



# Contents

- 1 What is CLUBB?
- 2 Introduction
- 3 Definitions
- 4 Definition of the trinormal distribution,  $P_{tmg}$
- 5 Formulas for higher-order moments
- 6 Asymptotics
- 7 Integration using SymPy

# Contents

## 1 What is CLUBB?

## 2 Introduction

- Motivation to add a third normal component
- Closing turbulence pdes by integration over a pdf
- Derivation of trinormal closures by transformation of binormal closures
- Goal
- Inputs and Outputs

## 3 Definitions

- Normal Distribution
- Variates of the pdf

## 4 Definition of the trinormal distribution, $P_{tmg}$

## 5 Formulas for higher-order moments

## 6 Asymptotics

## 7 Integration using SymPy

# Cloud **L**ayers **U**nified **B**y **B**inormals . . .

. . . is an atmospheric model that tries to predict the weather based on modeling a grid box with a sum of two normal distributions.

# Contents

## 1 What is CLUBB?

## 2 Introduction

- Motivation to add a third normal component
- Closing turbulence pdes by integration over a pdf
- Derivation of trinormal closures by transformation of binormal closures
- Goal
- Inputs and Outputs

## 3 Definitions

- Normal Distribution
- Variates of the pdf

## 4 Definition of the trinormal distribution, $P_{tmg}$

## 5 Formulas for higher-order moments

## 6 Asymptotics

## 7 Integration using SymPy

Nature does not look like this, this is too binormal.




Figure: Binormal plot for two strong up-/downdrafts

$$w_1 = 5, w_2 = -5, \theta_{l1} = 5, \theta_{l2} = -5, \alpha = 0.5, \sigma_w = 2, \sigma_{\theta_{l1}} = 2, \sigma_{\theta_{l2}} = 2.$$

To reduce the bimodality we can increase the widths.




Figure: Binormal plot for two strong up-/downdrafts with increased standard deviations

$$w_1 = 5, w_2 = -5, \theta_{l1} = 5, \theta_{l2} = -5, \alpha = 0.5, \sigma_w = 5, \sigma_{\theta_{l1}} = 5, \sigma_{\theta_{l2}} = 5.$$

# Adding a third normal is more realistic.




Figure: Trinormal plot for two strong up-/downdrafts with varying  $\delta$

$$w_1 = 5, w_2 = -5, \theta_{l1} = 5, \theta_{l2} = -5, \alpha = 0.5, \sigma_w = 2, \sigma_{\theta_{l1}} = 2, \sigma_{\theta_{l2}} = 2, \\ \sigma_{w3} = 2, \sigma_{3\theta_l} = 2, \rho_{w\theta_l} = 0.5.$$



A third normal even allows “weird” shapes.




Figure: Trinormal plot for two strong up-/downdrafts with a third peak in the middle

$$w_1 = 5, w_2 = -5, \theta_{l1} = 5, \theta_{l2} = -5, \alpha = 0.5, \delta = 0.5, \sigma_w = 2, \sigma_{\theta_{l1}} = 2, \\ \sigma_{\theta_{l2}} = 2, \sigma_{w3} = 2, \sigma_{\theta_{l3}} = 2, \rho_{w\theta_l} = 0.5.$$

Consider the following prognostic pde [Lar22, p. 21]:

$$\frac{\partial \overline{w'\theta'_l}}{\partial t} = -\overline{w} \frac{\partial \overline{w'\theta'_l}}{\partial z} - \frac{1}{\rho_s} \frac{\partial \rho_s \overline{w'^2 \theta'_l}}{\partial z} - \overline{w'^2} \frac{\partial \overline{\theta'_l}}{\partial z} - \overline{w'\theta'_l} \frac{\partial \overline{w}}{\partial z} + \dots$$

We need to close the third order moment ( $\overline{w'^2 \theta'_l}$ ) by integration over the pdf.

There already exist closures that assume a binormal pdf [LG05], e.g.

$$\overline{w'^2} = \alpha[(w_1 - \overline{w})^2 + \sigma_w^2] + (1 - \alpha)[(w_2 - \overline{w})^2 + \sigma_w^2]. \quad (2.1)$$

We can transform to a trinormal pdf using the following formulas:

$$\overline{w'^2} \frac{1 - \delta \lambda_w}{1 - \delta} = \overline{w'^2}_{dGn} \quad (2.2)$$

$$\overline{w'^3} \frac{1}{1 - \delta} = \overline{w'^3}_{dGn} \quad (2.3)$$

$$\frac{\overline{w'^3}}{\overline{w'^2}^{3/2}} \frac{(1 - \delta)^{1/2}}{(1 - \lambda_w \delta)^{3/2}} = \frac{\overline{w'^3}_{dGn}}{\overline{w'^2}_{dGn}^{3/2}} \quad (2.4)$$


$$\overline{\theta'_l{}^2} \frac{1 - \delta \lambda_\theta}{1 - \delta} = \overline{\theta'_l{}^2}_{dGn} \quad (2.5)$$

$$\overline{w' \theta'_l} \frac{1 - \delta \lambda_{w\theta}}{1 - \delta} = \overline{w' \theta'_l}_{dGn} \quad (2.6)$$

If we substitute in a formula for  $\lambda_w$  (4.6), which will be explained later on, we get

$$\overline{w'^2} \left( 1 - \delta \frac{\sigma_{w3}^2}{\overline{w'^2}} \right) = (1 - \delta) \overline{w'^2}_{dGn} \quad (2.7)$$

$$\Longleftrightarrow \overline{w'^2} = \overline{w'^2}_{dGn} - \delta \left( \overline{w'^2}_{dGn} - \sigma_{w3}^2 \right) \quad (2.8)$$

Table: 1D Plots for different  $\delta$  and  $\sigma_{w3}$ 

$w_1 = 5$ ,  $w_2 = -5$ ,  $\alpha = 0.2$ ,  $\sigma_w = 2$ . The blue plot represents the third normal, the orange/red one represents the binormal, and the green one represents the mixture. The  $x$  and  $y$  labels and ticks are omitted for clarity.

The goal of this thesis is to verify that all the transformations work out well.

# Forward run (weather forecast)

- Given:  $\overline{w}$ ,  $\overline{w'^2}$ ,  $\overline{w'^3}$ ,  $\overline{\theta_l}$ ,  $\overline{w'\theta'_l}$ ,  $\overline{r_t}$ ,  $\overline{w'r'_t}$ ,  $\overline{\theta_l'^2}$ ,  $\overline{r_t'^2}$ ,  $\overline{r_t'\theta'_l}$ .



# Forward run (weather forecast)

- Given:  $\overline{w}$ ,  $\overline{w'^2}$ ,  $\overline{w'^3}$ ,  $\overline{\theta_l}$ ,  $\overline{w'\theta'_l}$ ,  $\overline{r_t}$ ,  $\overline{w'r'_t}$ ,  $\overline{\theta'^2_l}$ ,  $\overline{r'^2_t}$ ,  $\overline{r'_t\theta'_l}$ .
- Find: Parameters, which describe the shape of the underlying pdf, for ultimately describing higher-order moments, e.g.  $\overline{w'^2\theta'_l}$  in terms of lower-order moments.

# Backward run (verification direction)

- Given: pdf parameters, e.g. mean, standard deviation

# Backward run (verification direction)

- Given: pdf parameters, e.g. mean, standard deviation
- Find: lower- and higher-order moments

# Contents

- 1 What is CLUBB?
- 2 Introduction
  - Motivation to add a third normal component
  - Closing turbulence pdes by integration over a pdf
  - Derivation of trinormal closures by transformation of binormal closures
  - Goal
  - Inputs and Outputs
- 3 Definitions
  - Normal Distribution
  - Variates of the pdf
- 4 Definition of the trinormal distribution,  $P_{tmg}$
- 5 Formulas for higher-order moments
- 6 Asymptotics
- 7 Integration using SymPy

# Multivariate

We say that a random vector  $\mathbf{X}$  is distributed according to a multivariate normal distribution when it has the following joint density function [Ize08, p. 59]:

Definition (pdf of a multivariate normal distribution)

$$f(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-\frac{r}{2}} |\boldsymbol{\Sigma}|^{-\frac{1}{2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right), \mathbf{x} \in \mathbb{R}^r, \quad (3.1)$$

where

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \vdots \\ \mu_r \end{pmatrix} \in \mathbb{R}^r, \text{ and } \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_1^2 & \rho_{12}\sigma_1\sigma_2 & \dots & \rho_{1r}\sigma_1\sigma_r \\ \rho_{12}\sigma_1\sigma_2 & \sigma_2^2 & \dots & \vdots \\ \vdots & \dots & \ddots & \vdots \\ \rho_{1r}\sigma_1\sigma_r & \dots & \dots & \sigma_r^2 \end{pmatrix} \in \mathbb{R}^{r \times r} \quad (3.2)$$

# Moments

We denote the skewness and kurtosis by the following:

$$\mathbb{E}[X^3] = \mathbb{E} \left[ \left( \frac{X - \mu}{\sigma} \right)^3 \right] = \frac{\mathbb{E}[(X - \mu)^3]}{(\mathbb{E}[(X - \mu)^2])^{3/2}} = \frac{\mu_3}{\sigma^3} \quad (3.3)$$

$$\mathbb{E}[X^4] = \mathbb{E} \left[ \left( \frac{X - \mu}{\sigma} \right)^4 \right] = \frac{\mathbb{E}[(X - \mu)^4]}{(\mathbb{E}[(X - \mu)^2])^2} = \frac{\mu_4}{\sigma^4} \quad (3.4)$$

- $w$  - upward wind (or up-/downdraft)
- $r_t$  - total water mixing ratio
- $\theta_l$  - liquid water potential temperature

The variables mostly appear in centered form, e.g.  $w' = w - \bar{w}$ .  
For example  $\overline{w'^2}$  is the centered variance.

# Contents

## 1 What is CLUBB?

## 2 Introduction

- Motivation to add a third normal component
- Closing turbulence pdes by integration over a pdf
- Derivation of trinormal closures by transformation of binormal closures
- Goal
- Inputs and Outputs

## 3 Definitions

- Normal Distribution
- Variates of the pdf

## 4 Definition of the trinormal distribution, $P_{tmg}$

## 5 Formulas for higher-order moments

## 6 Asymptotics

## 7 Integration using SymPy



# Normal Mixture

$$\begin{aligned} P_{tmg}(w, \theta_l, r_t) = & \alpha(1 - \delta)\mathcal{N}(\mu_1, \Sigma_1) \\ & + (1 - \alpha)(1 - \delta)\mathcal{N}(\mu_2, \Sigma_2) \\ & + \delta\mathcal{N}(\mu_3, \Sigma_3), \end{aligned} \tag{4.1}$$

where  $\mathcal{N}$  denotes the multivariate normal distribution,  $\alpha \in (0, 1)$  is the mixture fraction of the binormal, and  $\delta \in [0, 1)$  is the weight of the third normal.

# Mean of first and second component

$$\mu_1 = \begin{pmatrix} w_1 \\ \theta_{l1} \\ r_{t1} \end{pmatrix}, \mu_2 = \begin{pmatrix} w_2 \\ \theta_{l2} \\ r_{t2} \end{pmatrix} \quad (4.2)$$

# Covariance matrices of first and second component

$$\Sigma_1 = \begin{pmatrix} \sigma_w^2 & 0 & 0 \\ 0 & \sigma_{\theta_{l1}}^2 & \rho_{\theta_l r_t} \sigma_{\theta_{l3}} \sigma_{r_t 3} \\ 0 & \rho_{\theta_l r_t} \sigma_{\theta_{l3}} \sigma_{r_t 3} & \sigma_{r_{t1}}^2 \end{pmatrix} \quad (4.3)$$

$$\Sigma_2 = \begin{pmatrix} \sigma_w^2 & 0 & 0 \\ 0 & \sigma_{\theta_{l2}}^2 & \rho_{\theta_l r_t} \sigma_{\theta_{l3}} \sigma_{r_t 3} \\ 0 & \rho_{\theta_l r_t} \sigma_{\theta_{l3}} \sigma_{r_t 3} & \sigma_{r_{t2}}^2 \end{pmatrix} \quad (4.4)$$

# Placing of the third component

We place the third normal component at the mean in order to simplify the math.

$$\mu_3 = \begin{pmatrix} \overline{w} \\ \overline{\theta_l} \\ \overline{r_t} \end{pmatrix}, \text{ and } \Sigma_3 = \begin{pmatrix} \sigma_{w3}^2 & \rho_{w\theta_l3}\sigma_{w3}\sigma_{\theta_l3} & \rho_{wr_t3}\sigma_{w3}\sigma_{r_t3} \\ \rho_{w\theta_l3}\sigma_{w3}\sigma_{\theta_l3} & \sigma_{\theta_l3}^2 & \rho_{\theta_lr_t3}\sigma_{\theta_l3}\sigma_{r_t3} \\ \rho_{wr_t3}\sigma_{w3}\sigma_{r_t3} & \rho_{\theta_lr_t3}\sigma_{\theta_l3}\sigma_{r_t3} & \sigma_{r_t3}^2 \end{pmatrix} \quad (4.5)$$

# Additional definitions

$$\lambda_w \equiv \frac{\sigma_{w3}^2}{w'^2}, \quad \lambda_\theta \equiv \frac{\sigma_{\theta_l 3}^2}{\theta_l'^2}, \quad \lambda_r \equiv \frac{\sigma_{r_t 3}^2}{r_t'^2}, \quad (4.6)$$

$$\lambda_{\theta r} \equiv \frac{\rho_{\theta_l r_t} \sigma_{\theta_l 3} \sigma_{r_t 3}}{r_t' \theta_l'}, \quad \lambda_{w\theta} \equiv \frac{\rho_{w\theta_l} \sigma_{w3} \sigma_{\theta_l 3}}{w' \theta_l'}, \quad \lambda_{wr} \equiv \frac{\rho_{wr_t} \sigma_{w3} \sigma_{r_t 3}}{w' r_t'} \quad (4.7)$$

# Contents

- 1 What is CLUBB?
- 2 Introduction
  - Motivation to add a third normal component
  - Closing turbulence pdes by integration over a pdf
  - Derivation of trinormal closures by transformation of binormal closures
  - Goal
  - Inputs and Outputs
- 3 Definitions
  - Normal Distribution
  - Variates of the pdf
- 4 Definition of the trinormal distribution,  $P_{tmg}$
- 5 Formulas for higher-order moments
- 6 Asymptotics
- 7 Integration using SymPy

$$\begin{aligned}
\overline{w'^4} &= \left(\overline{w'^2}\right)^2 \frac{(1 - \delta\lambda_w)^2}{(1 - \delta)} \left(3\tilde{\sigma}_w^4 + 6(1 - \tilde{\sigma}_w^2) \tilde{\sigma}_w^2 + (1 - \tilde{\sigma}_w^2)^2\right) \\
&+ \frac{1}{(1 - \tilde{\sigma}_w^2)} \frac{1}{(1 - \delta\lambda_w)} \frac{\left(\overline{w'^3}\right)^2}{\overline{w'^2}} \\
&+ \delta 3\lambda_w^2 \left(\overline{w'^2}\right)^2
\end{aligned} \tag{5.1}$$

$$\overline{w'^2 \theta'_l} = \frac{1}{(1 - \tilde{\sigma}_w^2)} \frac{1 - \delta \lambda_{w\theta}}{1 - \delta \lambda_w} \frac{\overline{w'^3}}{\overline{w'^2}} \overline{w' \theta'_l} \quad (5.2)$$



$$\begin{aligned}
\overline{w' \theta_l'^2} &= \frac{2}{3} \frac{(1 - \delta \lambda_{w\theta})^2}{(1 - \delta \lambda_w)^2} \frac{1}{(1 - \tilde{\sigma}_w^2)^2} \frac{\overline{w'^3}}{(\overline{w'^2})^2} \left( \overline{w' \theta_l'} \right)^2 \\
&+ \frac{1}{3} \frac{(1 - \delta \lambda_w)}{(1 - \delta \lambda_{w\theta})} (1 - \tilde{\sigma}_w^2) \frac{\overline{w'^2} \overline{\theta_l'^3}}{\overline{w' \theta_l'}}
\end{aligned} \tag{5.3}$$

# Contents

- 1 What is CLUBB?
- 2 Introduction
  - Motivation to add a third normal component
  - Closing turbulence pdes by integration over a pdf
  - Derivation of trinormal closures by transformation of binormal closures
  - Goal
  - Inputs and Outputs
- 3 Definitions
  - Normal Distribution
  - Variates of the pdf
- 4 Definition of the trinormal distribution,  $P_{tmg}$
- 5 Formulas for higher-order moments
- 6 Asymptotics
- 7 Integration using SymPy

# Limit for $\overline{w'^4}$ as $\delta$ goes to 1

As skewness goes to zero we want the pdf to revert to a single normal ( $\delta \rightarrow 1$ ).

$$\lim_{\delta \rightarrow 1} \left( \overline{w'^4} \right) = \left( \frac{\left( \overline{w'^3} \right)^2}{(1 - \delta \lambda_w) \left( \overline{w'^2} - \frac{\sigma_w^2}{2 - c_1} \right)} \right) + 3 \left( \overline{w'^2} \right)^2 \quad (6.1)$$

# Limit for $\overline{w'^2\theta'_l}$ as $\delta$ goes to 1

As skewness goes to zero we want the pdf to revert to a single normal ( $\delta \rightarrow 1$ ).

$$\lim_{\delta \rightarrow 1} \left( \overline{w'^2\theta'_l} \right) = \frac{(c_1 - 2)^2 \overline{w'\theta'_l} \cdot \overline{w'^3}}{(c_2 - 2) \left( (c_1 - 2) \overline{w'^2} + \sigma_w^2 \right)} \quad (6.2)$$

# Limit for $\overline{w'^2 \theta'_l}$ as $\delta$ goes to 1

As skewness goes to zero we want the pdf to revert to a single normal ( $\delta \rightarrow 1$ ).

$$\lim_{\delta \rightarrow 1} \left( \overline{w'^2 \theta'_l} \right) = \lim_{\delta \rightarrow 1} \left( \frac{2}{3} \frac{(1 - \delta \lambda_{w\theta})^2}{(1 - \delta \lambda_w)^2} \frac{1}{(1 - \tilde{\sigma}_w^2)^2} \frac{\overline{w'^3}}{(\overline{w'^2})^2} \left( \overline{w' \theta'_l} \right)^2 \right. \\ \left. + \frac{1}{3} \frac{(1 - \delta \lambda_w)}{(1 - \delta \lambda_{w\theta})} (1 - \tilde{\sigma}_w^2) \frac{\overline{w'^2} \overline{\theta'_l{}^3}}{\overline{w' \theta'_l}} \right) \quad (6.3)$$

# Contents

## 1 What is CLUBB?

## 2 Introduction

- Motivation to add a third normal component
- Closing turbulence pdes by integration over a pdf
- Derivation of trinormal closures by transformation of binormal closures
- Goal
- Inputs and Outputs

## 3 Definitions

- Normal Distribution
- Variates of the pdf

## 4 Definition of the trinormal distribution, $P_{tmg}$

## 5 Formulas for higher-order moments

## 6 Asymptotics

## 7 Integration using SymPy

We have checked the higher-order moment formulas using SymPy.

**DEMONSTRATION**  
(Analytic integration using SymPy [Meu+17])

# Code to follow along the demonstration I

## Listing: Import statements

```
import sympy as sp
from IPython.display import display
from sympy import abc, oo, Symbol, Integral
from sympy.stats import Normal, density
```

## Listing: Defining symbols

```
sigma_w = Symbol('\sigma_w')
w_1 = Symbol('w_1')
w_2 = Symbol('w_2')
w_bar = Symbol('\overline{w}')
sigma_w_3 = Symbol('\sigma_{w3}')
w_prime_2_bar = Symbol('\overline{w}^{'2'})
```



# Code to follow along the demonstration II

## Listing: Defining the marginals

```
G_1_w = Normal(name='G_1_w', mean=w_1, std=sigma_w)
G_1_w_density = density(G_1_w)(sp.abc.w)
G_2_w = Normal(name='G_2_w', mean=w_2, std=sigma_w)
G_2_w_density = density(G_2_w)(sp.abc.w)
G_3_w = Normal(name='G_3_w', mean=w_bar, std=sigma_w_3)
G_3_w_density = density(G_3_w)(sp.abc.w)
G_w = ((1 - sp.abc.delta) * sp.abc.alpha * G_1_w_density +
        (1 - sp.abc.delta) * (1 - sp.abc.alpha) * G_2_w_density +
        sp.abc.delta * G_3_w_density)
```

## Listing: Defining and displaying the needed integral

```
w_prime_2_bar_int = sp.Integral((sp.abc.w - w_bar) ** 2 * G_w, [sp.abc.w, -oo,
↪ oo])
display(sp.Eq(w_prime_2_bar, w_prime_2_bar_int))
```

# Code to follow along the demonstration III

## Listing: Calculating and printing the integral

```
w_prime_2_bar_int_val = w_prime_2_bar_int.doit(conds='none').simplify()
display(sp.Eq(w_prime_2_bar, w_prime_2_bar_int_val))
```

## Listing: Python function for the second order moment

```
def w_prime_2_bar_check(delta=sp.abc.delta, alpha=sp.abc.alpha, w_1=w_1,
↪ w_2=w_2, w_bar=w_bar, sigma_w=sigma_w, sigma_w_3=sigma_w_3):
    return (((1 - delta) * alpha * ((w_1 - w_bar) ** 2 + sigma_w ** 2))
            + ((1 - delta) * (1 - alpha) * ((w_2 - w_bar) ** 2 + sigma_w ** 2))
            + (delta * sigma_w_3 ** 2))
```

## Listing: Printing the symbolic equation

```
display(sp.Eq(w_prime_2_bar, w_prime_2_bar_check()))
```

# Code to follow along the demonstration IV

Listing: Check if the integral and the given formula are the same

```
display(sp.factor(sp.Eq(w_prime_2_bar_int_val, w_prime_2_bar_check()),  
↪ sp.abc.alpha, sp.abc.delta))
```

# References

- [Ize08] Alan Julian Izenman. *Modern multivariate statistical techniques: regression, classification, and manifold learning*. Springer texts in statistics. OCLC: ocn225427579. New York ; London: Springer, 2008. ISBN: 9780387781884.
- [Lar22] Vincent E. Larson. *CLUBB-SILHS: A parameterization of subgrid variability in the atmosphere*. 2022. arXiv: 1711.03675 [physics.ao-ph].
- [LG05] Vincent E Larson and Jean-Christophe Golaz. “Using probability density functions to derive consistent closure relationships among higher-order moments”. In: *Monthly Weather Review* 133.4 (2005), pp. 1023–1042.
- [Meu+17] Aaron Meurer et al. “SymPy: symbolic computing in Python”. In: *PeerJ Computer Science* 3 (Jan. 2017), e103. ISSN: 2376-5992. DOI: 10.7717/peerj-cs.103. URL: <https://doi.org/10.7717/peerj-cs.103>.

We say that a random variable  $X$  is distributed according to a normal distribution ( $X \sim \mathcal{N}(\mu, \sigma^2)$ ) when it has the following pdf:

Definition (pdf of a normal distribution)

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2} \left(\frac{x - \mu}{\sigma}\right)^2\right) \quad (8.1)$$

Considering that the skewness goes to zero ( $Sk_w \rightarrow 0$ ), we want the pdf to revert to a single normal distribution. Therefore, we need

$$\delta, \lambda_w, \lambda_r, \lambda_\theta, \lambda_{wr}, \lambda_{w\theta}, \lambda_{\theta r} \rightarrow 1. \quad (8.2)$$

In this limit, there are no third-order moments anymore, so they have to go to 0 as well. Also, we want to have that the kurtosis is approaching 3 (value of the kurtosis of a standard normal distribution). To ensure those points, as well as no division by zero in the code, we can define the following properties:

$$\lim_{\delta \rightarrow 1} (1 - \delta) \propto |Sk_w|, \quad (8.3)$$

which means that in the limit as  $\delta \rightarrow 1$ ,  $(1 - \delta)$  should “behave as” the absolute value of the skewness of  $w$ ,

$$0 < \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta \lambda_x}{1 - \delta} \right) < \infty, \quad (8.4)$$

where  $x$  means any of  $w$ ,  $r_t$ , or  $\theta_l$ ,

$$0 < \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta \lambda_x}{1 - \delta \lambda_y} \right) < \infty \quad (8.5)$$

where  $x$  is the same as above and  $y$  means any of  $w$ ,  $r_t$ , or  $\theta_l$ ,  $x \neq y$ ,

$$0 < \lim_{\delta \rightarrow 1} (\tilde{\sigma}_w)^2 = \lim_{\delta \rightarrow 1} \left( \frac{\sigma_w^2}{w'^2} \frac{1 - \delta}{1 - \delta \lambda_w} \right) < 1. \quad (8.6)$$

To ensure (8.3), we can use a linear “fit”, which looks like

$$\lambda_w = \lambda_\theta = \lambda_q = (1 - c_1)\delta + c_1, \quad (8.7)$$

where  $c_1$  is some constant. The fit for the other  $\lambda$ 's is

$$\lambda_{w\theta} = \lambda_{q\theta} = \lambda_{wq} = (1 - c_2)\delta + c_2, \quad (8.8)$$

where again  $c_2$  is some constant. Note, that we already have a definition for the  $\lambda$ 's ((4.6)). This definition is just for the backward run though, because we actually have to choose  $\lambda$  in the forward direction.

If we now look at the limit with the proposed fit ((8.7)), where  $x$  is one of the three variates, we get:

$$\lim_{\delta \rightarrow 1} \left( \frac{1 - \delta \lambda_x}{1 - \delta} \right) = \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta((1 - c_1)\delta + c_1)}{1 - \delta} \right) = \lim_{\delta \rightarrow 1} \left( \frac{1 - ((1 - c_1)\delta^2 + c_1\delta)}{1 - \delta} \right) \quad (8.9)$$

$$= \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta^2 + c_1\delta^2 - c_1\delta}{1 - \delta} \right) = \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta^2 + c_1\delta(\delta - 1)}{1 - \delta} \right) \quad (8.10)$$

$$= \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta^2}{1 - \delta} \right) - \lim_{\delta \rightarrow 1} \left( \frac{c_1\delta(1 - \delta)}{1 - \delta} \right) \quad (8.11)$$

$$(\text{L'Hôpital}) \stackrel{\left[\frac{0}{0}\right]}{=} \lim_{\delta \rightarrow 1} \left( \frac{-2\delta}{-1} \right) - \lim_{\delta \rightarrow 1} (c_1\delta) \quad (8.12)$$

$$= 2 - c_1. \quad (8.13)$$



Then, we can also define the range of  $c_1$ , which should be  $(0, 2)$  because we want to have  $0 < \delta\lambda_w < 1$ . For the reciprocal, we then have

$$\lim_{\delta \rightarrow 1} \left( \frac{1 - \delta}{1 - \delta\lambda_x} \right) = \lim_{\delta \rightarrow 1} \left( \frac{1 - \delta}{1 - \delta^2 + c_1\delta^2 - c_1\delta} \right) \stackrel{[\frac{0}{0}]}{=} \lim_{\delta \rightarrow 1} \left( \frac{-1}{-2\delta + 2c_1\delta - c_1} \right) \quad (8.14)$$

$$= \frac{-1}{-2 + c_1} = \frac{1}{2 - c_1}. \quad (8.15)$$