

Problem Set 1
Math modeling of the atmosphere

Reading: Germano, M., 1992: Turbulence: The filtering approach. *J. Fluid Mech.*, **238**, 325–336..

Problem 1: Spatial filtering

Consider a 1D field

$$f(x'') = \frac{1}{2} (1 - \cos(x'')) \quad (1)$$

that extends indefinitely to the left and the right. In this problem, we'll see how running-mean filtering of f affects the perturbations of f , namely f'' , from the filtered version of f , namely \bar{f} .

Assume that the filter used is a simple box (uniform) filter, extending from $x'' = x - \pi/2$ to $x'' = x + \pi/2$.

Throughout this problem, when you are asked to plot functions, you may sketch the functions by hand or plot them using a computer.

- i) Calculate $\bar{f}(x)$. Show your work.
- ii) Overplot the original function $f(x'')$ and the filtered version \bar{f} .
- iii) Suppose we define $f'' \equiv f(x'') - \bar{f}(x)$. (Note that the definition here uses $\bar{f}(x)$ rather than $\bar{f}(x'')$.) Calculate $\overline{f''}(x)$, showing your work.
- iv) Now suppose we define $f'' \equiv f(x'') - \bar{f}(x'')$. (Note that this time the definition uses $\bar{f}(x'')$.) Calculate $\overline{f''}(x)$ using this new definition, showing your work. Plot $\overline{f''}(x)$.
- v) Explain in physical terms why $\overline{f''}(x)$ from part iii) is different from $\overline{f''}(x)$ from part iv). Which part obeys Reynolds' rules?