

Problem Set 2
Math modeling of the atmosphere

Problem 1: Cloud fraction for a “double delta function” PDF

Consider a volume of air that is half occupied by air with $r_t = r_{t1}$ and half occupied by air with $r_t = r_{t2}$. (Let $r_{t1} > r_{t2}$, and let $\Delta r_t \equiv r_{t1} - r_{t2}$.) Further suppose that the volume has uniform temperature everywhere and that the saturation mixing ratio is r_s . Assume that any vapor in excess of saturation is immediately converted to liquid, as we assumed earlier in the semester for non-precipitating, ice-free clouds.

- (i) Write an expression for cloud fraction, C , in terms of r_{t1} , r_{t2} , r_s , and the Heaviside step function.
- (ii) Write an expression for average liquid water mixing ratio, $\langle r_l \rangle$, in terms of r_{t1} , r_{t2} , r_s , and the Heaviside step function.

Problem 2: Cloud fraction for a triangular PDF

Consider now a volume of air that has a symmetric PDF of r_t , $P(r_t)$, with the shape of an isosceles triangle. The PDF is a piecewise linear function of r_t that becomes non-zero at $r_t = \bar{r}_t - \Delta r_t$, increases linearly to a peak at $r_t = \bar{r}_t$, and then decreases linearly to zero at $r_t = \bar{r}_t + \Delta r_t$. Assume that temperature is constant and that the saturation mixing ratio is r_s . You don't need to write in terms of Heaviside step functions if you don't wish to do so. Instead you can just leave the answer in terms of discrete cases.

- (i) Write the functional form of $P(r_t)$ in terms of \bar{r}_t , Δr_t , and r_t . Recall that the area under $P(r_t)$ must be 1.
- (ii) Find the cloud fraction, C , in terms of \bar{r}_t , Δr_t , and r_s .

Problem 3: Sketch cloud fractions for the double delta and triangular PDFs

Overplot the two cloud fraction formulas that you have found in problems 1 and 2 as a function of \bar{r}_t for fixed r_s and Δr_t . You may sketch by hand, if you wish.

Problem Set 2 Solution Outline

Math modeling of the atmosphere

Problem 1: Cloud fraction and liquid water for a “double delta function” PDF

Consider a volume that is half occupied by air with $r_t = r_{t1}$ and half occupied by air with $r_t = r_{t2}$. (Let $r_{t1} > r_{t2}$, and let $\Delta r_t \equiv r_{t1} - r_{t2}$.) Further suppose that the volume has uniform temperature everywhere and that the saturation mixing ratio is r_s .

(i) Write an expression for cloud fraction, C , in terms of r_{t1} , r_{t2} , r_s , and the Heaviside step function.

$$C = \frac{1}{2}H(r_{t1} - r_s) + \frac{1}{2}H(r_{t2} - r_s) \quad (1)$$

(ii) Write an expression for liquid water mixing ratio, r_l , in terms of r_{t1} , r_{t2} , r_s , and the Heaviside step function.

$$\langle r_l \rangle = \frac{1}{2}(r_{t1} - r_s)H(r_{t1} - r_s) + \frac{1}{2}(r_{t2} - r_s)H(r_{t2} - r_s) \quad (2)$$

Problem 2: Cloud fraction for a triangular PDF

Consider now a volume of air that has a symmetric PDF of r_t , $P(r_t)$, with the shape of an isosceles triangle. The PDF is piecewise linear function of r_t that becomes non-zero at $r_t = \bar{r}_t - \Delta r_t$, increases linearly to a peak at $r_t = \bar{r}_t$, and then decreases linearly to zero at $r_t = \bar{r}_t + \Delta r_t$. Assume that temperature is constant and that the saturation mixing ratio is r_s .

(i) Write the functional form of $P(r_t)$. Recall that the area under $P(r_t)$ must be 1.

$$P(r_t) = \begin{matrix} 0 & r_t < \bar{r}_t - \Delta r_t \end{matrix} \quad (3)$$

$$\frac{r_t - \bar{r}_t}{\Delta r_t^2} + \frac{1}{\Delta r_t} \quad \bar{r}_t - \Delta r_t \leq r_t < \bar{r}_t \quad (4)$$

$$-\frac{r_t - \bar{r}_t}{\Delta r_t^2} + \frac{1}{\Delta r_t} \quad \bar{r}_t \leq r_t < \bar{r}_t + \Delta r_t \quad (5)$$

$$0 \quad r_t \geq \bar{r}_t + \Delta r_t \quad (6)$$

(7)

(ii) Write an expression for cloud fraction, C , in terms of \bar{r}_t , Δr_t , and r_s .

In general,

$$C = \int P(r_t) H(r_t - r_s) dr_t \quad (8)$$

For the triangular PDF,

$$C = \begin{matrix} 0 & \bar{r}_t < r_s - \Delta r_t \end{matrix} \quad (9)$$

$$\frac{1}{2} \left[\frac{r_s - (\bar{r}_t + \Delta r_t)}{\Delta r_t} \right]^2 \quad r_s - \Delta r_t \leq \bar{r}_t < r_s \quad (10)$$

$$1 - \frac{1}{2} \left[\frac{-r_s + (\bar{r}_t - \Delta r_t)}{\Delta r_t} \right]^2 \quad r_s \leq \bar{r}_t < r_s + \Delta r_t \quad (11)$$

$$1 \quad \bar{r}_t \geq r_s + \Delta r_t \quad (12)$$

$$(13)$$

Problem 3: Sketch cloud fractions for the double delta and triangular PDFs

Overplot the two cloud fraction formulas that you have found as a function of \bar{r}_t for fixed r_s and Δr_t . You may sketch by hand, if you wish.

Both formulas for C must asymptote to 0 for low values of \bar{r}_t and asymptote to 1 for high values of \bar{r}_t . However, the triangle PDF yields a smooth function of C , whereas the double delta formula yields discrete steps from $C = 0, 0.5, 1$. These steps are rather unrealistic, which is a drawback of the double delta PDF.