

Multivariate Statistical Analysis

Homework 3

Lucas Fellmeth, Helen Kafka, Sven Bergmann

02/28/24

Problem 1

```
swissbanknotes <- read.csv("Data_csv/SwissBankNotes.csv")
swissbanknotes$class <- factor(swissbanknotes$class)
banknote_type <- swissbanknotes$class
```

(a)

```
swissbanknotes2 <- scale(swissbanknotes[, -7])
summary(swissbanknotes2)
```

```
##      length      height_left      height_right      inner_frame_lower
## Min.      :-2.91060      Min.      :-3.1064      Min.      :-2.3672      Min.      :-1.5350
## 1st Qu.: -0.78608      1st Qu.: -0.6135      1st Qu.: -0.6348      1st Qu.: -0.8428
## Median :  0.01062      Median :  0.2174      Median :  0.1077      Median : -0.2198
## Mean    :  0.00000      Mean    :  0.0000      Mean    :  0.0000      Mean    :  0.0000
## 3rd Qu.:  0.54175      3rd Qu.:  0.7714      3rd Qu.:  0.6645      3rd Qu.:  0.8186
## Max.    :  3.72855      Max.    :  2.4333      Max.    :  2.8299      Max.    :  2.2723
## inner_frame_upper      diagonal
## Min.      :-3.67459      Min.      :-2.32889
## 1st Qu.: -0.68560      1st Qu.: -0.85354
## Median : -0.06289      Median : -0.02907
## Mean    :  0.00000      Mean    :  0.00000
## 3rd Qu.:  0.68435      3rd Qu.:  0.88218
## Max.    :  2.05431      Max.    :  1.66324
```

```
pca_banknotes <- princomp(swissbanknotes2)
summary(pca_banknotes)
```

```
## Importance of components:
##              Comp.1      Comp.2      Comp.3      Comp.4      Comp.5
## Standard deviation      1.7119668  1.1276938  0.9298857  0.66896923  0.51704305
## Proportion of Variance  0.4909264  0.2130140  0.1448388  0.07496145  0.04477948
## Cumulative Proportion  0.4909264  0.7039403  0.8487791  0.92374054  0.96852002
##              Comp.6
## Standard deviation      0.43351526
## Proportion of Variance  0.03147998
## Cumulative Proportion  1.00000000
```

The cumulative proportion of the first two components is 0.704.

(b)

```
pca_banknotes$loadings[, 1]
```

##	length	height_left	height_right	inner_frame_lower
##	0.006987029	-0.467758161	-0.486678705	-0.406758327
##	inner_frame_upper	diagonal		
##	-0.367891118	0.493458317		

Component 1

The largest elements (in absolute value) correspond to height_left, height_right, diagonal length. So it's an index of height / diagonal length, where diagonal length has a negative sign. Observations with large negative component scores ksi_{i1} are shorter banknotes with longer diagonal whereas observations with large positive component scores ksi_{i1} are taller banknotes with a shorter diagonal.

```
pca_banknotes$loadings[, 2]
```

##	length	height_left	height_right	inner_frame_lower
##	0.81549497	0.34196711	0.25245860	-0.26622878
##	inner_frame_upper	diagonal		
##	-0.09148667	0.27394074		


Component 2

The largest element (in absolute value) corresponds to length. So it's an index of banknote length. Observations with large negative component scores ksi_{i2} are shorter banknotes whereas Observations with large positive component scores ksi_{i2} are longer banknotes.

(c)


```
plot(pca_banknotes$scores[, 1], main = "Component Scores of first component",  
     col = banknote_type, pch = 16)  
legend("topright", legend = levels(banknote_type), col = seq_along(banknote_type),  
      pch = 16)
```

Component Scores of first component



```
plot(pca_banknotes$scores[, 2], main = "Component Scores of second component",  
     col = banknote_type, pch = 16)  
legend("bottomleft", legend = levels(banknote_type), col = seq_along(banknote_type),  
       pch = 16)
```

Component Scores of second component



The two groups for counterfeit banknotes and genuine banknotes are pretty well separated for both components.

Problem 2

```
turtles <- read.csv("Data_csv/turtles.csv")
turtle_sex <- factor(turtles$sex)
turtles <- turtles[-1]
```

(a)

Before applying log we check if there are any zero values.

```
which(turtles == 0)
```

```
## integer(0)
```

```
turtles2 <- log(turtles)
turtles2 <- scale(turtles2)
pca_turtles <- princomp(turtles2)
summary(pca_turtles)
```

```
## Importance of components:
```

```
##               Comp.1      Comp.2      Comp.3
## Standard deviation    1.6942473 0.21117868 0.149765480
## Proportion of Variance 0.9771826 0.01518177 0.007635642
## Cumulative Proportion 0.9771826 0.99236436 1.000000000
```

The cumulative proportion of the first two components is 0.992.

(b)

```
pca_turtles$loadings[, 1]
```

```
##      length      width      height
## 0.5785513 0.5784099 0.5750829
```

Component 1

The largest elements (in absolute value) correspond to length, width and height. So it's a positive index of length, width and height. Observations with large negative component scores ksi_{i1} are turtles with smaller length, width and height whereas Observations with large positive component scores ksi_{i1} are turtles with larger length, width and height.

```
pca_turtles$loadings[, 2]
```

```
##      length      width      height
## 0.3936563 0.4195001 -0.8179575
```

Component 2


The largest element (in absolute value) corresponds to height. So it's a negative index of turtle height. Observations with large negative component scores ksi_{i2} are taller turtles whereas observations with large positive component scores ksi_{i2} are shorter turtles.

(c)

```
plot(pca_turtles$scores[, 1], main = "Component Scores of first component",
     col = turtle_sex, pch = 16)
```


```
legend("topright", legend = levels(turtle_sex), col = seq_along(turtle_sex),
      pch = 16)
```

Component Scores of first component



```
plot(pca_turtles$scores[, 2], main = "Component Scores of second component",
     col = turtle_sex, pch = 16)
legend("topleft", legend = levels(turtle_sex), col = seq_along(turtle_sex),
      pch = 16)
```

Component Scores of second component



The two groups for female turtles and male turtles are pretty well separated for both components.