

MTHSTAT 564/564G/764–Time Series Analysis

Spring 2024

Problem Solving Set 3

Please think about the following problems from the textbook in advance of our problem solving sessions on them:

Problem Solving 3

1. Suppose $Y_t = \beta_0 + \beta_1 t + X_t$, where $\{X_t\}$ is a zero-mean stationary series with autocovariance function γ_k and β_0 and β_1 are constants.
 - (a) Show that $\{Y_t\}$ is not stationary, but that $W_t = \nabla Y_t = Y_t - Y_{t-1}$ is stationary.
 - (b) In general, show that if $Y_t = \mu_t + X_t$, where $\{X_t\}$ is a zero-mean stationary series, and μ_t is a polynomial in t of degree d , then $\nabla^m Y_t = \nabla(\nabla^{m-1} Y_t)$ is stationary for $m \geq d$ and nonstationary for $0 \leq m < d$.
2. Let $\{X_t\}$ be a zero-mean, unit-variance stationary process with autocorrelation function ρ_k . Suppose that μ_t is a nonconstant function, and that σ_t is a positive-valued, nonconstant function. The observed series is formed as $Y_t = \mu_t + \sigma_t X_t$.
 - (a) Find the mean and covariance function for the $\{Y_t\}$ process.
 - (b) Show that the autocorrelation function for the $\{Y_t\}$ process depends only on the time lag. Is the $\{Y_t\}$ process stationary?
 - (c) Is it possible to have a time series with a constant mean and with $\text{Corr}(Y_t, Y_{t-k})$ free of t , but with $\{Y_t\}$ not stationary?
3. Suppose $\text{Cov}(X_t, X_{t-k}) = \gamma_k$ is free of t , but that $\mathbb{E}[X_t] = 3t$.
 - (a) Is $\{X_t\}$ stationary?
 - (b) Let $Y_t = 7 - 3t + X_t$. Is $\{Y_t\}$ stationary?
4. Evaluate the mean and covariance function for each of the following processes. In each case, determine whether or not the process is stationary.
 - (a) $Y_t = \theta_0 + te_t$.
 - (b) $W_t = \nabla Y_t$, where Y_t is given in part (a).
 - (c) $Y_t = e_t e_{t-1}$, where $\{e_t\}$ is normally distributed white noise.

5. Let $\{Y_t\}$ be stationary with autocovariance function γ_k . Define the sample variance as

$$S^2 = \frac{1}{n-1} \sum_{t=1}^{n-1} (Y_t - \bar{Y})^2.$$

- (a) First show that

$$\sum_{t=1}^n (Y_t - \mu)^2 = \sum_{t=1}^n (Y_t - \bar{Y})^2 + n(\bar{Y} - \mu)^2.$$

- (b) Use part (a) to show that

$$\mathbb{E}[S^2] = \frac{n}{n-1} \gamma_0 - \frac{n}{n-1} \text{Var}(\bar{Y}) = \gamma_0 - \frac{2}{n-1} \sum_{k=1}^{n-1} \left(1 - \frac{k}{n}\right) \gamma_k$$

using the results of Exercise 2.17 for the last expression.

- (c) If $\{Y_t\}$ is a white noise process with variance γ_0 , show that $\mathbb{E}[S^2] = \gamma_0$.
6. Let $Y_1 = \theta_0 + e_1$, and then for $t > 1$, define Y_t recursively by $Y_t = \theta_0 + Y_{t-1} + e_t$. Here, θ_0 is a constant. The process $\{Y_t\}$ is called a random walk with drift.
- (a) Show that Y_t may be rewritten as $Y_t = t\theta_0 + e_t + e_{t-1} + \dots + e_1$.
- (b) Find the mean function for Y_t .
- (c) Find the autocovariance function for Y_t .
7. Two processes $\{Z_t\}$ and $\{Y_t\}$ are said to be independent if for any time points t_1, t_2, \dots, t_m and s_1, s_2, \dots, s_n , the random variables $\{Z_{t_1}, \dots, Z_{t_m}\}$ are independent of the random variables $\{Y_{s_1}, \dots, Y_{s_n}\}$. Show that if $\{Z_t\}$ and $\{Y_t\}$ are independent stationary processes, then $W_t = Z_t + Y_t$ is stationary.
8. Suppose that $Y_t = R \cos(2\pi(ft + \Phi))$ for $t = 0, \pm 1, \pm 2, \dots$, where $0 < f < \frac{1}{2}$ is a fixed frequency and R and Φ are uncorrelated random variables and with Φ uniformly distributed on $(0, 1)$.
- (a) Show that $\mathbb{E}[Y_t] = 0$ all t .
- (b) Show that the process is stationary with $\gamma_k = \frac{1}{2} \mathbb{E}[R^2] \cos(2\pi fk)$. Use the calculations leading up to Eq 2.3.4 on Page 19.
9. Suppose that $Y_t = R \cos[2\pi(ft + \Phi)]$ for $t = 0, \pm 1, \pm 2, \dots$, where R and Φ are independent random variables and f is a fixed frequency. The phase Φ is assumed to be uniformly distributed on $(0, 1)$, and the amplitude R has a Rayleigh distribution with pdf $f(r) = re^{-\frac{r^2}{2}}$ for $r > 0$. Show that for each time point t , Y_t has a normal distribution. (Hint: Let $Y = R \cos[2\pi(ft + \Phi)]$ and $X = R \sin[2\pi(ft + \Phi)]$. Now find the joint distribution of X and Y . It can also be shown that all of the finite-dimensional distributions are multivariate normal, and hence the process is stationary.)