

More than just a line: Shining light on the flattening radial metallicity gradient of a NIHAO Milky-Way analogue simulation

S. Buder,^{1,2*} and T. Buck,^{3,4}

¹Research School of Astronomy & Astrophysics, Australian National University, ACT 2611, Australia

²Center of Excellence for Astrophysics in Three Dimensions (ASTRO-3D), Australia

³Universität Heidelberg, Interdisziplinäres Zentrum für Wissenschaftliches Rechnen, Im Neuenheimer Feld 205, D-69120 Heidelberg, Germany

⁴Universität Heidelberg, Zentrum für Astronomie, Institut für Theoretische Astrophysik, Albert-Ueberle-Straße 2, D-69120 Heidelberg, Germany

Accepted YYYY Month DD. Received YYYY Month DD

ABSTRACT

Radial metallicity gradients in galaxies are crucial for understanding the processes of galactic formation and its dynamical and chemical evolution. We use young stars and gas of a high-resolution simulation of a NIHAO Milky Way analogue galaxy to analyse four properties of the radial metallicity gradient, namely its linearity and scatter as well as its coherence with radial coverage, azimuth, and age. While a global linear fit with slope traces the gradient well, we find it to be too flat in the inner galaxy and too steep in the outer galaxy. An initially stepper but then flattening second order quadratic function provides a superior fit down to the smallest radial steps, where less regular effects take over. We identify these as **SB: spiral arms or warp?** when tracing them across the Galactic position. The simulation further exhibits a steadily increasing spread in [Fe/H] even among the young stars from 0.01 dex in the inner galaxy to 0.14 dex in at a radius 20 kpc coinciding with decreasing stellar density and increased flaring of stars towards higher heights with little correlation across the small age range of 1 Gyr. We conclude that future Galactic and extragalactic observational studies should test our theoretical prediction of a quadratic gradient function with their data to confirm or reject previous observational claims of a broken linear radial metallicity gradients proposed for the Milky Way and other spiral galaxies, as either function implies different dynamical and chemical galaxy formation and evolution scenarios.

Key words: cosmology: observations – Galaxy: structure – Galaxy: abundances – galaxies: structure – galaxies: abundances

1 INTRODUCTION

Radial metallicity gradients in galaxies, defined as the change in metal abundance with galactocentric radius, offer vital insights into the processes that shape the chemical and dynamical evolution of galaxies. The decrease in metallicity with increasing distance from the Galactic center is well-established both theoretically (Larson 1976; Tinsley 1980; Chiosi 1980) and observationally in the Milky Way (Searle 1971; Janes 1979; Twarog et al. 1997). However, the specific shape and characteristics of this gradient remain somewhat unclear.

The Milky Way, being the only galaxy where we can resolve millions of stars, provides a unique opportunity to study these gradients in detail. Over two decades ago, Chiappini (2002) highlighted the contradictions in the observed shape, magnitude, and evolution of abundance gradients. Recent advancements, such as the *Gaia* mission (Gaia Collaboration et al. 2016), have significantly expanded our observational data, allowing for more detailed studies of these gradients.

For instance, Hogg et al. (2019) created an extensive metallicity map of the Milky Way using APOGEE and *Gaia* data, while Poggio et al. (2022) mapped young stars and found a metallicity excess around spiral arms (see also Zari et al. 2018, 2021; Poggio et al. 2021; Hackshaw et al. 2024). Similarly, Imig et al. (2023, among

others) traced gradients across different stellar populations and ages, emphasizing the importance of considering radial migration effects (Binney & Tremaine 2008; Frankel et al. 2018, 2020).

Historically, radial metallicity gradients have been measured using various stellar populations and gas tracers. Early evidence by Janes (1979) suggested a gradient on the order of

$$\frac{d[\text{Fe}/\text{H}]}{dR} = -0.05 \pm 0.01, \text{dex, kpc}^{-1} \quad (1)$$

for the Milky Way which aligns with more recent measurements (Anders et al. 2017; Hayden et al. 2015). Estimated gradients also seem to be broadly consistent across different stellar tracers, such as young open clusters (e.g. Yong et al. 2012; Cunha et al. 2016; Magrini et al. 2017; Casamiquela et al. 2019; Donor et al. 2020; Spina et al. 2021; Myers et al. 2022), young hot (OB-type) stars (Zari et al. 2018, 2021; Poggio et al. 2021, 2022), field stars close to the Galactic plane (e.g. Bergemann et al. 2014) or Cepheids (Andrievsky et al. 2002a,b; Lemasle et al. 2007, 2013). However, these gradients are accompanied by significant spread in [Fe/H] of 0.1 – 0.15 dex, as noted by Twarog (1980), which may imply a fine structure of the metallicity gradient (see Genovali et al. 2014).

Despite extensive observational efforts, several challenges persist:

- The completeness (or patchiness) of observed datasets remains a fundamental issue (Bergemann et al. 2014).
- How robustly can we use the incomplete data to predict substructure in the gradients, such as the need to actually fit two linear

* E-mail: sven.buder@anu.edu.au

gradients with a break radius at corotation radius (Bresolin et al. 2012, and references therein) or further out (Yong et al. 2012; Donor et al. 2020) - or even more complicated functions (see e.g. Chiappini et al. 2001; Kubryk et al. 2015)?

- Different samples yield varying gradients, potentially due to biases in data or the inclusion of older stars (e.g. Boeche et al. 2013; Allende Prieto et al. 2006; Katz et al. 2011; Hayden et al. 2014; Anders et al. 2014; Vickers et al. 2021; Willett et al. 2023).
- The impact of spiral arm structures (Poggio et al. 2021), the Galactic warp (Lemasle et al. 2022) or bar-driven mixing (Di Matteo et al. 2013) on metallicity gradients is not fully understood.
- Methodologies for fitting linear models to scattered data need re-evaluation (Metha et al. 2021).

Understanding these gradients in the Milky Way is also crucial for extragalactic studies, where spatial resolution limits observations. New instruments like the MUSE integral field spectrograph has enabled a plethora of extragalactic studies to contrast the Milky Way to and techniques like the spectroscopy of HII regions and planetary nebulae have helped to infer gas metallicity distributions in external galaxies (Shaver et al. 1983; Vilchez & Esteban 1996; Rolleston et al. 2000; Bresolin et al. 2012). Recent examples include Sánchez et al. (2014) with CALIFA galaxy observations as well as Mun et al. (2024) and Chen et al. (2024) who use MAGPI galaxy observations to probe for example the effects of spiral arms. Notable is also the scatter that Chen et al. (2023) found for the gas metallicity across galactic radius with TYPHOON observations (see their Figs. 4-6).

From a modelling perspective, galactic chemical evolution models can both test understanding of radial metallicity gradients and make predictions beyond the limited volumes and tracers tested by Milky Way and extragalactic studies. Such galactic chemical evolution models include Chiappini et al. (2001); Minchev et al. (2014); Kubryk et al. (2015); Stanghellini et al. (2015); Matteucci (2001).

New suites of large-scale simulations now allow to gain insights into radial metallicity gradients across a range of simulated galaxies - including Milky Way analogues. While Pilkington et al. (2012) found that such gradients are established by inside-out formation in RaDES simulations, whereas Tissera et al. (2019) used EAGLE simulations to study gradients across different galaxy characteristics, such as mergers and stellar mass. Bellardini et al. (2021) and Bellardini et al. (2022); Graf et al. (2024) compared the radial metallicity gradient and its azimuthal scatter for different simulations of the FIRE suite for gas and stars, respectively. Focusing on the change of the radial metallicity gradient over time, Grand et al. (2015) assessed the scatter of radial metallicity distribution for different stellar ages. Both Ma et al. (2017, see their Fig. 6) with FIRE and Agertz et al. (see their Fig. 9 2021) with VINTERGATAN further assessed the evolution of the radial metallicity gradient for gas and stars. Khoperskov et al. (2023) again focused on the gas and found the scatter of the gas metallicity of $\approx 0.04 - 0.06$ dex at a given galactocentric distance.

This study aims to advance these theoretical insight by addressing four critical scientific questions using a high-resolution simulation of a Milky Way analogue galaxy:

- Linearity of the gradient:* Assess the extent to which the radial metallicity gradient of young stars is linear.
- Scatter in the gradient:* Quantify the expected scatter in the radial metallicity gradient of young stars.
- Coherence of the gradient with position:* Investigate the gradient's variation with radial coverage and azimuth.
- Coherence of the gradient with age:* Determine the reliability of stars as tracers of the gas disk over different ages.

The paper is structured as follows: Section 2 describes the data of our Milky Way analogue NIHAO simulation. Section 3 analyses the linearity of the radial metallicity gradients of the simulation under first of our four objectives. Section 4 then analyses both the scatter and local deviations from the gradient as well as the coherence of the gradient with vertical and azimuthal position as well as age (the remaining three objectives). Section 5 discusses them individually and Section 6 bundles them into an overarching conclusion.

2 DATA: A NIHAO MILKY WAY ANALOGUE SIMULATION

For this project, we use a cosmological zoom-in simulation of a Milky Way analogue (g8.26e11) from the *Numerical Investigation of a Hundred Astronomical Objects* (NIHAO, Wang et al. 2015). The model volume has a total mass (gas, stars and dark matter) of $9.1 \cdot 10^{11} M_{\odot}$. At redshift zero, it contains $6.4 \cdot 10^{11} M_{\odot}$ gas mass and $2.3 \cdot 10^{10} M_{\odot}$ stellar mass with a stellar mass resolution of around $7.5 \cdot 10^3 M_{\odot}$ and was calculated as part of the NIHAO-UHD project (Buck et al. 2020). SB: Tobi: Can we check the numbers here. This is what I get out of pynbody, but it's different to 8.26e11.

Simulations were carried out with the smoothed particle hydrodynamics code Gasoline2 (Wadsley et al. 2017) using cosmological parameters from Planck Collaboration et al. (2014) with initial conditions and energetic feedback descriptions from the NIHAO project (Wang et al. 2015). Zoom-in simulations were then performed as described in detail by Buck et al. (2021) with star formation following Stinson et al. (2006) and energetic feedback following Stinson et al. (2013). We note that this is a slightly different rerun of the same simulation than the one studied by Buder et al. (2024).

Because computational resources still limit the mass resolution of simulations, we are relying on tracer particles that represent simple stellar populations (SSPs) with the same age, overall metallicity and discrete initial mass function (IMF). Buck et al. (2021) have implemented the flexible chemical evolution code CHEMPY (Rybicki et al. 2017) to calculate the chemical yields for the SSPs. In particular, SB: we use the alternative (alt) setup of CHEMPY that assumes a Chabrier (2003) IMF with high-mass slope of $\alpha_{\text{IMF}} = -2.3$ over a mass range of $0.1 - 100 M_{\odot}$ for SSPs across a metallicity range of $Z/Z_{\odot} \in [10^{-5}, 2]$. The code calculates the contribution from asymptotic giant branch (AGB) stars, CCSN across a mass range of $8 - 40 M_{\odot}$, and SNIa with a an exponential function with exponent -1.12 , a delay time of 40 Myr, and a normalization of the SNI rate of -2.9. For each of these nucleosynthetic channels, yields from the following studies are used: Limongi & Chieffi (2018) for CCSN, Seitenzahl et al. (2013) for SNIa, and Karakas & Lugaro (2016) for AGB stars. -> CHECK, since I think we used the new_fit, not alt setup, so Chieffi & Limongi (2004) not Limongi & Chieffi (2018)! Contrary to a previous study by Buder et al. (2024), we take the elemental abundances at face value and do not apply any shifts.

For this project, we limit the simulation data to the main halo by applying PYNBODY's implementation of the Amiga Halo Finder (Knollmann & Knebe 2009) and then reposition and rotate this main halo to be face-on based on the angular momentum with PYNBODY's ANALYSIS.ANGMOM.FACEON module (Pontzen et al. 2013). The main halo has a total half mass radius of 66.0 kpc for all matter and a stellar half mass radius 2.4 kpc. We then further transform the resulting galactocentric Cartesian coordinate (X, Y, Z) and velocities (V_X, V_Y, V_Z) to Cylindrical ones as done in a previous study of this main halo by Buder et al. (2024).

To achieve a roughly similar selection as the observational data




Figure 1. Logarithmic spatial density distribution of stars (upper panels) and gas (lower panels) within $R < 20$ kpc of the NIHAO Milky Way analogue g8.26e11 in galactocentric cartesian and cylindric coordinates. Panel c) shows the influence of selecting only young stars with ages below 1 Gyr.

of the Milky Way (Genovali et al. 2014) and other galaxies (e.g. Chen et al. 2023), we restrict the simulation data to a galactocentric radius of $R_{\text{Gal}} \leq 20$ kpc, as shown in Fig. 1. Similar to the Milky Way (Poggio et al. 2018; Lemasle et al. 2022), we note a warp of the stellar and gaseous disk (see Figs. 1b and 1e, respectively).

To avoid too strong effects of radial migration (Binney & Tremaine 2008; Frankel et al. 2018) we further enforce stars to be younger than 1 Gyr. This selection de-facto limits the vertical range of 99% of stars to $|z| = 1.6$ kpc. The strong influence of this age cut on the vertical distribution of stars in the Milky Way analogue can best be appreciated from the difference of vertical density distributions of stars in Figs. 1b and 1c. We are applying these cuts for all following analyses of the radial metallicity gradient in Section 3.

3 THE LINEARITY OF THE RADIAL METALLICITY GRADIENT IN NIHAO

We show the logarithmic density distribution of star particle iron abundances [Fe/H] across different galactocentric radii R_{Gal} in Fig. 2a. This distribution strongly suggests that gradient is predominantly linear, similar to findings of the Milky Way.

The completeness and amount of simulated data points on a global scale allows us, however, to probe this linearity (or more complex behaviour) in more detail with observations.

Table 1. Global linear gradient fit results with different methods. LINEARREGRESSION is part of the SKLEARN package.

Method	Intercept ($a_0 \pm \sigma_{a_0}$)	Slope ($a_1 \pm \sigma_{a_1}$)
STATSMODELS.API.ODR	0.46229 ± 0.00033	-0.04114 ± 0.00004
SCIPY.ODR	0.46232 ± 0.00033	-0.04114 ± 0.00004
NP.POLYFIT	0.46229 ± 0.00033	-0.04114 ± 0.00004
LINEARREGRESSION	0.46222 ± 0.00019	-0.04114 ± 0.00006

3.1 Global gradient fits

Based on our visual impression of the gradient from Fig. 2a, we first fit a global linear radial metallicity gradient to the data. We report the gradient when using using the LINEARREGRESSION module of SKLEARN.LINEAR_MODEL (Pedregosa et al. 2011) in Fig. 2a as red dashed line, but have confirmed this result within the fitting uncertainties with other fitting methods (see Table 1).

We show the fit residuals in Fig. 2b as density distribution as well as in Fig. 2c as percentile distributions in radial bins of $\Delta R_{\text{Gal}} = 0.5$ kpc. While the density distribution shows a lot of substructure, that we are investigating later in Section 4, we note an increase of the median residuals in Fig. 2c towards inner and outer radii, especially for $R_{\text{Gal}} > 17$ kpc. This suggests a non-linear, that is, quadratic component to the gradient, which we will follow up in Section 3.3.

3.2 The impact of radial coverage on linear fits

While we have gained a useful insight into the global function, observational data will rarely cover the full extend of the stellar disk. Using smaller ranges, observational studies have found hints of piece-wise




Figure 2. Global fits and deviation to the radial metallicity gradient $R - [\text{Fe}/\text{H}]$. Functional forms of the linear (red) and quadratic (orange) lines are shown in the legend. Panel a) shows the underlying data of all data points as logarithmic density and the global fit to them as red dashed line. Panel b) shows the deviation of data from a linear gradient as logarithmic density plot, whereas panel c) shows the 16th and 84th percentile around the median deviation as error bars in $\Delta R_{\text{Gal.}} = 0.5 \text{ kpc}$ bins.

linear gradients with a break radius in them based on a limited radial coverage (e.g. Yong et al. 2012; Donor et al. 2020; Chen et al. 2023).

These results are intriguing, since a quadratic function can to first order be approximated by two linear functions with a break radius. We therefore want to use this simulation to test if the radial coverage may indeed delude us into identifying broken linear gradients.

For this purpose, we are fitting and comparing linear radial metallicity gradients to subsets in galactocentric radius. We test fits across piece-wise linear radial ranges of 0.25, 0.5, 1, 2, 5, 8, 10, 15, and 20 kpc. We show their difference with respect to a global linear fit in Fig. 3, with color-coding and indicating the local gradient slope. In this figure, a horizontal dashed line indicates the same slope as the global fit, whereas the offset of a line from said horizontal dashed line indicates the local deviation from the global gradient. Differences in line slopes are visualising the difference in gradient slopes. We see that all ranges are suggesting more or less significant deviations from a global linear fit. The innermost fit is suggesting a significantly different gradient than the outermost fit. We also note increasing slope differences towards the smallest scales, hinting at local deviations from a global pattern. We follow these up in Section 4, but for now focus on the larger-scale trends.

When directly comparing an inner and outer radius fit, such as between $R_{\text{Gal.}} = 5 - 10 \text{ kpc}$ (thick grey line in Fig. 3) and $R_{\text{Gal.}} = 10 - 15 \text{ kpc}$ (thick black line in Fig. 3), we note a significant change, similar to previous estimates of the Milky Way (e.g. Yong et al. 2012; Lemasle et al. 2008). In our case the gradient estimate changes from $[\text{Fe}/\text{H}](R_{\text{Gal.}}) = 0.459 - 0.043 \cdot R_{\text{Gal.}}$ to $[\text{Fe}/\text{H}](R_{\text{Gal.}}) = 0.383 - 0.035 \cdot R_{\text{Gal.}}$. When looking at linear gradient fits across radial coverage of $\Delta R_{\text{Gal.}} = 5 - 15 \text{ kpc}$ in Fig. 3, the gradient was




Figure 3. Impact of different coverage in galactocentric radius when fitting a linear radial metallicity gradient to young stars. Each horizontal segment is using a different running radial fitting range between 0.25 and 15 kpc as outlined on the right. For better contrast, the global linear fit is subtracted from the local gradient estimates and each line is colored by the gradient slope with a color scale centered around the global fit slope.

steeper (bluer color) for smaller radii and flatter (redder) for larger radii.

While such effects have been seen before for the Milky Way, previous studies have mostly suggested to fit piece-wise linear gradients with a break radius (e.g. Andrievsky et al. 2002a; Yong et al. 2012; Boeche et al. 2013; Hayden et al. 2014; Anders et al. 2017; Donor et al. 2020). Our results from Fig. 3 suggest, however, not a gradient with a break radius, but rather a smoothly evolving flattening gradient, in better agreement with a quadratic function rather than two linear ones. In this section, we are therefore trying to expand the gradient function by a higher order, that is, a second order one.

3.3 Higher-order gradients

A quadratic fit (see orange lines in Fig. 2) results in a slightly steeper linear component of the gradient (from -0.0411 to $-0.0497 \text{ dex kpc}^{-1}$), which is counteracted by the quadratic flattening term of $+0.0005 \text{ dex kpc}^{-2}$. The latter leads to an effective flattening of $-0.172 + 0.200 = 0.028 \text{ dex}$ (linear vs. quadratic terms) at $R_{\text{Gal.}} = 20 \text{ dex}$. While seemingly only a nuisance correction across the large extend of $[\text{Fe}/\text{H}]$ and $R_{\text{Gal.}}$, this quadratic function is in evidently superior to the linear fit. This can be best appreciated in Fig. 2c, where the orange line is tracing the median residuals from the linear function much better across the all radii.

When reproducing the local deviations from the linear global fit (Fig. 3) with a quadratic global fit (Fig. 4), we see that the deviations of the local fits no longer have significant offsets with respect to the horizontal lines, while still showing the increasing slope differences at the local levels (especially on scales below 1 kpc).




Figure 4. Same as Fig. 3, but comparing local deviations to a quadratic global fit. Overall deviations have decreased most notably for innermost and outermost radii.

4 SCATTER AND LOCAL DEVIATIONS FROM THE GRADIENT

Now that we are sufficiently satisfied that our flattening gradient function reproduces the overall shape of the radial metallicity gradient, we are concerned with both the scatter and local slope deviations across the galactocentric radii.

When investigating the change of scatter from the innermost radii to the outermost (see Fig. 2c), we see a steady increase in $1 - \sigma$ spread. This spread increases from

$$\begin{aligned} \sigma[\text{Fe}/\text{H}] &= 0.01 \text{ dex at } R_{\text{Gal.}} = 0.25 \pm 0.25 \text{ kpc to} \\ \sigma[\text{Fe}/\text{H}] &= 0.06 \text{ dex at } R_{\text{Gal.}} = 8.25 \pm 0.25 \text{ kpc and finally reaches} \\ \sigma[\text{Fe}/\text{H}] &= 0.14 \text{ dex at } R_{\text{Gal.}} = 19.75 \pm 0.25 \text{ kpc, the largest } R_{\text{Gal.}}. \end{aligned}$$

When reminding ourselves of the observed significant spread in metallicities of young open cluster at the solar radius beyond observational uncertainty (e.g. Donor et al. 2020; Spina et al. 2021) and our selection of only young (< 1 Gyr) old stars from the simulation, a strong correlation of this scatter with stellar age (introducing the possibility of significant radial migration) should be excluded. At this point, we can imagine that this chemical diversity might be caused by less well mixed gas or non-radial effects (such as vertical or azimuthal ones), which we investigate subsequently.

4.1 Coherence of the gradient with position

To further investigate the increase of scatter and local deviation from a global gradient, we are now looking into deviations with respect to the vertical and azimuthal dimensions.

In Fig. 5 we are looking at the previously identified gradient deviations (lines following the left axis label) on top of the vertical density distribution ($R - z$) of gas (Fig. 5a) and young stars (Fig. 5b).




Figure 5. Local gradient deviations from second lowest row of Fig. 4 for radial gradients in 0.5 kpc steps overlapped on top of the logarithmic density distribution in $R - z$ for $|z| < 3$ kpc of gas (panel a) and stars (panel b).

between $-3 < z < 3$ kpc. While the quickly decreasing amount of young stars (Fig. 5b) at outer radii does not show substructure in the density plots for reasonable bin sizes, we see more substructure for the gaseous component in Fig. 5a. In particular, we see rather minor deviations at small radii (where most stars and gas are close to the plane). At increasing radii, we notice an increase in both the vertical distribution of stars, as well as increasing local gradient deviations. In particular, we note a significant deviation of the slope around $R_{\text{Gal.}} \sim 15$ kpc, where the gradient deviation line is steep and blue (indicating a much steeper gradient at this radius), and we notice a significant overdensity of gas around $z \sim 1$ kpc.

While such an edge-on view of the galaxy may indeed be the only observable one for extra-galactic targets, for example of the GECKOS survey of edge-on galaxies (van de Sande et al. 2023), we have the luxury of being able to analyse the azimuthal direction of our simulated galaxy, too. This seems especially important, as we have seen a significant warping of the gaseous and stellar disk of our simulated galaxy (see Figs. 1b and 1e). While the warp of the stellar disk in Fig. 1b is not as clear, we confirm that both the gas disk and the youngest stars below 1 Gyr are tracing each other across the simulation in both galactocentric radius $R_{\text{Gal.}}$ and height $z_{\text{Gal.}}$ for different sectors in azimuthal direction.

SB: CONTINUE

- Density variation of the radial metallicity gradient $R - [\text{Fe}/\text{H}]$ across 8 different sectors in Fig. 6
- Deviation from global radial metallicity gradient across 8 different sectors in Fig. 7
- Fig. 7f shows impressively, that if you look at a rather narrow range of $R_{\text{gal.}}$, such as $7 < R_{\text{gal.}} < 13$ kpc, the radial metallicity gradient could actually look like it is damping towards larger radii. Actually one can see this effect even better in Fig. 6b,d,e,f,h and i! Is this possibly what is happening in local studies of our galaxy, where




Figure 6. Density variation across 8 different sectors (with color-code visualised in panel a) of the radial metallicity gradient $R - [\text{Fe}/\text{H}]$ across 8 different azimuth ranges (panels b-i). A rotating lighthouse-like GIF animation of the median age and median density of the $R - [\text{Fe}/\text{H}]$ -relation across different azimuths is freely available on a [repository](#).




Figure 7. Density variation of the radial metallicity gradient $R - [\text{Fe}/\text{H}]$ across full stellar disk (panel a) and 8 different sectors (same panel order as for Fig. 6). Linear radial metallicity gradients have been fit globally (red dashed line) and for each sector with colors following the same color code as in Fig. 6.




Figure 8. Tracing young stars' iron abundance $[Fe/H]$ across galactocentric radii $R_{Gal.}$ and height $z_{Gal.}$ across the whole galaxy (panel a) and different azimuthal ranges/sectors (panels b-i).




Figure 9. Tracing young stars' iron abundance $[Fe/H]$ residuals (relative to global gradient) across galactocentric radii $R_{Gal.}$ and height $z_{Gal.}$ across the whole galaxy (panel a) and different azimuthal ranges/sectors (panels b-i).

many researchers are actually fitting 2 linear gradients with a break radius?

- Fig. 8
- Fig. 9
- Idea: Compute gradient over a radius R and then plot this one in color in the xy direction (like hogg/eliers). Maybe 2kpc is the best radius to do that? Can also try 5kpc.

4.2 Coherence of the gradient with age

- Do the same as Fig. 6, but with bins colored by median age: Fig. 10
- We track the scatter across age in Fig. 11.
- The maximum scatter in radius bins varies between 0.073 dex and 0.195 dex across the 0..0.1..1.0 Gyr bins.
- Note the diagonal streaks in each of the small age bins (half a dynamic timescale) -> Could this be related to the spiral arms triggering star formation? How could we further trace this? color this by age or $\varphi_{Gal.}$?
- Test gradient for different ages above 1 Gyr? Maybe not necessary. We see that in Fig. 11k and especially 11l the gradient flattens for larger radii for larger ages, that is, the deviation from a line fitted with young stars becomes positive.




Figure 10. Same as Fig. 7, but colored by median age per bin. A rotating lighthouse-like GIF animation of the median age and median density of the $R - [Fe/H]$ -relation across different azimuths is freely available a [repository](#).

4.3 Young stars vs. gas

After having investigated azimuthal variations of the gradient in Section 4.1 and its change in smaller age steps in Section 4.2, we are now aiming to compare the gradient of the youngest stars to that of the gas in the galaxy at redshift zero.

- Fig. 5
- Follow up Fig. 12 by looking at the stellar and gas $[Fe/H]$ for these regions!
- Talk about significant overdensities in Fig. 12, e.g. 7kpc for 0-45deg, 7, 11, 16 kpc for 135-180 or 9-11 for 270-315. Should we follow up the radial gas metallicity gradient as well!?
- Comparing Fig. 12 with Fig. 7: overlap these figures? looks like sometimes the Δz tracks the $\Delta[Fe/H]$, e.g. panels f? Also in oposite direction with panels b and i. So maybe better trace $\Delta|z|$ vs. $\Delta[Fe/H]$! Can we use this to identify a warp?
- Also include gas metallicity gradient: Fig. 13. Gas gradient is interestingly more linear!

5 DISCUSSION

Comparison to [Minchev et al. \(2014\)](#), see their Fig. 10: They see significant decrease of stars towards larger radii (note to myself: their bins are not scatter, but number density in Fig. 10). How does this compare to our number densities?

5.1 Linearity of the gradient

SB: How are the linear metallicity gradients usually measured? Any binning? If so, can we study what effect that has? How are uncertainties treated?

To what extent is the radial metallicity gradient of young stars linear? Alternatively, would it be better described by two linear relations, with a break radius at corotation radius ([Bresolin et al. 2012](#), and references therein) or further out ([Donor et al. 2020](#)) or a more sophisticated function (see e.g. [Chiappini et al. 2001](#); [Kubryk et al. 2015](#))? Investigating the gradient's form helps in understanding the galactic processes influencing metallicity at different radii ([Minchev et al. 2014](#)).

SB: "However, a simple, linear gradient may not be the better way to describe the distribution of the elements in the Galactic disk. From




Figure 11. Density distribution and spread of $[Fe/H]$ across different galactocentric radii with respect to a global linear radial metallicity gradient across different age ranges. Panels a-j) show young stars and exhibit a rather similar trend, whereas the scatter increases significantly for stars above 1 Gyr in panels k) and l).

Cepheids, Andrievsky et al. (2002b), Pedicelli et al. (2009, 2010), and Genovali et al. (2013) found a steeper gradient in the very inner disk (≤ 7 kpc)." from Lemasle et al. (2013)

SB: "There are numerous abundance studies of open clusters from different groups: [...] They all reach the same conclusion: a linear gradient of approximately -0.06 dex/kpc extending quite far into the outer disk, and a flattening at a level of $[Fe/H] \approx -0.3$ dex, somewhere between 10 and 14 kpc." from Lemasle et al. (2013).

SB: Lemasle et al. (2008) find $d[Fe/H]/dR = -0.050 \pm 0.008$ dex kpc $^{-1}$ in the 5-17 kpc range, but $d[Fe/H]/dR = -0.012 \pm 0.014$ dex kpc $^{-1}$ in a smaller 10-15 kpc range.

SB: How significant is the difference of linear to quadratic when looking at 5-19 kpc, like MW studies (Genovali et al. 2014)?

SB: Put some of these results into context of the Milky Way in Fig. 14. Don't forget to cite all the people who contributed to the




Figure 12. Tracing young stars and gas across galactocentric radii $R_{Gal.}$ and height $z_{Gal.}$ across the whole galaxy (panel a) and different azimuthal ranges/sectors (panels b-i). Small rectangles with cool-warm colors along the horizontal axis indicate the local gradient slopes as in Fig. 5.




Figure 13. Same as Fig. 2, but for gas.

G+14 compilation, as well as Spina for GALAH and Myers et al. (2022); Donor et al. (2020) for APOGEE

SB: Point out the low $[Fe/H]$ outliers with ages below 200 Gyr in both observation (below $R_{Gal.} < 10$ kpc and $[Fe/H] < -0.25$ in Fig. 14a) and simulation (few red dots around $[Fe/H] \sim -0.15$ in Fig. 14c).

5.2 Scatter in the gradient

How much scatter in the radial metallicity gradient of young stars is expected based on simulations? Quantifying the scatter can provide insights into the effects of transient events like mergers, star formation bursts, and gas accretion.




Figure 14. Comparison of the Milky Way’s radial metallicity trend as traced by Cepheids (black triangles, compiled from literature by Genovali et al. 2014, G+14) as well as young (< 1 Gyr) open cluster of the Milky Way as traced by the literature compilation from Genovali et al. (2014, G+14 as squares), APOGEE DR17 from Myers et al. (2022, M+22 as crosses), and GALAH DR3 from Spina et al. (2021, S+21 as circles). The latter two are compiled based on the membership and age catalogue by Cantat-Gaudin & Anders (2020, CG+20).

SB: Put gas and star results into context of the Milky Way in Fig. 15, inspired by Poggio et al. (2021) and Hackshaw et al. (2024). We do not have enough star particles to really see substructure. But the gas shows similar substructure as the actual Milky Way.

SB: How well can we resolve the differences of $[Fe/H]$ in a similar regions as for the Milky Way? Fig. 13 shows this for gas and stars, respectively. Clear metallicity decrease in Fig. 13a towards left (away from galactic centre). Residuals from linear gradient fit are more scattered towards left (away from galactic centre). Low number statistics (typically less than 5 particles per bin)? Gas (Fig. 13c) also show lower iron abundances with increasing distance from centre. However, it is significantly more nuanced and shows step-like behaviour of $[Fe/H]$ changes around edges of gas overdensities. Significant deviations from global gradient expectation especially in the large void around $-12 < X_{Gal} < -10$ kpc. Follow-up would be interesting but is beyond scope of this paper.

5.3 Coherence of the gradient with position

How significant is the change in the radial metallicity of young stars gradient as a function of position, both in terms of radial coverage and azimuth? Understanding these variations can reveal the underlying processes affecting metallicity distribution, such as spiral arm dynamics and bar-driven mixing (see their Figs. 5-8 Di Matteo et al. 2013).

5.4 Coherence of the gradient with age

Until what stellar age can stars be used as reliable tracers of the gas disk in simulations, in terms of both chemical composition and position? This question is pivotal for interpreting observed gradients across different ages (e.g. Willett et al. 2023) and comparing them with model predictions, particularly concerning the migration and heating of stellar populations (Binney & Tremaine 2008; Frankel et al. 2018).

5.5 The connection of young stars and gas

Discuss both Fig. 12 and Fig. 13

5.6 Application of quadratic gradient function onto observations

- Can we apply our functional form onto MW data? e.g. Yong et al. (2012); Andrievsky et al. (2004); Genovali et al. (2014)?
- Can we apply our functional form onto extragalactic data? e.g. Chen et al. (2023); Bresolin et al. (2012)?

Sánchez et al. (2014) analysed a sample of 306 galaxies and found that breaks in metallicity gradients are common in both spiral and barred galaxies, often occurring at radii between 1 and 2 effective radii (R_e): "Beyond ~ 2 disk effective radii our data show evidence of a flattening in the abundance"

Using data from the MaNGA survey, Belfiore et al. (2017) found that many galaxies exhibit breaks in their metallicity gradients: "At large radii, a flattening of the metallicity gradient with respect to the value $0.5 < R/R_e < 2.0$ is detected for $\log 10(M_\star/M_\odot) > 9.5$, although with varying degrees of significance." SB: look into their text on the shape and description of the flattening!

Bresolin et al. (2012) break in gradient for M83, NGC 1512, NGC3621, and NGC4625. Vlajić et al. (2009) maybe most reasonable fit with 2 lines and a break radius for NGC 300.

SB: this NIHAO galaxy has a strong warp (compared to other NIHAO-UHD simulations Buck 2020). How does the metallicity gradient look like for other galaxies without a warp?!

6 CONCLUSIONS

6.1 Main Take-Away

6.2 Future Research

- Can we say what the linear gradient and flattening part is for R_{eff} for our NIHAO Milky Way analogue. Can we relate this to the MW effective radius and other galaxies?

ACKNOWLEDGEMENTS

We acknowledge the traditional owners of the land on which the ANU stands, the Ngunnawal and Ngambri people. We pay our respects to




Figure 15. Comparison of density distribution of young stars and gas in the Milky Way and the NIHAO Milky Way analogue simulation. Panel a) shows the measurements of the Solar vicinity within 5 kpc by Poggio et al. (2021). Panels b) and c) show young stars and gas NIHAO, respectively, for a selected region similar to panel a). Black and white contour lines in panel b) trace overdensities in the gas distribution of panel c).




Figure 16. Comparison of density distribution of young stars and gas NIHAO Milky Way analogue simulation for the same regions as Figs. 15b and 15c. Panels a) and c) trace median gas and young star iron abundances, respectively. Panels b) and d) plot the residual $[\text{Fe}/\text{H}]$ of gas and stars, respectively, when correcting with a radial metallicity gradient fit. Black and white contour lines in panel b) trace overdensities in the gas distribution of Fig. 15c).

elders past, and present and are proud to continue their tradition of surveying the night sky and its mysteries to better understand our Universe.

This work was supported by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013. SB acknowledges support from the Australian Research Council under grant number DE240100150 which enabled SB to continue researching at the end of a fixed-term position and finalising this study. TB acknowledges funding from the Carl Zeiss Stiftung and support from the European Research Council under ERC-CoG grant CRAGSMAN-646955. We gratefully acknowledge the Gauss Centre for Supercomputing e.V. (www.gauss-centre.eu) for funding this project by providing computing time on the GCS Supercomputer SuperMUC at Leibniz Supercomputing Centre (www.lrz.de). Simulations were partially computed with High Performance Computing resources at New York University, Abu Dhabi.

SOFTWARE

The research for this publication was coded in PYTHON (version 3.7.4) and included its packages ASTROPY (v. 3.2.2; [Astropy Collaboration et al. 2013, 2018](#)), IPYTHON (v. 7.8.0; [Pérez & Granger 2007](#)), MATPLOTLIB (v. 3.1.3; [Hunter 2007](#)), NUMPY (v. 1.17.2; [Walt et al. 2011](#)), PYNBODY (v. 1.1.0; [Pontzen et al. 2013](#)), SCIPY (v. 1.3.1; [Virtanen et al. 2020](#)), SKLEARN (v. 1.5.1 [Pedregosa et al. 2011](#)) STATSMODELS (v. 0.14.2 [Perktold et al. 2024](#)) We further made use of TOPCAT (version 4.7; [Taylor 2005](#));

DATA AVAILABILITY

All code and data to reproduce the analysis and figures can be publicly accessed via https://github.com/svenbuder/nihao_radial_metallicity_gradients.

REFERENCES

- Agertz O., et al., 2021, *MNRAS*, **503**, 5826
- Allende Prieto C., Beers T. C., Wilhelm R., Newberg H. J., Rockosi C. M., Yanny B., Lee Y. S., 2006, *ApJ*, **636**, 804
- Anders F., et al., 2014, *A&A*, **564**, A115
- Anders F., et al., 2017, *A&A*, **600**, A70
- Andrievsky S. M., et al., 2002a, *A&A*, **381**, 32
- Andrievsky S. M., Bersier D., Kovtyukh V. V., Luck R. E., Maciel W. J., Lépine J. R. D., Beletsky Y. V., 2002b, *A&A*, **384**, 140
- Andrievsky S. M., Luck R. E., Martin P., Lépine J. R. D., 2004, *A&A*, **413**, 159
- Astropy Collaboration et al., 2013, *A&A*, **558**, A33
- Astropy Collaboration et al., 2018, *AJ*, **156**, 123
- Belfiore F., et al., 2017, *MNRAS*, **469**, 151
- Bellardini M. A., Wetzel A., Loebman S. R., Faucher-Giguère C.-A., Ma X., Feldmann R., 2021, *MNRAS*, **505**, 4586
- Bellardini M. A., Wetzel A., Loebman S. R., Bailin J., 2022, *MNRAS*, **514**, 4270
- Bergemann M., et al., 2014, *A&A*, **565**, A89
- Binney J., Tremaine S., 2008, *Galactic Dynamics*: Second Edition. Princeton University Press
- Boeche C., et al., 2013, *A&A*, **559**, A59
- Bresolin F., Kennicutt R. C., Ryan-Weber E., 2012, *ApJ*, **750**, 122
- Buck T., 2020, *MNRAS*, **491**, 5435
- Buck T., Obreja A., Macciò A. V., Minchev I., Dutton A. A., Ostriker J. P., 2020, *MNRAS*, **491**, 3461
- Buck T., Rybizki J., Buder S., Obreja A., Macciò A. V., Pfrommer C., Steinmetz M., Ness M., 2021, *MNRAS*, **508**, 3365
- Buder S., Mijnarens L., Buck T., 2024, *MNRAS*, **532**, 1010
- Cantat-Gaudin T., Anders F., 2020, *A&A*, **633**, A99
- Casamiquela L., et al., 2019, *MNRAS*, **490**, 1821
- Chabrier G., 2003, *PASP*, **115**, 763
- Chen Q.-H., Grasha K., Battisti A. J., Kewley L. J., Madore B. F., Seibert M., Rich J. A., Beaton R. L., 2023, *MNRAS*, **519**, 4801
- Chen Q.-H., et al., 2024, *MNRAS*, **527**, 2991
- Chiappini C., 2002, *Ap&SS*, **281**, 253
- Chiappini C., Matteucci F., Romano D., 2001, *ApJ*, **554**, 1044
- Chieffi A., Limongi M., 2004, *ApJ*, **608**, 405
- Chiosi C., 1980, *A&A*, **83**, 206
- Cunha K., et al., 2016, *Astronomische Nachrichten*, **337**, 922
- Di Matteo P., Haywood M., Combes F., Semelin B., Snaith O. N., 2013, *A&A*, **553**, A102
- Donor J., et al., 2020, *AJ*, **159**, 199
- Frankel N., Rix H.-W., Ting Y.-S., Ness M., Hogg D. W., 2018, *ApJ*, **865**, 96
- Frankel N., Sanders J., Ting Y.-S., Rix H.-W., 2020, arXiv e-prints, p. [arXiv:2002.04622](https://arxiv.org/abs/2002.04622)
- Gaia Collaboration et al., 2016, *A&A*, **595**, A1
- Genovali K., et al., 2014, *A&A*, **566**, A37
- Graf R. L., Wetzel A., Bellardini M. A., Bailin J., 2024, arXiv e-prints, p. [arXiv:2402.15614](https://arxiv.org/abs/2402.15614)
- Grand R. J. J., Kawata D., Cropper M., 2015, *MNRAS*, **447**, 4018
- Hackshaw Z., Hawkins K., Filion C., Horta D., Laporte C. F. P., Carr C., Price-Whelan A. M., 2024, arXiv e-prints, p. [arXiv:2405.18120](https://arxiv.org/abs/2405.18120)
- Hayden M. R., et al., 2014, *AJ*, **147**, 116
- Hayden M. R., et al., 2015, *ApJ*, **808**, 132
- Hogg D. W., Eilers A.-C., Rix H.-W., 2019, *AJ*, **158**, 147
- Hunter J. D., 2007, *Comput Sci Eng*, **9**, 90
- Imig J., et al., 2023, *ApJ*, **954**, 124
- Janes K. A., 1979, *ApJS*, **39**, 135
- Karakas A. I., Lugaro M., 2016, *ApJ*, **825**, 26
- Katz D., Soubiran C., Cayrel R., Barbuy B., Friel E., Bienaymé O., Perrin M. N., 2011, *A&A*, **525**, A90
- Khoperskov S., Sivkova E., Saburova A., Vasiliev E., Shustov B., Minchev I., Walcher C. J., 2023, *A&A*, **671**, A56
- Knollmann S. R., Knebe A., 2009, *ApJS*, **182**, 608
- Kubryk M., Prantzos N., Athanassoula E., 2015, *A&A*, **580**, A126
- Larson R. B., 1976, *MNRAS*, **176**, 31
- Lemasle B., François P., Bono G., Mottini M., Primas F., Romaniello M., 2007, *A&A*, **467**, 283
- Lemasle B., François P., Piersimoni A., Pedicelli S., Bono G., Laney C. D., Primas F., Romaniello M., 2008, *A&A*, **490**, 613
- Lemasle B., et al., 2013, *A&A*, **558**, A31
- Lemasle B., et al., 2022, *A&A*, **668**, A40
- Limongi M., Chieffi A., 2018, *ApJS*, **237**, 13
- Ma X., Hopkins P. F., Wetzel A. R., Kirby E. N., Anglés-Alcázar D., Faucher-Giguère C.-A., Kereš D., Quataert E., 2017, *MNRAS*, **467**, 2430
- Magrini L., et al., 2017, *A&A*, **603**, A2
- Matteucci F., ed. 2001, *The chemical evolution of the Galaxy*. Astrophysics and Space Science Library Vol. 253, doi:10.1007/978-94-010-0967-6.
- Metha B., Trenti M., Chu T., 2021, *MNRAS*, **508**, 489
- Minchev I., Chiappini C., Martig M., 2014, *A&A*, **572**, A92
- Mun M., et al., 2024, *MNRAS*, **530**, 5072
- Myers N., et al., 2022, *AJ*, **164**, 85
- Pedregosa F., et al., 2011, *J Mach Learn Res*, **12**, 2825
- Pérez F., Granger B. E., 2007, *Comput Sci Eng*, **9**, 21
- Perktold J., et al., 2024, statsmodels/statsmodels: Release 0.14.2, doi:10.5281/zenodo.593847
- Pilkington K., et al., 2012, *A&A*, **540**, A56
- Planck Collaboration et al., 2014, *A&A*, **571**, A16
- Poggio E., et al., 2018, *MNRAS*, **481**, L21
- Poggio E., et al., 2021, *A&A*, **651**, A104
- Poggio E., et al., 2022, *A&A*, **666**, L4
- Pontzen A., Roškar R., Stinson G. S., Woods R., 2013, pynbody: Astrophysics Simulation Analysis for Python, Astrophysics Source Code Library, record ascl:1305.002
- Rolleston W. R. J., Smartt S. J., Dufton P. L., Ryans R. S. I., 2000, *A&A*, **363**, 537
- Rybicki J., Just A., Rix H.-W., 2017, *A&A*, **605**, A59
- Sánchez S. F., et al., 2014, *A&A*, **563**, A49
- Searle L., 1971, *ApJ*, **168**, 327
- Seitenzahl I. R., et al., 2013, *MNRAS*, **429**, 1156
- Shaver P. A., McGee R. X., Newton L. M., Danks A. C., Pottasch S. R., 1983, *MNRAS*, **204**, 53
- Spina L., et al., 2021, *MNRAS*, **503**, 3279
- Stanghellini L., Magrini L., Casasola V., 2015, *ApJ*, **812**, 39
- Stinson G., Seth A., Katz N., Wadsley J., Governato F., Quinn T., 2006, *MNRAS*, **373**, 1074
- Stinson G. S., Brook C., Macciò A. V., Wadsley J., Quinn T. R., Couchman H. M. P., 2013, *MNRAS*, **428**, 129
- Taylor M. B., 2005, *ASPC*, **347**, 29
- Tinsley M. B., 1980, *Fundamentals Cosmic Phys.*, **5**, 287
- Tissera P. B., Rosas-Guevara Y., Bower R. G., Crain R. A., del P Lagos C., Schaller M., Schaye J., Theuns T., 2019, *MNRAS*, **482**, 2208
- Twarog B. A., 1980, *ApJ*, **242**, 242
- Twarog B. A., Ashman K. M., Anthony-Twarog B. J., 1997, *AJ*, **114**, 2556
- Vickers J. J., Shen J., Li Z.-Y., 2021, *ApJ*, **922**, 189
- Vilchez J. M., Esteban C., 1996, *MNRAS*, **280**, 720
- Virtanen P., et al., 2020, *Nature Methods*, **17**, 261
- Vlajić M., Bland-Hawthorn J., Freeman K. C., 2009, *ApJ*, **697**, 361
- Wadsley J. W., Keller B. W., Quinn T. R., 2017, *MNRAS*, **471**, 2357
- Walt S. v. d., Colbert S. C., Varoquaux G., 2011, *Comput Sci Eng*, **13**, 22
- Wang L., Dutton A. A., Stinson G. S., Macciò A. V., Penzo C., Kang X., Keller B. W., Wadsley J., 2015, *MNRAS*, **454**, 83
- Willett E., et al., 2023, *MNRAS*, **526**, 2141
- Yong D., Carney B. W., Friel E. D., 2012, *AJ*, **144**, 95
- Zari E., Hashemi H., Brown A. G. A., Jardine K., de Zeeuw P. T., 2018, *A&A*, **620**, A172
- Zari E., Rix H. W., Frankel N., Xiang M., Poggio E., Drimmel R., Tkachenko A., 2021, *A&A*, **650**, A112
- van de Sande J., Fraser-McKelvie A., Fisher D. B., Martig M., Hayden M. R., the GECKOS Survey collaboration 2023, arXiv e-prints, p. [arXiv:2306.00059](https://arxiv.org/abs/2306.00059)

This paper has been typeset from a TeX/LaTeX file prepared by the author.