

Supplementary Data: Table S1. Prescribing Patterns according to WHO-ATC anatomical groups

	Dementia (n=147)	No Dementia (n=436)	Total (n=583)	P-value ²	T-value, χ^2 or Fishers exact test	Controlling for Age, Sex and co- morbidity ³
Number of patients prescribed ≥ 1 of the following medications according to the WHO-ATC anatomical groups ¹ , N (%)						
Alimentary Tract and Metabolism (WHO-ATC A)	110 (74.8)	296 (67.9)	406 (69.6)	0.11	$\chi^2 = 2.5$	aOR = 0.9, 95% CI = 0.6-1.5
Blood and Blood Forming Organs (WHO-ATC B)	105 (71.4)	302 (69.3)	407 (69.8)	0.62	$\chi^2 = 0.2$	aOR = 1.0, 95% CI = 0.6-1.6
Cardiovascular System (WHO-ATC C)	126 (85.7)	381 (87.4)	507 (87.0)	0.60	$\chi^2 = 0.3$	aOR = 0.7, 95% CI = 0.4-1.3
Genito-Urinary System and Sex Hormones (WHO-ATC G)	31 (21.1)	70 (16.1)	101 (17.3)	0.16	$\chi^2 = 1.9$	aOR = 1.2, 95% CI = 0.7-2.0
Systemic Hormonal Preparations (WHO-ATC H)	25 (17.0)	100 (22.9)	125 (21.4)	0.13	$\chi^2 = 2.3$	aOR = 0.6, 95% CI = 0.4-1.0
Anti-infectives for Systemic Use (WHO-ATC J)	22 (15.0)	41 (9.4)	63 (10.8)	0.06	$\chi^2 = 3.5$	aOR = 1.6, 95% CI = 0.8-2.9
Anti-neoplastic and Immunomodulating Agents (WHO-ATC L)	5 (3.4)	26 (6.0)	31 (5.3)	0.23	$\chi^2 = 1.4$	aOR = 0.7, 95% CI = 0.2-1.9
Musculo-Skeletal System (WHO-ATC M)	38 (25.9)	100 (22.9)	138 (23.7)	0.47	$\chi^2 = 0.5$	aOR = 1.2, 95% CI = 0.8-2.0
Nervous System (WHO-ATC N)	111 (75.5)	239 (54.8)	350 (60.0)	<0.001*	$\chi^2 = 19.6$	aOR = 2.0, 95% CI = 1.3-3.2 [†]
Anti-parasitic Products, Insecticides and Repellants (WHO-ATC P)	2 (1.4)	10 (2.3)	12 (2.1)	0.74	F = 0.74	aOR = 0.6, 95% CI = 0.1-3.3
Respiratory System (WHO-ATC R)	37 (25.2)	125 (28.7)	162 (27.8)	0.41	$\chi^2 = 0.7$	aOR = 0.8, 95% CI = 0.5-1.2
Sensory Organs (WHO-ATC S)	13 (8.8)	28 (6.4)	41 (7.0)	0.32	$\chi^2 = 1.0$	aOR = 1.0, 95% CI = 0.2-2.2
Various (WHO-ATC V)	1 (0.7)	4 (0.9)	5 (0.9)	1.0	F = 1.0	aOR = 0.5, 95% CI = 0.1-5.2

¹WHO-ATC D (Dermatologicals) excluded as emollients without any active ingredients were not coded.

²P-value for two-way table with measures of association

³P-value for dependent variable utilising linear or logistic regression as appropriate, with age, sex and CIRS-G as the independent variables.

*Statistically significant at p-level <0.05, utilising χ^2 test.

† Statistically significant at p-level <0.05, utilising logistic regression.

WHO-ATC = World Health Organization Anatomical Therapeutic Chemical, CIRS-G = Cumulative Illness Rating Scale in Geriatrics, aOR = Adjusted Odds Ratio, β = beta- coefficient, 95% CI = 95% Confidence Interval