

TABLE 1. ICD-9-CM and ICD-10 Coding Algorithms for Charlson Comorbidities

Comorbidities	Deyo's ICD-9-CM +	ICD-10 *	Enhanced ICD-9-CM *
Myocardial infarction	410.x, 412.x	I21.x, I22.x, I25.2	410.x, 412.x
Congestive heart failure	428.x	I09.9, I11.0, I13.0, I13.2, I25.5, I42.0, I42.5-I42.9, I43.x, I50.x, P29.0	398.91, 402.01, 402.11, 402.91, 404.01, 404.03, 404.11, 404.13, 404.91, 404.93, 425.4--425.9, 428.x
Peripheral vascular disease	443.9, 441.x, 785.4, V43.4 Procedure 38.48	I70.x, I71.x, I73.1, I73.8, I73.9, I77.1, I79.0, I79.2, K55.1, K55.8, K55.9, Z95.8, Z95.9	093.0, 437.3, 440.x, 441.x, 443.1--443.9, 447.1, 557.1, 557.9, V43.4
Cerebrovascular disease	430.x--438.x	G45.x, G46.x, H34.0, I60.x--I69.x	362.34, 430.x--438.x
Dementia	290.x	F00.x--F03.x, F05.1, G30.x, G31.1	290.x, 294.1, 331.2
Chronic pulmonary disease	490.x--505.x, 506.4	I27.8, I27.9, J40.x--J47.x, J60.x--J67.x, J68.4, J70.1, J70.3	416.8, 416.9, 490.x--505.x, 506.4, 508.1, 508.8
Rheumatic disease	710.0, 710.1, 710.4, 714.0--714.2, 714.81, 725.x	M05.x, M06.x, M31.5, M32.x--M34.x, M35.1, M35.3, M36.0	446.5, 710.0--710.4, 714.0--714.2, 714.8, 725.x
Peptic ulcer disease	531.x--534.x	K25.x--K28.x	531.x--534.x
Mild liver disease	571.2, 571.4--571.6	B18.x, K70.0--K70.3, K70.9, K71.3--K71.5, K71.7, K73.x, K74.x, K76.0, K76.2--K76.4, K76.8, K76.9, Z94.4	070.22, 070.23, 070.32, 070.33, 070.44, 070.54, 070.6, 070.9, 570.x, 571.x, 573.3, 573.4, 573.8, 573.9, V42.7
Diabetes without chronic complication	250.0--250.3, 250.7	E10.0, E10.1, E10.6, E10.8, E10.9, E11.0, E11.1, E11.6, E11.8, E11.9, E12.0, E12.1, E12.6, E12.8, E12.9, E13.0, E13.1, E13.6, E13.8, E13.9, E14.0, E14.1, E14.6, E14.8, E14.9	250.0--250.3, 250.8, 250.9
Diabetes with chronic complication	250.0--250.6	E10.2--E10.5, E10.7, E11.2, E11.5, E11.7, E12.2--E12.5, E12.7, E13.2--E13.5, E13.7, E14.2--E14.5, E14.7	250.4--250.7
Hemiplegia or paraplegia	344.1, 342.x	G04.1, G11.4, G80.1, G80.2, G81.x, G82.x, G83.0--G83.4, G83.9	334.1, 342.x, 343.x, 344.0--344.6, 344.9
Renal disease	582.x, 583--583.7, 585.x, 586.x, 588.x	I12.0, I13.1, N03.2--N03.7, N05.2--N05.7, N18.x, N19.x, N25.0, Z49.0--Z49.2, Z94.0, Z99.2	403.01, 403.11, 403.91, 404.02, 404.03, 404.12, 404.13, 404.92, 404.93, 582.x, 583.0--583.7, 585.x, 586.x, 588.0, V42.0, V45.1, V56.x
Any malignancy, including lymphoma and leukemia, except malignant neoplasm of skin	140.x--172.x, 174.x--195.8, 200.x--208.x	C00.x--C26.x, C30.x--C34.x, C37.x--C41.x, C43.x, C45.x--C58.x, C60.x--C76.x, C81.x--C85.x, C88.x, C90.x--C97.x	140.x--172.x, 174.x--195.8, 200.x--208.x, 238.6
Moderate or severe liver disease	456.0--456.21, 572.2--572.8	I85.0, I85.9, I86.4, I98.2, K70.4, K71.1, K72.1, K72.9, K76.5, K76.6, K76.7	456.0--456.2, 572.2--572.8
Metastatic solid tumor	196.x--199.1	C77.x--C80.x	196.x--199.x
AIDS/HIV	042.x--044.x	B20.x--B22.x, B24.x	042.x--044.x

Source:

- + Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol.* 1992; 45: 613-9.
- * Quan H, Sundararajan V, Halfon P, et al. Coding algorithms for defining Comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care.* 2005 Nov; 43(11): 1130-9.