

FMA-PG Notes

Shivam and Sharan

September 2021

1 Softmax PPO Closed Form Update

We will consider direct functional representation with tabular parameterization, i.e. $\pi \equiv p^\pi$ is essentially an $\mathcal{S} \times \mathcal{A}$ table satisfying the constraints

$$\begin{aligned} \sum_a p^\pi(a|s) &= 1, & \forall s \in \mathcal{S} \\ p^\pi(a|s) &\geq 0, & \forall s \in \mathcal{S}, \forall a \in \mathcal{A}. \end{aligned}$$

Our goal is to find the closed form solution to the following optimization problem (from Eq. 6, Sharan et al., 2021):

$$\pi_{t+1} = \arg \max_{\pi \in \Pi} \left[\sum_s d^{\pi_t}(s) \sum_a p^{\pi_t}(a|s) \left(A^{\pi_t}(s, a) + \frac{1}{\eta} \right) \log \frac{p^\pi(s, a)}{p^{\pi_t}(s, a)} \right], \quad (1)$$

subject to the above constraints on p^π . The above equation is obtained by setting the mirror map to the weighted exponential function.

We begin by formulating this problem using Lagrange multipliers $\lambda_s, \lambda_{s,a}$ for all states s and actions a (and with a slight abuse of the notation, i.e. we write λ_s for all the $\lambda_{s,s}$, etc.):

$$\begin{aligned} \mathcal{L}(p^\pi, \lambda_s, \lambda_{s,a}) &= \sum_s d^{\pi_t}(s) \sum_a p^{\pi_t}(a|s) \left(A^{\pi_t}(s, a) + \frac{1}{\eta} \right) \log \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)} \\ &\quad - \sum_{s,a} \lambda_{s,a} p^\pi(a|s) - \sum_s \lambda_s \left(\sum_a p^\pi(a|s) - 1 \right). \end{aligned} \quad (2)$$

KKT conditions for this problem are:

$$\nabla_{p^\pi(x,b)} \mathcal{L}(p^\pi, \lambda_s, \lambda_{s,a}) = 0, \quad \forall x \in \mathcal{S}, \forall b \in \mathcal{A} \quad (C1)$$

$$\sum_a p^\pi(a|s) = 1, \quad \forall s \in \mathcal{S} \quad (C2)$$

$$p^\pi(a|s) \geq 0, \quad \forall s \in \mathcal{S}, \forall a \in \mathcal{A} \quad (C3)$$

$$\lambda_s \geq 0, \quad \forall s \in \mathcal{S} \quad (C4)$$

$$\lambda_s \left(\sum_a p^\pi(a|s) - 1 \right) = 0 \quad \forall s \in \mathcal{S} \quad (C5)$$

$$\lambda_{s,a} p^\pi(a|s) = 0, \quad \forall s \in \mathcal{S}, \forall a \in \mathcal{A}. \quad (C6)$$

Let us now try to solve this system. Solving the first equation for arbitrary state-action pair (x, b) , gives us:

$$\begin{aligned} \nabla_{p^\pi(b|x)} \mathcal{L}(p^\pi, \lambda_s, \lambda_{s,a}) &= d^{\pi_t}(x) p^{\pi_t}(b|x) \left(A^{\pi_t}(x, b) + \frac{1}{\eta} \right) \frac{1}{p^\pi(b|x)} - \lambda_{x,b} - \lambda_x = 0 \\ \Rightarrow p^\pi(b|x) &= \frac{d^{\pi_t}(x) p^{\pi_t}(b|x) (1 + \eta A^{\pi_t}(x, b))}{\eta (\lambda_x + \lambda_{x,b})}. \end{aligned} \quad (3)$$

Let us set

$$\lambda_{s,a} = 0, \quad \forall s \in \mathcal{S}, \forall a \in \mathcal{A}. \quad (4)$$

Combining Eq. 3 with the second KKT condition gives us

$$\lambda_s = \frac{1}{\eta} \sum_a d^{\pi_t}(s) p^{\pi_t}(a|s) (1 + \eta A^{\pi_t}(s, a)). \quad (5)$$

Therefore, with the additional assumption $d^{\pi_t}(s) > 0$, $p^\pi(a|s)$ becomes

$$p^\pi(a|s) = \frac{p^{\pi_t}(a|s)(1 + \eta A^{\pi_t}(s, a))}{\sum_b p^{\pi_t}(b|s)(1 + \eta A^{\pi_t}(s, b))}. \quad (6)$$

Note that $d^{\pi_t}(s), p^{\pi_t}(a|s) \geq 0$ for any state-action pair, since they are proper measures. All that remains is to ensure that

$$1 + \eta A^{\pi_t}(s, a) \geq 0$$

to satisfy the third and fourth KKT conditions. But how to do that? One straightforward way is to define $p^\pi(a|s) = 0$ whenever $1 + \eta A^{\pi_t}(s, a) \leq 0$, and accordingly re-define λ_s . This gives us the final solution to our original optimization problem (Eq. 1):

$$\pi_{t+1} = p^\pi(s, a) = \frac{p^{\pi_t}(a|s) \max(1 + \eta A^{\pi_t}(s, a), 0)}{\sum_b p^{\pi_t}(b|s) \max(1 + \eta A^{\pi_t}(s, b), 0)}. \quad (7)$$

This leaves one last problem: Is it always true that given any state s , there exists atleast one action a , such that $1 + \eta A^{\pi_t}(s, a) \geq 0$? Because otherwise, we would fail to satisfy the second KKT condition.

2 MDPO Closed Form Update

The paper (Sharan et al., 2021) considers the direct representation along with tabular parameterization of the policy, albeit with a small change in notation as compared to the previous section: $\pi(a|s) \equiv p^\pi(a|s, \theta)$. However, since this notation is more cumbersome, we will stick to our old one: $\pi(a|s) \equiv p^\pi(a|s)$. The constraints on these parameters are the same as before: $\sum_a p^\pi(a|s) = 1, \forall s \in \mathcal{S}$; and $p^\pi(a|s) \geq 0, \forall s \in \mathcal{S}, \forall a \in \mathcal{A}$. Our goal, this time, is to solve the following optimization problem (see Eq. 9, Sharan et al., 2021)

$$\pi_{t+1} = \arg \max_{\pi \in \Pi} \left[\sum_s d^{\pi_t}(s) \sum_a p^{\pi_t}(a|s) \left(Q^{\pi_t}(s, a) \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)} - \frac{1}{\eta} D_\phi(p^\pi(\cdot|s), p^{\pi_t}(\cdot|s)) \right) \right], \quad (8)$$

with the mirror map as the negative entropy (Eq. 5.27, Beck and Teboulle, 2002). This particular choice results in

$$D_\phi(p^\pi(\cdot|s), p^{\pi_t}(\cdot|s)) = \text{KL}(p^\pi(\cdot|s) \| p^{\pi_t}(\cdot|s)) := \sum_a p^\pi(a|s) \log \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)}. \quad (9)$$

The optimization problem then simplifies to

$$\pi_{t+1} = \arg \max_{\pi \in \Pi} \left[\sum_s d^{\pi_t}(s) \sum_a p^{\pi_t}(a|s) \left(Q^{\pi_t}(s, a) \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)} - \frac{1}{\eta} \sum_a p^\pi(a|s) \log \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)} \right) \right]. \quad (10)$$

Proceeding analogously to the previous section, we use Lagrange multipliers λ_s , $\lambda_{s,a}$ for all states s and actions a to obtain the function

$$\begin{aligned} \mathcal{L}(p^\pi, \lambda_s, \lambda_{s,a}) = & \sum_s d^{\pi_t}(s) \sum_a p^{\pi_t}(a|s) \left(Q^{\pi_t}(s, a) \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)} - \frac{1}{\eta} \sum_a p^\pi(a|s) \log \frac{p^\pi(a|s)}{p^{\pi_t}(a|s)} \right) \\ & - \sum_{s,a} \lambda_{s,a} p^\pi(a|s) - \sum_s \lambda_s \left(\sum_a p^\pi(a|s) - 1 \right). \end{aligned} \quad (11)$$

The KKT conditions are exactly the same as before (Eq. C1 to Eq. C6).

Again, we begin by solving the first KKT condition:

$$\begin{aligned} \nabla_{p^\pi(b|x)} \mathcal{L}(p^\pi, \lambda_s, \lambda_{s,a}) = & d^{\pi_t}(x) p^{\pi_t}(b|x) \left(\frac{Q^{\pi_t}(x, b)}{p^{\pi_t}(b|x)} - \frac{1}{\eta} \left[\log \frac{p^\pi(b|x)}{p^{\pi_t}(b|x)} + 1 \right] \right) - \lambda_{x,b} - \lambda_x \\ = & d^{\pi_t}(x) p^{\pi_t}(b|x) \left(\frac{Q^{\pi_t}(x, b)}{p^{\pi_t}(b|x)} - \frac{1}{\eta} \left[\log \frac{p^\pi(b|x)}{p^{\pi_t}(b|x)} + 1 \right] - \frac{\lambda_{x,b} + \lambda_x}{d^{\pi_t}(x) p^{\pi_t}(b|x)} \right) \\ = & 0 \\ \Rightarrow \quad \log \frac{p^\pi(b|x)}{p^{\pi_t}(b|x)} = & \frac{\eta Q^{\pi_t}(x, b)}{p^{\pi_t}(b|x)} - \frac{\eta(\lambda_{x,b} + \lambda_x)}{d^{\pi_t}(x) p^{\pi_t}(b|x)} - 1 \\ \Rightarrow \quad p^\pi(b|x) = & p^{\pi_t}(b|x) \cdot \exp \left(\frac{\eta Q^{\pi_t}(x, b)}{p^{\pi_t}(b|x)} \right) \cdot \exp \left(-\frac{\eta(\lambda_{x,b} + \lambda_x)}{d^{\pi_t}(x) p^{\pi_t}(b|x)} - 1 \right). \end{aligned} \quad (12)$$