

True-implies-false, true-implies-true and **nonseparable** Kochen-Specker sets

Adán Cabello*

Departamento de Física Aplicada II, Universidad de Sevilla, E-41012 Sevilla, Spain

José R. Portillo†

*Departamento de Matemática Aplicada I – **IMUS**, Universidad de Sevilla, E-41012 Sevilla, Spain*

Alberto Solís‡

Departamento de Matemática Aplicada I, Universidad de Sevilla, E-41012 Sevilla, Spain

Karl Svozil§

*Institute for Theoretical Physics, University of Technology Vienna,
Wiedner Hauptstrasse 8-10/136, 1040 Vienna, Austria*

(Dated: January 22, 2018)

The essential ingredient in Hardy-like proofs of Bell and Kochen-Specker (KS) theorems is a set of propositions about the outcomes of quantum measurements such that, when outcome noncontextuality is assumed, if proposition A is true then, due to exclusiveness and completeness, a nonexclusive proposition B must be false. We call such a set a *true-implies-false set* (TIFS). The KS theorem can be proved for any quantum state of a system of a given dimension by concatenating a set of propositions **containing a TIFS** such that, if proposition A is true then a nonexclusive proposition C must be true. We call such a set a *true-implies-true set* (TITS). **A nonseparating set (NSS) is one which contains propositions A and C which cannot be separated by any valid truth assignment.** Here we identify the **TIFSes, TITSes, and NSSes** that have the smallest number of propositions for every d -dimensional quantum system with $d \geq 3$.

PACS numbers: 03.65.Ta, 03.65.Ud

I. INTRODUCTION

Every noncontextuality (NC) [?] and **Boole-Bell type** inequality [?] can be represented by a graph in which vertices represent the propositions appearing when the correlations are expressed as a positive linear combination of probabilities of propositions, and exclusive propositions are represented by adjacent nodes [1, 2]. Quantum theory (QT) violates NC and Bell inequalities *if and only* if their exclusivity graphs are *imperfect*, i.e., contain, as induced subgraphs, odd cycles of length five or more (i.e., pentagons, heptagons, etc.), or their complements [1?, 2]. Moreover, every proof, with or without inequalities, of the Kochen-Specker (KS) [3?, 4] and Bell [?] theorems can be associated to an imperfect graph. Reciprocally, every imperfect graph can be used to prove that QT cannot be explained with noncontextual hidden variable theories [1, 2].

Some imperfect graphs allow us to present the conflict between QT and hidden variables in very appealing way: by pointing out a contradiction between a prediction *with certainty* of hidden variables and the corresponding prediction of QT. Proofs of this type are the ones by Stairs [5], Hardy, and others []. In addition,

these imperfect graphs play a fundamental role in quantum state-independent proofs of the KS theorem. See the proof of Bell [?] and Kochen and Specker [4]). In this article we will obtain the simplest of these imperfect graphs.

Hereafter by propositions we will mean statements the type “outcomes a and b will be respectively obtained when observables A and B will be jointly measured on the same physical system”, where A and B are assumed to be observables represented in QT by rank-one projectors that commute. We will say that two propositions are *compatible* if and only if all the observables involved in both propositions are jointly measurable, i.e., represented in QT by operators that commute. Two propositions are *exclusive* when both cannot simultaneously true. A set of mutually exclusive compatible propositions is *complete* when one of the propositions must be true. A *context* is a **complete set** (~~complete or not~~) of mutually exclusive compatible propositions. (Greechie) **orthogonality diagrams** [6] compactly represent contexts as single smooth lines (such as circles or straight unbroken lines) connecting mutually compatible (atomic) **exclusive propositions** which are represented as small circles; contexts intertwining at a single proposition are represented as non-smoothly connected lines, broken at that proposition.

The assumption of *outcome noncontextuality* assigns the same truth value (true or false) to any proposition with independence of which other compatible propositions are considered simultaneously. QT is in conflict with the assumption of outcome noncontextuality [? ? ?

* adan@us.es

† josera@us.es

‡ solisencina@us.es

§ svozil@tuwien.ac.at




FIG. 1. (Color online) Orthogonality diagram of the simplest known [4, 7–11] (a) TIFS, (b) TITS, and (c) NSS in $d = 3$. Points represent propositions, smooth lines represent complete sets (i.e., one and only one of the propositions must be true); in particular, they indicate that any pair of propositions connected by a smooth line cannot both be true (exclusiveness). (a) If A is true then B is false [?, Fig. 1, p. 182]. (b) If A is true then C is true [4, Γ_1]. (c) A and C can only be both true or both false [4, Γ_3]. These sets are realizable in S^2 by taking, for instance [11, p. 206, Fig. 1], $A = (1, \sqrt{2}, 0)/\sqrt{3}$, $v_1 = (\sqrt{2}, -1, 1)/2$, $v_2 = (\sqrt{2}, -1, -1)/2$, $v_3 = (0, 1, 1)/\sqrt{2}$, $v_4 = (0, 1, -1)/\sqrt{2}$, $v_5 = (\sqrt{2}, 1, -1)/2$, $v_6 = (\sqrt{2}, 1, 1)/2$, $B = (-1, \sqrt{2}, 0)/\sqrt{3}$, $v_7 = (0, 0, 1)$, $C = (\sqrt{2}, 1, 0)/\sqrt{3}$, $v_{1'} = (-1, \sqrt{2}, -1)/2$, $v_{2'} = (-1, \sqrt{2}, 1)/2$, $v_{3'} = (1, 0, -1)/\sqrt{2}$, $v_{4'} = (1, 0, 1)/\sqrt{2}$, $v_{5'} = (1, \sqrt{2}, 1)/2$, $v_{6'} = (1, \sqrt{2}, -1)/2$. In QT, the proposition v_i is represented by the projector $|v_i\rangle\langle v_i|$. To obtain a TIFS, TITS and NSS in $d = 4$ it is enough to add $\langle v| = (0, 0, 0, 1)$, and similarly to obtain TITSes in higher dimensions.

?. This conflict can be manifested in many ways, and in the following we shall investigate some which are particularly appealing as they involve smaller sets of propositions than configurations for proofs of the KS theorem.

We define (i) a *true-implies-false set* (TIFS), (ii) a *true-implies-true set* (TITS), and (iii) a *nonseparating set* (NSS) as a set S of propositions represented in QT by rank-one projection operators (or the vectors onto which they project) such that, due to the exclusiveness and completeness of some of the elements of S , when outcome noncontextuality is assumed, if proposition $A \in S$ is true, then a nonexclusive proposition (i) $B \in S$ must be false, (ii) $C \in S$ must be true as well, and (iii) $C \in S$ must be true, as well as *vice versa*, whenever $C \in S$ is true, then A must be true; i.e., A and C cannot be separated with classical probabilities. This is not the same as in (ii), as for (ii) $v(C) = 1$ does not imply $v(A) = 1$ (in particular, $v(A) = 0$ is allowed). Explicit examples for the cases (i–iii) are shown in Fig. 1(a–c), respectively. A TIFS, TITS, or NSS S is said to be *critical* if the set resulting from removing any element of S is not a TIFS,

TITS, or NSS, respectively.

HISTORY? Upon going through Clifton’s paper cited I found two early instances of TITS different from KS: Belinfante [12, Fig. C.I. p. 67], Pitowsky [13, p. 394]. Early examples of TIFS have been discussed in Refs. [14, Fig. 1, p. 182], [12, Fig. B.I. p. 64], [?, p. 588–589], [15, Sects. IV, Fig. 2], [9, p. 39, Fig. 2.4.6], and [7, 8]. Early examples of TITS can be found in Refs. [16, Γ_1 , p. 86], [12, Fig. C.I. p. 67], [13, p. 394], [15, 17, 18], and [19, Lemma 1]. A NSS is discussed in the KS paper [16, Γ_3 , p. 70].

Any TIFS, TITS, or NSS proves quantum contextuality (i.e., the impossibility of explaining QT with models satisfying outcome noncontextuality) because for a system prepared in the quantum state in which proposition A is true there is a nonzero probability of finding proposition B or C true, false, and different, respectively. This is the method followed in the proofs of quantum contextuality proposed by Stairs [?], Clifton [?], and Cabello et al. [? ?]). All these proofs can be converted into experimental tests of whether nature can be described with noncontextual hidden variable theories [?].

TITSes also serve to prove state-independent quantum contextuality, i.e., quantum contextuality for any initial quantum state of a quantum system of a given dimension $d \geq 3$. A proof of this can be obtained by concatenating several TITSes. This is the method followed by Bell [?] and Kochen and Specker (KS) [?] to prove state-independent quantum contextuality in $d = 3$. The same method can be extended to any $d \geq 3$ [?]. State-independent proofs of quantum contextuality have been recently converted into quantum state-independent experimental tests of contextuality [? ? ? ? ? ? ?].

TITS in which proposition A corresponds to an entangled state and the rest corresponds to product states can be used to prove quantum nonlocality (i.e., the impossibility of explaining QT with local hidden variable theories). This is exactly what is behind Hardy-like proofs of quantum nonlocality [? ?] (for a detailed explanation, see Refs. [? ?]).

These sets, and, in particular, NSS, are interesting because they demonstrate a growing deviation from nonclassicality below the KS theorem: there still exist classical valuations and truth tables, but they get “more pathologic” up to the point where propositional structures with NSS cannot be “embedded” [4] into any kind of hidden parameter model, such as partition logics [20], and their model realizations as Wright’s generalized urn model [21] or automaton logic [22] (still allowing logics with TIFS or TITS).

TITSes are known for any physical system described by a Hilbert space of dimension $d \geq 2$ [? ? ?]. In $d = 3$, Bell found one with $n = 13$ propositions [?], and KS found one with $n = 10$ [? ? ?], which is illustrated in Fig. 1. Both Bell’s and KS’s sets belong to a broader family with $n = 10 + 3m$, with $m = 0, 1, \dots$ [?]. For $d > 3$, TITS with $n = 7 + d$ are easy to construct from the set of Fig. 1 by adding the vector with all components

zero but the one corresponding to the new dimension [?].

An important question is which are the *simplest TIFSes, TITSes and NSSes* for any $d \geq 3$, i.e., those with a minimum number of propositions. This is the problem we address in this paper.

II. METHOD TO OBTAIN TITS WITH MINIMUM NUMBER OF PROPOSITIONS

QUIZAS HAYA QUE RECONSTRUIR TODA LA ESTRUCTURA DE SECCIONES

A TITS can be represented by a graph in which vertices represent propositions, edges connect exclusive propositions, and d -cliques (d mutually connected vertices) represent complete contexts. A graph is said to be *nonrealizable* in dimension d if it represents a set of rays (i.e., unit vectors) which is not realizable in S^{d-1} .

Lemma 1: [23?] The simplest nonrealizable graph in $d = 1$ consists of two vertices. The simplest nonrealizable graph in $d = 2$ has three vertices with one of them connected to the other two. From these to nonrealizable graphs one can recursively construct nonrealizable graphs in any dimension d by starting from the nonrealizable graph in dimension $d - 2$ and adding to it two vertices connected with all vertices of the nonrealizable graph in $d - 2$.

~~THIS PROOF IS IN THE CITES *Proof:* One cannot have two different ways in S^0 because there is only one ray in S^0 . In S^1 , if a ray is orthogonal to a second ray, and the second is orthogonal to the third, then the first and third ray should be the same. If we add two dimensions and two different rays, both orthogonal to those of previous set I , then these two new rays span a two dimensional subspace orthogonal to I . Therefore, if I was nonrealizable, then the resulting set is also nonrealizable.~~

Lemma 2: Every n -vertex graph corresponding to a critical TITS in dimension d contains an $(n + 1 - d)$ -vertex graph corresponding to a TIFS.

Proof: Let be G a graph corresponding to a TITS, then $b \in K_d$ is true and any adjacent v must be false. Then the subgraph taken from the other true a to false v forms a TIFS. Moreover, any $v_i \in K_d - 1$ can be taken as false to get a TIFS.

Lemma 3: The graph of a critical TIFS must be biconnected (i.e., when removing any vertex the resulting graph must be still connected).

Proof: Suppose it is not. Cut vertex split graph in two connected components, if true and false vertices are in the same component, then the graph is not critical and if true and false vertices are in different components then cut vertex can substitute anyone of them getting a smaller TIFS.

Corollary 1: Every vertex of a graph corresponding to a TIFS must be connected to, at least, two vertices (i.e., the graph must have minimal valency two).

Proof: Directly from Lemma 3.

Lemma 4: Every TIFS graph in dimension d contains, at least, two K_d graphs.

Proof: Let be G a TIFS graph with true vertex a and false vertex b . G is not trivial TIFS, so there is another true vertex x linked to b and other $d - 1$ vertices. ~~Let be C the rest of vertices in G (except a and b).~~ We consider two different cases. a) Every vertex in $V(G) - \{a, b, x\}$ belongs to $N(a)$ and b is linked to C . Note that in this case $a \cup N(a)$ and $b \cup N(b)$ form two K_d 's. b) Only several vertices in $V(G) - \{a, b, x\}$ belongs to $N(a)$. We have a set of vertices in $V(G) - \{a, b, x\}$ with value false, but such vertices are not linked directly to a , so their value must be forced through another true vertex ~~out C~~ . This vertex is not linked to a , so it has to belong to a context, i.e. a K_d .

~~*Lemma 5:* The graph of any 8-vertex TIFS in $d = 3$ must contain at least two triangles.~~

~~*Proof:* From Lemma 4. Otherwise the graph can be colored as follows: first vertex a true and everything else false.~~

III. DIMENSION 3 - SPECKER BUG

CHANGED SECTIONS ORDER

Therefore, we have a method to obtain the minimal TITS in $d = 3$ **which proceeds as follows:**

Step 1: We generate all nonisomorphic n -vertex biconnected graphs (Lemma 3) of minimal valence two (Corollary 1), not containing cycles on length four (Lemma 1) and containing at least two triangles (Lemma 4) for n less or equal than 8. This can be efficiently done using the computer program *nauty* [?]. We obtain that there are 2 **such** graphs for $n = 7$, and 8 graphs for $n = 8$. All of them are illustrated in Fig. 2.

Step 2: For every graph obtained after step 1, consider all possible pairs of vertices (v_i, v_j) . If, for one (v_i, v_j) , the graph does not admit a noncontextual assignment when $v_i = 1$ and $v_j = 1$ (i.e., then both are true), then the graph is a TIFS in which $A = v_i$ and $B = v_j$. The test of whether or not a graph admits a noncontextual assignment can be done using a simple computer program (e.g., [?]).

After step 2, we find that only the last graph in Fig. 2 corresponds to a TIFS. This graph, **also depicted in Fig. 1(a)**, was first introduced by Kochen and Specker [? , Fig. 1, p. 182]; and later used as a subgraph of the graph Γ_1 of Kochen and Specker [4], as depicted in Fig. 1(b).

This proves that, in $d = 3$, there is no TIFS with less than or equal number of propositions than the one introduced by Kochen and Specker in 1965 [? , Fig. 1, p. 182] (and then used in Ref. [?]), which means that there is no TITS with less than or equal number of propositions than the one in Fig. 1.

Theorem 1: Minimal TITS in dimension $d = 3$ have 10 vertices.




FIG. 2. (Color online) Orthogonality diagrams of all non-isomorphic 7-vertex (first row) and 8-vertex (the rest) biconnected graphs of minimal valence two, not containing cycles of length four, and containing at least two triangles. The last graph was a 9 vertex graph - FIXED




FIG. 3. (Color online) Two equivalent, isomorphic representations of the orthogonality diagram of the simplest known [14, Fig. 1, p. 182] TIFS in $d = 3$, with 7 contexts and 8 atomic propositions. It is realizable in S^2 by taking, for instance, $v_1 = (0, -1, \sqrt{2})/\sqrt{3}$, $v_2 = (1, 0, 0)$, $v_3 = (1, \sqrt{2}, 1)/2$, $v_4 = (0, 1, 0)$, $v_5 = (1, 0, -1)/\sqrt{2}$, $v_6 = (0, 0, 1)$, $v_7 = (-1, \sqrt{2}, -1)/2$, $v_8 = (\sqrt{2}, 1, 0)/\sqrt{3}$. Values on v_1 and v_8 cannot both be 1, as this would imply the values on $v_2 = v_3 = v_6 = v_7 = v_9$ to vanish, which in turn would require both values v_4 as well as v_5 to be 1; which contradicts exclusivity because those observables belong to the same context and are orthogonal. In QT, the proposition v_i is represented by the projector $|v_i\rangle\langle v_i|$.

Proof: This show gets a construction method from the TIFS to obtain a minimal TITS in the same dimension. It is sufficient to add two vertices, one true vertex b and other support vertex v_3 with value no. Both vertices must belong to K_d by the constraints of the problem, as much as the rest of vertices connected to b have value no by TIFS included, simply we must force the value of v_3 to be no, which is achieved by simply adding an edge from the true vertex a in the TIFS to vertex v_3 .

It will be seen later that this method is applicable to higher dimensions.

Do these arguments also apply for NSS? Not exactly. Searching with 17 vertices is computationally very hard. I am using alternative techniques, but without results (2017-12-02)

For $d = 3$, Minimum angle = $\arccos(1/3)$ approx 1.23096 radian approx 19.0986 (not minutes, decimal point) Orthonormal representation: $(0, -1/\sqrt{3}, \sqrt{2}/\sqrt{3})$, $(1, 0, 0)$, $(1/2, 1/\sqrt{2}, 1/2)$, $(0, 1, 0)$, $(1/\sqrt{2}, 0, -1/\sqrt{2})$, $(0, 0, 1)$, $(-1/2, 1/\sqrt{2}, -1/2)$, $(\sqrt{2}/\sqrt{3}, 1/\sqrt{3}, 0)$. An orthonormal representation using only 0 and 1 was done in Pentagons and Paradoxes and gives the minimal angle too $(1, 1, -1)$, $(0, 1, 1)$, $(1, 0, 1)$, $(1, 0, 0)$, $(0, 1, 0)$, $(0, 1, -1)$, $(1, 0, -1)$, $(1, 1, 1)$.

THE ORTHOGONAL REPRESENTATION OF THE PENTAGON DETERMINES UNIVOQUELY THE ORTHOGONAL REPRESENTATION OF THE ENTIRE BUDGE. SEE DRAFT IN ACTUAL PAPER SHEET.

IV. HIGHER DIMENSIONS

SOME PARAGRAPHS CHANGED ORDER IN THIS SECTION

In this section we generalize the results of the previous section to higher dimensions than 3.

Theorem 2: Let be G a critical TIFS graph in dimension d with two K_d , then $V(G) \geq d + 5$.

Proof: For this proof we use the notion of forbidden graph in dimension d for the orthogonal rank of a graph [?]. This concept is related to the impossible representation of a graph with several structure.

For a non trivial TIFS we have to use a context, so the first valid linked structure in dimension d is K_{d-1} linked to true vertex y and false vertex x such that $N(y) = N(x)$. But this graph ($|V(G)| = d + 1$) is not realizable because it form a forbidden graph in dimension d .

Suppose that we have a graph with one more vertex c ($|V(G)| = d + 2$). Is evident that $y \approx c$, so c have to belong to a context with other vertex from K_{d-1} (for now call Q). We consider two cases: a) $N(x) = \{v/v \in Q\}$, Then we have another forbidden graph between c and x . b) Choose $N(x) \cap N(y) = \emptyset$, in that case we have a forbidden structure between y and x .

We must consider a new vertex w ($|V(G)| = d + 3$). Previous results shows that neither y or x can belong to the two K_d necessary for the lemma 5, so the structure between y and x must have two K_d , but we only have

c and w as internal vertex different of Q . Is not enough to evade a forbidden structure again (both will create forbidden graph).

We introduce another vertex t ($|V(G)| = d + 4$). This new vertex will support to anyone internal vertex w or c . w.l.o.g. We consider that t support w , i.e. They belong to the same context. Now we have two K_d from $c \cup N(c) = Q$ and $w \cup t \cup Q - q$ where $q \in N(c)$. Here we can observe that again appear forbidden structure in dimension d between t and q because $N(t) = N(q)$.

Then we must consider $|V(G)| \geq d + 5$ to get a minimal TIFS. In fact, we know a structure of TIFS with $d + 5$ vertices obtained from the Fig. 4.

Example: The graph G of Fig. 1(a) ,with $V(G) = \{A, v_1, v_2, v_4, v_5, v_6, v_7, v_8\}$, is a critical TIFS graph in dimension 3.

The number of TIFSes in dimension 3, 4, 5, 6... is 1, 3, 4, 8, 13, 19... In general, the number of TIFSes in dimension d is $(d^2 - d - 4)/3$. We know all the corresponding graphs and its orthogonal representation. In the further sections, we will explain these results.

Theorem 3: Let be G a critical TITS that contain a TIFS graph with $d + k$ vertices, then G have, at least, $d + k + 2$ vertices.

Proof: We dismiss two different cases: a) Suppose the critical TITS graph has $d + k$ vertices with yes vertices a and c . Then the vertex c must belong to one of two K_d in TIFS with false vertex b , in that case reaches vertex b not be affected by the obtained colored TITS with vertices a and c that, for the Lemma 2, contains a smaller TIFS than the original. b) Suppose the critical TITS with $d + k + 1$ vertices and true vertex a, c . We add a vertex x to TIFS from true vertex a to false vertex b , is clear that the new vertex must be yes, i.e., it must be equal to vertex c , because if c was an internal vertex of TIFS with a yes and b no then we have the previous case.

Theorem 4: Minimal TITS have $d + 7$ vertices in dimension $d \geq 3$.

Proof: The addition of two vertices of the TIFS in dimension d with true vertex a and false vertex b by addition of necessary edges in the TITS with yes vertices a and c can not be a forbidden graph (you can check that the structure with 3 independent triangles full linked to K_{d-3} has dimension d), so this obtained a TITS realizable in dimension d . The general structure of the graph is in Fig. 4.

Corollary 2: The family of minimal TITSes with $d + 7$ vertices contains exactly 3 K_d .

Proof: Consider the structure of minimal TITS (Fig. 4), where $\{a\} \cup C$ is TIFS in dimension d and $|D| = d - 2$. There exist some edges between $C \in \text{TIFS}_{d=3}$ and D that form one more K_d . Furthermore, $\forall v \in D$ verifies that $v \in N(a)$.




FIG. 4. Schema for minimal TITS with $d + k + 2$ vertices in dimension d . The arrow represents connection between a and some vertices in C such that they form a minimal TIFS, also vertex a is linked with all vertices in D .

V. DIMENSION 4 - HARDY BUG & OTHER BUGS

There are three TIFS with a minimum number of propositions in dimension 4, as depicted in Fig 5.

TIPS: (a) The orthogonal representation il (almost) unique (b) minimal angle (c) Explain the three graphs

For $d = 4$, Minimum angle = $\arccos((1 - \varepsilon^2)/3)$ por the graphs (a) and (b) in Fig. 5 and $\arccos(1/3)$ por graph (c)

A. Specific realizations

FIGURES TO ONE COLUMN, BIBTEX ITEMS TO BIBTEX FILE CNLG.BIB

1. Pentagrams and Paradoxes

[26]

2. No-hidden-variables proof for two spin-particles preselected and postselected in unentangled states

[27]

3. Bell-Kochen-Specker theorem: A proof with 18 vectors

[25]

MAYBE A FIGURE WITH CABELLO18 GRAPH AND THE BUGDET INSIDE?




FIG. 5. (Color online) Orthogonality diagram of the three simplest (partially known [24–26]) TIFS in $d = 4$, with 8(9) contexts and 9 atomic propositions. It is realizable in S^3 by taking, for instance, (a) $v_1 = (0, -1, \sqrt{2}, 0)/\sqrt{3}$, $v_2 = (1, 0, 0, 0)$, $v_3 = (1, \sqrt{2}, 1, 0)/2$, $v_4 = (0, 1, 0, 0)$, $v_5 = (1, 0, -1, 0)/\sqrt{2}$, $v_6 = (0, 0, 1, 0)$, $v_7 = (-1, \sqrt{2}, -1, 0)/2$, $v_8 = (\sqrt{1 - \varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0) + \varepsilon(0, 0, 0, 1)$, $v_9 = (0, 0, 0, 1)$, (b) $v_1 = (\sqrt{1 - \varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0) + \varepsilon(0, 0, 0, 1)$, $v_8 = (\sqrt{2}, 1, 0, 0)/\sqrt{3}$, (c) $v_1 = (0, -1, \sqrt{2}, 0)/\sqrt{3}$, $v_8 = (\sqrt{2}, 1, 0, 0)/\sqrt{3}$. Values on v_1 and v_8 cannot both be 1, as this would imply the values on $v_2 = v_3 = v_6 = v_7 = v_9$ to vanish, which in turn would require both values v_4 as well as v_5 to be 1; which contradicts exclusivity because those observables belong to the same context and are orthogonal. In QT, the proposition v_i is represented by the projector $|v_i\rangle\langle v_i|$. **FIGURE IN ONE COLUMN. \texttt{\textbackslashwidetext COMMAND REMOVED}**




FIG. 6. Pentagrams and Paradoxes[26]




FIG. 7. No-hidden-variables proof for two spin-particles pre-selected and postselected in unentangled states [27]




FIG. 8. Bell-Kochen-Specker theorem: A proof with 18 vectors [25]

VI. DIMENSION 5 - GENERALIZED HARDY BUGS

There are four TIFS with a minimum number of propositions in dimension 5. They are shown in Fig 9.

TIPS: (a) The orthogonal representation is (almost) unique (b) Explain the four graphs

For $d = 5$, Minimum angle = $\arccos((1 - \varepsilon^2)/3)$ for all

graphs (a) and (b) in Fig. 5 except $\arccos(1/3)$ por graph (d)

VII. DIMENSION 6

There are eight TIFS with a minimum number of propositions in dimension 6. They are shown in Fig 10.

TIPS: (a) The orthogonal representation il (almost) unique (b) Explain the eight graphs and (c) generalize to higher dimensions

THIS PARAGRAPH CAN BE FULL REWRITTEN. WRONG FORMULAE. MUST BE $\binom{d-1}{2} = (d-1)(d-2)/2$ FOR $d \geq 5$ Para contar los grafos TITS con menor número de vértices, basta con contar las posibles conexiones (excepto isomorfismos) entre los vértices del K_{d-3} interior y los vértices TRUE y FALSE. (NOTA: ¿se puede aprovechar alguna de las figuras?). Como cada vértice del K_{d-3} puede tener 3 estados diferentes e incompatibles (adyacente sólo a TRUE, adyacente sólo a FALSE o adyacente a ambos), tenemos combinaciones con repetición de 3 elementos tomados en grupos de $d-3$. Fórmulas: $\binom{3}{d-3} = \binom{d-1}{d-3} = \binom{d-1}{2} = (d-1)(d-2)/2$.

Orthogonality diagram of the the four simplest known TIFS in $d = 6$, with 9 contexts and 11 atomic propositions. It is realizable in S^5 by taking, for instance, (a) $v_1 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0, 0) + \varepsilon(0, 0, 0, 0, 0, 1)$, $v_2 = (1, 0, 0, 0, 0, 0)$, $v_3 = (1, \sqrt{2}, 1, 0, 0, 0)/2$, $v_4 = (0, 1, 0, 0, 0, 0)$, $v_5 = (1, 0, -1, 0, 0, 0)/\sqrt{2}$, $v_6 = (0, 0, 1, 0, 0, 0)$, $v_7 = (-1, \sqrt{2}, -1, 0, 0, 0)/2$, $v_8 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0, 0) + (\varepsilon/\sqrt{2})(0, 0, 0, 1, 1, 0)$, $v_9 = (0, 0, 0, 1, 0, 0)$, $v_{10} = (0, 0, 0, 0, 1, 0)$, $v_{11} = (\sqrt{1-\varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0, 0) + (\varepsilon/\sqrt{2})(0, 0, 0, 0, 1, 1)$, (b) $v_1 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0, 0) + \varepsilon(0, 0, 0, 1, 0, 0)$, $v_2 = (1, 0, 0, 0, 0, 0)$, $v_3 = (1, \sqrt{2}, 1, 0, 0, 0)/2$, $v_4 = (0, 1, 0, 0, 0, 0)$, $v_5 = (1, 0, -1, 0, 0, 0)/\sqrt{2}$, $v_6 = (0, 0, 1, 0, 0, 0)$, $v_7 = (-1, \sqrt{2}, -1, 0, 0, 0)/2$, $v_8 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0, 0) + \varepsilon(0, 0, 0, 1, 0, 0)$, $v_9 = (0, 0, 0, 1, 0, 0)$, $v_{10} = (0, 0, 0, 0, 1, 0)$, $v_{11} = (\sqrt{1-\varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0, 0) + (\varepsilon/\sqrt{2})(0, 0, 0, 0, 1, 1)$.

(c) $v_1 = (0, -1, \sqrt{2}, 0, 0, 0)/\sqrt{3}$, $v_8 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0, 0) + (\varepsilon/\sqrt{2})(0, 0, 0, 1, 1, 0)$, (d) $v_1 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0, 0) + (\varepsilon/\sqrt{2})(0, 0, 0, 0, 1, 1)$, $v_8 = (\sqrt{2}, 1, 0, 0, 0, 0)/\sqrt{3}$, (e) $v_1 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0, 0) + \varepsilon(0, 0, 0, 0, 0, 1)$, $v_8 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0, 0) + \varepsilon(0, 0, 0, 1, 0, 0)$, (f) $v_1 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0, 0) + \varepsilon(0, 0, 0, 0, 0, 1)$, $v_8 = (\sqrt{2}, 1, 0, 0, 0, 0)/\sqrt{3}$, (g) $v_1 = (0, -1, \sqrt{2}, 0, 0, 0)/\sqrt{3}$, $v_8 = (\sqrt{1-\varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0, 0) + \varepsilon(0, 0, 0, 1, 0, 0)$, (h) $v_1 = (0, -1, \sqrt{2}, 0, 0, 0)/\sqrt{3}$, $v_8 = (\sqrt{2}, 1, 0, 0, 0, 0)/\sqrt{3}$. Values on v_1 and v_8 cannot both be 1, as this would imply the values on $v_2 = v_3 = v_6 = v_7 = v_9 = v_{10} = v_{11}$ to vanish, which in turn would require both values $v_4 = v_5$ to be 1; which contradicts exclusivity because those observables belong to the same context and are orthogonal. In QT, the proposition v_i is represented by the projector $|v_i\rangle\langle v_i|$.

For $d = 6$, Minimum angle = $\arccos((1-\varepsilon^2)/3)$ for all graphs (a) and (b) in Fig. 5 except $\arccos(1/3)$ for graph (h). GENERALIZE FOR HIGHER DIMENSIONS

VIII. CONCLUSIONS AND OPEN PROBLEMS

Peres conjectured that the simplest possible state-independent proof is one in [?], which requires 18 propositions in $d = 4$ [?]. The method we have developed in this article, and the obtained minimal TITS, can be helped to prove Peres' conjecture [?]. Other related open problem which can benefit from these results is which is the simplest state-independent proof in $d = 3$.

ACKNOWLEDGMENTS

This work was supported by the Project No. FIS2011-29400 (MINECO, Spain) with FEDER funds, the FQXi large grant project "The Nature of Information in Sequential Quantum Measurements," and the Brazilian program Science without Borders.

-
- [1] A. Cabello, S. Severini, and A. Winter, in *American Mathematical Society, 2010 Fall Southeastern Section Meeting (Richmond, Virginia, Nov 6-7, 2010)* (2010).
 - [2] A. Cabello, S. Severini, and A. Winter, *Phys. Rev. Lett.* **112**, 040401 (5 pages) (2014), URL <http://link.aps.org/doi/10.1103/PhysRevLett.112.040401>.
 - [3] S. Kochen and E. P. Specker, in *Theory of Models: Proceedings of the 1963 International Symposium [on the Theory of Models, held] at Berkeley, [from June 25 to July 11 1963]*, edited by J. W. Addison, L. Henkin, and A. Tarski (North-Holland, Amsterdam, 1965), pp. 177–189.
 - [4] S. Kochen and E. P. Specker, *Journal of Mathematics and Mechanics* **17**, 59 (1967).
 - [5] A. Stairs, *Philosophy of Science* **50**, 578 (1983), URL <http://doi.org/10.1086/289140>.
 - [6] R. J. Greechie, *Journal of Combinatorial Theory. Series A* **10**, 119 (1971), URL [http://doi.org/10.1016/0097-3165\(71\)90015-X](http://doi.org/10.1016/0097-3165(71)90015-X).
 - [7] I. Pitowsky, *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* **34**, 395 (2003), ISSN 1355-2198, quantum Information and Computation, arXiv:quant-ph/0208121, URL [http://doi.org/10.1016/S1355-2198\(03\)00035-2](http://doi.org/10.1016/S1355-2198(03)00035-2).
 - [8] I. Pitowsky, in *Physical Theory and its Interpretation*, edited by W. Demopoulos and I. Pitowsky (Springer Netherlands, 2006), vol. 72 of *The Western Ontario Series in Philosophy of Science*, pp. 213–240, ISBN 978-1-4020-4875-3, arXiv:quant-ph/0510095, URL http://doi.org/10.1007/1-4020-4876-9_10.
 - [9] P. Pták and S. Pulmannová, *Orthomodular Structures as Quantum Logics. Intrinsic Properties, State Space and*




FIG. 9. (Color online) Orthogonality diagram of the the four simplest known \square TIFS in $d = 5$, with 9 contexts and 10 atomic propositions. It is realizable in S^4 by taking, for instance, (a) $v_1 = (\sqrt{1 - \varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0) + \varepsilon(0, 0, 0, 0, 1)$, $v_2 = (1, 0, 0, 0, 0)$, $v_3 = (1, \sqrt{2}, 1, 0, 0)/2$, $v_4 = (0, 1, 0, 0, 0)$, $v_5 = (1, 0, -1, 0, 0)/\sqrt{2}$, $v_6 = (0, 0, 1, 0, 0)$, $v_7 = (-1, \sqrt{2}, -1, 0, 0)/2$, $v_8 = (\sqrt{1 - \varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0) + \varepsilon(0, 0, 0, 1, 0)$, $v_9 = (0, 0, 0, 1, 0)$, $v_{10} = (0, 0, 0, 0, 1)$, (b) $v_1 = (0, -1, \sqrt{2}, 0, 0)/\sqrt{3}$, $v_8 = (\sqrt{1 - \varepsilon^2}/\sqrt{3})(\sqrt{2}, 1, 0, 0, 0) + \varepsilon(0, 0, 0, 1, 0)$, (c) $v_1 = (\sqrt{1 - \varepsilon^2}/\sqrt{3})(0, -1, \sqrt{2}, 0, 0) + \varepsilon(0, 0, 0, 1, 0)$, $v_8 = (\sqrt{2}, 1, 0, 0, 0)/\sqrt{3}$, (d) $v_1 = (0, -1, \sqrt{2}, 0, 0)/\sqrt{3}$, $v_8 = (\sqrt{2}, 1, 0, 0, 0)/\sqrt{3}$. Values on v_1 and v_8 cannot both be 1, as this would imply the values on $v_2 = v_3 = v_6 = v_7 = v_9 = v_{10} = v_{11}$ to vanish, which in turn would require both values $v_4 = v_5$ to be 1; which contradicts exclusivity because those observables belong to the same context and are orthogonal. In QT, the proposition v_i is represented by the projector $|v_i\rangle\langle v_i|$. **CORRECTED FIGURE (a) FOR SIMILAR COLORS TO OTHER FIGURES**

- Probabilistic Topics*, vol. 44 of *Fundamental Theories of Physics* (Kluwer Academic Publishers, Springer Netherlands, Dordrecht, 1991), ISBN 978-0-7923-1207-9.
- [10] K. Svozil and J. Tkadlec, *Journal of Mathematical Physics* **37**, 5380 (1996), URL <http://doi.org/10.1063/1.531710>.
- [11] J. Tkadlec, *International Journal of Theoretical Physics* **37**, 203 (1998), URL <http://doi.org/10.1023/A:1026646229896>.
- [12] F. J. Belinfante, *A Survey of Hidden-Variables Theories*, vol. 55 of *International Series of Monographs in Natural Philosophy* (Pergamon Press, Elsevier, Oxford, New York, 1973), ISBN 978-0-08-017032-9, 0080170323, URL <http://doi.org/10.1016/B978-0-08-017032-9.50001-7>.
- [13] Itamar, *Philosophy of Science* **49** (1982), URL <http://doi.org/10.2307/187281>.
- [14] S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley* (North Holland, Amsterdam, New York, Oxford, 1965), pp. 177–189, ISBN 9781483275345, reprinted in Ref. [28], pp. 209–221], URL <https://www.elsevier.com/books/the-theory-of-models/addison/978-0-7204-2233-7>.
- [15] *American Journal of Physics* **61**, 443 (1993), URL <http://doi.org/10.1119/1.17239>.
- [16] S. Kochen and E. P. Specker, *Journal of Mathematics and Mechanics* (now *Indiana University Mathematics Journal*) **17**, 59 (1967), ISSN 0022-2518, URL <http://doi.org/10.1512/iumj.1968.17.17004>.
- [17] H. B. Johansen, *American Journal of Physics* **62** (1994), URL <http://doi.org/10.1119/1.17551>.
- [18] P. E. Vermaas, *American Journal of Physics* **62** (1994), URL <http://doi.org/10.1119/1.17488>.
- [19] A. Cabello and G. García-Alcaine, *Journal of Physics A: Mathematical and General Physics* **29** (1996), URL <http://doi.org/10.1088/0305-4470/29/5/016>.
- [20] K. Svozil, *International Journal of Theoretical Physics* **44**, 745 (2005), arXiv:quant-ph/0209136, URL <http://doi.org/10.1007/s10773-005-7052-0>.
- [21] R. Wright, *Foundations of Physics* **20**, 881 (1990), URL <http://doi.org/10.1007/BF01889696>.
- [22] M. Schaller and K. Svozil, *International Journal of Theoretical Physics* **35** (1996), URL <http://doi.org/10.1007/BF02302381>.
- [23] A. Solís-Encina and J. Portillo, arXiv [math.CO] **1504.03662**, 1 (2015), URL <http://arxiv.org/abs/1504.03662>.
- [24] L. Hardy, *Physical Review Letters* **71**, 1665 (1993), URL <http://dx.doi.org/10.1103/PhysRevLett.71.1665>.
- [25] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, *Physics Letters A* **212**, 183 (1996), arXiv:quant-




FIG. 10. (Color online) see text

- ph/9706009, URL [http://doi.org/10.1016/0375-9601\(96\)00134-X](http://doi.org/10.1016/0375-9601(96)00134-X).
- [26] P. Badziąg, I. Bengtsson, A. Cabello, H. Granström, and J.-A. Larsson, *Foundations of Physics* **41** (2011), URL <http://doi.org/10.1007/s10701-010-9433-3>.
- [27] A. Cabello, *Physical Review A* **55**, 4109 (1997), URL <http://doi.org/doi/10.1103/PhysRevA.55.4109>.
- [28] E. Specker, *Selecta* (Birkhäuser Verlag, Basel, 1990), ISBN 978-3-0348-9966-6, 978-3-0348-9259-9, URL <http://doi.org/10.1007/978-3-0348-9259-9>.