

# Functional Epistemology “Nullifies” Dyson’s Rebuttal of Perturbation Theory

Karl Svozil <sup>1</sup> 

<sup>1</sup> Institute for Theoretical Physics, TU Wien, Wiedner Hauptstrasse 8-10/136, 1040 Vienna, Austria; [svozil@tuwien.ac.at](mailto:svozil@tuwien.ac.at); <http://tph.tuwien.ac.at/~svozil>

**Abstract:** Functional epistemology is about ways to access functional objects, either conceptualized by ontology or nominalism, by some operational algorithmic means. Not all such means are equally capable of reproducing these functions in the desired consistency and resolution. Dyson’s argument against the perturbative expansion of quantum field theoretic terms, in a radical form (never pursued by Dyson), is an example of epistemology taken as ontology.

**Keywords:** Asymptotic divergence, perturbation series, partial function

## 1. Functional epistemology

Regardless of one’s metaphysical preferences about ontological realism or Platonism—claiming that “some [[mathematical or physical objects or]] entities sometimes exist without being experienced by any finite mind” [1,2]—versus mathematical nominalism claiming that mathematical entities such as numbers and functions do not exist, quasi “a subject with no object” [3]—every application of mathematical formalism requires some operational access to these objects and entities. In a broader perspective this can also be seen as semantics in need for a syntactical formalization.

One important aspect of access is a representation of functional objects and entities that in some formal form correspond to important aspects of those objects and entities. Nevertheless, although representations vary—spanning a wide range of efficacies and deficiencies—they should not be confused with the respective mathematical objects or entities.

The original informal conception of function  $y = f(x)$  was that of a unique association of an output “value”  $y$  given an input “argument”  $x$ . More recent conceptions consider ordered pairs  $(x, y)$ , where again  $x$  stands for argument(s) and  $y$  for unique value(s); in particular, there must not be two pairs  $(x, y)$  and  $(x, y')$  with  $y \neq y'$ .

This naive functional conception was challenged by Gödel’s, Kleene’s, and Turing’s formalization of what functional “access” means; for instance, in the form of paper and pencil operations on a “paper machine” [4]. These developments closely followed the paradigm change from Cantor’s naive set theory [5] to axiomatic set theories [6]; for instance, Zermelo-Fraenkel set theory. Indeed, from a foundational perspective, it might not get worse: Rice’s theorem, usually proven by reduction to the halting problem, states that any nontrivial semantic property of a computable function (evaluated by a Turing machine) is undecidable.

Therefore, due to incompleteness and related theorems, for the sake of formalization, these earlier intuitive and heuristic perceptions of functional performance had to be modified and restricted. The current formalization of functions is in terms of “desirable” properties, such as, in particular, effective computability. This has resulted in the abandonment of functional totality—the pretension that any (every) arbitrary argument  $x$  can be associated with a (unique) value  $y$ —in favor of partiality: certain functions, such as predictors of the large-scale performance of deterministic systems, need not have a value accessible by some algorithm—in short, there is a difference between determinism and

**Citation:** Svozil, K. Functional Epistemology “Nullifies” Dyson’s Rebuttal of Perturbation Theory.

*Axioms* **2021**, *1*, 0. <https://doi.org/>

Received:

Accepted:

Published:

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Copyright:** © 2023 by the author. Submitted to *Axioms* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

predictability [7]. In such a regime, it can no longer be maintained that the value  $y$  exists—that is, can be obtained or accessed by some algorithm or computation.

Typical examples of such “critical” functions are Turing’s halting function, or Specker’s theorems of recursive analysis [8–10], or Chaitin’s “halting probability”  $\Omega$  in terms of its bitwise expansion [11]. Physical realizations have been, for instance, by reduction to the halting problem [12], suggested in terms of undecidable classical dynamics [13,14],  $N$ -body problems [15], or spectral gaps [16].

In particular, it can happen that a program implementable on a computer “computes a limit”, but this limit is uncomputable and algorithmically inaccessible: Because some resources, such as computing time or space, that are necessary to compute this limit with, say, precision up to its  $n$ th bit, grow faster asymptotically than any computable function of  $n$  [17]. In intuitive algorithmic terms the difference between a total versus a partial function may be imagined as the distinction between a DO-loop (with fixed finite beginning, ending and increments) and a WHILE-loop. The latter WHILE-loop may or may not “take forever”, depending on the respective termination condition.

Another area of partial value assignments is quantum mechanics. Extensions of the Kochen-Specker theorem suggest that, relative to the assumption of noncontextuality, only a “star-shaped” [18, Fig. 5] (in terms of hypergraphs [19] representing individual contexts by smooth curves [20]) context can have definite value assignments. Observables in all other contexts must be value indefinite [21–23].

Still another issue of functional epistemology is the means relativity of functional representation. The same function can have very different representations and encodings; some exhibiting more or less problematic issues. For the sake of an example, we shall later represent the same function in five different forms.

The selection of particular means is often not a matter of choice but one of pragmatism or even desperation. In particular, theoretical physicists are often criticized for their “relaxed” stance on formal rigor. Dirac’s introduction of the needle-shaped delta function is often quoted as an example. Heaviside, in another instance, responded to criticism for his use of the “highly nonsmooth” unit step function [24, p. 9, § 225]: “*But then the rigorous logic of the matter is not plain! Well, what of that? Shall I refuse my dinner because I do not fully understand the process of digestion? No, not if I am satisfied with the result.*”

This, in a nutshell, seems to be the attitude of field theorists regarding the use of perturbation series: It is well documented [25–27] that the commonly used power series expansion which can be rewritten as inverse power series expansion

$$\begin{aligned} f(\alpha) &= \alpha a_0 + a_1 \alpha + a_2 \alpha^2 + \cdots \\ &= \sum_{n=0}^{\infty} a_n \alpha^n = \frac{1}{\alpha} \sum_{n=0}^{\infty} \frac{a_n}{\left(\frac{1}{\alpha}\right)^{n+1}}. \end{aligned} \quad (1)$$

in terms of the fine structure constant  $\alpha$ —that is, the square of the) coupling constant—is divergent.

For this power series to converge, there has to be a finite radius of convergence centered at the origin at (fictitious) value  $\alpha = 0$ , thereby including (fictitious) nonvanishing negative values  $\alpha < 0$  within which  $F$  has to be analytic. However, because if the (fictitious) coupling between like charges becomes negative, and because by tunneling this cannot be “contained”, the vacuum becomes unstable due to pair creation, and (fictitiously) disintegrates explosively. Hence, Dyson concludes, the power series  $f(-\alpha)$  cannot converge and thus cannot be analytic—a complete contradiction to the assumption.

An immediate reaction would be to perceive these coincidences as “bordering on the mysterious” [28]. This spirit is corroborated [29] by statements like Carrier’s Rule, pointing out that “divergent series converge faster than convergent series because they don’t have to converge.” However, as already surmised by Dyson, quantitative considerations from partial sum show [30, p. 4] that convergent series “initially”—that is, with only “a few orders” added—may “largely deviate” from the true value of the function it encodes (eg,

Let  $S(x) = \int_0^\infty e^{-t}/(1+tx)dt$  stand for the Stieltjes function (cf. [48, formula 5.1.28, page 230] but with  $x \mapsto \frac{1}{x}$ ),  $\gamma = \lim_{n \rightarrow \infty} \left( \sum_{j=1}^n \frac{1}{j} - \log n \right) \approx 0.5772$ , be the Euler-Mascheroni constant [49],  $\Gamma(z, x)$  represent the upper incomplete gamma function [48, formula 6.5.3, page 260],  $\mathfrak{B}[s(x)]$  be the Borel transform of  $s(x)$ ,  $(x)_n = \Gamma(x+n)/\Gamma(x) = x(x+1) \cdots (x+n-1)$  and  $(x)_0 = 1$  be Pochhammer symbols [48, Section 24.1.3, page 824], also known as the

rising factorial power, and  $S_j^{(k)}$  be Sterling numbers of the first kind that are the polynomial coefficients of the Pochhammer symbol  $(z - j + 1)_j$  [48, Section 24.1.3, page 824]; that is,  $\sum_{k=0}^j S_j^{(k)} z^k = (z - j + 1)_j = (-1)^j (-z)_j$  for  $j \in \mathbb{N} \cup \{0\}$ . Then

$$\begin{aligned} s(x) &= xS(x) = e^{\frac{1}{x}} \Gamma\left(0, \frac{1}{x}\right) \\ &= -e^{\frac{1}{x}} \left[ \gamma - \log x + \sum_{n=1}^{\infty} \frac{(-1)^n}{n! n x^n} \right] \end{aligned} \quad (4)$$

$$\begin{aligned} &= \int_0^{\infty} \mathcal{BS}(y) e^{-\frac{y}{x}} dy \\ &= \int_0^{\infty} \frac{e^{-\frac{y}{x}}}{1+y} dy = \int_0^{\infty} \frac{x e^{-t}}{1+xt} dt \end{aligned} \quad (5)$$

$$= \int_0^x \frac{e^{\frac{1}{x}-\frac{1}{t}}}{t} dt \quad (6)$$

$$= \sum_{j=0}^{\infty} (-1)^j j! x^{j+1} \quad (7)$$

$$= x \sum_{j=0}^{\infty} \frac{(-1)^j}{\left(\frac{1}{x}\right)_{j+1}} \sum_{k=0}^j S_j^{(k)} k! \quad (8)$$

What we can learn from this prototypic example is the wide variety of mathematical representations associated with some function. Not everybody might agree with all the equality signs in (4)–(8) and the legality of the respective methods though, thereby reflecting a variety of metamathematical stances.

### 3. Quantum field theoretical perturbation series need not diverge

Let us discuss two critical aspects in the derivation of the power series expansion of (1). One critical step in the derivation of  $f$  amounts to interchanging a sum with an integral in the case of nonuniform convergence of the former [50, Sect. II.A]. One may perceive asymptotic divergence as a “penalty” for such manipulations. It may come as a surprise that those calculations performed well for empirical predictions.

Ritt’s theorem inspires one strategy to cope with such issues [51,52] by stating that any (not necessarily asymptotic) divergent power series with arbitrary coefficients can be converted into nonunique analytic functions. Thereby, every summand is multiplied with a suitable nonunique “convergence factor.” (Conversely, every analytic function can be approximated by a unique asymptotic series.)

A general regularisation of divergent series using such convergence factors, also called *cutoff functions*, has been recently introduced by Tao [53, Section 3.7]. The resulting smoothed sums may become uniformly convergent, thereby allowing interchanging a sum with an integral and avoiding the aforementioned issues while preserving inherent properties of the original divergent series. This is not dissimilar to the use of test functions in the theory of distributions.

A second critical aspect is the expansion of (1) in terms of a power series, and the resulting vanishing of the radius of convergence. Dyson already mentioned a possible remedy, his “*Alternative A: There may be discovered a new method of carrying through the renormalization program, not making use of power series expansions.*” One such candidate expansion that does not necessarily share the catastrophic fate of the power series caused by the “explosive disintegration” of the vacuum state for negative arguments, is an expansion

of  $f(\alpha)$  in terms of inverse factorial series [54,55] and recently investigated by Weniger [47] as well as O. Costin, R. D. Costin and Dunne [56,57]:

$$f(\alpha) = b_0 \frac{0!}{\alpha} + b_1 \frac{1!}{\alpha(\alpha+1)} + \dots = \sum_{n=0}^{\infty} b_n \frac{n!}{(\alpha)_{n+1}}, \quad (9)$$

where again  $(\alpha)_{n+1}$  are Pochhammer symbols.

Stirling numbers of the first kind  $S_j^{(k)}$  mentioned earlier serve as “translations”—that is, as expansions from an inverse power  $1/\alpha^{n+1}$  in terms of inverse factorial series  $(\alpha)_{n+j+1}$ : for  $k \in \mathbb{N} \cup \{0\}$  [58, Equation (6), § 30, p. 78],

$$\frac{1}{\alpha^{n+1}} = \sum_{j=0}^{\infty} \frac{(-1)^j}{(\alpha)_{n+j+1}} S_{n+j}^{(n)}. \quad (10)$$

The respective “reverse” expansion of a Pochhammer symbol  $(\alpha)_{k+1}$  in terms of an inverse power series  $1/\alpha^{n+j+1}$  for  $k \in \mathbb{N} \cup \{0\}$  and  $|\alpha| > 0$  [58, Equation (9), § 26, p. 68] is given by

$$\frac{1}{(\alpha)_{n+1}} = \sum_{j=0}^{\infty} \frac{(-1)^j}{\alpha^{n+j+1}} S_{n+j}^{(n)}. \quad (11)$$

Insertion of (10) into (1), rearranging the order of the summations through an index shift  $m = n + j$  with  $n \geq 0$  and  $j \geq 0$ , hence  $m \geq 0$  and  $j = m - n \geq 0$  and  $n \leq m$  yields

$$\begin{aligned} f(\alpha) &= \frac{1}{\alpha} \sum_{n=0}^{\infty} a_n \sum_{j=0}^{\infty} \frac{(-1)^j}{\left(\frac{1}{\alpha}\right)_{n+j+1}} S_{n+j}^{(n)} \\ &= \frac{1}{\alpha} \sum_{j=0}^{\infty} \sum_{n=0}^{\infty} a_n \frac{(-1)^j}{\left(\frac{1}{\alpha}\right)_{n+j+1}} S_{n+j}^{(n)} \\ &= \frac{1}{\alpha} \sum_{m=0}^{\infty} \frac{(-1)^m}{\left(\frac{1}{\alpha}\right)_{m+1}} \sum_{n=0}^m (-1)^{\pm n} S_m^{(n)} a_n. \end{aligned} \quad (12)$$

Therefore, if we define the inverse power series  $f'(\beta) = (1/\beta)f(1/\beta) = \sum_{m=0}^{\infty} a'_m/\beta^{m+1} = \sum_{m=0}^{\infty} b'_m m! / (\beta)_{m+1}$  with  $\beta = 1/\alpha$ , then, by comparison,

$$b'_m = \frac{1}{m!} \sum_{n=0}^m (-1)^{m \pm n} S_m^{(n)} a'_n. \quad (13)$$

For the sake of studying the fascinating convergence [29,47,55,58] of the inverse factorial series, note that terms of the form  $n!/(z)_{n+1}$  can, for large  $z \rightarrow \infty$ , be estimated with the help of [48, Formula 6.1.47]  $\Gamma(z+a)/\Gamma(z+b) = z^{a-b} \left[1 + O\left(\frac{1}{z}\right)\right]$  and for large  $n \rightarrow \infty$  as follows:

$$\begin{aligned} \frac{n!}{(\alpha)_{n+1}} &= \frac{\Gamma(n+1)}{[\Gamma(\alpha+n+1)/\Gamma(\alpha)]} = \frac{\Gamma(n+1)}{\Gamma(n+1+\alpha)} \Gamma(\alpha) \\ &= (n+1)^{-\alpha} \left[1 + O\left(\frac{1}{n+1}\right)\right] (\alpha-1)! = O(n^{-\alpha}). \end{aligned} \quad (14)$$

Therefore, the inverse factorial series (9) converges with the possible exception of the points  $\alpha = -m$  with  $m \in \mathbb{N} \cup \{0\}$  (where the Pochhammer symbols in the denominator might vanish) if and only if the associated Dirichlet series  $\sum_{n=1}^{\infty} c_n n^{-\alpha}$  converges.

Unlike a power series that has a radius of convergence, a Dirichlet series has an abscissa of convergence  $\Re(\alpha) > \lambda$ , that is, it converges on this half-plane [59, § 58, 255,

page 456]. Therefore, the inverse factorial series may converge for all positive coupling constants  $\alpha > 0$  although it may diverge for negative values  $\alpha < 0$ . The physically relevant region lies within the abscissa of convergence. Even if the inverse power series diverges factorially, the respective inverse factorial series may converge, but this has to be checked explicitly.

Thus, as already suggested by Dyson [25], representing quantum field theoretical entities in terms of the Tomonaga-Schwinger-Feynman-Dyson power series expansion in the coupling constant [60] may suffice for all practical purposes [61] so far, although their divergencies may cause uneasiness for a variety of (pragmatic and formal) reasons. One should not confuse these field-theoretic entities with their actual representations; that is, functional ontology with epistemology. We have intended to present a positive outlook for an improved theory of convergent perturbation series. A candidate for such a theory might be inverse factorial expansions exhibiting an abscissa rather than a radius of convergence.

However, a convergence issue encountered in inverse factorial series is the Stokes phenomenon [56,57]: the asymptotic behavior of functions need not be uniform in different regions of the complex plane, bounded by (anti-)Stokes lines. In particular, inverse factorial series may not be suitable for the study of Stokes phenomena if Stokes lines are present in the right complex half-plane  $\Re(\alpha) > \lambda$  because of the singularities on these Stokes lines. One may conjecture that inverse factorials might converge in regions where the associated power series are Borel summable; yet convergence fails in the presence of Stokes lines. This would mean that quantum field theories have convergent inverse factorial expansions only in less than four dimensions; and that this expansion might fail for four dimensions. Nevertheless, Berry, O Costin, R. D. Costin and Howls have pointed out [62, p. 10] that, although “typically, classical factorial series have two major limitations: slow convergence, at best power-like, and limited domain of convergence (a half plane which cannot be centered on the asymptotically important Stokes line)  $\dots$  for resurgent functions these limitations can be overcome. Ecalle-Borel summable series can be summed by rapidly convergent factorial series.”

Transseries from Borel-Ecalle summations of divergent (power) series offer a method to “recover” nonperturbative information from such power series [63,64], thereby indicating that the divergent perturbative power series expansion contains information of the nonperturbative kind. The situation is not totally dissimilar from tempered distributions, with test functions with unbounded (noncompact) support, allowing the representation and reconstruction of generalized functions by Fourier transforms.

A further method for alternative representations of functions are Padé approximations by rational functions (of given order) near a specific point. Padé approximants offer practical methods of definition and the computation of the value of a power series even if such series diverge [65].

#### 4. Summary

In this brief exposé I have bundled together two ideas: first, that mathematical objects or entities may have very varied representations and realizations [66]. Not all of them might require comparable means to access them—think of convergence or (asymptotic) divergence, or of partial functions. Different means might not be equally appropriate or sufficient and necessary for different purposes.

Second, in particular and more specifically, as conjectured already by Dyson, arguments against the existence or convergence of power expansions of Tomonaga-Schwinger-Feynman-Dyson perturbative quantum field theory might be “liftable” by using other expansion techniques.

For the sake of illustration, suppose for a moment that Gödel’s “unadulterated” Platonism [2,67] is acceptable. (An analogous argument can be made within nominalism.) Then mathematical objects or entities such as functions can be conceptualized by their ontological existence.



However, on second thought, it is an entirely different, highly nontrivial, issue to “touch” or to epistemically access those objects or entities. We have presented some examples of such access which analytically spread over a wide variety of (asymptotic) divergent and convergent expressions.

We have, in particular, argued that (asymptotic) divergence of a particular type of perturbation series based on power series expansions, could be overcome by other methods of perturbative series; in particular, by inverse factorial series. This still leaves open the consistent existence of quantum field theory, but at least it indicates conceivable convergent access to quantum field theoretical objects and functions.

**Funding:** This research was funded in whole, or in part, by the Austrian Science Fund (FWF), Project No. I 4579-N. For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

**Acknowledgments:** I kindly acknowledge discussions with and suggestions by Thomas Sommer.

#### Conflicts of Interest:

The author declares no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

#### References

1. Stace, W.T. The Refutation of Realism. *Mind* **1934**, *43*, 145–155. <https://doi.org/10.1093/mind/XLIII.170.145>.
2. Parsons, C. Platonism and Mathematical Intuition in Kurt Gödel’s Thought. *Bulletin of Symbolic Logic* **1995**, *1*, 44–74. <https://doi.org/10.2307/420946>.
3. Burgess, J.P.; Rosen, G. *A Subject With No Object*; Oxford University Press, 1999. <https://doi.org/10.1093/0198250126.001.0001>.
4. Turing, A.M. Intelligent Machinery. In *Cybernetics. Key Papers*; Evans, C.R.; Robertson, A.D.J., Eds.; Butterworths: London, 1968; pp. 27–52.
5. Halmos, P.R. *Naïve Set Theory*; Undergraduate Texts in Mathematics, Springer: New York, NY, USA, 1974. <https://doi.org/10.1007/978-1-4757-1645-0>.
6. Hrbacek, K.; Jech, T. *Introduction to Set Theory*, third ed.; CRC Press, 1999. <https://doi.org/10.1201/9781315274096>.
7. Suppes, P. The Transcendental Character of Determinism. *Midwest Studies In Philosophy* **1993**, *18*, 242–57. <https://doi.org/10.1111/j.1475-4975.1993.tb00266.x>.
8. Specker, E. Der Satz vom Maximum in der rekursiven Analysis. In *Proceedings of the Constructivity in mathematics: proceedings of the colloquium held at Amsterdam, 1957*; Heyting, A., Ed.; North-Holland Publishing Company: Amsterdam, 1959; pp. 254–265. Reprinted in Ref. [9], pp. 148–159]; English translation: *Theorems of Analysis which cannot be proven constructively*, [https://doi.org/10.1007/978-3-0348-9259-9\\_12](https://doi.org/10.1007/978-3-0348-9259-9_12).
9. Specker, E. *Selecta*; Birkhäuser Verlag: Basel, 1990. <https://doi.org/10.1007/978-3-0348-9259-9>.
10. Kreisel, G. A notion of mechanistic theory. *Synthese* **1974**, *29*, 11–26. <https://doi.org/10.1007/BF00484949>.
11. Calude, C.S.; Dinneen, M.J. Exact Approximations of Omega Numbers. *International Journal of Bifurcation and Chaos* **2007**, *17*, 1937–1954. CDMTCS report series 293, <https://doi.org/10.1142/S0218127407018130>.
12. Yanofsky, N.S. Paradoxes, Contradictions, and the Limits of Science. *American Scientist* **2016**, *104*, 166. <https://doi.org/10.1511/2016.120.166>.
13. Moore, C.D. Unpredictability and undecidability in dynamical systems. *Physical Review Letters* **1990**, *64*, 2354–2357. Cf. Ch. Bennett, *Nature*, **346**, 606 (1990), <https://doi.org/10.1103/PhysRevLett.64.2354>.
14. Bennett, C.H. Undecidable dynamics. *Nature* **1990**, *346*, 606–607. <https://doi.org/10.1038/346606a0>.
15. Svozil, K. Omega and the time evolution of the *N*-body problem. In *Proceedings of the Randomness and Complexity, from Leibniz to Chaitin*; Calude, C.S., Ed.; World Scientific: Singapore, 2007; pp. 231–236, [arXiv:physics/0703031]. [https://doi.org/10.1142/9789812770837\\_0013](https://doi.org/10.1142/9789812770837_0013).
16. Cubitt, T.S.; Perez-Garcia, D.; Wolf, M.M. Undecidability of the spectral gap. *Nature* **2015**, *528*, 207–211, [arXiv:1502.04135,arXiv:1502.04573]. <https://doi.org/10.1038/nature16059>.

17. Rado, T. On Non-Computable Functions. *The Bell System Technical Journal* **1962**, *XLI*(41), 877–884. <https://doi.org/10.1002/j.1538-7305.1962.tb00480.x>. 256
18. Abbott, A.A.; Calude, C.S.; Svozil, K. Value-indefinite observables are almost everywhere. *Physical Review A* **2014**, *89*, 032109, [arXiv:1309.7188]. <https://doi.org/10.1103/PhysRevA.89.032109>. 257
19. Bretto, A. *Hypergraph theory*; Mathematical Engineering, Springer: Cham, Heidelberg, New York, Dordrecht, London, 2013; pp. xiv+119. <https://doi.org/10.1007/978-3-319-00080-0>. 258
20. Greechie, R.J. Orthomodular lattices admitting no states. *Journal of Combinatorial Theory. Series A* **1971**, *10*, 119–132. [https://doi.org/10.1016/0097-3165\(71\)90015-X](https://doi.org/10.1016/0097-3165(71)90015-X). 259
21. Pitowsky, I. Infinite and finite Gleason's theorems and the logic of indeterminacy. *Journal of Mathematical Physics* **1998**, *39*, 218–228. <https://doi.org/10.1063/1.532334>. 260
22. Hrushovski, E.; Pitowsky, I. Generalizations of Kochen and Specker's theorem and the effectiveness of Gleason's theorem. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* **2004**, *35*, 177–194, [arXiv:quant-ph/0307139]. <https://doi.org/10.1016/j.shpsb.2003.10.002>. 261
23. Abbott, A.A.; Calude, C.S.; Svozil, K. A variant of the Kochen-Specker theorem localising value indefiniteness. *Journal of Mathematical Physics* **2015**, *56*, 102201(1–17), [arXiv:1503.01985]. <https://doi.org/10.1063/1.4931658>. 262
24. Heaviside, O. *Electromagnetic theory*; "The Electrician" Printing and Publishing Corporation: London, 1894–1912. 263
25. Dyson, F.J. Divergence of Perturbation Theory in Quantum Electrodynamics. *Physical Review* **1952**, *85*, 631–632. <https://doi.org/10.1103/PhysRev.85.631>. 264
26. Le Guillou, J.C.; Zinn-Justin, J. *Large-Order Behaviour of Perturbation Theory*; Vol. 7, *Current Physics-Sources and Comments*, North Holland, Elsevier: Amsterdam, 1990,2013. 265
27. Vainshtein, A.I. Decaying Systems and Divergence of the Series of Perturbation Theory. In *Proceedings of the Continuous Advances in QCD 2002*; World Scientific Publishing Co. Pte. Ltd.: Singapore, 2002. Novosibirsk Institute of Nuclear Physics Report, December 1964; reprinted in the proceedings of the Conference "2002 Arkadyfest-Honoring the 60th Birthday of Arkady Vainshtein" held 17-23 May 2002 at the University of Minnesota; foreword and translation by M. A. Shifman, [https://doi.org/10.1142/9789812776310\\_others01](https://doi.org/10.1142/9789812776310_others01). 266
28. Wigner, E.P. The unreasonable effectiveness of mathematics in the natural sciences. Richard Courant Lecture delivered at New York University, May 11, 1959. *Communications on Pure and Applied Mathematics* **1960**, *13*, 1–14. <https://doi.org/10.1002/cpa.3160130102>. 267
29. Landau, E. Über die Grundlagen der Theorie der Fakultätenreihen. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften* **1906**, *36*, 151–218,482. 268
30. Bleistein, N.; Handelsman, R.A. *Asymptotic Expansions of Integrals*; Dover Books on Mathematics, Dover, 1975, 1986. 269
31. Aoyama, T.; Hayakawa, M.; Kinoshita, T.; Nio, M. Tenth-Order QED Contribution to the Electron  $g - 2$  and an Improved Value of the Fine Structure Constant. *Physical Review Letters* **2012**, *109*, 111807. <https://doi.org/10.1103/PhysRevLett.109.111807>. 270
32. Hanneke, D.; Fogwell, S.; Gabrielse, G. New Measurement of the Electron Magnetic Moment and the Fine Structure Constant. *Physical Review Letters* **2008**, *100*, 120801. <https://doi.org/10.1103/PhysRevLett.100.120801>. 271
33. Hanneke, D.; Fogwell Hoogerheide, S.; Gabrielse, G. Cavity control of a single-electron quantum cyclotron: Measuring the electron magnetic moment. *Phys. Rev. A* **2011**, *83*, 052122. <https://doi.org/10.1103/PhysRevA.83.052122>. 272
34. Aoyama, T.; Hayakawa, M.; Kinoshita, T.; Nio, M. Complete Tenth-Order QED Contribution to the Muon  $g - 2$ . *Physical Review Letters* **2012**, *109*, 111808. <https://doi.org/10.1103/PhysRevLett.109.111808>. 273
35. Keshavarzi, A.; Khaw, K.S.; Yoshioka, T. Muon  $g - 2$ : A review. *Nuclear Physics B* **2022**, *975*, 115675. <https://doi.org/10.1016/j.nuclphysb.2022.115675>. 274
36. Janka, G.; Ohayon, B.; Crivelli, P. Muonium Lamb shift: theory update and experimental prospects. *EPJ Web of Conferences* **2022**, *262*, 01001. <https://doi.org/10.1051/epjconf/202226201001>. 275
37. Bezginov, N.; Valdez, T.; Horbatsch, M.; Marsman, A.; Vutha, A.C.; Hessels, E.A. A measurement of the atomic hydrogen Lamb shift and the proton charge radius. *Science* **2019**, *365*, 1007–1012. <https://doi.org/10.1126/science.aau7807>. 276



38. Ohayon, B.; Janka, G.; Cortinovis, I.; Burkley, Z.; Borges, L.d.S.; Depero, E.; Golovizin, A.; Ni, X.; Salman, Z.; Suter, A.; et al. Precision Measurement of the Lamb Shift in Muonium. *Physical Review Letters* **2022**, *128*, 011802. <https://doi.org/10.1103/PhysRevLett.128.011802>. 313–315
39. Boyd, J.P. The Devil's Invention: Asymptotic, Superasymptotic and Hyperasymptotic Series. *Acta Applicandae Mathematica* **1999**, *56*, 1–98. <https://doi.org/10.1023/A:1006145903624>. 316–317
40. Rousseau, C. Divergent series: Past, present, future. *Mathematical Reports – Comptes rendus mathématiques* **2016**, *38*, 85–98, [arXiv:1312.5712]. 318–319
41. Flory, M.; Helling, R.C.; Sluka, C. How I Learned to Stop Worrying and Love QFT, 2012, [arXiv:1201.2714]. course presented by Robert C. Helling at the Ludwig-Maximilians-Universität München in the summer of 2011, notes by Mario Flory and Constantin Sluka. 320–322
42. Costin, O. *Asymptotics and Borel Summability*; Vol. 141, *Monographs and surveys in pure and applied mathematics*, Chapman & Hall/CRC, Taylor & Francis Group: Boca Raton, FL, 2009. 323–324
43. Zinn-Justin, J. Summation of divergent series: Order-dependent mapping. *Applied Numerical Mathematics* **2010**, *60*, 1454–1464, [arXiv:1001.0675]. <https://doi.org/10.1016/j.apnum.2010.04.002>. 325–327
44. Costin, O.; Dunne, G.V. Convergence from divergence. *Journal of Physics A: Mathematical and Theoretical* **2017**, *51*, 04LT01, [arXiv:1904.11593]. <https://doi.org/10.1088/1751-8121/aa9e30>. 328–329
45. Brüning, E.A.K. How to reconstruct an analytic function from its asymptotic expansion? *Complex Variables, Theory and Application: An International Journal* **1996**, *30*, 199–220. <https://doi.org/10.1080/17476939608814924>. 330–332
46. Euler, L. De seriebus divergentibus. *Novi Commentarii Academiae Scientiarum Petropolitanae* **1760**, *5*, 205–237, [arXiv:1202.1506]. In *Opera Omnia*: Series 1, Volume 14, pp. 585–617. Available on the Euler Archive as E247. 333–335
47. Weniger, E.J. Summation of divergent power series by means of factorial series. *Applied Numerical Mathematics* **2010**, *60*, 1429–1441, [arXiv:1005.0466]. <https://doi.org/10.1016/j.apnum.2010.04.003>. 336–338
48. Abramowitz, M.; Stegun, I.A., Eds. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*; Number 55 in National Bureau of Standards Applied Mathematics Series, U.S. Government Printing Office, Washington, D.C., 1964; pp. xiv+1046. 339–340
49. Sloane, N.J.A. A001620 Decimal expansion of Euler's constant (or the Euler-Mascheroni constant), gamma. (Formerly M3755 N1532). The On-Line Encyclopedia of Integer Sequences, 2019. accessed on July 17rd, 2019. 341–344
50. Pernice, S.A.; Oleaga, G. Divergence of perturbation theory: Steps towards a convergent series. *Physical Review D* **1998**, *57*, 1144–1158, [arXiv:hep-th/9609139]. <https://doi.org/10.1103/PhysRevD.57.1144>. 345–347
51. Pittnauer, F. *Vorlesungen über asymptotische Reihen*; Vol. 301, *Lecture Notes in Mathematics*, Springer Verlag: Berlin Heidelberg, 1972. <https://doi.org/10.1007/BFb0059524>. 348–349
52. Remmert, R. *Theory of Complex Functions*, 1 ed.; Vol. 122, *Graduate Texts in Mathematics*, Springer-Verlag: New York, NY, 1991. <https://doi.org/10.1007/978-1-4612-0939-3>. 350–351
53. Tao, T. *Compactness and contradiction*; American Mathematical Society: Providence, RI, 2013. 352
54. Watson, G.N. The transformation of an asymptotic series into a convergent series of inverse factorials [Memoir crowned by the Danish Royal Academy of Science]. *Rendiconti del Circolo Matematico di Palermo* **1912**, *34*, 41–88. <https://doi.org/10.1007/bf03015008>. 353–355
55. Doetsch, G. *Handbuch der Laplace-Transformation: Band II Anwendungen der Laplace-Transformation*; Springer Basel AG (Birkhäuser): Basel, 1972. <https://doi.org/10.1007/978-3-0348-5956-1>. 356–357
56. Costin, O.; Costin, R.D. A new type of factorial series expansions and applications, 2016, [arXiv:1608.01010]. <https://doi.org/10.48550/arXiv.1608.01010>. 358–359
57. Costin, O.; Dunne, G.V. Convergence from divergence. *Journal of Physics A: Mathematical and Theoretical* **2017**, *51*, 04LT01, [arXiv:1705.09687]. <https://doi.org/10.1088/1751-8121/aa9e30>. 360–361
58. Nielsen, N. *Die Gammafunktion*; AMS Chelsea Publishing: Bronx, New York, NY, 1965. reprint of “Handbuch der Theorie der Gammafunktion”, first published in 1906, and “Theorie des Integrallogarithmus und verwandter Transzendenten”, first published in 1906. 362–364
59. Knoop, K. *Theorie und Anwendung der unendlichen Reihen*; Springer: Berlin, Heidelberg, 1996. fünfte verbesserte Auflage, <https://doi.org/10.1007/978-3-642-61406-4>. 365–366
60. Dyson, F.J. The Radiation Theories of Tomonaga, Schwinger, and Feynman. *Physical Review* **1949**, *75*, 486–502. <https://doi.org/10.1103/PhysRev.75.486>. 367–368
61. Bell, J.S. Against ‘measurement’. *Physics World* **1990**, *3*, 33–41. <https://doi.org/10.1088/2058-7058/3/8/26>. 369–370

62. Berry, M.V.; Costin, O.; Costin, R.D.; Howls, C.J. Borel Plane Resurgence in Hyperasymptotics and Factorial Series, 2016. Talk at the Resurgence Meeting, presented by Ovidiu Costin on July 19, 2016, 10:00 to 11:00 at Amphitheatre Ea1, North Tower, IST, Lisboa; accessed July 7th, 2022. 371
63. Costin, O. Exponential asymptotics, transseries, and generalized Borel summation for analytic, nonlinear, rank-one systems of ordinary differential equations. *International Mathematics Research Notices* **1995**, 1995, 377, [arXiv:math/0608414]. <https://doi.org/10.1155/s1073792895000286>. 372
64. Edgar, G.A. Transseries for Beginners. *Real Analysis Exchange* **2009**, 35, 253–310, [arXiv:0801.4877]. <https://doi.org/10.48550/arXiv.0801.4877>. 373
65. Baker, G.A.; Graves-Morris, P. *Pad'e Approximants*, 2 ed.; Encyclopedia of Mathematics and its Applications, Cambridge University Press: Cambridge, UK, 1996. <https://doi.org/10.1017/CBO9780511530074>. 374
66. Aigner, M.; Ziegler, G.M. *Proofs from THE BOOK*, four ed.; Springer: Heidelberg, 1998–2010. <https://doi.org/10.1007/978-3-642-00856-6>. 375
67. Kreisel, G. Kurt Gödel. 28 April 1906–14 January 1978. *Biographical memoirs of Fellows of the Royal Society* **1980**, 26, 148–224. corrections *Ibid.* **27**, 697; *ibid.* **28**, 718, <https://doi.org/10.1098/rsbm.1980.0005>. 376