Four-Dimensional Context Hypergraphs with Separating Two-Valued States and a Chromatic Gap

Karl Svozil [1]

[1] *Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria* *

(Dated: October 22, 2026)

We construct context hypergraphs in four-dimensional real Hilbert space that admit a unital, separating family of two-valued states while their orthogonality graphs require five colors and have clique number four. The main example is a linear hypergraph with 65 rays and 29 contexts, obtained by completing a known 17-ray Peres subgraph. We give an injective integer realization with exactly the prescribed orthogonalities, prove the chromatic gap by a parity argument, and determine the fractional chromatic number and the full two-valued state space. A family of 48 states already provides a partition representation of the vertices and listed contexts. An elementary apex example and a hyperplane obstruction explain the role of the four-dimensional completion.

I. INTRODUCTION AND OVERVIEW

Let $H = (V, E)$, with $E \neq \emptyset$, be a 4-uniform context hypergraph: each context $e \in E$ is a 4-element subset of V, and its orthogonality graph $\Gamma(H)$ has vertex set V and edge set

$$E(\Gamma(H)) = \big\{\{u, v\} \subseteq V : u \neq v,\ \exists\, e \in E,\ \{u, v\} \subseteq e\big\}. \tag{1}$$

Writing K_t for the complete graph on t vertices, every context $e \in E$ forms a 4-clique (K_4) in $\Gamma(H)$, so the clique number satisfies $\omega(\Gamma(H)) \geq 4$, and equality $\omega(\Gamma(H)) = 4$ holds whenever $\Gamma(H)$ is K_5-free. A *strong k-coloring* of H is a proper k-coloring $c : V \to \{0, 1, \ldots, k-1\}$ of $\Gamma(H)$, so that every context $e \in E$ receives four distinct colors; the minimum such k is the chromatic number $\chi(H) = \chi(\Gamma(H)) \geq \omega(\Gamma(H))$.

In four-dimensional Hilbert space, a 4-uniform hypergraph $H = (V, E)$ is *edge-faithfully coordinatizable in* $\mathbb{R}^4$ if there is an injective ray assignment $i \mapsto [v_i] \in \mathbb{P}(\mathbb{R}^4)$, the projective space of rays in real Euclidean four-space ($v_i \in \mathbb{R}^4 \setminus \{0\}$) such that

$$v_i \cdot v_j = 0 \iff \{i, j\} \in E(\Gamma(H)) \qquad (i \neq j \in V). \tag{2}$$

In particular, every context $e \in E$ is mapped to an orthogonal basis of $\mathbb{R}^4$, no two vertices share a ray, and no extra orthogonalities occur outside $E(\Gamma(H))$.

Independently of quantum coordinatization, we call H *set-representable on its vertices and listed contexts* if its collection $\mathcal{S}(H)$ of *two-valued states*—maps $s : V \to \{0, 1\}$ satisfying $\sum_{i \in e} s(i) = 1$ for every $e \in E$—is both:

1. *unital*: for every $i \in V$, there exists $s \in \mathcal{S}(H)$ with $s(i) = 1$; and

2. *separating*: for every pair of distinct vertices $i \neq j \in V$, there exists $s \in \mathcal{S}(H)$ with $s(i) \neq s(j)$.

* Corresponding author: karl.svozil@tuwien.ac.at; https://tph.tuwien.ac.at/~svozil

For any unital and separating family $\Omega \subseteq \mathcal{S}(H)$, set

$$A_i = \{s \in \Omega : s(i) = 1\} \qquad (i \in V). \tag{3}$$

The sets A_i are nonempty and pairwise distinct, and $\{A_i : i \in e\}$ is a partition of Ω for each $e \in E$. This is the precise partition representation meant here [1, 2]; it does not require injectivity on all compound Hilbert-space events. The support of a state is $\mathrm{supp}(s) = \{i \in V : s(i) = 1\}$.

The *independence number* $\alpha(G)$ is the largest size of a vertex set containing no adjacent pair. The *fractional chromatic number* $\chi_f(G)$ is the minimum total nonnegative weight assigned to independent sets so that the sets containing each vertex have total weight at least one.

Because the color classes of any strong 4-coloring of a 4-uniform hypergraph define four two-valued states that partition V, chromatic completeness ($\chi(H) = 4$) implies unitality, whereas Kochen–Specker (KS) configurations such as the 24-ray Peres set in $\mathbb{R}^4$ [3] satisfy $\chi > 4$ because $\mathcal{S}(H) = \emptyset$. The subscript YO abbreviates Yu–Oh. Conversely, the existence of unital and separating states need not supply a strong 4-coloring: such a coloring requires four states whose supports are pairwise disjoint and cover V.

The purpose of this note is to exhibit explicit 4-uniform context hypergraphs H with $\omega(\Gamma(H)) = 4$ that are simultaneously set-representable (unital and separating two-valued states), edge-faithfully coordinatizable in $\mathbb{R}^4$, and chromatically incomplete ($\chi(H) = 5 > 4 = \omega(\Gamma(H))$). Table I summarizes the configurations analyzed below.

II. NON-LINEAR WARM-UP $H_{\mathrm{YO}}^{(4)}$ AND THE HYPERPLANE OBSTRUCTION

Before constructing a linear 4-uniform example, we record an elementary 26-ray, 16-context non-linear example obtained by lifting the 3-uniform completed Yu–Oh hypergraph $H_{\mathrm{YO}}^{+} = (V_{\mathrm{YO}}^{+}, E_{\mathrm{YO}}^{+})$ [4–6] from $\mathbb{R}^3$ to $\mathbb{R}^4$. The obstruction below rules out retaining linearity when two intersecting complete orthogonal triads in one hyperplane are lifted to distinct four-element contexts.

TABLE I. Comparison of 4-uniform context hypergraphs in $\mathbb{R}^4$ with clique number $\omega(\Gamma(H)) = 4$ and chromatic number $\chi(H) = 5 > 4$, showing the number of pairwise non-collinear vectors $|V|$, the number of 4-element contexts $|E|$, and the number of orthogonality edges $|E(\Gamma(H))|$.

Hypergraph	$\lvert V\rvert$	$\lvert E\rvert$	$\lvert E(\Gamma(H))\rvert$	Linear	$\omega(\Gamma(H))$	$\chi(H)$	Edge-faithful	$\lvert\mathcal{S}(H)\rvert$	Set-representable
Peres 24-ray set [3]	24	24	108	No ($\lvert e\cap e'\rvert \le 2$)	4	5	Yes ($\mathbb{R}^4$)	0	No (KS)
Apex suspension $H^{(4)}_{\mathrm{YO}}$ (Prop. 1)	26	16	73	No ($\lvert e\cap e'\rvert \le 2$)	4	5	Yes ($\mathbb{R}^4$)	25	Yes
Canonical linear H_{65} (Thm. 4)	65	29	174	Yes ($\lvert e\cap e'\rvert \le 1$)	4	5	Yes ($\mathbb{R}^4$)	393,216	Yes (48 canonical)

Proposition 1 (Non-linear apex suspension $H^{(4)}_{\mathrm{YO}}$)**.** *Let $H^{+}_{\mathrm{YO}} = (V^{+}_{\mathrm{YO}}, E^{+}_{\mathrm{YO}})$ be the 25-vertex, 16-context completed Yu–Oh hypergraph in $\mathbb{R}^3$, with vectors $r_1, \dots, r_{25} \in \mathbb{R}^3$ and contexts $E^{+}_{\mathrm{YO}} = \{e_1, \dots, e_{16}\}$. Define $H^{(4)}_{\mathrm{YO}} = (V_{26}, E^{(4)}_{16})$ on $V_{26} = \{1, \dots, 26\}$ by appending an apex vertex 26 to every context:*

$$E^{(4)}_{16} = \big\{e \cup \{26\} : e \in E^{+}_{\mathrm{YO}}\big\}, \tag{4}$$

and assign the vectors $v_i = (r_i, 0)^T \in \mathbb{R}^4$ for $1 \le i \le 25$ and $v_{26} = (0, 0, 0, 1)^T \in \mathbb{R}^4$ (enumerated explicitly in Appendix A 2 and depicted as a 4-uniform hypergraph diagram in Fig. 1). Then:

1. *$H^{(4)}_{\mathrm{YO}}$ is 4-uniform with clique number $\omega(\Gamma(H^{(4)}_{\mathrm{YO}})) = 4$ and chromatic number $\chi(H^{(4)}_{\mathrm{YO}}) = \chi(H^{+}_{\mathrm{YO}}) + 1 = 5 > 4$.*
2. *The vectors $v_1, \dots, v_{26} \in \mathbb{R}^4$ form an injective, edge-faithful orthogonal representation of $H^{(4)}_{\mathrm{YO}}$ in $\mathbb{R}^4$.*
3. *$H^{(4)}_{\mathrm{YO}}$ admits exactly 25 two-valued states (enumerated in Appendix B 1), which form a unital and separating collection on V_{26}.*

Proof. (1)–(2) Since $v_{26} = (0, 0, 0, 1)^T$ is orthogonal to $\mathbb{R}^3 \times \{0\}$ and $r_1, \dots, r_{25}$ is an injective, edge-faithful representation of H^{+}_{YO} in $\mathbb{R}^3$, the vectors $v_1, \dots, v_{26}$ are pairwise non-proportional and satisfy $v_i \cdot v_j = 0$ if and only if $\{i, j\} \in E(\Gamma(H^{(4)}_{\mathrm{YO}}))$. Thus $\Gamma(H^{(4)}_{\mathrm{YO}})$ is the graph join $\Gamma(H^{+}_{\mathrm{YO}}) * K_1$, which immediately gives $\omega(\Gamma(H^{(4)}_{\mathrm{YO}})) = \omega(\Gamma(H^{+}_{\mathrm{YO}})) + 1 = 3 + 1 = 4$ and $\chi(H^{(4)}_{\mathrm{YO}}) = \chi(H^{+}_{\mathrm{YO}}) + 1 = 4 + 1 = 5$.

(3) In any two-valued state $s : V_{26} \to \{0, 1\}$ on $H^{(4)}_{\mathrm{YO}}$, either $s(26) = 1$, which forces $s(1) = \cdots = s(25) = 0$ (giving one apex state s_{apex}), or $s(26) = 0$, in which case $\sum_{i \in e} s(i) = 1$ for all $e \in E^{+}_{\mathrm{YO}}$ so that $s|_{V^{+}_{\mathrm{YO}}}$ is one of the 24 two-valued states of H^{+}_{YO}. Since the 24 states of H^{+}_{YO} are unital and separating on $\{1, \dots, 25\}$ and all vanish at 26, while $s_{\mathrm{apex}}(26) = 1$ and $s_{\mathrm{apex}}(i) = 0$ for $i \le 25$, the 25 states of $H^{(4)}_{\mathrm{YO}}$ are unital and separating on all 26 vertices. □

Although $H^{(4)}_{\mathrm{YO}}$ satisfies all three requirements of set-representability, edge-faithfulness in $\mathbb{R}^4$, and $\chi = 5 > 4 = \omega$, it is *non-linear*: the apex vertex 26 lies in all 16 contexts, so $|(e \cup \{26\}) \cap (e' \cup \{26\})| = 2$ whenever $e, e' \in E^{+}_{\mathrm{YO}}$ share a vertex in $\mathbb{R}^3$ (giving Berge girth 2). A Berge cycle of length r consists of r distinct contexts and r distinct vertices, with successive contexts meeting at the corresponding vertices cyclically; the Berge girth is the shortest such length. The following proposition shows that this non-linearity is unavoidable whenever two contexts have intersecting three-dimensional restrictions inside a common hyperplane.

Proposition 2 (Hyperplane obstruction to linearity in dimension four)**.** *Let $H = (V, E)$ be a 4-uniform context hypergraph with an injective orthogonal representation $i \mapsto [v_i] \in \mathbb{P}(\mathbb{F}^4)$ ($\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, with $\mathbb{C}$ the complex numbers). Orthogonality over $\mathbb{C}$ is defined by the Hermitian product $\langle x, y\rangle = \sum_{j=1}^{4} \overline{x_j} y_j$. For a subspace W, write $e_W = \{i \in e : v_i \in W\}$. If two distinct contexts $e, e' \in E$ each contain at least three vertices whose rays lie in a common three-dimensional hyperplane $W = z^{\perp} \subset \mathbb{F}^4$ ($z \in \mathbb{F}^4 \setminus \{0\}$), then both e and e' must contain the unique ray $[z]$ orthogonal to W. Consequently, if $|e_W \cap e'_W| \ge 1$, then $|e \cap e'| \ge 2$, violating linearity.*

Proof. Let $e = \{u_1, u_2, u_3, u_4\}$ with $v_{u_1}, v_{u_2}, v_{u_3} \in W$. Because $v_{u_1}, v_{u_2}, v_{u_3}$ are mutually orthogonal nonzero vectors in the 3-dimensional subspace W, they form an orthogonal basis of W. Since $v_{u_4} \perp v_{u_1}, v_{u_2}, v_{u_3}$, we have $v_{u_4} \in W^{\perp} = \mathrm{span}(z)$, so $[v_{u_4}] = [z]$. The same argument applied to e' shows that its fourth ray must also equal $[z]$; by injectivity, the same vertex $z_0 \in V$ with $[v_{z_0}] = [z]$ belongs to both e and e'. If e and e' also share any vertex inside W, then $|e \cap e'| \ge 2$. □

III. CONSTRUCTION OF THE CANONICAL LINEAR HYPERGRAPH H_{65} IN $\mathbb{R}^4$

By Proposition 2, two intersecting complete orthogonal triads in one hyperplane cannot be extended to distinct contexts of a linear 4-uniform hypergraph with an injective orthogonal representation in $\mathbb{R}^4$. The assertion concerns triads, not arbitrary smaller restrictions of contexts. Instead, we start from a four-dimensional core of 17 rays in $\{-1, 0, 1\}^4$ and complete its 24 uncovered orthogonal pairs into 4-contexts via the following general lemma. All completion vectors are integer vectors, with $\mathbb{Z}$ denoting the integers.

A. The two-plane completion lemma in $\mathbb{R}^4$

In $\mathbb{R}^3$, two orthogonal rays $\{v_u, v_v\}$ uniquely determine their third orthogonal basis ray $[v_u \times v_v]$, which may inadvertently be orthogonal or parallel to other rays of the configuration. In $\mathbb{R}^4$, by contrast, the orthogonal complement $\{v_u, v_v\}^\perp$ of an orthogonal pair is a *two-dimensional plane*, providing an infinite one-parameter family of orthogonal completion pairs $\{w_1, w_2\}$.

Lemma 3 (Simultaneous plane-completion faithfulness and state extension in $\mathbb{R}^4$)**.** *Let V_{core} be a finite vertex set carrying pairwise non-collinear vectors $v_i \in \mathbb{Q}^4 \setminus \{0\}$, where $\mathbb{Q}$ denotes the rational numbers, let E_{base} be a collection of 4-element orthogonal bases of $\mathbb{R}^4$ drawn from V_{core} with $|C \cap C'| \le 1$ for all $C \neq C' \in E_{\text{base}}$, and let*

$$E_{\text{bridge}} = \big\{\{u,v\} \subset V_{\text{core}} : v_u \cdot v_v = 0\big\} \setminus \bigcup_{C \in E_{\text{base}}} \binom{C}{2} \quad (5)$$

be the set of orthogonal pairs in V_{core} not already covered by E_{base}, with $E_{\text{bridge}} = \{e_1, \dots, e_M\}$ ($e_m = \{u_m, v_m\}$). For each $m \in \{1, \dots, M\}$, set $L_m = \operatorname{span}(v_{u_m}, v_{v_m})$ and $P_m = L_m^\perp$.

1. Span conditions for simultaneous edge-faithful completion: *The pairs $e_1, \dots, e_M \in E_{\text{bridge}}$ can be simultaneously completed to 4-element orthogonal bases $E_m = \{u_m, v_m, w_{m,1}, w_{m,2}\}$ by fresh integer vectors $w_{m,1}, w_{m,2} \in P_m \cap \mathbb{Z}^4$ having* no *extra orthogonalities or collinearities with any other vectors in the completed hypergraph $H = (V_{\text{core}} \cup \bigcup_{m=1}^M \{w_{m,1}, w_{m,2}\},\ E_{\text{base}} \cup \{E_1, \dots, E_M\})$ if and only if both*

$$v_z \notin L_m \quad \text{for every } z \in V_{\text{core}} \setminus \{u_m, v_m\} \quad (1 \le m \le M) \quad (6)$$

and

$$P_\ell \neq L_m \ (\text{i.e., } L_\ell \not\perp L_m) \quad (1 \le \ell \neq m \le M). \quad (7)$$

2. Linearity, clique number, and chromatic preservation: *The completed 4-uniform hypergraph H is linear ($|e \cap e'| \le 1$ for all $e \neq e' \in E(H)$). If $E(H) \neq \emptyset$ and the core orthogonality graph G_{core} is K_5-free, then $\omega(\Gamma(H)) = 4$, and for every $k \ge 4$, H is strongly k-colorable if and only if G_{core} is properly k-colorable.*

3. Canonical state extension: *Let $S_1, \dots, S_N \subset V_{\text{core}}$ be all independent sets in G_{core} satisfying $|S_\alpha \cap C| = 1$ for every $C \in E_{\text{base}}$. For each $\alpha \in \{1, \dots, N\}$ and $\varepsilon \in \{0,1\}$, define $s_{\alpha,\varepsilon} : V(H) \to \{0,1\}$ by $s_{\alpha,\varepsilon}(i) = \mathbf{1}_{i \in S_\alpha}$ for $i \in V_{\text{core}}$ and, for each $m \in \{1, \dots, M\}$,*

$$\big(s_{\alpha,\varepsilon}(w_{m,1}),\ s_{\alpha,\varepsilon}(w_{m,2})\big) = \begin{cases} (0,0), & |S_\alpha \cap \{u_m, v_m\}| = 1, \\ (1-\varepsilon,\ \varepsilon), & S_\alpha \cap \{u_m, v_m\} = \emptyset. \end{cases} \quad (8)$$

Then every $s_{\alpha,\varepsilon}$ is a valid two-valued state on H; here $\mathbf{1}_{i \in S_\alpha}$ is the indicator of membership in S_α. The two indices $\varepsilon = 0, 1$ can give the same state when S_α hits every bridge edge. If $S_1, \dots, S_N$ are unital and separating on V_{core}, and the M edge zero-profiles

$$Z(e_m) = \big\{\alpha \in \{1, \dots, N\} : u_m \notin S_\alpha \text{ and } v_m \notin S_\alpha\big\} \quad (9)$$

are nonempty and pairwise distinct for $m = 1, \dots, M$, then the indexed canonical family $\{s_{\alpha,\varepsilon}\}$, containing at most $2N$ distinct states, is unital and separating on all $|V_{\text{core}}| + 2M$ vertices of H.

Proof. (1) For necessity, if $v_z \in L_m$ for some $z \in V_{\text{core}} \setminus \{u_m, v_m\}$, then $P_m = L_m^\perp \subset v_z^\perp$, so every completion vector in P_m is orthogonal to v_z; similarly, if $P_\ell = L_m$ for $\ell \neq m$, then $P_\ell \perp P_m$, forcing $w_{\ell,a} \perp w_{m,b}$ for all $a, b \in \{1,2\}$.

For sufficiency, assume Eqs. (6)–(7). Because $v_{u_m}, v_{v_m} \in \mathbb{Q}^4$, each P_m is a rational 2-plane and therefore admits an integer orthogonal basis $\{\eta_A, \eta_B\} \subset P_m \cap \mathbb{Z}^4$ (obtained by rational Gram–Schmidt and clearing denominators) with $n_A = \|\eta_A\|^2$ and $n_B = \|\eta_B\|^2$. We choose $\{w_{m,1}, w_{m,2}\} \subset P_m \cap \mathbb{Z}^4$ inductively for $m = 1, \dots, M$ so as to maintain the invariant that, for every $r \ge m$, no previously placed vector $z \notin \{v_{u_r}, v_{v_r}\}$ lies in L_r (which holds at $m = 1$ by Eq. (6)). At step m, the invariant for $r = m$ ensures that every previously placed vector $z \notin \{v_{u_m}, v_{v_m}\}$ satisfies $z \notin L_m = P_m^\perp$, so $z^\perp \cap P_m$ and $\mathbb{R}z \cap P_m$ are at most 1-dimensional lines in P_m. Moreover, for every future $r > m$, Eq. (7) gives $P_m \neq L_r$, so $P_m \cap L_r$ is also at most a 1-dimensional line in P_m. The integer parameterization

$$\begin{aligned} w_{m,1}(a,b) &= a\,\eta_A + b\,\eta_B, \\ w_{m,2}(a,b) &= -n_B b\,\eta_A + n_A a\,\eta_B \end{aligned} \quad (10)$$

produces infinitely many projectively distinct orthogonal pairs in $P_m \cap \mathbb{Z}^4$ as (a,b) ranges over $\mathbb{Z}^2 \setminus \{(0,0)\}$, and can avoid every forbidden line for both vectors: the inverse image of each such line under either invertible linear map in Eq. (10) is a line in the (a,b)-parameter plane, and finitely many lines cannot cover all integer slopes. This avoids the finite union of forbidden lines in P_m, preventing any extra orthogonalities or collinearities with previously placed vectors while preserving $w_{m,1}, w_{m,2} \notin L_r$ for all $r > m$.

(2) Each E_m shares at most one vertex with any $C \in E_{\text{base}}$ (since $\binom{C}{2} \cap E_{\text{bridge}} = \emptyset$) and at most one vertex with any $E_{m'}$ ($m' \neq m$), while $w_{m,1}, w_{m,2}$ have degree 1 in H. In any proper k-coloring ($k \ge 4$) of G_{core}, $c(u_m) \neq c(v_m)$, leaving $k - 2 \ge 2$ colors in $\{0, \dots, k-1\} \setminus \{c(u_m), c(v_m)\}$ to color $w_{m,1}$ and $w_{m,2}$.

(3) Since S_α is independent in G_{core}, $|S_\alpha \cap \{u_m, v_m\}| \in \{0,1\}$, so Eq. (8) gives $s_{\alpha,\varepsilon}(u_m) + s_{\alpha,\varepsilon}(v_m) + s_{\alpha,\varepsilon}(w_{m,1}) + s_{\alpha,\varepsilon}(w_{m,2}) = 1$ on every E_m. Across the $2N$ indexed states $(s_{1,0}, s_{1,1}, \dots, s_{N,0}, s_{N,1})$, the profile of $w_{m,1}$ has $(1,0)$ in block α for $\alpha \in Z(e_m)$ and $(0,0)$ otherwise; the

profile of $w_{m,2}$ has $(0,1)$ in block α for $\alpha \in Z(e_m)$ and $(0,0)$ otherwise; and the profile of $i \in V_{\text{core}}$ has $(1,1)$ for $i \in S_\alpha$ and $(0,0)$ otherwise. Since $Z(e_m) \neq \emptyset$ and $Z(e_m) \neq Z(e_{m'})$ for $m \neq m'$, all $|V_{\text{core}}|+2M$ state profiles are nonzero and pairwise distinct. □

B. Explicit definition of H_{65} and provenance of the 17-ray core

We define the 17-vertex core $V_{\text{core}} = \{1, \ldots, 17\}$ in $\mathbb{R}^4$ using four structural groups of vectors in $\{-1, 0, 1\}^4$:

1. *Standard coordinate basis* $C_1 = \{1, 2, 3, 4\}$*:*

$$v_1 = (1,0,0,0), \qquad v_2 = (0,1,0,0),$$
$$v_3 = (0,0,1,0), \qquad v_4 = (0,0,0,1).$$

2. *Even-parity Hadamard basis* $C_2 = \{5, 6, 7, 8\}$ (the ray representatives with first coordinate $+1$ and an even number of minus signs):

$$v_5 = (1,1,1,1), \qquad v_6 = (1,1,-1,-1),$$
$$v_7 = (1,-1,1,-1), \qquad v_8 = (1,-1,-1,1).$$

3. *Odd-parity Hadamard hub triad* $\{9, 10, 11\}$ (three mutually orthogonal ± 1 vectors with an odd number of minus signs):

$$v_9 = (1,1,-1,1), \qquad v_{10} = (1,-1,1,1),$$
$$v_{11} = (1,-1,-1,-1).$$

4. *Three binary bridge pairs* $B_1 = \{12, 13\}$, $B_2 = \{14, 15\}$, $B_3 = \{16, 17\}$ completing the three pairs of $\{9, 10, 11\}$ into 4-element orthogonal bases $C_3 = \{9, 10, 12, 13\}$, $C_4 = \{9, 11, 14, 15\}$, and $C_5 = \{10, 11, 16, 17\}$:

$$v_{12} = (1,0,0,-1), \qquad v_{13} = (0,1,1,0),$$
$$v_{14} = (1,0,1,0), \qquad v_{15} = (0,1,0,-1),$$
$$v_{16} = (1,1,0,0), \qquad v_{17} = (0,0,1,-1).$$

As a 17-ray orthogonality graph in $\mathbb{R}^4$, G_{core} is orthogonally equivalent to the 17-ray subgraph of the 24-ray Peres configuration introduced by Haynes *et al.* [7, Sec. 1.2, pp. 8–9] as an example with orthogonal dimension 4 and chromatic number 5. Explicitly, the orthogonal matrix

$$T = \frac{1}{2}\begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & -1 & -1 & -1 \end{pmatrix}, \qquad T^T T = I_4, \tag{11}$$

where I_4 is the 4×4 identity matrix. The matrix T maps the hub triad $\{v_9, v_{10}, v_{11}\}$ to the coordinate rays $\{e_2, e_3, e_4\}$, the bases $C_1 \cup C_2 = \{v_1, \ldots, v_8\}$ to the eight rays $(1, \pm 1, \pm 1, \pm 1)^T$, and the six bridge rays $\{v_{12}, \ldots, v_{17}\}$ to the six binary rays $e_1 \pm e_j$ $(j = 2, 3, 4)$ of Ref. [7]. Choosing the coordinate-Hadamard frame above makes the bipartite parity structure between C_1, C_2, and $B_1 \cup B_2 \cup B_3$ transparent, allowing us to complete G_{core} into the linear 4-uniform context hypergraph H_{65} and analyze its two-valued state space by hand.

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
$x_{i,1}$	1	0	0	0	1	1	1	1	1	1	1	1	0	1	0	1	0
$x_{i,2}$	0	1	0	0	1	1	−1	−1	1	−1	−1	0	1	0	1	1	0
$x_{i,3}$	0	0	1	0	1	−1	1	−1	−1	1	−1	0	1	1	0	0	1
$x_{i,4}$	0	0	0	1	1	−1	−1	1	1	1	−1	−1	0	0	−1	0	−1
$c(i)$	0	1	2	3	0	1	2	4	0	1	3	3	4	2	1	0	2

i	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41
$x_{i,1}$	1	3	1	3	0	0	2	3	1	3	1	3	0	0	2	3	1	3	1	3	0	0	2	3
$x_{i,2}$	0	0	3	−2	1	3	−3	1	0	0	−3	2	1	3	3	−1	−1	−3	−1	−3	2	3	0	0
$x_{i,3}$	−3	2	0	0	−1	−3	3	−1	−1	−3	−1	−3	3	−2	0	0	0	0	3	−2	3	−1	−3	1
$x_{i,4}$	1	3	1	3	3	−2	0	0	−3	2	0	0	1	3	3	−1	3	−2	0	0	3	−1	−3	1
$c(i)$	0	2	0	1	1	2	0	1	0	3	0	1	2	3	0	3	1	3	1	2	1	3	0	3

i	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65
$x_{i,1}$	1	2	1	2	1	5	3	3	1	2	1	2	3	3	1	5	1	2	1	2	1	5	3	3
$x_{i,2}$	−5	−1	−3	3	−2	−1	−1	2	3	−3	5	1	−2	1	2	1	−1	−2	−1	−2	3	−3	−5	1
$x_{i,3}$	3	−3	5	1	2	1	1	−2	−1	−2	−1	−2	1	−5	3	−3	−5	−1	−3	3	−2	−1	−1	2
$x_{i,4}$	1	2	1	2	−3	3	5	−1	5	1	3	−3	−2	1	2	1	−3	3	−5	−1	−2	−1	−1	2
$c(i)$	1	2	0	1	0	2	0	1	0	3	0	1	2	3	0	3	1	3	1	2	1	3	0	3

(12)

Here the weight of a vector is its number of nonzero coordinates, and $N(i) = \{j \in V_{\text{core}} \setminus \{i\} : v_i \cdot v_j = 0\}$ denotes its neighborhood in G_{core}. Because every vector in C_1 has weight 1 and every vector in $C_2 \cup \{9, 10, 11\}$ has weight 4 with odd/even parity, there are no orthogonalities among $\{1, \ldots, 11\}$ outside C_1, C_2, and the triad $\{9, 10, 11\}$. Each of the six binary bridge vectors $b \in \{12, \ldots, 17\}$, having weight 2, is orthogonal to exactly two coordinate rays in C_1 and two even-parity Hadamard rays in C_2, creating $6 \times 2 + 6 \times 2 = 24$ bridge edges $E_{\text{bridge}} = \{e_1, \ldots, e_{24}\}$. Completing each $e_m = \{u_m, v_m\}$ via Lemma 3 with fresh integer rays $\{2m + 16,\ 2m + 17\} \subset \{18, \ldots, 65\}$ yields the 29 contexts of $H_{65} = (V_{65}, E_{29})$:

$$
\begin{aligned}
E_{29} = \big\{ & \{1,2,3,4\},\ \{5,6,7,8\},\ \{9,10,12,13\},\\
& \{9,11,14,15\},\ \{10,11,16,17\},\ \{2,12,18,19\},\\
& \{3,12,20,21\},\ \{1,13,22,23\},\ \{4,13,24,25\},\\
& \{2,14,26,27\},\ \{4,14,28,29\},\ \{1,15,30,31\},\\
& \{3,15,32,33\},\ \{3,16,34,35\},\ \{4,16,36,37\},\\
& \{1,17,38,39\},\ \{2,17,40,41\},\ \{5,12,42,43\},\\
& \{8,12,44,45\},\ \{6,13,46,47\},\ \{7,13,48,49\},\\
& \{6,14,50,51\},\ \{8,14,52,53\},\ \{5,15,54,55\},\\
& \{7,15,56,57\},\ \{7,16,58,59\},\ \{8,16,60,61\},\\
& \{5,17,62,63\},\ \{6,17,64,65\}\big\}.
\end{aligned}
\tag{13}
$$

The complete coordinate table $v_i = (x_{i,1}, x_{i,2}, x_{i,3}, x_{i,4})^T \in \mathbb{R}^4$ and a strong 5-coloring $c(i) \in \{0, 1, 2, 3, 4\}$ for all 65 vertices are given in Eq. (12) (and enumerated vertex-by-vertex and context-by-context in Appendix A 1). Figure 1 illustrates the 26-vertex, 16-context non-linear warm-up apex suspension $H_{\text{YO}}^{(4)}$ of Proposition 1 (with the common apex vertex 26 at the center surrounded by 16 concentric colored disks), while Fig. 2 presents the full 65-vertex, 29-context four-uniform hypergraph diagram of the canonical linear hypergraph H_{65}.

Both drawings represent 4-uniform context hypergraphs. They differ from Greechie diagrams in the strict sense of Ref. [1, Definition 2.4]: the warm-up has contexts sharing two vertices, whereas H_{65} contains the three-context Berge cycle C_3, C_4, C_5, with successive intersection vertices $9, 11, 10$. Strict Greechie diagrams permit neither feature. The smooth-curve drawing convention is used only to display the listed incidences.

IV. MAIN THEOREM AND PROOFS

Theorem 4 (Properties of the canonical linear 4-uniform hypergraph H_{65})**.** *The 65-vertex, 29-context 4-uniform hypergraph $H_{65} = (V_{65}, E_{29})$ defined by Eqs. (12)–(13) satisfies:*

1. Linearity and clique number: *H_{65} is linear ($|e \cap e'| \le 1$ for all distinct $e, e' \in E_{29}$), and the 4-cliques of $\Gamma(H_{65})$ are precisely the 29 contexts of E_{29}; in particular, $\omega(\Gamma(H_{65})) = 4$.*

2. Edge-faithful realization in $\mathbb{R}^4$: *The 65 integer vectors $v_1, \ldots, v_{65} \in \mathbb{R}^4$ in Eq. (12) are pairwise non-collinear and satisfy $v_i \cdot v_j = 0$ if and only if $\{i, j\} \in E(\Gamma(H_{65}))$ (174 orthogonal pairs).*

3. Chromatic–clique gap $\chi(H_{65}) = 5 > 4 = \omega(\Gamma(H_{65}))$: *The 17-vertex core G_{core} has independence number $\alpha(G_{\text{core}}) = 4$, fractional chromatic number $\chi_f(\Gamma(H_{65})) = 17/4 = 4.25 > 4$, and $\chi(H_{65}) = 5$.*

4. Set-representability: *The core $(V_{\text{core}}, \{C_1, \ldots, C_5\})$ admits exactly 24 two-valued state supports $S_1, \ldots, S_{24}$ compatible with G_{core} (listed in Table II), and the resulting 48 canonical two-valued states $s_{\alpha,\varepsilon}$ on H_{65} via Eq. (8) form a unital and separating collection on all 65 vertices (within a full state space of $24 \times 2^{14} = 393{,}216$ states, enumerated in Appendix B 2).*

Proof. (1)–(2) By inspection of Eq. (13), the five base contexts $C_1, \ldots, C_5$ satisfy $|C_a \cap C_b| \le 1$ (C_1, C_2 are disjoint from everything else, while $C_3 \cap C_4 = \{9\}$, $C_3 \cap C_5 = \{10\}$, $C_4 \cap C_5 = \{11\}$), and the 24 pairs $e_m = \{u_m, v_m\}$ in $E_1, \ldots, E_{24}$ are precisely the 24 orthogonal pairs between $C_1 \cup C_2$ and $B_1 \cup B_2 \cup B_3$.

To determine the triangles and 4-cliques of G_{core} and $\Gamma(H_{65})$, let $e_m = \{u_m, v_m\}$ be a bridge edge with $u_m \in C_a$ ($a \in \{1, 2\}$) and $v_m \in B_k$. Because u_m is non-orthogonal to the other basis C_{3-a} and to $\{9, 10, 11\}$, while v_m has no neighbors in $(B_1 \cup B_2 \cup B_3) \setminus B_k$ and its partner $v'_m \in B_k$ is non-orthogonal to u_m (Table III), the *unique* common third neighbor of u_m and v_m in G_{core} is the second vertex of $N(v_m) \cap C_a$. Consequently, G_{core} has exactly 13 triangles not contained in $C_1, \ldots, C_5$: the 12 bridge triangles

$$
\begin{aligned}
&\{1,2,17\},\ \{1,3,15\},\ \{1,4,13\},\\
&\{2,3,12\},\ \{2,4,14\},\ \{3,4,16\},\\
&\{5,6,17\},\ \{5,7,15\},\ \{5,8,12\},\\
&\{6,7,13\},\ \{6,8,14\},\ \{7,8,16\},
\end{aligned}
$$

together with the hub triad $\{9, 10, 11\}$. Since each bridge edge has only one common third neighbor in V_{core}, *no 4-clique (K_4) in G_{core} contains a bridge edge.* This has two immediate consequences:

- First, for any two distinct bridge edges e_ℓ, e_m, the four rays $\{u_\ell, v_\ell, u_m, v_m\}$ cannot form a K_4 in G_{core}, so $L_\ell \not\perp L_m$ ($P_\ell \neq L_m$), verifying Eq. (7). Likewise, any nonzero vector in $L_m = \text{span}(v_{u_m}, v_{v_m})$ outside $\mathbb{R}v_{u_m} \cup \mathbb{R}v_{v_m}$ either has support size 3 (when $u_m \in C_1$, whereas no vector in V_{core} has support size 3) or has components outside $\{-1, 0, 1\}$ up to scaling (when $u_m \in C_2$), verifying Eq. (6). The reproducible completion rule and exact integer verification in Appendix C confirm that




FIG. 1. Four-uniform hypergraph diagram of the 26-vertex, 16-context non-linear warm-up apex suspension $H^{(4)}_{\mathrm{YO}} = (V_{26}, E^{(4)}_{16})$ in $\mathbb{R}^4$ of Proposition 1 (26 pairwise non-collinear integer vectors $v_1, \ldots, v_{26} \in \mathbb{Z}^4$ enumerated in Eq. (A2), 16 four-element contexts $e^{(4)}_1, \ldots, e^{(4)}_{16}$ listed in Table VI, and 73 orthogonality edges). Every context $e^{(4)}_r = e_r \cup \{26\}$ $(r = 1, \ldots, 16)$ is drawn as a circular arc of a distinct color passing through its four vertices. Each vertex number is surrounded by concentric filled disks of successive radii matching the colors of its incident context arcs, with a white-filled center behind the vertex label: the 12 completion vertices $14, \ldots, 25$ have 1 colored disk, the 3 coordinate vertices $2, 5, 8$ have 2 nested disks, the 10 vertices $1, 3, 4, 6, 7, 9, 10, 11, 12, 13$ have 3 nested disks, and the common apex vertex 26 at the center is surrounded by 16 nested colored disks corresponding to all 16 contexts containing it. Unmarked arc crossings are not vertices.

TABLE II. The 24 core state supports $S_\alpha \subset V_{\mathrm{core}}$ $(\alpha = 1, \ldots, 24)$ of H_{65}, grouped by their hub ray $h \in \{9, 10, 11\}$ and opposite bridge ray $b \in B_1 \cup B_2 \cup B_3$.

Hub h	Bridge b	Allowed $(i_1, i_2) \in C_1 \times C_2$	Supports $S_\alpha = \{i_1, i_2, h, b\}$
9	16	$\{1,2\} \times \{5,6\}$	$S_1 = \{1,5,9,16\}$, $S_3 = \{1,6,9,16\}$, $S_7 = \{2,5,9,16\}$, $S_9 = \{2,6,9,16\}$
9	17	$\{3,4\} \times \{7,8\}$	$S_{15} = \{3,7,9,17\}$, $S_{17} = \{3,8,9,17\}$, $S_{21} = \{4,7,9,17\}$, $S_{23} = \{4,8,9,17\}$
10	14	$\{1,3\} \times \{5,7\}$	$S_2 = \{1,5,10,14\}$, $S_5 = \{1,7,10,14\}$, $S_{13} = \{3,5,10,14\}$, $S_{16} = \{3,7,10,14\}$
10	15	$\{2,4\} \times \{6,8\}$	$S_{10} = \{2,6,10,15\}$, $S_{11} = \{2,8,10,15\}$, $S_{19} = \{4,6,10,15\}$, $S_{24} = \{4,8,10,15\}$
11	12	$\{1,4\} \times \{6,7\}$	$S_4 = \{1,6,11,12\}$, $S_6 = \{1,7,11,12\}$, $S_{20} = \{4,6,11,12\}$, $S_{22} = \{4,7,11,12\}$
11	13	$\{2,3\} \times \{5,8\}$	$S_8 = \{2,5,11,13\}$, $S_{12} = \{2,8,11,13\}$, $S_{14} = \{3,5,11,13\}$, $S_{18} = \{3,8,11,13\}$

$v_i \cdot v_j = 0$ holds exactly for the $5 \times 6 + 24 \times 6 = 174$ edges of $\Gamma(H_{65})$.

- Second, deleting all 24 bridge edges from G_{core} leaves only the edges of $C_1, \ldots, C_5$, whose only K_4 cliques are $C_1, \ldots, C_5$ themselves. Since every fresh completion vertex $w \in \{18, \ldots, 65\}$ has degree 3 in $\Gamma(H_{65})$ with neighborhood equal to the remaining three vertices of its unique completion context E_m, the 4-cliques of $\Gamma(H_{65})$ are precisely the 29 contexts of E_{29}, and $\omega(\Gamma(H_{65})) = 4$ (which also follows directly from the injective orthogonal realization in $\mathbb{R}^4$).

(3) To prove $\alpha(G_{\mathrm{core}}) = 4$ by hand, let $I \subseteq V_{\mathrm{core}}$ be an independent set in G_{core}, and write $W_9 = C_3 \cup C_4 \cup C_5 = \{9, 10, 11\} \cup B_1 \cup B_2 \cup B_3$. Since C_1 and C_2 are 4-cliques, $|I \cap C_1| \leq 1$ and $|I \cap C_2| \leq 1$. Moreover, W_9 is partitioned into the triangle $\{9, 10, 11\}$ and the three edges B_1, B_2, B_3, so $|I \cap W_9| \leq 4$; in fact, if $I \cap \{9, 10, 11\} \neq \emptyset$, say $9 \in I$, then $9 \in C_3 \cap C_4$ excludes all




FIG. 2. Complete four-uniform hypergraph diagram of the canonical linear 4-uniform hypergraph $H_{65} = (V_{65}, E_{29})$ in $\mathbb{R}^4$ of Theorem 4 (65 pairwise non-collinear integer vectors $v_1, \ldots, v_{65} \in \mathbb{Z}^4$ enumerated in Eq. (12) and Table IV, 29 four-element contexts listed in Eq. (13) and Table V, and $29 \times \binom{4}{2} = 174$ orthogonality edges). All 29 four-element contexts have distinct colors, matched by the concentric filled disk layers of successive radii surrounding each vertex number, with a white center behind every label: the five core contexts $C_1 = \{1, 2, 3, 4\}$, $C_2 = \{5, 6, 7, 8\}$, $C_3 = \{9, 10, 12, 13\}$, $C_4 = \{9, 11, 14, 15\}$, and $C_5 = \{10, 11, 16, 17\}$ are drawn as bold smooth curves, while the 24 bridge-completion contexts $E_1, \ldots, E_{24}$ (such as the straight line $E_{21} = \{7, 16, 58, 59\}$) connect the binary bridge vertices $12, \ldots, 17$ (each surrounded by 5 nested colored disks) to the basis vertices $1, \ldots, 8$ (4 nested disks) via the degree-1 completion pairs $18, \ldots, 65$ (1 colored disk), with the hub vertices $9, 10, 11$ surrounded by 2 nested disks. Unmarked crossings are not vertices.

TABLE III. Orthogonal neighbors in $C_1 = \{1, 2, 3, 4\}$ and $C_2 = \{5, 6, 7, 8\}$ for each of the six binary bridge rays $b \in B_1 \cup B_2 \cup B_3$.

Pair	Ray b	$N(b) \cap C_1$	$N(b) \cap C_2$
B_1	$12 = (1, 0, 0, -1)$	$\{2, 3\}$	$\{5, 8\}$
B_1	$13 = (0, 1, 1, 0)$	$\{1, 4\}$	$\{6, 7\}$
B_2	$14 = (1, 0, 1, 0)$	$\{2, 4\}$	$\{6, 8\}$
B_2	$15 = (0, 1, 0, -1)$	$\{1, 3\}$	$\{5, 7\}$
B_3	$16 = (1, 1, 0, 0)$	$\{3, 4\}$	$\{7, 8\}$
B_3	$17 = (0, 0, 1, -1)$	$\{1, 2\}$	$\{5, 6\}$

of $\{10, 11, 12, 13, 14, 15\}$ and leaves at most one vertex in $B_3 = \{16, 17\}$, so $|I \cap W_9| \leq 2$ and hence $|I| \leq 1+1+2 = 4$.

The only remaining way to have $|I| \geq 5$ is for $I \cap \{9, 10, 11\} = \emptyset$ and $|I \cap W_9| = 3$, which requires $I \cap W_9 = \{b_1, b_2, b_3\}$ with $(b_1, b_2, b_3) \in B_1 \times B_2 \times B_3$. Table III lists the orthogonal neighbors of each $b \in B_1 \cup B_2 \cup B_3$ in C_1 and C_2. Call a triple $(b_1, b_2, b_3) \in B_1 \times B_2 \times B_3$ even or odd according to the parity of the number of its entries in $\{13, 15, 17\}$, the second elements of the three ordered bridge pairs. Observe the complementary parity of the $2 + 2$ partitions of C_1 and C_2:

- For the four *even-parity* triples $(b_1, b_2, b_3) \in \{(12, 14, 16), (12, 15, 17), (13, 14, 17), (13, 15, 16)\}$, the union of their orthogonal neighbors in C_2 is all of $C_2 = \{5, 6, 7, 8\}$ (leaving only one non-neighbor in C_1 and *zero* non-neighbors in C_2), so $|I| \leq 3 + 1 + 0 = 4$.
- For the four *odd-parity* triples $(b_1, b_2, b_3) \in$

$\{(12,14,17),(12,15,16),(13,14,16),(13,15,17)\}$, the union of their orthogonal neighbors in C_1 is all of $C_1 = \{1,2,3,4\}$ (leaving *zero* non-neighbors in C_1 and only one non-neighbor in C_2), so again $|I| \le 3+0+1 = 4$.

Thus $\alpha(G_{\text{core}}) = 4$ (attained, e.g., by $S_1 = \{1,5,9,16\}$). In any proper k-coloring of G_{core}, each color class is an independent set of size at most 4, so $4k \ge |V_{\text{core}}| = 17$, forcing $k \ge \lceil 17/4 \rceil = 5$ and $\chi_f(\Gamma(H_{65})) \ge \chi_f(G_{\text{core}}) \ge 17/4 = 4.25$. Together with the explicit 5-coloring in Eq. (12), this establishes $\chi(H_{65}) = 5 > 4 = \omega(\Gamma(H_{65}))$.

For the matching fractional upper bound, the independent four-sets of G_{core} consist of the 24 sets in Table II, each with one hub, one bridge vertex, and one vertex from each of C_1, C_2; 12 sets with two bridge vertices from different pairs and one vertex from each basis; and 8 sets with three bridge vertices and one basis vertex. These counts follow from the neighbor lists in Table III, which also show that assigning weight 1/8 to each set of the first type and 1/16 to each of the other types covers every vertex with total weight one. The total weight is $24/8 + 12/16 + 8/16 = 17/4$.

Equivalently, take two copies of each of the first 24 independent sets and one copy of each of the other 20, obtaining 68 color classes in which every core vertex occurs 16 times. For any bridge edge, its adjacent endpoints occupy 32 disjoint classes, leaving 36 classes available. Assign 16 available classes to its first completion vertex and 16 other available classes to its second. Completion vertices from different contexts are nonadjacent, so these choices are independent. This extends the coloring to all 65 vertices and proves $\chi_f(\Gamma(H_{65})) \le 68/16 = 17/4$. Together with the lower bound, this proves equality.

(4) A subset $S \subset V_{\text{core}}$ is the support of a core state compatible with G_{core} if and only if S is independent in G_{core} and $|S \cap C_a| = 1$ for $a = 1,\dots,5$. (Without the 24 bridge-edge constraints, the five-context hypergraph $(V_{\text{core}}, \{C_1,\dots,C_5\})$ alone has $4\times4\times(3\times2+2^3) = 224$ two-valued states.) By the parity analysis in Part (3), no triple $(b_1,b_2,b_3) \in B_1\times B_2\times B_3$ can be extended to include both a vertex of C_1 and a vertex of C_2. Consequently, S must select:

- exactly one hub vertex $h \in \{9,10,11\}$ (3 choices), which satisfies the two contexts among C_3, C_4, C_5 containing h;
- one binary bridge vertex b from the unique remaining pair B_k (2 choices), satisfying the third context among C_3, C_4, C_5;
- one of the two vertices $i_1 \in C_1 \setminus N(b)$ (2 choices); and
- one of the two vertices $i_2 \in C_2 \setminus N(b)$ (2 choices).

This yields precisely $3\times2\times2\times2 = 24$ graph-compatible core state supports $S_1,\dots,S_{24}$, listed in Table II.

To count the full state space $\mathcal{S}(H_{65})$, note that in the 24-edge bipartite bridge graph between $C_1 \cup C_2$ and $B_1 \cup B_2 \cup B_3$, every coordinate vertex $i_1 \in C_1$ has bridge degree 3, every Hadamard vertex $i_2 \in C_2$ has bridge degree 3, every bridge vertex $b \in B_1 \cup B_2 \cup B_3$ has bridge degree 4, and every hub vertex $h \in \{9,10,11\}$ has bridge degree 0. Because each $S_\alpha = \{i_1, i_2, h, b\}$ is independent in G_{core}, no bridge edge has both endpoints in S_α; hence *every* S_α meets $3+3+0+4 = 10$ bridge edges and misses $24-10 = 14$ bridge edges. Each of the 14 missed completion contexts $E_m = \{u_m, v_m, w_{m,1}, w_{m,2}\}$ admits 2 independent binary extensions $(s(w_{m,1}), s(w_{m,2})) \in \{(1,0),(0,1)\}$, while the 10 hit completion contexts force $(0,0)$, giving $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$.

Finally, to see that the 24 edge zero-profiles $Z(e_m)$ in Eq. (9) are nonempty and pairwise distinct, partition $S_1,\dots,S_{24}$ into three blocks $\mathcal{B}_9, \mathcal{B}_{10}, \mathcal{B}_{11}$ of 8 supports each according to their hub ray $h \in \{9,10,11\}$ (the three horizontal sections of Table II). Every vertex $u \in C_1 \cup C_2$ occurs in 2 supports in each of the three blocks $\mathcal{B}_9, \mathcal{B}_{10}, \mathcal{B}_{11}$ (total weight 6, block weights $(2,2,2)$), whereas each bridge vertex $v \in B_k$ occurs in 4 supports concentrated entirely in the single block $\mathcal{B}_{h_k}$ of the hub h_k opposite B_k (block weights $(4,0,0)$ up to permutation). Since $u_m \perp v_m$, the indicators $\mathbf{1}_{u_m}$ and $\mathbf{1}_{v_m}$ are disjoint, so the complement of $Z(e_m)$ has indicator $f_m = \mathbf{1}_{u_m} + \mathbf{1}_{v_m}$ of weight $6+4 = 10$ (so $|Z(e_m)| = 24-10 = 14 > 0$) and block weights $(6,2,2)$ up to permutation. The unique block of weight 6 identifies $\mathcal{B}_{h_k}$ and hence B_k; within $\mathcal{B}_{h_k}$, f_m takes value 1 on all 4 supports containing v_m and on only 2 of the 4 supports containing its partner $v'_m \in B_k$, which uniquely identifies v_m; and subtracting $\mathbf{1}_{v_m}$ from f_m recovers $\mathbf{1}_{u_m}$, which uniquely identifies $u_m \in C_1 \cup C_2$ since all 17 core vertex profiles are distinct. Thus all 24 zero-profiles $Z(e_1),\dots,Z(e_{24})$ are nonempty and pairwise distinct, and Lemma 3(3) guarantees that the 48 canonical states $s_{\alpha,\varepsilon}$ are unital and separating on all 65 vertices of H_{65}. □

A. Scope of the partition representation

For either the canonical family or the full state space, the sets A_i of Eq. (3) separate all named vertices and turn the listed contexts into partitions, while remaining weaker than an order-determining or compound-event-faithful representation (as already seen for the warm-up $H_{\text{YO}}^{(4)}$ in Appendix B 1, where the four non-orthogonal triad atoms $10,11,12,13$ have pairwise disjoint state sets $A_{10}, A_{11}, A_{12}, A_{13}$ and satisfy strict inclusions such as $A_{11} \subsetneq A_{25}$). For H_{65}, the classification of its core supports similarly implies

$$A_9 = A_{16} \dot\cup A_{17}, \quad A_{10} = A_{14} \dot\cup A_{15}, \quad A_{11} = A_{12} \dot\cup A_{13}. \tag{14}$$

Here $\dot\cup$ denotes disjoint union. For example, Eq. (14) gives the same set image to the ray $[v_9]$ and the plane $\text{span}(v_{16}, v_{17})$, which are distinct Hilbert-space events. Thus the vertex-and-context representation is not an in-

jective representation of all compound Hilbert events.

This distinction also appears at the probability level. Let $q = (1,1,1,-1)^T$, the missing fourth odd-parity Hadamard ray, and define $\Pi_i = v_i v_i^T/\|v_i\|^2$ and $\Pi_q = qq^T/4$. Direct summation gives

$$\sum_{i=1}^{17} \Pi_i = 4I_4 + \Pi_q. \tag{15}$$

Every two-valued state has exactly four active core vertices, so every convex mixture of such states gives total core probability 4. By Eq. (15), quantum preparation along q, represented by the unit vector $q/2$, gives total core probability 5. The existence of separating two-valued states therefore does not imply a classical model for all quantum preparations.

Appendix A: Explicit enumeration of unnormalized vectors, contexts, and non-collinearity certificates

For direct human and computational verification, this appendix enumerates all unnormalized integer vectors in $\mathbb{Z}^4$ and all four-element contexts for both hypergraphs studied in this paper:

1. The *canonical linear hypergraph* $H_{65} = (V_{65}, E_{29})$ (Theorem 4, Appendix A 1), which has exactly 65 **pairwise non-collinear integer vectors** (17 core vectors $v_1, \ldots, v_{17}$ and 48 degree-1 completion vectors $v_{18}, \ldots, v_{65}$), 29 **four-element contexts** (5 core contexts $C_1, \ldots, C_5$ and 24 bridge-completion contexts $E_1, \ldots, E_{24}$), and 174 **orthogonality edges** ($29 \times \binom{4}{2} = 174$).

2. The *non-linear warm-up apex suspension* $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$ (Proposition 1, Appendix A 2), which has exactly 26 **pairwise non-collinear integer vectors** (13 Yu–Oh rays $v_1, \ldots, v_{13}$, 12 completion rays $v_{14}, \ldots, v_{25}$ in $\mathbb{R}^3 \times \{0\}$, and 1 apex ray v_{26}), 16 **four-element contexts**, and 73 **orthogonality edges** ($16 \times \binom{3}{2} = 48$ edges in $\mathbb{R}^3 \times \{0\}$ plus 25 apex edges).

Canonical primitive normal form and human-verifiable non-collinearity. Every vector $v_i = (x_{i,1}, x_{i,2}, x_{i,3}, x_{i,4})^T \in \mathbb{Z}^4$ listed below in both H_{65} and $H_{\mathrm{YO}}^{(4)}$ is written in *canonical primitive projective form*:

$$\begin{gathered}\gcd\big(|x_{i,1}|, |x_{i,2}|, |x_{i,3}|, |x_{i,4}|\big) = 1, \\ \text{first nonzero entry of } v_i > 0.\end{gathered} \tag{A1}$$

Because every projective ray $[v] \in \mathbb{P}(\mathbb{Q}^4)$ possesses a *unique* representative in $\mathbb{Z}^4$ satisfying Eq. (A1), two vectors v_i, v_j in canonical primitive form are collinear (proportional over $\mathbb{R}$ or $\mathbb{C}$) *if and only if they are identical* 4-*tuples* ($v_i = v_j$) and, in particular, have the same squared Euclidean norm $\|v_i\|^2 = \|v_j\|^2$.

1. The canonical linear 65-vertex, 29-context hypergraph H_{65}

Table IV enumerates all 65 unnormalized integer vectors $v_1, \ldots, v_{65} \in \mathbb{Z}^4$ of H_{65}, together with their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0,1,2,3,4\}$ in the strong 5-coloring of Eq. (12), and the contexts in E_{29} to which each vertex belongs. Table V enumerates all 29 contexts $C_1, \ldots, C_5, E_1, \ldots, E_{24} \in E_{29}$, displaying side by side both the vertex indices $\{i,j,k,\ell\}$ and the four mutually orthogonal integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

To verify by hand that all 65 vectors in Table IV are pairwise non-collinear across all $\binom{65}{2} = 2{,}080$ pairs, observe that all 65 tuples satisfy Eq. (A1) and partition by squared norm $\|v_i\|^2$ into seven small, disjoint norm shells ($4 + 6 + 7 + 12 + 12 + 12 + 12 = 65$):

- $\|v_i\|^2 = 1$ (4 vectors): indices $\{1,2,3,4\}$, the 4 standard coordinate basis vectors;
- $\|v_i\|^2 = 2$ (6 vectors): indices $\{12, \ldots, 17\}$, one vector for each of the $\binom{4}{2} = 6$ coordinate-pair supports;
- $\|v_i\|^2 = 4$ (7 vectors): indices $\{5, \ldots, 11\}$, seven distinct ± 1 vectors with $x_{i,1} = +1$;
- $\|v_i\|^2 = 11$ (12 vectors): indices $\{18, 20, 22, 25, 26, 28, 30, 33, 34, 36, 39, 41\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct sign/entry patterns of $\{0, \pm 1, \pm 1, \pm 3\}$;
- $\|v_i\|^2 = 18$ (12 vectors): indices $\{43, 45, 46, 49, 51, 53, 54, 56, 59, 61, 62, 65\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 2, \pm 2, \pm 3\}$;
- $\|v_i\|^2 = 22$ (12 vectors): indices $\{19, 21, 23, 24, 27, 29, 31, 32, 35, 37, 38, 40\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct patterns of $\{0, \pm 2, \pm 3, \pm 3\}$;
- $\|v_i\|^2 = 36$ (12 vectors): indices $\{42, 44, 47, 48, 50, 52, 55, 57, 58, 60, 63, 64\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 1, \pm 3, \pm 5\}$.

Because no two 4-tuples within any of the seven norm shells coincide, no two of the 65 vectors are collinear; quantitatively, the maximum squared cosine over all $\binom{65}{2} = 2{,}080$ pairs is $\max_{1 \le i < j \le 65} (v_i \cdot v_j)^2/(\|v_i\|^2 \|v_j\|^2) = 19^2/(11 \times 36) = 361/396 \approx 0.9116 < 1$, attained by (v_{41}, v_{47}), whose squared norms are 11 and 36 and whose inner product is 19.

2. The non-linear warm-up 26-vertex, 16-context apex suspension $H_{\mathrm{YO}}^{(4)}$

For completeness, the 26 unnormalized integer vectors $v_1, \ldots, v_{26} \in \mathbb{Z}^4$ of the non-linear warm-up hypergraph

TABLE IV. Complete enumeration of the 65 unnormalized integer vectors $v_i \in \mathbb{Z}^4$ of H_{65}, their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0,1,2,3,4\}$, and the contexts in E_{29} containing vertex i.

i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i	i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i
1	$(1,0,0,0)$	1	0	C_1, E_3, E_7, E_{11}	34	$(1,-1,0,3)$	11	1	$E_9 = \{3,16,34,35\}$
2	$(0,1,0,0)$	1	1	C_1, E_1, E_5, E_{12}	35	$(3,-3,0,-2)$	22	3	$E_9 = \{3,16,34,35\}$
3	$(0,0,1,0)$	1	2	C_1, E_2, E_8, E_9	36	$(1,-1,3,0)$	11	1	$E_{10} = \{4,16,36,37\}$
4	$(0,0,0,1)$	1	3	C_1, E_4, E_6, E_{10}	37	$(3,-3,-2,0)$	22	2	$E_{10} = \{4,16,36,37\}$
5	$(1,1,1,1)$	4	0	$C_2, E_{13}, E_{19}, E_{23}$	38	$(0,2,3,3)$	22	1	$E_{11} = \{1,17,38,39\}$
6	$(1,1,-1,-1)$	4	1	$C_2, E_{15}, E_{17}, E_{24}$	39	$(0,3,-1,-1)$	11	3	$E_{11} = \{1,17,38,39\}$
7	$(1,-1,1,-1)$	4	2	$C_2, E_{16}, E_{20}, E_{21}$	40	$(2,0,-3,-3)$	22	0	$E_{12} = \{2,17,40,41\}$
8	$(1,-1,-1,1)$	4	4	$C_2, E_{14}, E_{18}, E_{22}$	41	$(3,0,1,1)$	11	3	$E_{12} = \{2,17,40,41\}$
9	$(1,1,-1,1)$	4	0	C_3, C_4	42	$(1,-5,3,1)$	36	1	$E_{13} = \{5,12,42,43\}$
10	$(1,-1,1,1)$	4	1	C_3, C_5	43	$(2,-1,-3,2)$	18	2	$E_{13} = \{5,12,42,43\}$
11	$(1,-1,-1,-1)$	4	3	C_4, C_5	44	$(1,-3,5,1)$	36	0	$E_{14} = \{8,12,44,45\}$
12	$(1,0,0,-1)$	2	3	$C_3, E_1, E_2, E_{13}, E_{14}$	45	$(2,3,1,2)$	18	1	$E_{14} = \{8,12,44,45\}$
13	$(0,1,1,0)$	2	4	$C_3, E_3, E_4, E_{15}, E_{16}$	46	$(1,-2,2,-3)$	18	0	$E_{15} = \{6,13,46,47\}$
14	$(1,0,1,0)$	2	2	$C_4, E_5, E_6, E_{17}, E_{18}$	47	$(5,-1,1,3)$	36	2	$E_{15} = \{6,13,46,47\}$
15	$(0,1,0,-1)$	2	1	$C_4, E_7, E_8, E_{19}, E_{20}$	48	$(3,-1,1,5)$	36	0	$E_{16} = \{7,13,48,49\}$
16	$(1,1,0,0)$	2	0	$C_5, E_9, E_{10}, E_{21}, E_{22}$	49	$(3,2,-2,-1)$	18	1	$E_{16} = \{7,13,48,49\}$
17	$(0,0,1,-1)$	2	2	$C_5, E_{11}, E_{12}, E_{23}, E_{24}$	50	$(1,3,-1,5)$	36	0	$E_{17} = \{6,14,50,51\}$
18	$(1,0,-3,1)$	11	0	$E_1 = \{2,12,18,19\}$	51	$(2,-3,-2,1)$	18	3	$E_{17} = \{6,14,50,51\}$
19	$(3,0,2,3)$	22	2	$E_1 = \{2,12,18,19\}$	52	$(1,5,-1,3)$	36	0	$E_{18} = \{8,14,52,53\}$
20	$(1,3,0,1)$	11	0	$E_2 = \{3,12,20,21\}$	53	$(2,1,-2,-3)$	18	1	$E_{18} = \{8,14,52,53\}$
21	$(3,-2,0,3)$	22	1	$E_2 = \{3,12,20,21\}$	54	$(3,-2,1,-2)$	18	2	$E_{19} = \{5,15,54,55\}$
22	$(0,1,-1,3)$	11	1	$E_3 = \{1,13,22,23\}$	55	$(3,1,-5,1)$	36	3	$E_{19} = \{5,15,54,55\}$
23	$(0,3,-3,-2)$	22	2	$E_3 = \{1,13,22,23\}$	56	$(1,2,3,2)$	18	0	$E_{20} = \{7,15,56,57\}$
24	$(2,-3,3,0)$	22	0	$E_4 = \{4,13,24,25\}$	57	$(5,1,-3,1)$	36	3	$E_{20} = \{7,15,56,57\}$
25	$(3,1,-1,0)$	11	1	$E_4 = \{4,13,24,25\}$	58	$(1,-1,-5,-3)$	36	1	$E_{21} = \{7,16,58,59\}$
26	$(1,0,-1,-3)$	11	0	$E_5 = \{2,14,26,27\}$	59	$(2,-2,-1,3)$	18	3	$E_{21} = \{7,16,58,59\}$
27	$(3,0,-3,2)$	22	3	$E_5 = \{2,14,26,27\}$	60	$(1,-1,-3,-5)$	36	1	$E_{22} = \{8,16,60,61\}$
28	$(1,-3,-1,0)$	11	0	$E_6 = \{4,14,28,29\}$	61	$(2,-2,3,-1)$	18	2	$E_{22} = \{8,16,60,61\}$
29	$(3,2,-3,0)$	22	1	$E_6 = \{4,14,28,29\}$	62	$(1,3,-2,-2)$	18	1	$E_{23} = \{5,17,62,63\}$
30	$(0,1,3,1)$	11	2	$E_7 = \{1,15,30,31\}$	63	$(5,-3,-1,-1)$	36	3	$E_{23} = \{5,17,62,63\}$
31	$(0,3,-2,3)$	22	3	$E_7 = \{1,15,30,31\}$	64	$(3,-5,-1,-1)$	36	0	$E_{24} = \{6,17,64,65\}$
32	$(2,3,0,3)$	22	0	$E_8 = \{3,15,32,33\}$	65	$(3,1,2,2)$	18	3	$E_{24} = \{6,17,64,65\}$
33	$(3,-1,0,-1)$	11	3	$E_8 = \{3,15,32,33\}$					

$H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$ in Proposition 1 (written in canonical primitive form satisfying Eq. (A1)) are:

$$
\begin{aligned}
v_1 &= (0,1,1,0), & v_2 &= (1,0,0,0),\\
v_3 &= (0,1,-1,0), & v_4 &= (1,1,0,0),\\
v_5 &= (0,0,1,0), & v_6 &= (1,-1,0,0),\\
v_7 &= (1,0,1,0), & v_8 &= (0,1,0,0),\\
v_9 &= (1,0,-1,0), & v_{10} &= (1,-1,1,0),\\
v_{11} &= (1,-1,-1,0), & v_{12} &= (1,1,1,0),\\
v_{13} &= (1,1,-1,0), & v_{14} &= (2,1,1,0),\\
v_{15} &= (1,2,-1,0), & v_{16} &= (1,-2,1,0),\\
v_{17} &= (1,1,-2,0), & v_{18} &= (1,-1,-2,0),\\
v_{19} &= (2,1,-1,0), & v_{20} &= (2,-1,1,0),\\
v_{21} &= (1,-2,-1,0), & v_{22} &= (1,1,2,0),\\
v_{23} &= (1,2,1,0), & v_{24} &= (1,-1,2,0),\\
v_{25} &= (2,-1,-1,0), & v_{26} &= (0,0,0,1).
\end{aligned}
\tag{A2}
$$

These 26 canonical primitive vectors partition by squared norm $\|v_i\|^2$ into four disjoint norm shells ($4+6+4+12 = 26$):

- $\|v_i\|^2 = 1$ (4 vectors): $\{v_2, v_5, v_8, v_{26}\}$, the four coordinate basis vectors of $\mathbb{R}^4$;
- $\|v_i\|^2 = 2$ (6 vectors): $\{v_1, v_3, v_4, v_6, v_7, v_9\}$, the six binary rays in $\mathbb{R}^3 \times \{0\}$;
- $\|v_i\|^2 = 3$ (4 vectors): $\{v_{10}, v_{11}, v_{12}, v_{13}\}$, the four triad rays $(1, \pm 1, \pm 1, 0)^T$;
- $\|v_i\|^2 = 6$ (12 vectors): $\{v_{14}, \ldots, v_{25}\}$, all $3 \times 4 = 12$ canonical primitive permutations and sign patterns of $(\pm 2, \pm 1, \pm 1, 0)^T$.

Since all 26 canonical 4-tuples are distinct, all $\binom{26}{2} = 325$ pairs of vectors are strictly non-collinear, with maximum squared cosine $\max_{1 \le i < j \le 26} (v_i \cdot v_j)^2 / (\|v_i\|^2 \|v_j\|^2) = 4^2/(3 \times 6) = 8/9 \approx 0.8889 < 1$. Table VI lists all 16 four-element contexts of $E_{16}^{(4)}$ side by side with their four mutually orthogonal integer vectors.

TABLE V. Complete enumeration of the 29 four-element contexts of $H_{65} = (V_{65}, E_{29})$, listing for each context both its vertex indices $\{i, j, k, \ell\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

Context	Vertex set	v_i	v_j	v_k	v_ℓ
Five core 4-contexts $C_1, \ldots, C_5$ on $V_{\text{core}} = \{1, \ldots, 17\}$:					
C_1	$\{1, 2, 3, 4\}$	$v_1 = (1, 0, 0, 0)$	$v_2 = (0, 1, 0, 0)$	$v_3 = (0, 0, 1, 0)$	$v_4 = (0, 0, 0, 1)$
C_2	$\{5, 6, 7, 8\}$	$v_5 = (1, 1, 1, 1)$	$v_6 = (1, 1, -1, -1)$	$v_7 = (1, -1, 1, -1)$	$v_8 = (1, -1, -1, 1)$
C_3	$\{9, 10, 12, 13\}$	$v_9 = (1, 1, -1, 1)$	$v_{10} = (1, -1, 1, 1)$	$v_{12} = (1, 0, 0, -1)$	$v_{13} = (0, 1, 1, 0)$
C_4	$\{9, 11, 14, 15\}$	$v_9 = (1, 1, -1, 1)$	$v_{11} = (1, -1, -1, -1)$	$v_{14} = (1, 0, 1, 0)$	$v_{15} = (0, 1, 0, -1)$
C_5	$\{10, 11, 16, 17\}$	$v_{10} = (1, -1, 1, 1)$	$v_{11} = (1, -1, -1, -1)$	$v_{16} = (1, 1, 0, 0)$	$v_{17} = (0, 0, 1, -1)$
Twelve coordinate-to-bridge completion contexts $E_1, \ldots, E_{12}$ (between C_1 and $B_1 \cup B_2 \cup B_3$):					
E_1	$\{2, 12, 18, 19\}$	$v_2 = (0, 1, 0, 0)$	$v_{12} = (1, 0, 0, -1)$	$v_{18} = (1, 0, -3, 1)$	$v_{19} = (3, 0, 2, 3)$
E_2	$\{3, 12, 20, 21\}$	$v_3 = (0, 0, 1, 0)$	$v_{12} = (1, 0, 0, -1)$	$v_{20} = (1, 3, 0, 1)$	$v_{21} = (3, -2, 0, 3)$
E_3	$\{1, 13, 22, 23\}$	$v_1 = (1, 0, 0, 0)$	$v_{13} = (0, 1, 1, 0)$	$v_{22} = (0, 1, -1, 3)$	$v_{23} = (0, 3, -3, -2)$
E_4	$\{4, 13, 24, 25\}$	$v_4 = (0, 0, 0, 1)$	$v_{13} = (0, 1, 1, 0)$	$v_{24} = (2, -3, 3, 0)$	$v_{25} = (3, 1, -1, 0)$
E_5	$\{2, 14, 26, 27\}$	$v_2 = (0, 1, 0, 0)$	$v_{14} = (1, 0, 1, 0)$	$v_{26} = (1, 0, -1, -3)$	$v_{27} = (3, 0, -3, 2)$
E_6	$\{4, 14, 28, 29\}$	$v_4 = (0, 0, 0, 1)$	$v_{14} = (1, 0, 1, 0)$	$v_{28} = (1, -3, -1, 0)$	$v_{29} = (3, 2, -3, 0)$
E_7	$\{1, 15, 30, 31\}$	$v_1 = (1, 0, 0, 0)$	$v_{15} = (0, 1, 0, -1)$	$v_{30} = (0, 1, 3, 1)$	$v_{31} = (0, 3, -2, 3)$
E_8	$\{3, 15, 32, 33\}$	$v_3 = (0, 0, 1, 0)$	$v_{15} = (0, 1, 0, -1)$	$v_{32} = (2, 3, 0, 3)$	$v_{33} = (3, -1, 0, -1)$
E_9	$\{3, 16, 34, 35\}$	$v_3 = (0, 0, 1, 0)$	$v_{16} = (1, 1, 0, 0)$	$v_{34} = (1, -1, 0, 3)$	$v_{35} = (3, -3, 0, -2)$
E_{10}	$\{4, 16, 36, 37\}$	$v_4 = (0, 0, 0, 1)$	$v_{16} = (1, 1, 0, 0)$	$v_{36} = (1, -1, 3, 0)$	$v_{37} = (3, -3, -2, 0)$
E_{11}	$\{1, 17, 38, 39\}$	$v_1 = (1, 0, 0, 0)$	$v_{17} = (0, 0, 1, -1)$	$v_{38} = (0, 2, 3, 3)$	$v_{39} = (0, 3, -1, -1)$
E_{12}	$\{2, 17, 40, 41\}$	$v_2 = (0, 1, 0, 0)$	$v_{17} = (0, 0, 1, -1)$	$v_{40} = (2, 0, -3, -3)$	$v_{41} = (3, 0, 1, 1)$
Twelve Hadamard-to-bridge completion contexts $E_{13}, \ldots, E_{24}$ (between C_2 and $B_1 \cup B_2 \cup B_3$):					
E_{13}	$\{5, 12, 42, 43\}$	$v_5 = (1, 1, 1, 1)$	$v_{12} = (1, 0, 0, -1)$	$v_{42} = (1, -5, 3, 1)$	$v_{43} = (2, -1, -3, 2)$
E_{14}	$\{8, 12, 44, 45\}$	$v_8 = (1, -1, -1, 1)$	$v_{12} = (1, 0, 0, -1)$	$v_{44} = (1, -3, 5, 1)$	$v_{45} = (2, 3, 1, 2)$
E_{15}	$\{6, 13, 46, 47\}$	$v_6 = (1, 1, -1, -1)$	$v_{13} = (0, 1, 1, 0)$	$v_{46} = (1, -2, 2, -3)$	$v_{47} = (5, -1, 1, 3)$
E_{16}	$\{7, 13, 48, 49\}$	$v_7 = (1, -1, 1, -1)$	$v_{13} = (0, 1, 1, 0)$	$v_{48} = (3, -1, 1, 5)$	$v_{49} = (3, 2, -2, -1)$
E_{17}	$\{6, 14, 50, 51\}$	$v_6 = (1, 1, -1, -1)$	$v_{14} = (1, 0, 1, 0)$	$v_{50} = (1, 3, -1, 5)$	$v_{51} = (2, -3, -2, 1)$
E_{18}	$\{8, 14, 52, 53\}$	$v_8 = (1, -1, -1, 1)$	$v_{14} = (1, 0, 1, 0)$	$v_{52} = (1, 5, -1, 3)$	$v_{53} = (2, 1, -2, -3)$
E_{19}	$\{5, 15, 54, 55\}$	$v_5 = (1, 1, 1, 1)$	$v_{15} = (0, 1, 0, -1)$	$v_{54} = (3, -2, 1, -2)$	$v_{55} = (3, 1, -5, 1)$
E_{20}	$\{7, 15, 56, 57\}$	$v_7 = (1, -1, 1, -1)$	$v_{15} = (0, 1, 0, -1)$	$v_{56} = (1, 2, 3, 2)$	$v_{57} = (5, 1, -3, 1)$
E_{21}	$\{7, 16, 58, 59\}$	$v_7 = (1, -1, 1, -1)$	$v_{16} = (1, 1, 0, 0)$	$v_{58} = (1, -1, -5, -3)$	$v_{59} = (2, -2, -1, 3)$
E_{22}	$\{8, 16, 60, 61\}$	$v_8 = (1, -1, -1, 1)$	$v_{16} = (1, 1, 0, 0)$	$v_{60} = (1, -1, -3, -5)$	$v_{61} = (2, -2, 3, -1)$
E_{23}	$\{5, 17, 62, 63\}$	$v_5 = (1, 1, 1, 1)$	$v_{17} = (0, 0, 1, -1)$	$v_{62} = (1, 3, -2, -2)$	$v_{63} = (5, -3, -1, -1)$
E_{24}	$\{6, 17, 64, 65\}$	$v_6 = (1, 1, -1, -1)$	$v_{17} = (0, 0, 1, -1)$	$v_{64} = (3, -5, -1, -1)$	$v_{65} = (3, 1, 2, 2)$

Appendix B: Explicit enumeration of two-valued states

This appendix provides the complete enumeration of all two-valued states on both hypergraphs studied in the main text: the 25 two-valued states of the nonlinear warm-up hypergraph $H_{\text{YO}}^{(4)}$ (Appendix B 1), and the complete branch-by-branch parametrization of all $24 \times 2^{14} = 393{,}216$ two-valued states—together with the 48 canonical unital and separating states $s_{\alpha,\varepsilon}$ and the 65 vertex state-profiles—of the canonical linear hypergraph H_{65} (Appendix B 2).

1. Complete enumeration of the 25 two-valued states on $H_{\text{YO}}^{(4)}$

Table VII lists all 25 two-valued states $s_1, \ldots, s_{25} : V_{26} \to \{0, 1\}$ on $H_{\text{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$. States $s_1, \ldots, s_{24}$ have $s_r(26) = 0$ and correspond to the 24 two-valued states of the completed Yu–Oh hypergraph H_{YO}^{+} in $\mathbb{R}^3$ (grouped into 12 pairs of states sharing the same support on the three coordinate contexts $\{1, 2, 3\}$, $\{4, 5, 6\}$, $\{7, 8, 9\}$), while s_{25} is the unique apex state with support $\{26\}$. Table VIII displays the transposed 26×25 vertex-by-state binary matrix, listing for each vertex $i \in \{1, \ldots, 26\}$ its state weight $\sum_{r=1}^{25} s_r(i) \in \{1, 3, 8, 13\}$, the set of states $A_i = \{s_r : s_r(i) = 1\}$, and its 25-bit profile $(s_1(i), \ldots, s_{25}(i))$.

Inspection of Table VIII highlights three structural properties of the state space $\mathcal{S}(H_{\text{YO}}^{(4)})$:

1. *Unitality and atom separation:* Every row of Table VIII has positive weight (1, 3, 8, or 13) and all 26 binary rows are distinct ($A_i \neq A_j$ for all $1 \leq i < j \leq 26$), so $\mathcal{S}(H_{\text{YO}}^{(4)})$ is nonempty, unital, and separating on atoms.

2. *Exclusion of simultaneous* 1*s on the non-orthogonal triad atoms* $\{10, 11, 12, 13\}$*:* Although the four Yu–Oh triad rays $v_{10}, v_{11}, v_{12}, v_{13}$ are pairwise non-orthogonal ($v_i \cdot v_j = \pm 1 \neq 0$ for $10 \leq i < j \leq 13$), their active state sets in Table VIII are pairwise

TABLE VI. Complete enumeration of the 16 four-element contexts of the non-linear warm-up apex suspension $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, listing for each context both its vertex indices $\{i, j, k, 26\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_{26}\} \subset \mathbb{Z}^4$.

Context	Vertex set	v_i	v_j	v_k	v_{26}
$e_1^{(4)}$	$\{1,2,3,26\}$	$v_1 = (0,1,1,0)$	$v_2 = (1,0,0,0)$	$v_3 = (0,1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_2^{(4)}$	$\{7,8,9,26\}$	$v_7 = (1,0,1,0)$	$v_8 = (0,1,0,0)$	$v_9 = (1,0,-1,0)$	$v_{26} = (0,0,0,1)$
$e_3^{(4)}$	$\{4,5,6,26\}$	$v_4 = (1,1,0,0)$	$v_5 = (0,0,1,0)$	$v_6 = (1,-1,0,0)$	$v_{26} = (0,0,0,1)$
$e_4^{(4)}$	$\{2,5,8,26\}$	$v_2 = (1,0,0,0)$	$v_5 = (0,0,1,0)$	$v_8 = (0,1,0,0)$	$v_{26} = (0,0,0,1)$
$e_5^{(4)}$	$\{3,12,25,26\}$	$v_3 = (0,1,-1,0)$	$v_{12} = (1,1,1,0)$	$v_{25} = (2,-1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_6^{(4)}$	$\{9,12,16,26\}$	$v_9 = (1,0,-1,0)$	$v_{12} = (1,1,1,0)$	$v_{16} = (1,-2,1,0)$	$v_{26} = (0,0,0,1)$
$e_7^{(4)}$	$\{6,12,17,26\}$	$v_6 = (1,-1,0,0)$	$v_{12} = (1,1,1,0)$	$v_{17} = (1,1,-2,0)$	$v_{26} = (0,0,0,1)$
$e_8^{(4)}$	$\{3,11,14,26\}$	$v_3 = (0,1,-1,0)$	$v_{11} = (1,-1,-1,0)$	$v_{14} = (2,1,1,0)$	$v_{26} = (0,0,0,1)$
$e_9^{(4)}$	$\{7,11,15,26\}$	$v_7 = (1,0,1,0)$	$v_{11} = (1,-1,-1,0)$	$v_{15} = (1,2,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{10}^{(4)}$	$\{4,11,24,26\}$	$v_4 = (1,1,0,0)$	$v_{11} = (1,-1,-1,0)$	$v_{24} = (1,-1,2,0)$	$v_{26} = (0,0,0,1)$
$e_{11}^{(4)}$	$\{9,10,23,26\}$	$v_9 = (1,0,-1,0)$	$v_{10} = (1,-1,1,0)$	$v_{23} = (1,2,1,0)$	$v_{26} = (0,0,0,1)$
$e_{12}^{(4)}$	$\{1,10,19,26\}$	$v_1 = (0,1,1,0)$	$v_{10} = (1,-1,1,0)$	$v_{19} = (2,1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{13}^{(4)}$	$\{4,10,18,26\}$	$v_4 = (1,1,0,0)$	$v_{10} = (1,-1,1,0)$	$v_{18} = (1,-1,-2,0)$	$v_{26} = (0,0,0,1)$
$e_{14}^{(4)}$	$\{1,13,20,26\}$	$v_1 = (0,1,1,0)$	$v_{13} = (1,1,-1,0)$	$v_{20} = (2,-1,1,0)$	$v_{26} = (0,0,0,1)$
$e_{15}^{(4)}$	$\{7,13,21,26\}$	$v_7 = (1,0,1,0)$	$v_{13} = (1,1,-1,0)$	$v_{21} = (1,-2,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{16}^{(4)}$	$\{6,13,22,26\}$	$v_6 = (1,-1,0,0)$	$v_{13} = (1,1,-1,0)$	$v_{22} = (1,1,2,0)$	$v_{26} = (0,0,0,1)$

disjoint:

$$A_{10} = \{s_{13}, s_{19}, s_{23}\}, \quad A_{11} = \{s_5, s_7, s_{15}\}, \\ A_{12} = \{s_1, s_3, s_9\}, \quad A_{13} = \{s_{11}, s_{17}, s_{21}\}. \tag{B1}$$

Consequently, $\sum_{i=10}^{13} s(i) \le 1$ holds for every $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$, and no two-valued state assigns $(1,1)$ to any of the $\binom{4}{2} = 6$ non-orthogonal pairs $\{10,11\}$, $\{10,12\}$, $\{10,13\}$, $\{11,12\}$, $\{11,13\}$, $\{12,13\}$.

3. *Failure of order-determinacy:* Even though $A_i \neq A_j$ for all $i \neq j$, the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is not order-determining on atoms or compound propositions [1]. In particular, each triad atom $t \in \{10,11,12,13\}$ (*weight* 3) is strictly dominated across all 25 states ($A_t \subsetneq A_w$, i.e., $s(t) \le s(w)$ for all $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$) by four non-orthogonal completion atoms $w \in \{14, \dots, 25\}$ (*weight* 13):

$$A_{10} \subsetneq A_{16}, A_{20}, A_{24}, A_{25}, \quad A_{11} \subsetneq A_{18}, A_{21}, A_{22}, A_{25}, \\ A_{12} \subsetneq A_{14}, A_{22}, A_{23}, A_{24}, \quad A_{13} \subsetneq A_{15}, A_{17}, A_{19}, A_{25}, \tag{B2}$$

together with compound-event inclusions such as $A_{12} \subsetneq A_3 \dot\cup A_{14}$, $A_{11} \subsetneq A_3 \dot\cup A_{25}$, and their unions with the apex set $A_{26} = \{s_{25}\}$.

2. Complete parametrization of all 393,216 two-valued states and the 48 canonical separating states on H_{65}

Every two-valued state $s \in \mathcal{S}(H_{65})$ on the canonical linear hypergraph $H_{65} = (V_{65}, E_{29})$ is uniquely indexed by a pair $(\alpha, \boldsymbol{\varepsilon})$ consisting of:

1. a branch index $\alpha \in \{1, \dots, 24\}$ selecting one of the 24 four-element core state supports $S_\alpha \subset V_{\mathrm{core}} = \{1, \dots, 17\}$; and
2. a 14-bit binary vector $\boldsymbol{\varepsilon} = (\varepsilon_m)_{m \in \mathcal{F}_\alpha} \in \{0,1\}^{14}$ indexed by the 14 *free completion contexts*

$$\mathcal{F}_\alpha = \big\{ m \in \{1, \dots, 24\} : S_\alpha \cap \{u_m, v_m\} = \emptyset \big\},$$

whose complement is the set of 10 *hit completion contexts* $\mathcal{H}_\alpha = \{1, \dots, 24\} \setminus \mathcal{F}_\alpha$.

Explicitly, the state $s_{\alpha,\boldsymbol{\varepsilon}} : V_{65} \to \{0,1\}$ takes values

$$s_{\alpha,\boldsymbol{\varepsilon}}(i) = \mathbf{1}_{i \in S_\alpha} \qquad (1 \le i \le 17), \\ \big(s_{\alpha,\boldsymbol{\varepsilon}}(2m+16),\, s_{\alpha,\boldsymbol{\varepsilon}}(2m+17)\big) \\ = \begin{cases} (0,0), & m \in \mathcal{H}_\alpha, \\ (1-\varepsilon_m,\, \varepsilon_m), & m \in \mathcal{F}_\alpha. \end{cases} \tag{B3}$$

Because $|\mathcal{F}_\alpha| = 14$ for every $\alpha \in \{1, \dots, 24\}$, each branch α yields $2^{14} = 16{,}384$ states $s_{\alpha,\boldsymbol{\varepsilon}}$, giving $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$ two-valued states in total. Setting all 14 free bits equal to $\varepsilon \in \{0,1\}$ ($\varepsilon_m = \varepsilon$ for all $m \in \mathcal{F}_\alpha$) yields the 48 *canonical states* $s_{\alpha,0}, s_{\alpha,1}$ of Eq. (8), which have support size $|S_\alpha| + 14 = 18$:

$$\mathrm{supp}(s_{\alpha,0}) = S_\alpha \cup \big\{2m+16 : m \in \mathcal{F}_\alpha\big\}, \\ \mathrm{supp}(s_{\alpha,1}) = S_\alpha \cup \big\{2m+17 : m \in \mathcal{F}_\alpha\big\}. \tag{B4}$$

Table IX enumerates all 24 branches $\alpha = 1, \dots, 24$, specifying S_α, the 10 hit context indices $\mathcal{H}_\alpha \subset \{1, \dots, 24\}$, the 14 free context indices $\mathcal{F}_\alpha \subset \{1, \dots, 24\}$, and the 24-bit free-context indicator string. Table X lists the resulting state profiles and weights for all 65 vertices of H_{65} across both the 48 canonical states $\{s_{\alpha,\varepsilon}\}$ and the full space of 393,216 states: across the 48 canonical

TABLE VII. Complete enumeration of the 25 two-valued states $s_1, \ldots, s_{25} : V_{26} \to \{0,1\}$ on the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, showing for each state its support on the 13-ray Yu–Oh core $\{1,\ldots,13\}$, its forced support on $\{14,\ldots,26\}$, its total support size $|S_r|$, and its 26-bit binary vector $(s_r(1\ldots13)\, s_r(14\ldots25)\, s_r(26))$.

State	Core support $S_r \cap \{1,\ldots,13\}$	Completion/apex support $S_r \cap \{14,\ldots,26\}$	$\lvert S_r\rvert$	Binary vector $(s_r(1\ldots13)\, s_r(14\ldots25)\, s_r(26))$
s_1	$\{1,4,8,12\}$	$\{14,15,21,22,23\}$	9	1001000100010 1100000111000
s_2	$\{1,4,8\}$	$\{14,15,16,17,21,22,23,25\}$	11	1001000100000 1111000111010
s_3	$\{1,5,7,12\}$	$\{14,18,22,23,24\}$	9	1000101000010 1000100011100
s_4	$\{1,5,7\}$	$\{14,16,17,18,22,23,24,25\}$	11	1000101000000 1011100011110
s_5	$\{1,5,9,11\}$	$\{17,18,21,22,25\}$	9	1000100010100 0001100110010
s_6	$\{1,5,9\}$	$\{14,15,17,18,21,22,24,25\}$	11	1000100010000 1101100110110
s_7	$\{1,6,8,11\}$	$\{16,18,21,23,25\}$	9	1000010100100 0010100101010
s_8	$\{1,6,8\}$	$\{14,15,16,18,21,23,24,25\}$	11	1000010100000 1110100101110
s_9	$\{2,4,7,12\}$	$\{14,19,20,22,23\}$	9	0101001000010 1000011011000
s_{10}	$\{2,4,7\}$	$\{14,16,17,19,20,22,23,25\}$	11	0101001000000 1011011011010
s_{11}	$\{2,4,9,13\}$	$\{14,15,17,19,25\}$	9	0101000010001 1101010000010
s_{12}	$\{2,4,9\}$	$\{14,15,17,19,20,21,22,25\}$	11	0101000010000 1101011110010
s_{13}	$\{2,6,7,10\}$	$\{14,16,20,24,25\}$	9	0100011001000 1010001000110
s_{14}	$\{2,6,7\}$	$\{14,16,18,19,20,23,24,25\}$	11	0100011000000 1010111001110
s_{15}	$\{2,6,9,11\}$	$\{18,19,20,21,25\}$	9	0100010010100 0000111100010
s_{16}	$\{2,6,9\}$	$\{14,15,18,19,20,21,24,25\}$	11	0100010010000 1100111100110
s_{17}	$\{3,4,8,13\}$	$\{15,16,17,19,23\}$	9	0011000100001 0111010001000
s_{18}	$\{3,4,8\}$	$\{15,16,17,19,20,21,22,23\}$	11	0011000100000 0111011111000
s_{19}	$\{3,5,7,10\}$	$\{16,17,20,22,24\}$	9	0010101001000 0011001010100
s_{20}	$\{3,5,7\}$	$\{16,17,18,19,20,22,23,24\}$	11	0010101000000 0011111011100
s_{21}	$\{3,5,9,13\}$	$\{15,17,18,19,24\}$	9	0010100010001 0101110000100
s_{22}	$\{3,5,9\}$	$\{15,17,18,19,20,21,22,24\}$	11	0010100010000 0101111101000
s_{23}	$\{3,6,8,10\}$	$\{15,16,20,21,24\}$	9	0010010101000 0110001100100
s_{24}	$\{3,6,8\}$	$\{15,16,18,19,20,21,23,24\}$	11	0010010100000 0110111011000
s_{25}	$\emptyset$	$\{26\}$	1	0000000000000 0000000000001

states $(s_{1,0}, s_{1,1}, \ldots, s_{24,0}, s_{24,1})$, each '1' in row i of Table X expands to '(1,1)' for core vertices $i \in \{1,\ldots,17\}$, to '(1,0)' for first completion vertices $w_{m,1} = 2m{+}16$, and to '(0,1)' for second completion vertices $w_{m,2} = 2m+17$ (and each '0' expands to '(0,0)'), making all 65 canonical 48-bit vertex profiles nonzero and pairwise distinct by inspection.

Appendix C: Reproducible completion rule and exact verification

The coordinates in Eq. (12) can be reproduced by the following finite selection rule. Normalize an integer vector by dividing by the greatest common divisor of its entries and choosing the sign so that its first nonzero entry is positive. Form the set of normalized nonzero vectors obtained from $\{-3,-2,\ldots,3\}^4$, ordered by squared norm, then maximum absolute entry, then lexicographically. Process bridge edges in the order $e_{13},\ldots,e_{24},e_1,\ldots,e_{12}$. For each pair, choose η_A to be the first vector in this ordered pool orthogonal to its endpoints, and η_B the first such vector also orthogonal to η_A.

For all $(a,b) \in \{-7,-6,\ldots,7\}^2 \setminus \{(0,0)\}$, form the pair in Eq. (10) and normalize both vectors. Discard pairs that repeat an already placed ray or have an orthogonality to any already placed vertex other than the two endpoints. Among the remaining ordered pairs choose the lexicographically smallest score

$$\left(\max_{j,k} |(w_{m,j})_k|,\ \|w_{m,1}\|^2 + \|w_{m,2}\|^2,\ w_{m,1}, w_{m,2} \right). \tag{C1}$$

In Eq. (C1), the order on the last two entries is the usual lexicographic order on their integer coordinates. This rule yields exactly the 48 completion vectors in Table IV. It is a recipe for this specific example; the general existence assertion is established independently by Lemma 3.

ACKNOWLEDGMENTS

This research was funded in part by the Austrian Science Fund (FWF), Grant DOI 10.55776/PIN5424624, and the Czech Science Foundation (GAČR), Grant No. 25-20013L. The author acknowledges FWF and TU Wien Bibliothek for financial support through its Open Access Funding Programme, as well as support from Google via Gemini 4 Argon High and contributions from SOL 6.2 xhigh and Sonnet 5.5 high.

TABLE VIII. Vertex state profiles across the 25 two-valued states $s_1, \ldots, s_{25}$ of $H_{\mathrm{YO}}^{(4)}$, showing for each vertex $i \in \{1, \ldots, 26\}$ its state weight $\sum_{r=1}^{25} s_r(i)$, the active state indices $\{r : s_r(i) = 1\}$, and the 25-bit profile $(s_{1\ldots12}(i)\, s_{13\ldots24}(i)\, s_{25}(i))$. All 26 rows are nonzero and pairwise distinct.

Vertex i	Weight $\sum_r s_r(i)$	Active states $\{r \in \{1, \ldots, 25\} : s_r(i) = 1\}$	25-bit profile $(s_{1\ldots12}(i)\, s_{13\ldots24}(i)\, s_{25}(i))$
1	8	$\{1, 2, 3, 4, 5, 6, 7, 8\}$	111111110000 000000000000 0
2	8	$\{9, 10, 11, 12, 13, 14, 15, 16\}$	000000001111 111100000000 0
3	8	$\{17, 18, 19, 20, 21, 22, 23, 24\}$	000000000000 000011111111 0
4	8	$\{1, 2, 9, 10, 11, 12, 17, 18\}$	110000001111 000011000000 0
5	8	$\{3, 4, 5, 6, 19, 20, 21, 22\}$	001111000000 000000111100 0
6	8	$\{7, 8, 13, 14, 15, 16, 23, 24\}$	000000110000 111100000011 0
7	8	$\{3, 4, 9, 10, 13, 14, 19, 20\}$	001100001100 110000110000 0
8	8	$\{1, 2, 7, 8, 17, 18, 23, 24\}$	110000110000 000011000011 0
9	8	$\{5, 6, 11, 12, 15, 16, 21, 22\}$	000011000011 001100001100 0
10	3	$\{13, 19, 23\}$	000000000000 100000100010 0
11	3	$\{5, 7, 15\}$	000010100000 001000000000 0
12	3	$\{1, 3, 9\}$	101000001000 000000000000 0
13	3	$\{11, 17, 21\}$	000000000010 000010001000 0
14	13	$\{1, 2, 3, 4, 6, 8, 9, 10, 11, 12, 13, 14, 16\}$	111101011111 110100000000 0
15	13	$\{1, 2, 6, 8, 11, 12, 16, 17, 18, 21, 22, 23, 24\}$	110001010011 000111001111 0
16	13	$\{2, 4, 7, 8, 10, 13, 14, 17, 18, 19, 20, 23, 24\}$	010100110100 110011110011 0
17	13	$\{2, 4, 5, 6, 10, 11, 12, 17, 18, 19, 20, 21, 22\}$	010111000111 000011111100 0
18	13	$\{3, 4, 5, 6, 7, 8, 14, 15, 16, 20, 21, 22, 24\}$	001111110000 011100011101 0
19	13	$\{9, 10, 11, 12, 14, 15, 16, 17, 18, 20, 21, 22, 24\}$	000000001111 011111011101 0
20	13	$\{9, 10, 12, 13, 14, 15, 16, 18, 19, 20, 22, 23, 24\}$	000000001101 111101110111 0
21	13	$\{1, 2, 5, 6, 7, 8, 12, 15, 16, 18, 22, 23, 24\}$	110011110001 001101000111 0
22	13	$\{1, 2, 3, 4, 5, 6, 9, 10, 12, 18, 19, 20, 22\}$	111111001101 000001110100 0
23	13	$\{1, 2, 3, 4, 7, 8, 9, 10, 14, 17, 18, 20, 24\}$	111100111100 010011010001 0
24	13	$\{3, 4, 6, 8, 13, 14, 16, 19, 20, 21, 22, 23, 24\}$	001101010000 110100111111 0
25	13	$\{2, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16\}$	010111110111 111100000000 0
26	1	$\{25\}$	000000000000 000000000000 1

[1] K. Svozil and J. Tkadlec, Greechie diagrams, nonexistence of measures in quantum logics and Kochen–Specker type constructions, Journal of Mathematical Physics **37**, 5380 (1996).

[2] M. H. Shekarriz and K. Svozil, Noncontextual coloring of orthogonality hypergraphs, Journal of Mathematical Physics **63**, 032104 (2022), arXiv:2105.08520.

[3] A. Peres, Two simple proofs of the Kochen-Specker theorem, Journal of Physics A: Mathematical and General **24**, L175 (1991).

[4] S. Yu and C. H. Oh, State-independent proof of Kochen-Specker theorem with 13 rays, Physical Review Letters **108**, 030402 (2012), arXiv:1109.4396.

[5] M. Pavičić, Hypergraph contextuality, Entropy **21**, 1107 (2019).

[6] K. Svozil, Chromatic quantum contextuality, Entropy **27**, 387 (2025), arXiv:2501.15261.

[7] G. Haynes, C. Park, A. Schaeffer, J. Webster, and L. H. Mitchell, Orthogonal vector coloring, The Electronic Journal of Combinatorics **17**, R55 (2010).

TABLE IX. Complete branch-by-branch parametrization of the $24 \times 2^{14} = 393{,}216$ two-valued states $s_{\alpha,\boldsymbol{\varepsilon}}$ and the 48 canonical states $s_{\alpha,0}, s_{\alpha,1}$ on H_{65} via Eqs. (B3)–(B4). For each $\alpha \in \{1, \ldots, 24\}$, the table lists the 4-element core support S_α, the 10 hit completion context indices $m \in \mathcal{H}_\alpha$ where $(s(2m+16), s(2m+17)) = (0,0)$ is forced on E_m, the 14 free completion context indices $m \in \mathcal{F}_\alpha$ where $(s(2m+16), s(2m+17)) = (1-\varepsilon_m, \varepsilon_m)$ with $\varepsilon_m \in \{0,1\}$ on E_m, and the 24-bit indicator string $(\mathbf{1}_{1\in\mathcal{F}_\alpha} \ldots \mathbf{1}_{12\in\mathcal{F}_\alpha}\, \mathbf{1}_{13\in\mathcal{F}_\alpha} \ldots \mathbf{1}_{24\in\mathcal{F}_\alpha})$.

Branch	Core support S_α	Hit contexts $\mathcal{H}_\alpha$ (forced $(0,0)$)	Free contexts $\mathcal{F}_\alpha$ (2^{14} choices)	Free indicator $m = 1 \ldots 24$
S_1	$\{1,5,9,16\}$	$\{3,7,9,10,11,13,19,21,22,23\}$	$\{1,2,4,5,6,8,12,14,15,16,17,18,20,24\}$	110111010001 011111010001
S_2	$\{1,5,10,14\}$	$\{3,5,6,7,11,13,17,18,19,23\}$	$\{1,2,4,8,9,10,12,14,15,16,20,21,22,24\}$	110100011101 011100011101
S_3	$\{1,6,9,16\}$	$\{3,7,9,10,11,15,17,21,22,24\}$	$\{1,2,4,5,6,8,12,13,14,16,18,19,20,23\}$	110111010001 110101110010
S_4	$\{1,6,11,12\}$	$\{1,2,3,7,11,13,14,15,17,24\}$	$\{4,5,6,8,9,10,12,16,18,19,20,21,22,23\}$	000111011101 000101111110
S_5	$\{1,7,10,14\}$	$\{3,5,6,7,11,16,17,18,20,21\}$	$\{1,2,4,8,9,10,12,13,14,15,19,22,23,24\}$	110100011101 111000100111
S_6	$\{1,7,11,12\}$	$\{1,2,3,7,11,13,14,16,20,21\}$	$\{4,5,6,8,9,10,12,15,17,18,19,22,23,24\}$	000111011101 001011100111
S_7	$\{2,5,9,16\}$	$\{1,5,9,10,12,13,19,21,22,23\}$	$\{2,3,4,6,7,8,11,14,15,16,17,18,20,24\}$	011101110010 011111010001
S_8	$\{2,5,11,13\}$	$\{1,3,4,5,12,13,15,16,19,23\}$	$\{2,6,7,8,9,10,11,14,17,18,20,21,22,24\}$	010001111110 010011011101
S_9	$\{2,6,9,16\}$	$\{1,5,9,10,12,15,17,21,22,24\}$	$\{2,3,4,6,7,8,11,13,14,16,18,19,20,23\}$	011101110010 110101110010
S_{10}	$\{2,6,10,15\}$	$\{1,5,7,8,12,15,17,19,20,24\}$	$\{2,3,4,6,9,10,11,13,14,16,18,21,22,23\}$	011101001110 110101001110
S_{11}	$\{2,8,10,15\}$	$\{1,5,7,8,12,14,18,19,20,22\}$	$\{2,3,4,6,9,10,11,13,15,16,17,21,23,24\}$	011101001110 101110001011
S_{12}	$\{2,8,11,13\}$	$\{1,3,4,5,12,14,15,16,18,22\}$	$\{2,6,7,8,9,10,11,13,17,19,20,21,23,24\}$	010001111110 100010111011
S_{13}	$\{3,5,10,14\}$	$\{2,5,6,8,9,13,17,18,19,23\}$	$\{1,3,4,7,10,11,12,14,15,16,20,21,22,24\}$	101100100111 011100011101
S_{14}	$\{3,5,11,13\}$	$\{2,3,4,8,9,13,15,16,19,23\}$	$\{1,5,6,7,10,11,12,14,17,18,20,21,22,24\}$	100011100111 010011011101
S_{15}	$\{3,7,9,17\}$	$\{2,8,9,11,12,16,20,21,23,24\}$	$\{1,3,4,5,6,7,10,13,14,15,17,18,19,22\}$	101111100100 111011100100
S_{16}	$\{3,7,10,14\}$	$\{2,5,6,8,9,16,17,18,20,21\}$	$\{1,3,4,7,10,11,12,13,14,15,19,22,23,24\}$	101100100111 111000100111
S_{17}	$\{3,8,9,17\}$	$\{2,8,9,11,12,14,18,22,23,24\}$	$\{1,3,4,5,6,7,10,13,15,16,17,19,20,21\}$	101111100100 101110111000
S_{18}	$\{3,8,11,13\}$	$\{2,3,4,8,9,14,15,16,18,22\}$	$\{1,5,6,7,10,11,12,13,17,19,20,21,23,24\}$	100011100111 100010111011
S_{19}	$\{4,6,10,15\}$	$\{4,6,7,8,10,15,17,19,20,24\}$	$\{1,2,3,5,9,11,12,13,14,16,18,21,22,23\}$	111010001011 110101001110
S_{20}	$\{4,6,11,12\}$	$\{1,2,4,6,10,13,14,15,17,24\}$	$\{3,5,7,8,9,11,12,16,18,19,20,21,22,23\}$	001010111011 000101111110
S_{21}	$\{4,7,9,17\}$	$\{4,6,10,11,12,16,20,21,23,24\}$	$\{1,2,3,5,7,8,9,13,14,15,17,18,19,22\}$	111010111000 111011100100
S_{22}	$\{4,7,11,12\}$	$\{1,2,4,6,10,13,14,16,20,21\}$	$\{3,5,7,8,9,11,12,15,17,18,19,22,23,24\}$	001010111011 001011100111
S_{23}	$\{4,8,9,17\}$	$\{4,6,10,11,12,14,18,22,23,24\}$	$\{1,2,3,5,7,8,9,13,15,16,17,19,20,21\}$	111010111000 101110111000
S_{24}	$\{4,8,10,15\}$	$\{4,6,7,8,10,14,18,19,20,22\}$	$\{1,2,3,5,9,11,12,13,15,16,17,21,23,24\}$	111010001011 101110001011

TABLE X. Vertex state profiles and weights for all 65 vertices of H_{65}, showing for each core vertex $i \in \{1,\dots,17\}$ and each completion pair $(w_{m,1}, w_{m,2}) = (2m+16, 2m+17)$ $(m = 1,\dots,24)$ the number of active branches W_{24}, the weight $W_{48} = \sum_{\alpha,\varepsilon} s_{\alpha,\varepsilon}(i)$ across the 48 canonical states, the weight $W_{393,216} = \sum_{s\in\mathcal{S}(H_{65})} s(i)$ across all 393,216 states, the active branch indices $\alpha \in \{1,\dots,24\}$ (equal to $Z(e_m)$ for $(w_{m,1}, w_{m,2})$), and the 24-bit branch profile across $S_1,\dots,S_{24}$ (in which each `1` expands to `11` for core vertices i, to `10` for $w_{m,1}$, and to `01` for $w_{m,2}$ across the 48 canonical states).

Vertex / completion pair	W_{24}	W_{48}	$W_{393,216}$	Active branches $\alpha \in \{1,\dots,24\}$	24-bit branch profile ($1\dots8 \mid 9\dots16 \mid 17\dots24$)
Core 1	6	12	98,304	$\{1,2,3,4,5,6\}$	11111100 00000000 00000000
Core 2	6	12	98,304	$\{7,8,9,10,11,12\}$	00000011 11110000 00000000
Core 3	6	12	98,304	$\{13,14,15,16,17,18\}$	00000000 00001111 11000000
Core 4	6	12	98,304	$\{19,20,21,22,23,24\}$	00000000 00000000 00111111
Core 5	6	12	98,304	$\{1,2,7,8,13,14\}$	11000011 00001100 00000000
Core 6	6	12	98,304	$\{3,4,9,10,19,20\}$	00110000 11000000 00110000
Core 7	6	12	98,304	$\{5,6,15,16,21,22\}$	00001100 00000011 00001100
Core 8	6	12	98,304	$\{11,12,17,18,23,24\}$	00000000 00110000 11000011
Core 9	8	16	131,072	$\{1,3,7,9,15,17,21,23\}$	10100010 10000010 10001010
Core 10	8	16	131,072	$\{2,5,10,11,13,16,19,24\}$	01001000 01101001 00100001
Core 11	8	16	131,072	$\{4,6,8,12,14,18,20,22\}$	00010101 00010100 01010100
Core 12	4	8	65,536	$\{4,6,20,22\}$	00010100 00000000 00010100
Core 13	4	8	65,536	$\{8,12,14,18\}$	00000001 00010100 01000000
Core 14	4	8	65,536	$\{2,5,13,16\}$	01001000 00001001 00000000
Core 15	4	8	65,536	$\{10,11,19,24\}$	00000000 01100000 00100001
Core 16	4	8	65,536	$\{1,3,7,9\}$	10100010 10000000 00000000
Core 17	4	8	65,536	$\{15,17,21,23\}$	00000000 00000010 10001010
E_1 : (18, 19)	14	14	114,688	$\{1,2,3,5,13,14,15,16,17,18,19,21,23,24\}$	11101000 00001111 11101011
E_2 : (20, 21)	14	14	114,688	$\{1,2,3,5,7,8,9,10,11,12,19,21,23,24\}$	11101011 11110000 00101011
E_3 : (22, 23)	14	14	114,688	$\{7,9,10,11,13,15,16,17,19,20,21,22,23,24\}$	00000010 11101011 10111111
E_4 : (24, 25)	14	14	114,688	$\{1,2,3,4,5,6,7,9,10,11,13,15,16,17\}$	11111110 11101011 10000000
E_5 : (26, 27)	14	14	114,688	$\{1,3,4,6,14,15,17,18,19,20,21,22,23,24\}$	10110100 00000110 11111111
E_6 : (28, 29)	14	14	114,688	$\{1,3,4,6,7,8,9,10,11,12,14,15,17,18\}$	10110111 11110110 11000000
E_7 : (30, 31)	14	14	114,688	$\{7,8,9,12,13,14,15,16,17,18,20,21,22,23\}$	00000011 10011111 11011110
E_8 : (32, 33)	14	14	114,688	$\{1,2,3,4,5,6,7,8,9,12,20,21,22,23\}$	11111111 10010000 00011110
E_9 : (34, 35)	14	14	114,688	$\{2,4,5,6,8,10,11,12,19,20,21,22,23,24\}$	01011101 01110000 00111111
E_{10} : (36, 37)	14	14	114,688	$\{2,4,5,6,8,10,11,12,13,14,15,16,17,18\}$	01011101 01111111 11000000
E_{11} : (38, 39)	14	14	114,688	$\{7,8,9,10,11,12,13,14,16,18,19,20,22,24\}$	00000011 11111101 01110101
E_{12} : (40, 41)	14	14	114,688	$\{1,2,3,4,5,6,13,14,16,18,19,20,22,24\}$	11111100 00001101 01110101
E_{13} : (42, 43)	14	14	114,688	$\{3,5,9,10,11,12,15,16,17,18,19,21,23,24\}$	00101000 11110011 11101011
E_{14} : (44, 45)	14	14	114,688	$\{1,2,3,5,7,8,9,10,13,14,15,16,19,21\}$	11101011 11001111 00101000
E_{15} : (46, 47)	14	14	114,688	$\{1,2,5,6,7,11,13,15,16,17,21,22,23,24\}$	11001110 00101011 10001111
E_{16} : (48, 49)	14	14	114,688	$\{1,2,3,4,7,9,10,11,13,17,19,20,23,24\}$	11110010 11101000 10110011
E_{17} : (50, 51)	14	14	114,688	$\{1,6,7,8,11,12,14,15,17,18,21,22,23,24\}$	10000111 00110110 11001111
E_{18} : (52, 53)	14	14	114,688	$\{1,3,4,6,7,8,9,10,14,15,19,20,21,22\}$	10110111 11000110 00111100
E_{19} : (54, 55)	14	14	114,688	$\{3,4,5,6,9,12,15,16,17,18,20,21,22,23\}$	00111100 10010011 11011110
E_{20} : (56, 57)	14	14	114,688	$\{1,2,3,4,7,8,9,12,13,14,17,18,20,23\}$	11110011 10011100 11010010
E_{21} : (58, 59)	14	14	114,688	$\{2,4,8,10,11,12,13,14,17,18,19,20,23,24\}$	01010001 01111100 11110011
E_{22} : (60, 61)	14	14	114,688	$\{2,4,5,6,8,10,13,14,15,16,19,20,21,22\}$	01011101 01001111 00111100
E_{23} : (62, 63)	14	14	114,688	$\{3,4,5,6,9,10,11,12,16,18,19,20,22,24\}$	00111100 11110001 01110101
E_{24} : (64, 65)	14	14	114,688	$\{1,2,5,6,7,8,11,12,13,14,16,18,22,24\}$	11001111 00111101 01000101