

Archimedean Resolution of the Bondi Runaway: Holes in a Background Substratum Gravitates Repulsively without Paradox

Karl Svozil ^{1,*}

¹*Institute for Theoretical Physics, TU Wien, Wiedner Hauptstrasse 8-10/136, 1040 Vienna, Austria*
(Dated: April 17, 2026)

A localized absence of substratum in a positive-energy background acts, to leading order, as an effective negative-mass source whose pair dynamics with ordinary matter evades the Bondi runaway. A hole near a positive mass M_+ is pushed away from M_+ by Archimedean buoyancy in the substratum's infall, while M_+ is simultaneously repelled by the hole's negative effective source; both accelerations are aligned in the sense of mutual separation. No fundamental field violates the energy conditions: the apparent negativity is a background-subtraction artifact, structurally identical to Casimir energy, band-hole effective mass, and the peculiar gravity of cosmic voids.

Introduction.—The negative-mass Schwarzschild solution has been regarded since Bondi [1] as physically unrealizable. Treated as an autonomous point particle with intrinsic negative inertia, a negative mass placed near a positive one produces a runaway: the pair accelerates indefinitely in a common direction, conserving energy and momentum but violating dynamical stability. The standard energy conditions [2, 3] forbid fundamental matter with $\rho < 0$, reinforcing the view that repulsive solutions of the Einstein equations are mathematical curiosities.

We show that this conclusion does not survive once space is modeled as a *background substratum*: a positive-energy-density medium $\rho_0 > 0$ whose local absence defines what we call a hole. The picture is an old one, going back to Einstein's 1920 reinterpretation of the gravitational ether [4] and to Sakharov's induced gravity [5], and it has a directly observed realization in cosmic voids [6–8]. What is new here is a mechanical observation: in such a picture, the Bondi runaway is replaced by Archimedean mutual repulsion. The negative-mass degree of freedom is not autonomous; it is a buoyant displacement whose dynamics are set by the medium, and the medium's own gravitational response to a nearby positive mass drives the hole away from, not toward, that mass.

Setup.—The substratum stress-energy $T_{\mu\nu}^{(0)}$ with $\rho_0 > 0$ is assumed to fill space outside the hole. For a Lorentz scalar, $T_{\mu\nu}^{(0)} = -\rho_0 g_{\mu\nu}$ ($p_0 = -\rho_0$, equivalent to a cosmological constant $\Lambda = 8\pi G\rho_0/c^2$); for a pressureless dust, $T_{\mu\nu}^{(0)} = \rho_0 u_\mu u_\nu$. Both choices support the central mechanism; the dust realization is empirically instantiated by cold dark matter on cosmological scales.

The hole occupies a bounded region $\Omega \subset \mathbb{R}^3$ in which $T_{\mu\nu} = 0$. Only deviations from the uniform background source observable gravity; we define the background-subtracted effective stress-energy

$$T_{\mu\nu}^{\text{eff}} = T_{\mu\nu} - T_{\mu\nu}^{(0)}, \quad (1)$$

which vanishes outside Ω and equals $-T_{\mu\nu}^{(0)}$ inside. The underlying field is positive everywhere; the negativity of $T_{\mu\nu}^{\text{eff}}$ in Ω is a renormalization, not a fundamental violation of the energy conditions.

Newtonian and linearized limit.—In the weak-field limit, the shell theorem applied to $\rho^{\text{eff}} = -\rho_0 \chi_\Omega$ yields a potential

$$\Phi(\mathbf{r}) = \frac{G\rho_0 V(\Omega)}{r} + \mathcal{O}(\mathcal{R}^2/r^3) \quad (2)$$

outside the bounding sphere $r = \mathcal{R}$ of Ω , where $V(\Omega)$ is its Lebesgue measure. The leading field is positive and repels test bodies with an effective source

$$M_{\text{eff}} = -\rho_0 V(\Omega). \quad (3)$$

Equation (3) depends only on the volume of the absence: the shape of Ω —spherical cavity, cubic cavity, foam, or fractal—enters only through subleading multipoles suppressed by $(\mathcal{R}/r)^2$. This shape-universality is exact in the linearized regime.

Exact exterior geometry.—For a spherical hole in a Lorentz-scalar substratum, Birkhoff's theorem applies only to the source-free interior, which is therefore Minkowski. The exterior is not vacuum: it carries $T_{\mu\nu}^{(0)} \propto g_{\mu\nu}$, and the correct spherically symmetric solution is the Schwarzschild–de Sitter (Kottler) metric [9],

$$F(r) = 1 - \frac{2GM_{\text{ext}}}{c^2 r} - \frac{\Lambda r^2}{3}. \quad (4)$$

Matching Minkowski interior to SdS exterior at $r = \mathcal{R}$ requires, by the Israel junction conditions [10, 11], a thin boundary shell whose surface tension balances the bulk pressure jump $\Delta p = \rho_0$. The shell carries positive surface energy and ordinary tensile stress—not exotic matter. The asymptotic mass parameter is

$$M_{\text{ext}} = -\frac{4\pi}{3}\rho_0 \mathcal{R}^3 + M_{\text{shell}}, \quad (5)$$

and in the sub-cosmological regime $\Lambda \mathcal{R}^2 \ll 1$ the bulk deficit dominates, $M_{\text{ext}} < 0$, in agreement with Eq. (3). Radial timelike geodesics in $F(r)$ accelerate outward: holes repel. For a dust substratum, the exterior is FLRW and the hole is a Lemaître–Tolman–Bondi void [12, 13]; no shell is needed and the boundary evolves smoothly.

Archimedean dynamics.—We now turn to the main claim: the pair dynamics of a hole and a nearby positive mass M_+ does not exhibit the Bondi runaway. Let $\hat{\mathbf{n}}$ denote the unit vector from the hole’s centroid toward M_+ , and let r be their separation.

Two effects operate simultaneously (Fig. 1). The positive mass, treated as a test body in the hole’s exterior field Eq. (4), experiences a gravitational acceleration directed away from the hole,

$$\mathbf{a}_{M_+} = +\frac{G|M_{\text{ext}}|}{r^2} \hat{\mathbf{n}}, \quad (6)$$

with the $+\hat{\mathbf{n}}$ sign inherited from $M_{\text{ext}} < 0$. Independently, the substratum surrounding the hole is itself pulled toward M_+ by ordinary gravitational attraction, acquiring an acceleration $\mathbf{g}_{\text{at hole}}(M_+) = +(GM_+/r^2)\hat{\mathbf{n}}$ at the hole’s location. The hole, being the *absence* of substratum within a flowing medium, is Archimedean: its acceleration is opposite to that of the displaced fluid,

$$\mathbf{a}_{\text{hole}} = -\mathbf{g}_{\text{at hole}}(M_+) = -\frac{GM_+}{r^2} \hat{\mathbf{n}}. \quad (7)$$

The positive mass accelerates along $+\hat{\mathbf{n}}$ (away from the hole), the hole along $-\hat{\mathbf{n}}$ (away from M_+). Both objects accelerate *away from each other*. The pair separates; there is no runaway.

Two features distinguish this from Bondi’s analysis. First, the hole has no intrinsic inertia. Its dynamical response is inherited entirely from the surrounding substratum and follows from Archimedes’ principle applied to a non-uniform acceleration field. The paradoxical “chasing” behavior that Bondi derived requires treating the negative-mass object as an autonomous Newtonian particle with $m < 0$; the hole is not such an object. Second, the sign of $\mathbf{a}_{\text{hole}}$ is fixed by fluid kinematics, not by M_{ext} : it would be the same for any substratum flow directed toward M_+ , regardless of what the hole’s effective gravitational charge might be. Momentum conservation is respected, with the missing momentum supplied by the

substratum’s gravitational response to M_+ ; the medium, not the hole, is the reservoir.

The same mechanism governs well-known non-gravitational analogues. A semiconductor hole subjected to an electric field accelerates with the sign appropriate to negative effective mass, but no runaway instability arises because the underlying electrons, all of positive mass, obey ordinary dynamics within the adiabatic band [14, 15]. Dirac’s hole-theoretic positron [16] is the relativistic analogue of the same construction. The present result is the gravitational analogue.

Energy conditions.—The fundamental stress-energy $T_{\mu\nu}$ is nowhere negative. Inside the hole $T_{\mu\nu} = 0$, outside $T_{\mu\nu} = T_{\mu\nu}^{(0)}$ with $\rho_0 > 0$. The Lorentz-scalar realization has $p_0 = -\rho_0$ and thereby violates the strong energy condition—but in exactly the same sense and to the same degree as the cosmological-constant component driving the observed cosmic acceleration [17, 18], a violation universally accepted as empirically required. The dust realization satisfies all standard energy conditions. Negativity enters only in the bookkeeping object $T_{\mu\nu}^{\text{eff}}$ of Eq. (1), constructed to isolate the local gravitational effect of the deviation from baseline. This is the same structural move that renders the Casimir energy negative [19, 20], the band hole’s inertial mass effectively negative [14], and the peculiar gravity of a cosmic void repulsive [6, 7]. In none of these cases does the negativity of the subtracted quantity signal a pathology of the underlying theory.

Discussion.—The result is narrow but sharp. Within a background-substratum framework, gravitational repulsion between an underdense region and ordinary matter is generic, the Bondi pair-runaway objection does not apply, and the standard energy conditions of the fundamental field are preserved. What the framework does *not* establish is the existence of such a substratum on subcosmological scales, which remains a hypothesis. The closest empirical realization is cosmic void cosmology, in which the dust-like substratum is simply cold dark matter and the repulsive peculiar gravity of voids relative to the cosmic mean is an observed phenomenon, measured through redshift-space distortions in the void-galaxy cross-correlation and other probes [7, 8].

The Archimedean mechanism suggests where to look for a compact-object realization. A candidate signature is a localized object of negative gravitational charge whose trajectory tracks, at leading order, the negative gradient of the Newtonian potential produced by surrounding matter—opposite to the trajectory of an ordinary self-gravitating body. For such an object, lensing is defocusing, Shapiro delays become advances, and orbital dynamics are inverted. None has been detected at subcosmological scales, and the framework presented here is consistent with that null result: it predicts only that *if* a ponderable substratum exists and *if* it admits localized




FIG. 1. Archimedean repulsion. The substratum (wavy band) flows gravitationally toward M_+ with $\hat{\mathbf{n}}$ the unit vector from hole to M_+ . The hole, as a buoyant absence in this flow, is displaced in the $-\hat{\mathbf{n}}$ direction, $\mathbf{a}_{\text{hole}}$ [Eq. (7)]. Simultaneously, M_+ is gravitationally repelled along $+\hat{\mathbf{n}}$ by the hole’s effective negative source, $\mathbf{a}_{M_+}$ [Eq. (6)]. Both accelerations point away from the other body; the pair separates rather than running away in a common direction.

absences on astrophysical scales, such signatures should appear. The framework moves the question from “can negative-mass sources exist at all” to “does a substratum exist in which they can be realized as absences,” which is an empirical question we do not claim to settle.

Conclusion.—The Bondi runaway is an artifact of treating a negative-mass source as an autonomous dynamical particle. In a substratum picture, the negative effective source is a buoyant absence with no intrinsic inertia, and its dynamics, set by the surrounding medium, produce mutual repulsion with any nearby positive mass rather than runaway pursuit. The result requires neither exotic fundamental matter nor violation of the underlying energy conditions, and it places gravitational repulsion conceptually alongside Casimir energy, band holes, and void peculiar gravity—familiar, unexotic consequences of background subtraction in a positive-energy medium.

The author thanks an anonymous referee of an earlier version for incisive criticism concerning the non-vacuum exterior, the Israel junction conditions, the substratum equation of state, and the Archimedean mechanism itself. This text was partially created and revised with assistance from large language models; all content, ideas, and prompts were provided by the author. This research was funded in whole or in part by the Austrian Science Fund (FWF) Grant DOI: 10.55776/PIN5424624. The author acknowledges TU Wien Bibliothek for financial support through its Open Access Funding Programme.

* karl.svozil@tuwien.ac.at

- [1] H. Bondi, Negative mass in general relativity, *Reviews of Modern Physics* **29**, 423 (1957).
- [2] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, 1973).
- [3] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).
- [4] A. Einstein, Ether and the theory of relativity, in *The Genesis of General Relativity. Sources and Interpretations*, Boston Studies in the Philosophy and History of Science, Vol. 250 (Springer, Dordrecht, The Netherlands, Netherlands, 2007) pp. 1537–1542.
- [5] A. D. Sakharov, Vacuum quantum fluctuations in curved space and the theory of gravitation, *General Relativity and Gravitation* **32**, 365 (2000), translated from *Doklady Akademii Nauk SSSR*, vol. 177, No. 1, pp. 70–71, November 1967.
- [6] R. K. Sheth and R. van de Weygaert, A hierarchy of voids: Much ado about nothing, *Monthly Notices of the Royal Astronomical Society* **350**, 517 (2004), arXiv:astro-ph/0311260.
- [7] N. Hamaus, P. M. Sutter, and B. D. Wandelt, Universal density profile for cosmic voids, *Physical Review Letters* **112**, 251302 (2014), arXiv:1403.5499.
- [8] S. Nadathur, P. M. Carter, W. J. Percival, H. A. Winther, and J. E. Bautista, Beyond BAO: Improving cosmological constraints from BOSS data with measurement of the void-galaxy cross-correlation, *Physical Review D* **100**, 023504 (2019), arXiv:1904.01030.
- [9] F. Kottler, Über die physikalischen Grundlagen der Einsteinschen Gravitationstheorie, *Annalen der Physik* **361**, 401 (1918).
- [10] W. Israel, Singular hypersurfaces and thin shells in general relativity, *Il Nuovo Cimento B* **44**, 1 (1966), erratum: *Nuovo Cimento B* **48**, 463 (1967), DOI: 10.1007/BF02712210.
- [11] E. Poisson, *A Relativist’s Toolkit: The Mathematics of Black-Hole Mechanics* (Cambridge University Press, Cambridge, 2004).
- [12] R. C. Tolman, Effect of inhomogeneity on cosmological models, *Proceedings of the National Academy of Sciences of the USA* **20**, 169 (1934).
- [13] H. Bondi, Spherically symmetrical models in general relativity, *Monthly Notices of the Royal Astronomical Society* **107**, 410 (1947), reprinted in *Gen. Relativ. Gravit.* **31**, 1783–1805 (1999), DOI: 10.1023/A:1026726520289.
- [14] C. Kittel, *Introduction to Solid State Physics*, 8th ed. (John Wiley & Sons, New York, 2005).
- [15] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Holt, Rinehart and Winston, New York, 1976).
- [16] P. A. M. Dirac, A theory of electrons and protons, *Proceedings of the Royal Society of London A* **126**, 360 (1930).
- [17] A. G. Riess, A. V. Filippenko, P. Challis, A. Clocchiatti, A. Diercks, P. M. Garnavich, R. L. Gilliland, C. J. Hogan, S. Jha, R. P. Kirshner, B. Leibundgut, M. M. Phillips, D. Reiss, B. P. Schmidt, R. A. Schommer, R. C. Smith, J. Spyromilio, C. Stubbs, N. B. Suntzeff, and J. Tonry, Observational evidence from supernovae for an accelerating universe and a cosmological constant, *The Astronomical Journal* **116**, 1009 (1998), arXiv:astro-ph/9805201.
- [18] S. Perlmutter, G. Aldering, G. Goldhaber, R. A. Knop, P. Nugent, P. G. Castro, S. Deustua, S. Fabbro, A. Goobar, D. E. Groom, I. M. Hook, A. G. Kim, M. Y. Kim, J. C. Lee, N. J. Nunes, R. Pain, C. R. Pennyacker, R. Quimby, C. Lidman, R. S. Ellis, M. Irwin, R. G. McMahon, P. Ruiz-Lapuente, N. Walton, B. Schaefer, B. J. Boyle, A. V. Filippenko, T. Matheson, A. S. Fruchter, N. Panagia, H. J. M. Newberg, and W. J. Couch, Measurements of Ω and Λ from 42 high-redshift supernovae, *The Astrophysical Journal* **517**, 565 (1999), arXiv:astro-ph/9812133.
- [19] H. B. G. Casimir, On the attraction between two perfectly conducting plates, *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* **51**, 793 (1948).
- [20] K. A. Milton, *The Casimir Effect: Physical Manifestations of Zero-Point Energy* (World Scientific, Singapore, 2001).