# Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments

Karl Svozil [a)]
*Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria*

(Dated: November 5, 2026)

Suppose intertwined complete experiments each measure a maximal observable with one common spectrum, and the eigenvalue of a shared outcome is an element of physical reality, the same in every experiment. Such noncontextual eigenvalues exist exactly when the context hypergraph can be colored with as many colors as the dimension. They can fail even when the two-valued states separate all outcomes, so that the logic is classically realizable; completed Yu–Oh is an example. When no such assignment exists, any classical model needs more hidden states than the dimension, just to tell apart preparations along orthogonal rays. Where noncontextual eigenvalues exist, they can force two distinct rays to share an eigenvalue, a chromatic form of Kochen–Specker inseparability; for one quantum preparation, every classical model must put a third of its weight on truth assignments that no eigenvalue assignment produces. We also correct an earlier inference from coloring to geometry.

## I. INTRODUCTION

An experimental context with $d$ rank-one outcomes admits a nondegenerate observable with $d$ distinct eigenvalues. These eigenvalues distinguish all $d$ outcomes and provide the maximal resolution of a complete projective experiment within that context. Such an observable is maximal in the sense that every self-adjoint observable commuting with it is a function of it, so the statistics of the latter can be obtained by processing its measurement outcomes. Noninjective functions merge outcomes and reduce this resolution. This is the finite-dimensional setting of von Neumann's functional representation of commuting observables[1] (Satz 8, p. 221); see also Halmos[2] (§84, pp. 171–172).

When complete experiments are intertwined, they share outcomes represented by the same rank-one projectors. We ask whether the eigenvalues of the maximal observables measured in such intertwined experiments can be assigned to their outcomes noncontextually, in the spirit of the elements of physical reality of Einstein, Podolsky, and Rosen[3]: whether every outcome carries a definite eigenvalue that belongs to the outcome itself, rather than to the experiment in which it is recorded. Three assumptions make this precise.

(i) *Maximality.* Each complete experiment measures a maximal observable, so its $d$ outcomes are never merged and carry $d$ distinct eigenvalues. A degenerate observable would discard part of what the experiment resolves.

(ii) *Uniformity.* All maximal observables share one spectrum of $d$ eigenvalues $\Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$. The eigenvalues are nothing but labels of the $d$ outcomes of a complete experiment. Since self-adjoint operators with a common nondegenerate spectrum are unitarily equivalent, uniformity says that every complete experiment measures one and the same quantity in a different orientation, $A_e = U_e A U_e^\dagger$.

(iii) *Noncontextuality of eigenvalues.* An outcome shared by several complete experiments carries the same eigenvalue in each of them, just as Kochen and Specker require a shared outcome to carry the same truth value[4].

[a)] Electronic mail: karl.svozil@tuwien.ac.at

No constraint is placed on how the $d$ eigenvalues are distributed over the outcomes of different contexts; any permutation within a context is allowed. A noncontextual eigenvalue assignment in this sense labels the outcomes, the most elementary record a complete experiment produces, and assumption (iii) treats these labels as elements of physical reality. For spin one, an example is the observable $aS_x^2+bS_y^2+cS_z^2$ with distinct real $a, b, c$ considered by Kochen and Specker[4]: every orthogonal triad of directions yields one such observable, with the same eigenvalues $b+c$, $a+c$, and $a+b$. A noncontextual eigenvalue assignment then gives each direction the same coefficient, $a$, $b$, or $c$, in every triad that contains it.

These assumptions concern eigenvalues rather than which outcome occurs, yet they have immediate consequences for truth assignments. A two-valued state selects one outcome in every context and retains only the distinction between that outcome and the others. A noncontextual eigenvalue assignment yields $d$ such states at once, one for each eigenvalue $\lambda_a$: "the outcome with eigenvalue $\lambda_a$ occurs in every experiment." These $d$ truth assignments have pairwise disjoint supports covering every outcome. A noncontextual eigenvalue assignment is thus a hidden-variable model whose hidden variable is the eigenvalue itself; its $d$ equally weighted values reproduce the maximally mixed state. Because this requirement is more demanding than Kochen–Specker noncontextuality, which asks for a single truth assignment, its failure—*chromatic contextuality*—is a weaker form of contextuality that already appears in configurations retaining many classical truth assignments. Even when a noncontextual eigenvalue assignment exists, its consistency across intertwined contexts can force relations between distinct outcomes, such as equal eigenvalues for nonorthogonal rays, that are absent from the full family of individual truth assignments. We study both the existence of noncontextual eigenvalue assignments and the event distinctions and logical relations that survive when such assignments are available[5–7].

To express these questions, let a $d$-uniform context hypergraph $H = (V, E)$ encode the complete experiments. An ordinary two-valued state $s : V \to \{0, 1\}$ selects exactly one outcome in every context, so $\sum_{v\in e} s(v) = 1$ for every $e \in E$. The Kochen–Specker (KS) theorem[4] establishes the nonexistence of such a global truth assignment for suitable finite families of intertwined projective contexts in dimension $d \geq 3$.

A strong $d$-coloring $c : V \to \{0, \ldots, d-1\}$ assigns all $d$ distinct labels bijectively within every context and independently of context at shared vertices. It is the formal counterpart of a noncontextual eigenvalue assignment: $c(i) = a$ means that outcome $i$ carries the eigenvalue $\lambda_a$ in every context containing it, so assumptions (i)–(iii) hold if and only if $\chi(H) \leq d$. For an injective ray realization $i \mapsto [v_i] \in \mathbb{P}(\mathbb{F}^d)$, with $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, rank-one projectors $P_i$, and a common nondegenerate real spectrum $\Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$, it defines the maximal context observables

$$A_e = \sum_{i\in e} \lambda_{c(i)} P_i. \tag{1}$$

Because the indicator functions $s_a(v) = \mathbf{1}_{\{c(v)=a\}}$ $(a \in \{0, \ldots, d-1\})$ of any strong $d$-coloring form $d$ two-valued states whose supports partition $V$, one has the strict one-way implication

$$\mathcal{S}(H) = \varnothing \quad \Longrightarrow \quad \chi(H) > d, \tag{2}$$

whose converse fails even when $\mathcal{S}(H)$ is unital and separates every pair of vertices. The earlier introduction in Ref.[6] inferred absence of global two-valued assignments from $\chi > d$. Equation (2) is the valid implication; completed Yu–Oh refutes the converse. Both the implication and the failure of its converse, for the original 13-ray Yu–Oh set, were noted by Cabello[8]. In terms of eigenvalues, if no noncontextual truth assignment exists, no noncontextual eigenvalue assignment exists either; but noncontextual eigenvalues can be impossible while truth assignments, even a unital separating family of them, survive. The distinct Appendix B inference about geometry is corrected below. We say that $H$ has an *exact palette*, or is *chromatically complete*, when $\chi(H) = d$, with no additional density or geometric assumption. This asymmetry separates chromatic contextuality into two distinct regimes that are unified in this work:

1. *Regime I, common-spectrum chromatic obstructions ($\chi(H) > d$, possibly with $\mathcal{S}(H) \neq \varnothing$):* Here no strong $d$-coloring exists ($\mathcal{C}_d(H) = \varnothing$), so the outcomes admit no noncontextual assignment of $d$ eigenvalues: every family of context observables with one common spectrum of $d$ eigenvalues must either merge outcomes, and so fail to be maximal, or let a shared outcome change its eigenvalue from context to context; maximal observables with noncontextual eigenvalues need more than $d$ eigenvalues. This happens even in examples in which classical truth assignments survive. A critical foundational clarification—correcting an interpretation in Appendix B of the earlier work with Shekarriz[6]—is that the chromatic obstruction $\chi(H) > d$ is an operator spectral-value obstruction and *does not* by itself obstruct geometric ray realizability in $\mathbb{F}^d$. Indeed, for finite 3-uniform linear hypergraphs of Berge girth five with unital separating state spaces, strong 3-colorability and injective $\mathbb{C}^3$ realizability are logically independent (Proposition 10). Ruling out geometric ray assignments for Greechie's Petersen-derived $G_{32}$[9] requires a direct Hermitian projection-identity argument (Theorem 11) rather than $\chi(G_{32}) = 4$.

2. *Regime II, exact-palette chromatic separability and logical gadgets ($\chi(H) = d$ with $\mathcal{S}_{\text{ext}}(H) \subsetneq \mathcal{S}(H)$):* In this regime, $\mathcal{C}_d(H) \neq \varnothing$, so noncontextual eigenvalue assignments exist, but some two-valued states $s \in \mathcal{S}(H)$ cannot be completed to a strong $d$-coloring: they are truth assignments that no noncontextual eigenvalue assignment produces. We characterize the extendable states $\mathcal{S}_{\text{ext}}(H)$ by the residual chromatic condition $\chi(\Gamma(H)[V \setminus \operatorname{supp} s]) \leq d-1$ (Theorem 2), which in dimension $d = 3$ yields an odd-cycle nonextendability certificate. In the specified subhypergraphs—such as the faithful real 17-ray and 21-ray restrictions $H_{17,10}$ and $H_{21,13}$ of completed Yu–Oh[7,10] (Theorem 15) and our faithful 3- and 4-dimensional logical gadgets (Secs. IV C and IV D)—ordinary states separate all vertices while every admissible $d$-coloring forces nonadjacent rays to share a color ($u \sim_c v$, chromatic inseparability / implicit identity[11]) or obeys purely chromatic true-implies-false (TIFS) or ternary exclusions violated by nonextendable states in $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$.

These two regimes are governed by a shared structural mechanism. Lemma 3 ($\chi(H) = d$ if and only if $H$ admits $d-1$ pairwise disjoint states) is the global unanchored form of Theorem 2. Likewise, the eight normalized 3-colorings of the four base contexts of completed Yu–Oh $H_{\text{YO}}^{+}$ partition into four antipodal pairs that each make one of the four diagonal neighborhoods rainbow (Theorem 13): retaining all four diagonal sectors forces $\chi(H_{\text{YO}}^{+}) = 4$ (Regime I), whereas retaining two or three sectors leaves four or two normalized 3-colorings and forces two or 21 chromatically inseparable pairs in $H_{17,10}$ and $H_{21,13}$ (Regime II), which our exhaustive 65,535-subset inventory classification (Proposition 16) proves minimal among the faithful restrictions of $H_{\text{YO}}^{+}$ with inherited coordinates.

For truth values, elements of reality fail as in the Kochen–Specker theorem: the logic has no two-valued states, or too few to separate its outcomes, and then it has no classical realization. For eigenvalues the situation is different. If the two-valued states of $H$ are unital and separating, $H$ can be realized classically as a partition logic[4,12,13]: outcome $i$ becomes the set of states with $s(i) = 1$, and each context becomes a partition of this set. Noncontextual eigenvalues can still be impossible, because they require a strong $d$-coloring, in the classical realization as much as in the quantum one. Greechie's $G_{32}$ is an example: it has a classical realization, no orthogonal ray assignment in $\mathbb{C}^3$ (Theorem 11), and $\chi(G_{32}) = 4$. Thus $\chi(H) > d$ is a condition on the spectra of maximal observables, not on truth values. It does, however, bound the size of every classical model. Preparations along orthogonal rays can be told apart with certainty, so in any hidden-variable model they have disjoint sets of hidden states; picking one hidden state for each ray then colors $\Gamma(H)$, and the model needs at least $\chi(H)$ hidden states (Sec. VIII works this out for Yu–Oh). A noncontextual eigenvalue assignment keeps orthogonal rays apart with only $d$ hidden states, the eigenvalues themselves. Thus, when $H$ is realized in $\mathbb{C}^d$, $\chi(H) > d$ means that a classical model needs more hidden states than the quantum system has dimensions merely to tell apart preparations along orthogonal rays, a gap that Prakash[14] calls representational contextuality and turns into a

memory advantage of quantum automata. A stronger separation, which excludes every local classical model, arises when an experiment fixes the label of each outcome, as the $d$-party supersinglet of the next paragraph does.

Noncontextual eigenvalues and strong colorings were first related by Cabello[8]. The first version of that preprint assigned noncontextual values $-1, 0, 1$ to the spin-one components $S_u$ along the rays $u$ and concluded that $\chi > d$ characterizes state-independent contextuality. For an orthogonal triad $u, v, w$, however, the eigenvectors of $S_u$, $S_v$, $S_w$ for the eigenvalue 1 (or $-1$) are not mutually orthogonal, so quantum mechanics fixes only which of them takes the value 0; the necessity of $\chi > d$ and the minimality of the 13-ray Yu–Oh set were later published in Ref.[15]. The context observables $A_e$ of Eq. (1) are diagonal in their contexts, so assumptions (i)–(iii) are equivalent to $\chi(H) \leq d$ without such a gap. The second version of the preprint distributes the $d$ particles of the totally antisymmetric supersinglet among $d$ parties. If all parties measure bases containing a ray $v$, exactly one of them obtains $v$; if $d-1$ parties measure the same basis, the ray of that basis which none of them obtains is obtained with certainty by the last party in any basis containing it. A local hidden-variable model of these perfect correlations exists exactly when the hypergraph of the measured bases has a strong $d$-coloring, the color of a ray being the party that obtains it; for the Yu–Oh set this hypergraph is $H_{\mathrm{YO}}^{+}$. Here the label of an outcome is the party that records it, and its noncontextuality, assumption (iii), follows from locality and the EPR criterion of reality[3] instead of being postulated. Quantum mechanics produces these correlations whenever $H$ has an orthogonal ray assignment in $\mathbb{C}^d$, while local classical models need a strong $d$-coloring, so here $\chi(H) > d$ does separate quantum from classical physics. $G_{32}$ has no ray assignment and gives no such separation. For a single system in the maximally mixed state the threshold is fractional: $I_d/d$ has a noncontextual truth-assignment model exactly when $\chi_f(\Gamma(H)) = d$[16] (Proposition 6). Contrary to the claim of Ref.[8], $\chi > d$ is therefore not sufficient for the contextuality of $I_d/d$: the faithful complex Hesse hypergraph $H_{37}$ has $\chi = 4$ and $\chi_f = 3$ (Theorem 19). The gap between $\chi$ and the least dimension of an orthogonal ray assignment has also been studied in connection with the quantum chromatic number[17].

We further connect these combinatorial and geometric layers to orthomodular event representations and quantum probability polytopes $\mathcal{P}_{\mathrm{ext}}(H) \subseteq \mathcal{P}(H) = \operatorname{conv} \mathcal{S}(H)$, where $\mathcal{P}_{\mathrm{ext}}(H)$ collects the probability vectors of hidden-variable models that choose a noncontextual eigenvalue assignment and one of its $d$ eigenvalues. On $H_{17,10}$, an explicit quantum ray preparation lies in $\mathcal{P}(H_{17,10}) \setminus \mathcal{P}_{\mathrm{ext}}(H_{17,10})$ and requires nonextendable state weight exactly $1/9$; the preparation along the coordinate ray $z_3$ requires exactly $1/3$, the largest weight that a single chromatic identity of $H_{17,10}$ can certify (Theorem 24). In complex dimension three, the 37-ray Hesse restriction $H_{37}$[18] has no injective $\mathbb{R}^3$ realization, pastes into a 76-event orthomodular lattice with exact minimum state-family sizes 6, 7, and 18 for vertex separation, event separation, and order determination (Theorem 19), and demonstrates that exact event-order determination does not guarantee classical mixtures for all Born preparations (Corollary 25). In real dimension four, our two-plane completion lemma (Lemma 21) produces the canonical linear 65-ray hypergraph $H_{65}$ completing the 17-ray core of Haynes *et al.*[19] with $\chi = 5$, $\chi_f = 17/4$, and 393,216 states that fail atom-order determination and collapse a ray with a plane (Theorem 22). Finally, we formulate and prove the existence form of the KS ancestry hypothesis (Proposition 26), give an abstract transverse chromatic-to-stateless construction (Proposition 27), establish exact inclusions into specified KS parents (Bub's 49-ray set[20], the 165-ray MUB inventory, Peres's 24-ray geometry, and 68-ray extensions containing the 18-ray parity proof $C_{18,9}$ of Cabello *et al.*[21] and a copy $C_{20,11}$ of Kernaghan's 20-ray parity proof[22]), and contrast abstract context states $\mathcal{S}_{\mathrm{ctx}}$ with geometric states $\mathcal{S}_{\mathrm{geom}}$ on the 36-ray completion and 45-ray vertex deletion of $C_{18,9}$. Complete vector, state, and coloring enumerations and finite verification certificates are provided in the supplementary material, which additionally cites Refs.[23–25].

## II. UNIFIED FRAMEWORK: STATES, COLORINGS, SEPARABILITY, AND COMPLETION

### A. Context hypergraphs, orthogonal realizations, and spectral regimes

All hypergraphs and graphs considered here are finite. A $d$-uniform *context hypergraph* $H = (V, E)$ ($d \geq 2$) consists of a nonempty collection $E$ of $d$-element subsets of $V$, called *contexts*, with $V = \bigcup_{e \in E} e$. Its *two-section* (or *context*) *graph* $\Gamma(H)$ has vertex set $V$ and edge set $E(\Gamma(H)) = \bigcup_{e \in E} \binom{e}{2}$. A hypergraph $H$ is *linear* if $|e \cap e'| \leq 1$ for all distinct $e, e' \in E$. A *Berge cycle* of length $\ell \geq 2$ is an alternating sequence of distinct vertices $x_1, \ldots, x_\ell$ and distinct contexts $e_1, \ldots, e_\ell$ with $\{x_j, x_{j+1}\} \subseteq e_j$ cyclically; the least such $\ell$ is the *Berge girth* $g(H)$ (or $\infty$ if acyclic). In Greechie's pasting of Boolean blocks[9,26], linearity ($g(H) \geq 3$) with $g(H) \geq 4$ yields an *orthomodular poset*, and $g(H) \geq 5$ yields an *orthomodular lattice* (OML).

With the Hermitian inner product $\langle x, y \rangle = \sum_j \overline{x_j} y_j$ on $\mathbb{F}^d$ ($\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$), we distinguish three levels of geometric realization:

1. An *orthogonal ray assignment* $\rho : V \to \mathbb{P}(\mathbb{F}^d)$, $i \mapsto [v_i]$ ($v_i \neq 0$), requires $\langle v_i, v_j \rangle = 0$ whenever $i \sim j$ in $\Gamma(H)$; nonadjacent vertices may share a ray.
2. An *injective orthogonal representation* additionally requires $[v_i] \neq [v_j]$ for all $i \neq j$.
3. An injective representation is *edge-faithful* (abbreviated FOR) if $\langle v_i, v_j \rangle = 0 \iff i \sim j$ in $\Gamma(H)$ for all $i \neq j$[27].

A *strong $k$-coloring* of $H$ is a map $c : V \to \{0, \ldots, k-1\}$ injective on every context $e \in E$, i.e., a proper $k$-coloring of $\Gamma(H)$; the least such $k$ is $\chi(H) = \chi(\Gamma(H))$. We call $H$ *chromatically complete* (equivalently, $H$ has an exact palette) when $\chi(H) = d$ and write $\mathcal{C}_d(H)$ for the set of strong $d$-colorings of $H$. For a realized $d$-uniform hypergraph with rank-one projectors $P_i = v_i v_i^\dagger / \langle v_i, v_i \rangle$, three spectral labelings arise: (a) *context-dependent common spectrum*, where bijections $\ell_e : e \to \Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$ define $A_e = \sum_{i \in e} \ell_e(i) P_i$ with shared projectors carrying context-dependent eigenvalues; (b) *context-independent vertex labeling* via a strong $k$-coloring $c$ and distinct real scalars $\lambda_0, \ldots, \lambda_{k-1}$ in Eq. (1), where spectra vary across contexts whenever $|c(V)| > d$ (unavoidable when $\chi(H) > d$); and (c) *chromatic completeness* ($k = d$), which provides both context-independent eigenvalues and a single $d$-element spectrum $\Sigma$ for all $A_e$. Only labeling (c) meets all three assumptions of the Introduction: labeling (a) gives up noncontextual eigenvalues, and labeling (b) keeps them at the price of more than $d$ eigenvalues, so that different contexts have different spectra.

### B. Two-valued states, color extendability, and the separation hierarchy

A *two-valued context state* is a function $s : V \to \{0, 1\}$ satisfying

$$\sum_{v \in e} s(v) = 1 \qquad (e \in E). \tag{3}$$

We write $\mathcal{S}(H) = \mathcal{S}_{\mathrm{ctx}}(H)$ for the set of context states. For a ray realization $\rho$, the *geometric state space*

$$\mathcal{S}_{\mathrm{geom}}(H, \rho) = \{s \in \mathcal{S}(H) : s(i) + s(j) \leq 1 \text{ whenever } i \neq j,\ v_i \perp v_j\} \tag{4}$$

additionally enforces orthogonality exclusions on uncovered pairs[28]; $\mathcal{S}_{\mathrm{geom}}(H, \rho) = \mathcal{S}(H)$ when $\rho$ is edge-faithful. A state family $\Omega \subseteq \mathcal{S}(H)$ is *unital* if every $v \in V$ has $s(v) = 1$ for some $s \in \Omega$, and *vertex separating* if distinct $u, v \in V$ have $s(u) \neq s(v)$ for some $s \in \Omega$, inducing the partition set representation $A_i = \{s \in \Omega : s(i) = 1\}$. On an event poset $(L, \leq)$, $\Omega$ is *event separating* if distinct $x, y \in L$ have distinct profiles on $\Omega$, and *order determining* if $x \leq y \iff (\forall s \in \Omega,\ s(x) \leq s(y))$.

**Definition 1** (Chromatic separability and extendable states)**.** For chromatically complete $H$ ($\mathcal{C}_d(H) \neq \varnothing$), vertices $u, v \in V$ are *chromatically inseparable*, written $u \sim_c v$, if

$$c(u) = c(v) \qquad \text{for every } c \in \mathcal{C}_d(H), \tag{5}$$

an *implicit identity* in graph coloring[11] (Def. 3.8). We call $H$ *chromatically separating* if $u \sim_c v \implies u = v$. A state $s \in \mathcal{S}(H)$ is *extendable*, written $s \in \mathcal{S}_{\text{ext}}(H)$, if $\operatorname{supp} s = \{v \in V : s(v) = 1\}$ is a color class of some $c \in \mathcal{C}_d(H)$[5–7].

Physically, $u \sim_c v$ means that every noncontextual eigenvalue assignment gives $u$ and $v$ the same eigenvalue, and an extendable state is exactly a truth assignment of the form "the outcome with eigenvalue $\lambda_a$ occurs" for some noncontextual eigenvalue assignment $c$ and eigenvalue $\lambda_a$.

**Theorem 2** (Completion criterion)**.** *For $s \in \mathcal{S}(H)$ with $S = \operatorname{supp} s$, the following are equivalent:*

1. *$s \in \mathcal{S}_{\text{ext}}(H)$;*
2. *$\chi(\Gamma(H)[V \setminus S]) \leq d - 1$ (so $S$ is a critical independent set[11] (Def. 3.3));*
3. *$s$ belongs to $d$ states in $\mathcal{S}(H)$ with pairwise disjoint supports partitioning $V$;*
4. *$s$ belongs to $d - 1$ states in $\mathcal{S}(H)$ with pairwise disjoint supports.*

*For $d = 3$, $s \notin \mathcal{S}_{\text{ext}}(H)$ if and only if $\Gamma(H)[V \setminus S]$ contains an odd cycle.*

*Proof.* A strong $d$-coloring with color class $S$ leaves a $(d-1)$-coloring on $V \setminus S$. Conversely, any $(d-1)$-coloring of $\Gamma(H)[V \setminus S]$ combined with a $d$th color on $S$ is a strong $d$-coloring because each context meets $S$ once and forms a $(d-1)$-clique in $V \setminus S$; its color indicators yield (3). Conversely, $d$ states with disjoint supports $S_1, \ldots, S_d$ partitioning $V$ define the strong $d$-coloring $c(v) = k$ for $v \in S_k$, since each context meets each $S_k$ exactly once; hence (3) $\implies$ (1). Trivially (3) $\implies$ (4), and given $d-1$ disjoint state supports $S_1, \ldots, S_{d-1}$, the complement $s_d(v) = 1 - \sum_{k=1}^{d-1} s_k(v)$ is binary and sums to $d - (d-1) = 1$ on every context, giving (3). For $d = 3$, $\chi(\Gamma(H)[V \setminus S]) \leq 2$ holds if and only if $\Gamma(H)[V \setminus S]$ is bipartite (odd-cycle free). □

Applying the equivalence (3) $\iff$ (4) of Theorem 2 globally without anchoring a state $s$ yields the characterization of chromatic completeness:

**Lemma 3** (Disjoint states and strong coloring)**.** *A finite $d$-uniform hypergraph $H$ satisfies $\chi(H) = d$ if and only if $\mathcal{S}(H)$ contains $d - 1$ states with pairwise disjoint supports.*

**Corollary 4** (The separation hierarchy)**.** *For chromatically complete $H$ and $u, v \in V$,*

$$u \sim_c v \quad \iff \quad s(u) = s(v) \text{ for all } s \in \mathcal{S}_{\text{ext}}(H). \tag{6}$$

*Thus inseparability by $\mathcal{S}(H)$ implies chromatic inseparability ($\sim_c$), equivalently chromatic separation implies separation by $\mathcal{S}(H)$, whereas the converse fails even for faithful real ray configurations. By Theorem 0 of Kochen and Specker[4], separation of all outcomes by $\mathcal{S}(H)$ is what allows an embedding into a Boolean algebra; chromatic inseparability is the same failure of separation, with $\mathcal{S}(H)$ replaced by the states that noncontextual eigenvalue assignments produce. Moreover, $\mathcal{S}_{\text{ext}}(H)$ is always unital whenever $\mathcal{C}_d(H) \neq \varnothing$.*

*Proof.* The indicator states $s_a(v) = \mathbf{1}_{\{c(v)=a\}}$ over all $c \in \mathcal{C}_d(H)$ and $a \in \{0, \ldots, d-1\}$ are precisely $\mathcal{S}_{\text{ext}}(H) \subseteq \mathcal{S}(H)$. For each fixed coloring, the $d$ supports partition $V$. The inclusion need not be strict: for a single $d$-element context every state is extendable. Strict inclusion is an additional property established for the specified examples. □

**Proposition 5** (Canonical chromatic quotient and closure test)**.** *Let $H$ be chromatically complete.*

1. *The relation $\sim_c$ is an equivalence relation with no two equivalent vertices in a common context. The quotient $\overline{H} = (V/\!\sim_c, \overline{E})$ is $d$-uniform and chromatically separating, and pullback along $V \to V/\!\sim_c$ induces a bijection $\mathcal{C}_d(\overline{H}) \to \mathcal{C}_d(H)$.*

2. *For $u \neq v$, adjoining one context $\{u, v, w_1, \ldots, w_{d-2}\}$ with fresh vertices $w_i$ yields chromatic number $d+1$ if and only if $u \sim_c v$, and $d$ otherwise. Since recoloring one vertex with a new color separates any pair, the* separable chromatic number $\chi_{\mathrm{sep}}$ *of a finite graph, the least $t$ for which some set of proper colorings with at most $t$ colors separates every pair of distinct vertices*[5]*, satisfies $\chi_{\mathrm{sep}} \in \{\chi, \chi+1\}$.*

*Proof.* (1) Adjacent vertices receive distinct colors, so each context projects injectively to $V/\!\sim_c$, and every $c \in \mathcal{C}_d(H)$ factors uniquely through $V/\!\sim_c$. (2) A coloring $c \in \mathcal{C}_d(H)$ extends to the fresh context if and only if $c(u) \neq c(v)$; if $u \sim_c v$, recoloring $u$ with a $(d+1)$th color separates $(u, v)$ while preserving all old context inequalities. □

The quotient preserves colorings, not an injective realization. Closing an inseparable pair is likewise abstract: faithful coordinates make the pair nonorthogonal, so the added context cannot retain those rays.

Put $\mathcal{P}_{\mathrm{ext}} = \operatorname{conv} \mathcal{S}_{\mathrm{ext}}$ and $\mathcal{P} = \operatorname{conv} \mathcal{S}$. Thus $\mathcal{P}$ is the set of probability vectors of noncontextual truth-assignment models, and $\mathcal{P}_{\mathrm{ext}}$ is the subset produced by models that choose a noncontextual eigenvalue assignment $c$ and an eigenvalue $\lambda_a$ at random and report, in every context, the outcome with eigenvalue $\lambda_a$. A binary vector cannot be a convex combination of distinct binary vectors, since each positive-weight term must agree at every coordinate. Consequently,

$$\mathcal{P}_{\mathrm{ext}} = \mathcal{P} \quad \Longleftrightarrow \quad \mathcal{S}_{\mathrm{ext}} = \mathcal{S}. \tag{7}$$

This is stronger than equality of vertex-separation kernels: the pentagon has separating colors yet a nonextendable state. Pair identities alone do not describe all completion constraints.

### C. Fractional coloring and fresh-vertex context completion

Write $\alpha(G)$ for the independence number and $\chi_f(G)$ for the fractional chromatic number of $G = \Gamma(H)$.

**Proposition 6** (Fractional threshold and uniform state mixture)**.** *For any finite $d$-uniform context hypergraph $H$,*

$$\chi_f(\Gamma(H)) = d \quad \Longleftrightarrow \quad \frac{1}{d}\mathbf{1} \in \mathcal{P}(H) = \operatorname{conv} \mathcal{S}(H). \tag{8}$$

*Proof.* A state mixture $\sum_{s \in \mathcal{S}(H)} \mu_s s = \frac{1}{d}\mathbf{1}$ assigns weight $d\mu_s$ to the independent set $\operatorname{supp} s$, covering each vertex with weight 1 at total weight $d$, so $\chi_f(\Gamma(H)) \leq d$; since every context is a $d$-clique, $\chi_f(\Gamma(H)) \geq \omega(\Gamma(H)) \geq d$. Conversely, a fractional coloring $\{w_I\}$ of total weight $d$ obeys $d \leq \sum_{v \in e} \sum_{I \ni v} w_I = \sum_I w_I |I \cap e| \leq \sum_I w_I = d$ for each $e \in E$, forcing $|I \cap e| = 1$ whenever $w_I > 0$, so $\{w_I/d\}$ is a state mixture equal to $\frac{1}{d}\mathbf{1}$. □

Equation (8) restates, for context states, the fractional-chromatic criterion of Ramanathan and Horodecki[16] for the contextuality of the maximally mixed state.

**Proposition 7** (Fresh-vertex context completion)**.** *Let $G$ be a finite core graph with designated $d$-clique base contexts ($d \geq 3$). Complete selected uncovered edges $\{u, v\}$ into $d$-cliques using $d-2$ degree-one fresh vertices per edge, covering all core vertices and edges. For the resulting hypergraph $H$ and $G' = \Gamma(H)$,*

$$\begin{aligned} \chi(G') &= \max\{d, \chi(G)\}, \\ \chi_f(G') &= \max\{d, \chi_f(G)\}, \\ |\mathcal{S}(H)| &= \sum_S (d-2)^{m(S)}, \end{aligned} \tag{9}$$

TABLE I. Principal configurations. FOR means injective and edge-faithful; $g$ is Berge girth. Counts refer to all context states and all color-class states.

| Configuration | $d$ | $\lvert V\rvert/\lvert E\rvert$ | $g$ | $\chi$ | $\chi_f$ | $\lvert\mathcal{S}\rvert$ | $\lvert\mathcal{S}_{\text{ext}}\rvert$ | Geometry and principal distinction |
|---|---|---|---|---|---|---|---|---|
| One triad | 3 | 3/1 | $\infty$ | 3 | 3 | 3 | 3 | Real FOR; colors separate. |
| $H_{6,3}$ | 3 | 6/3 | 3 | 3 | 3 | 4 | 3 | No injective complex realization. |
| $P_5$ | 3 | 10/5 | 5 | 3 | 3 | 11 | 10 | Real FOR; colors separate; ternary exclusion. |
| $H_8$ | 3 | 14/8 | 5 | 3 | 3 | 11 | 7 | Repeated-ray assignments; no complex injection. |
| $H_9$ | 3 | 15/9 | 5 | 4 | 3 | 8 | 0 | No complex ray assignment; context-critical. |
| $G_{32}$ | 3 | 15/10 | 5 | 4 | 3 | 6 | 0 | No complex ray assignment. |
| $H_{17,10}$ | 3 | 17/10 | 5 | 3 | 3 | 18 | 10 | Real FOR; two color-identical pairs. |
| $H_{21,13}$ | 3 | 21/13 | 5 | 3 | 3 | 21 | 6 | Real FOR; 21 color-identical pairs. |
| Completed Yu–Oh | 3 | 25/16 | 5 | 4 | 35/11 | 24 | 0 | Real FOR; context-critical. |
| $H_{37}$ | 3 | 37/24 | 5 | 4 | 3 | 153 | 0 | Complex FOR; no real injection; OML minima 6/7/18. |
| $H_{\text{YO}}^{(4)}$ | 4 | 26/16 | 2 | 5 | 46/11 | 25 | 0 | Real FOR; common-apex suspension, nonlinear. |
| $C_{36,18}$ | 4 | 36/18 | 3 | 5 | 9/2 | 0 | 0 | Faithful linear KS completion. |
| $H_{65}$ | 4 | 65/29 | 3 | 5 | 17/4 | 393,216 | 0 | Real FOR; false atom order and compound collapse. |

*where $S$ ranges over independent core supports meeting each base context once and $m(S)$ counts completed edges missed by $S$.*

*Proof.* Any $k$-coloring ($k \geq d$) or measurable fractional palette of size $k = \max\{d, \chi_f(G)\}$ assigns disjoint colors/sets of measure 1 to $u$ and $v$, leaving $k-2 \geq d-2$ colors (or measure $k-2 \geq d-2$) for the $d-2$ fresh vertices. A core support $S$ hitting $\{u, v\}$ forces its fresh vertices to 0, whereas a missed edge allows $d-2$ choices of a single fresh vertex set to 1. □

Table I summarizes the principal configurations studied below.

The logical situation has two distinct branches. If a hypergraph has no admissible two-valued context state, it cannot admit a strong $d$-coloring, because the color classes of such a coloring would supply $d$ admissible states. The converse implication fails: a noncontextual assignment of $d$ eigenvalues to the outcomes may be impossible while ordinary states, including a unital separating family, remain available.

When a noncontextual eigenvalue assignment does exist, its color classes provide a nonempty unital family of extendable states inside the full context-state space. A particular ordinary state belongs to this restricted family precisely when deleting its support leaves a $(d-1)$-colorable graph. Chromatic identities describe pairs having equal eigenvalues in every noncontextual eigenvalue assignment, and hence equal values in every state of that restricted family; they need not hold throughout the ordinary state space. These assignment questions have to be distinguished from the separate requirements of an injective geometric realization, a representation of event order, and a classical mixture reproducing a specified Born probability vector.

## III. INDEPENDENCE OF COLORING AND COORDINATIZATION, PETERSEN OBSTRUCTIONS, AND COMPLETED YU–OH

### A. Small abstract examples and four-quadrant independence

The non-linear hypergraph $H_{4,2}$ with contexts $\{1,2,3\}$ and $\{1,2,4\}$ has three states $\{1\}$, $\{2\}$, $\{3,4\}$ (all extendable), forces $3 \sim_c 4$ and $s(3) = s(4)$, and has no injective $\mathbb{C}^3$ realization because two orthogonal rays determine a unique orthogonal-complement ray.

**Proposition 8** (Smallest linear chromatically inseparable hypergraph)**.** *The linear 3-uniform hypergraph $H_{6,3}$ with contexts $\{1,2,3\}$, $\{3,4,5\}$, $\{5,6,1\}$ of Berge girth three has four unital separating states $\{1,4\}$, $\{2,5\}$, $\{3,6\}$, $\{2,4,6\}$, a unique strong 3-coloring up to permutation with inseparable classes $\{1,4\}$, $\{2,5\}$, $\{3,6\}$, and three extendable states (as $V \setminus \{2,4,6\} = \{1,3,5\}$ is a triangle). No linear 3-uniform hypergraph with $|V| < 6$ or $|E| < 3$ has a chromatically inseparable pair, and $H_{6,3}$ admits no injective realization in $\mathbb{C}^3$.*

*Proof.* The triangle $\{1,3,5\}$ in $\Gamma(H_{6,3})$ forces three distinct colors on $1,3,5$, whence $c(4) = c(1)$, $c(2) = c(5)$, and $c(6) = c(3)$. Similarly, in $\mathbb{C}^3$, $\{v_1, v_3, v_5\}$ forms an orthonormal basis, forcing $[v_4] = [v_1]$, $[v_2] = [v_5]$, and $[v_6] = [v_3]$. With $|E| \leq 2$, independent permutations of unshared vertices separate all pairs; by inclusion–exclusion, three linear 3-contexts require $|V| \geq 6$. □

**Proposition 9** (Clique number at Berge girth at least four)**.** *If $H = (V, E)$ is an $n$-uniform linear hypergraph ($n \geq 3$) of Berge girth $g(H) \geq 4$, every clique $K \subseteq V$ of size $|K| \geq 3$ in $\Gamma(H)$ is contained in a unique context $e \in E$. Hence $\omega(\Gamma(H)) = n$, and chromatic incompleteness ($\chi(H) > n$) is equivalent to the chromatic–clique gap $\chi(\Gamma(H)) > \omega(\Gamma(H))$.*

*Proof.* For any three vertices $u, v, w \in K$, pairwise adjacency gives contexts $e_1 \supseteq \{u, v\}$, $e_2 \supseteq \{v, w\}$, $e_3 \supseteq \{w, u\}$. Distinct $e_1, e_2, e_3$ would form a Berge 3-cycle, contradicting $g(H) \geq 4$. By linearity, the shared context $e$ containing $\{u, v, w\}$ is unique for $\{u, v\}$ and thus contains all of $K$. □

**Proposition 10** (Logical independence of coloring and coordinatization)**.** *For finite 3-uniform context hypergraphs, all four combinations of chromatic completeness ($\chi(H) = 3$) and injective coordinatizability in $\mathbb{C}^3$ occur even within the orthomodular-lattice admissible class of linear hypergraphs of Berge girth $g(H) \geq 5$[9,26] with unital separating state spaces:*

1. *$\chi = 3$, injectively realizable in $\mathbb{R}^3$: a single context $C_1 = \{\{1,2,3\}\}$ (or $P_5$, $H_{17,10}$, $H_{21,13}$);*
2. *$\chi = 3$, not injectively realizable in $\mathbb{C}^3$: the 8-context Petersen restriction $H_8$ (Theorem 11);*
3. *$\chi = 4$, edge-faithfully realizable in $\mathbb{R}^3$: completed Yu–Oh $H^+_{\mathrm{YO}}$ (Theorem 13);*
4. *$\chi = 4$, admitting no orthogonal ray assignment in $\mathbb{C}^3$: Greechie's $G_{32}$ and $H_9$ (Theorem 11).*

Injectivity is essential to quadrant (2): any strong $n$-coloring $c$ yields a noninjective ray assignment $i \mapsto [e_{c(i)+1}] \in \mathbb{P}(\mathbb{F}^n)$. At Berge girth four (an orthomodular poset that is not a lattice), $E_{\square} = \{\{1,2,3\}, \{3,4,5\}, \{5,6,7\}, \{7,8,1\}\}$ also satisfies $\chi = 3$ while forcing $[v_3] = [v_7] \perp \mathrm{span}(v_1, v_5)$ whenever $[v_1] \neq [v_5]$.

### B. Greechie's $G_{32}$, multipole forcing, and correction of the JMP spectral argument

Greechie's $G_{32}$[9] (Fig. 6) has 15 vertices and ten contexts

$$\begin{aligned} E(G_{32}) = \{&123, 345, 567, 789, 9\,10\,11, \\ &11\,12\,1, 4\,10\,13, 6\,12\,14, \\ &8\,2\,15, 13\,14\,15\}. \end{aligned} \tag{10}$$

Each vertex lies in two contexts, so $G_{32}$ is dual to the Petersen graph $P$, $\Gamma(G_{32}) = L(P)$, $\mathrm{Aut}(G_{32}) \cong S_5$, and $\chi(G_{32}) = \chi'(P) = 4$. Its six two-valued state supports satisfy

$$|\mathcal{S}(G_{32})| = 6, \qquad |S_\alpha| = 5, \qquad |S_\alpha \cap S_\beta| = 1 \ (\alpha \neq \beta). \tag{11}$$

These supports yield the $\binom{6}{2} = 15$ two-element vertex profiles $B_i = \{\alpha : i \in S_\alpha\}$. Their nonempty pairwise intersections exclude even two disjoint states, confirming $\chi(G_{32}) = 4$ by Lemma 3.

In Appendix B of Shekarriz and Svozil[6], the chromatic obstruction $\chi(G_{32}) = 4 > 3$ was invoked to argue against a 3-dimensional realization of $G_{32}$. As Proposition 10 demonstrates via $H^+_{\mathrm{YO}}$ ($\chi = 4$, yet faithfully realizable in $\mathbb{R}^3$), $\chi(H) > 3$ rules out only a context-independent 3-value spectral assignment—a noncontextual assignment of the eigenvalues of uniform maximal observables to their outcomes—*not* geometric rays. A genuine geometric exclusion requires the projection identity: for an orthonormal basis $\{x, y, z\} \subset \mathbb{C}^3$ with $u \perp x$ and $w \perp z$,

$$\langle u, w\rangle = \langle u, y\rangle\langle y, w\rangle. \tag{12}$$

**Theorem 11** (Multipole forcing and minimal complex obstructions)**.** *Let* $H_8 = G_{32} \setminus \{\{8, 2, 15\}, \{13, 14, 15\}\}$ *(*14 *vertices,* 8 *contexts) and* $H_9 = G_{32} \setminus \{\{13, 14, 15\}\}$ *(*15 *vertices,* 9 *contexts).*

1. *In every orthogonal ray assignment of* $H_8$ *in* $\mathbb{C}^3$ *(allowing repeated rays), the terminal rays satisfy*

$$[v_2] = [v_8] \qquad \textit{and} \qquad [v_{13}] = [v_{14}], \tag{13}$$

*and either* $[v_1] = [v_4] = [v_9]$ *or* $[v_3] = [v_7] = [v_{12}]$*. Thus* $H_8$ *has no injective realization in* $\mathbb{C}^3$*.*

2. $H_8$ *is chromatically complete (*$\chi(H_8) = 3$*) with* 11 *unital separating states, of which* 7 *are extendable (the* 4 *nonextendable states leave odd cycles in* $\Gamma(H_8)$*; see Table S38). Its* 3 *normalized* 3*-colorings (fixing* $(c(1), c(2), c(3)) = (0, 1, 2)$*) are listed in Table S39, forcing the* 6 *chromatically inseparable pairs* $(1, 9)$*,* $(2, 8)$*,* $(3, 7)$*,* $(4, 6)$*,* $(10, 12)$*,* $(13, 14)$*; deleting any single context of* $H_8$ *removes all forced color identities (leaving* $6, 20, 20, 6, 20, 20, 6, 6$ *normalized colorings on the remaining active vertices, up to global color permutation). Isolated vertices are dropped: retaining the isolated vertex created by deleting* $\{1, 2, 3\}$*,* $\{7, 8, 9\}$*,* $\{4, 10, 13\}$*, or* $\{6, 12, 14\}$ *multiplies the corresponding count by three, from* 6 *to* 18*.*

3. *Neither* $H_9$*,* $G_{32}$*, nor the* 12*-vertex,* 21*-edge subgraph* $\Gamma_0$ *of* $\Gamma(H_9)$ *on* $\{1, \ldots, 12\}$ *admits any orthogonal ray assignment in* $\mathbb{C}^3$*. Moreover,* $H_9$ *is context-minimal and* $\Gamma_0$ *is edge-minimal: deleting any context of* $H_9$ *or edge of* $\Gamma_0$ *yields a* 3*-colorable structure realizable in* $\mathbb{R}^3$ *by standard basis rays.*

*Proof.* (1) Let $v_1, \ldots, v_{14} \in \mathbb{C}^3$ be unit vectors for $H_8$. In $\{v_1, v_2, v_3\}$, $v_{11}, v_{12} \perp v_1$ and $v_4, v_5 \perp v_3$, so Eq. (12) gives $\langle x, y\rangle = \langle x, v_2\rangle\langle v_2, y\rangle$ for $x \in \{v_{11}, v_{12}\}, y \in \{v_4, v_5\}$. Expanding $v_{12}$ in $\{v_5, v_6, v_7\}$ ($v_{12} \perp v_6$) and $v_4$ in $\{v_9, v_{10}, v_{11}\}$ ($v_4 \perp v_{10}$) yields $v_7\langle v_7, v_{12}\rangle = v_{12} - v_5\langle v_5, v_{12}\rangle$ and $v_9\langle v_9, v_4\rangle = v_4 - v_{11}\langle v_{11}, v_4\rangle$. Since $v_7 \perp v_9$, $v_{12} \perp v_{11}$, and $v_5 \perp v_4$, their inner product vanishes:

$$\begin{aligned} 0 &= \langle v_{12}, v_4\rangle + \langle v_{12}, v_5\rangle\langle v_5, v_{11}\rangle\langle v_{11}, v_4\rangle \\ &= \langle v_{12}, v_2\rangle\langle v_2, v_4\rangle\big(1 + |\langle v_2, v_5\rangle|^2|\langle v_2, v_{11}\rangle|^2\big). \end{aligned} \tag{14}$$

Thus $\langle v_2, v_4\rangle = 0$ or $\langle v_{12}, v_2\rangle = 0$. If $\langle v_2, v_4\rangle = 0$, set $(v_1, v_3, v_2) = (e_1, e_2, e_3)$; then $v_4 \perp v_2, v_3 \implies [v_4] = [e_1]$, $\{3, 4, 5\} \implies [v_5] = [e_3] = [v_2]$, $\{9, 10, 11\}$ with $v_{10}, v_{11} \perp e_1$ forces $[v_9] = [e_1]$, $\{5, 6, 7\}$ and $\{7, 8, 9\}$ force $[v_7] = [e_2], [v_6] = [e_1], [v_8] = [e_3] = [v_2]$, and $\{4, 10, 13\}$, $\{9, 10, 11\}$, $\{11, 12, 1\}$, $\{6, 12, 14\}$ in $\mathrm{span}(e_2, e_3)$ force $[v_{13}] = [v_{11}] = [v_{14}]$ and $[v_1] = [v_4] = [v_9]$. Reflection symmetry gives the second branch when $\langle v_{12}, v_2\rangle = 0$.

(2) In $H_8$, fixing $(c(1), c(2), c(3)) = (0, 1, 2)$ leaves 16 choices along $345, 567, 789, 9\,10\,11$, of which 13 fail on $11\,12\,1, 4\,10\,13, 6\,12\,14$, leaving $c_1, c_2, c_3$ in Table S39. Equal columns identify the 6 pairs.

(3) Both $H_9$ (hence $G_{32}$) and $\Gamma_0$ add the orthogonality $v_2 \perp v_8$ to $H_8$, contradicting $[v_2] = [v_8]$. For minimality, deleting $\{8, 2, 15\}$ from $H_9$ leaves $H_8$ ($\chi = 3$); deleting $\{4, 10, 13\}$ or $\{6, 12, 14\}$ is isomorphic by $D_6$ symmetry to deleting $\{8, 2, 15\}$; and deleting any ring context (say $\{1, 2, 3\}$) allows the 3-coloring $(c(1), \ldots, c(15)) = (2, 1, 2, 0, 1, 0, 2, 0, 1, 2, 0, 1, 1, 2, 2)$. On $\Gamma_0$, the following deletions cover all 21 edges up to $D_6$ symmetry: $\{1, 2\}$ (colored by 001020120121), $\{2, 3\}$ (011020120121), $\{1, 3\}$ (010120102012), and $\{4, 10\}$ (012010201021), each yielding a standard-basis $\mathbb{R}^3$ assignment $i \mapsto [e_{c(i)+1}]$. For the minimum $\Delta_3$ of $\sum |\langle v_i, v_j \rangle|^2$ over the 30 edges $\{i, j\}$ of $\Gamma(G_{32})$ and all unit vectors $v_1, \ldots, v_{15} \in \mathbb{C}^3$, the robustness bounds $1/14400 < \Delta_3 < 0.927496$ are proved in Sec. S3 of the supplementary material. □

In his proof that state-independent contextuality in $\mathbb{C}^3$ requires 13 rays, Cabello[8] found $\Gamma_0$ to be the only square-free graph with fewer than 13 vertices and chromatic number above 3, and excluded it by showing, in real coordinates, that ten of its vertices (in our labels, all but 2 and 8) cannot be distinct rays. Part (1) proves this over $\mathbb{C}^3$: the rays $2, 8, 13, 14$ of $H_8$ are then fixed as orthogonal completions, and the ten rays satisfy $[v_1] = [v_4] = [v_9]$ or $[v_3] = [v_7] = [v_{12}]$. Part (3) shows more: $\Gamma_0$ admits no orthogonal ray assignment in $\mathbb{C}^3$ at all, even with repeated rays.

The complete normalized coloring lists for $H_8$ and $H_{17,10}$ are in the supplement.

**Corollary 12** (Blanuša and Isaacs dot-product snarks)**.** *If a cubic graph $G$ admits no orthogonal vector edge coloring in $\mathbb{C}^3$, neither does any Isaacs dot product $G \cdot P$[29], obtained by deleting two nonadjacent edges of $G$ and two adjacent vertices $u, v$ of the Petersen graph $P$, and joining the two dangling edges of $u$ to the ends of one deleted edge and those of $v$ to the ends of the other. In particular, both 18-vertex Blanuša snarks $B_1, B_2 = P \cdot P$[30] and all iterated dot products $X = P \cdot P \cdots P$ satisfy $\chi(H(X)) = 4 > 3 = \omega(L(X))$ and admit no orthogonal ray assignment in $\mathbb{C}^3$.*

*Proof.* In $G \cdot P$, the embedded copy of $P \setminus \{u, v\}$ has dual hypergraph $H_8$, so Theorem 11(1) forces $[v_2] = [v_8]$ and $[v_{13}] = [v_{14}]$ on the two pairs of cut edges, inducing a valid orthogonal ray assignment on $G$. □

Silva and Ibiapina[23] (Question 6) asked whether a cubic graph with chromatic index 4 can have an orthogonal vector edge coloring in $\mathbb{R}^3$. For the Petersen graph, the Blanuša snarks, and these dot products the answer is no, even in $\mathbb{C}^3$.

### C. Completed Yu–Oh $H^+_{\mathrm{YO}}$, base-coloring partition, and context criticality

Completing each of the 12 uncovered orthogonal pairs of the 13-ray Yu–Oh set[10] ($r_1, \ldots, r_{13} \in \mathbb{R}^3$, consisting of $z_1 = r_2, z_2 = r_8, z_3 = r_5$, $y_1^{\pm} = r_{1,3}, y_2^{\pm} = r_{7,9}, y_3^{\pm} = r_{4,6}$, and diagonals $h_0 = r_{12}, h_1 = r_{11}, h_2 = r_{10}, h_3 = r_{13}$) by cross-product rays $r_{14}, \ldots, r_{25}$ (Table S2) yields the 25-ray, 16-context linear hypergraph $H^+_{\mathrm{YO}}$ of Berge girth five[7,31]:

$$\begin{aligned} E^+_{\mathrm{YO}} &= \{123, 789, 456, 285\} \cup \bigcup_{h=10}^{13} \mathcal{E}_h, \\ \mathcal{E}_h &= \{\{x, h, w_{xh}\} : x \in N(h)\}, \end{aligned} \tag{15}$$

All 25 vectors, 16 contexts, and restriction-membership flags are tabulated in the supplement. Here the neighborhoods are $N(10) = \{1, 9, 4\}$, $N(11) = \{3, 7, 4\}$, $N(12) = \{3, 9, 6\}$ and $N(13) = \{1, 7, 6\}$, and each $w_{xh}$ is the unique completing ray. It has a real FOR in $\mathbb{R}^3$ (48 orthogonal pairs) and 24 unital separating states (each of the 12 binary assignments on the four base contexts leaves uniquely one diagonal $h_j$ with all three neighbors zero, allowing

$s(h_j) \in \{0,1\}$ and hence $12 + 4 \times 3 = 24$ states). Completed Yu–Oh is therefore a faithfully realized real configuration in which noncontextual truth assignments survive, enough of them to separate every pair of rays, while by the following theorem no noncontextual assignment of three eigenvalues to its outcomes exists.

**Theorem 13** (Base-coloring partition and context criticality of $H_{\mathrm{YO}}^{+}$)**.** *The completed Yu–Oh hypergraph $H_{\mathrm{YO}}^{+}$ satisfies $\omega(\Gamma(H_{\mathrm{YO}}^{+})) = 3$, $\chi_f(\Gamma(H_{\mathrm{YO}}^{+})) = 35/11 > 3 = \chi_f(\Gamma(G_{32}))$, and $\chi(H_{\mathrm{YO}}^{+}) = 4$, and is context-critical for $\chi > 3$:*

1. *Fixing $(c(2), c(8), c(5)) = (0, 1, 2)$, the four base contexts $\{1,2,3\}$, $\{7,8,9\}$, $\{4,5,6\}$, $\{2,8,5\}$ admit eight strong 3-colorings indexed by $(b_1, b_2, b_3) \in \{0,1\}^3$:*

$$\begin{aligned} (c(1), c(3)) &= (2 - b_1, 1 + b_1), \\ (c(7), c(9)) &= (2 - 2b_2, 2b_2), \\ (c(4), c(6)) &= (1 - b_3, b_3). \end{aligned} \tag{16}$$

   *Each antipodal pair of $(b_1, b_2, b_3)$ makes the neighborhood $N(h_j)$ of* uniquely one *diagonal ray $h_j \in \{10, 11, 12, 13\}$ rainbow ($\{0,1,2\}$) while the other three neighborhoods are 2-colored:*

| *Antipodal binary patterns* | *Blocked diagonal* |
|---|---|
| $101, 010$ | 12 |
| $110, 001$ | 11 |
| $000, 111$ | 10 |
| $011, 100$ | 13 |

(17)

2. *Consequently, $\chi(H_{\mathrm{YO}}^{+}) = 4$, whereas deleting any single context $e \in E_{\mathrm{YO}}^{+}$ yields $\chi(H_{\mathrm{YO}}^{+} \setminus \{e\}) = 3$.*

*Proof.* Direct check of Eqs. (16)–(17) shows that every base 3-coloring blocks one $h_j$, so $\chi(H_{\mathrm{YO}}^{+}) \geq 4$, matched by the four-coloring

$$(c(1), \ldots, c(25)) = (2,0,1,1,2,0,2,1,0,3,0,2,1,2,1,1,1,0,0,0,0,2,1,2,0),$$

in which only $h_2 = 10$ receives the fourth color. Deleting a sector context $\{x, h_j, w\}$ frees $h_j$ in its antipodal base colorings by setting $c(h_j) = c(x)$; deleting $\{2,8,5\}$ allows $c(2) = c(8) = c(5) = 0$; deleting $\{1,2,3\}$ allows $(c(1), c(2), c(3)) = (0,0,1)$, $(c(7), c(8), c(9)) = (2,1,0)$, $(c(4), c(5), c(6)) = (1,2,0)$, and $(c(10), c(11), c(12), c(13)) = (2,0,2,1)$; the cases $\{7,8,9\}$ and $\{4,5,6\}$ follow by the threefold cyclic coordinate symmetry of the Yu–Oh set. For $\chi_f$, weight $1/2$ on the 6 states of $G_{32}$ gives $\chi_f(G_{32}) = 3$, while Proposition 7 preserves the 13-ray value $\chi_f = 35/11$[15]. □

## IV. EXACT-PALETTE CHROMATIC SEPARABILITY AND LOGICAL GADGETS IN DIMENSIONS THREE AND FOUR

From now on we assume that noncontextual assignments of $d$ eigenvalues exist ($\chi(H) = d$) and ask what they imply for the relations between distinct outcomes. Such assignments may force outcomes to share or to avoid an eigenvalue although ordinary truth assignments permit otherwise; these forced relations are their logical content.

### A. Kochen–Specker antecedents and palette-invariant implications

Three historical Kochen–Specker constructions clarify the hierarchy of logical gadgets:

1. *1965 abstract TIFS precursor:* Identifying repeated labels $v_1$ and $v_2$ in Kochen and Specker's 1965 Fig. 1[32] (p. 182) yields an 11-vertex, 7-context hypergraph $\{134, 325, 2\,9\,11, 11\,1\,10, 479, 5\,8\,10, 678\}$ implementing $s(1) = 1 \implies s(2) = 0$. It has only four states ($\{1, 5, 6, 9\}$, $\{2, 4, 6, 10\}$, $\{3, 6, 9, 10\}$, $\{4, 5, 6, 11\}$), all with $s(6) = 1$ and $s(7) = s(8) = 0$, so $\chi > 3$; moreover, 1 and 9 are both orthogonal to 4 and 11, so the distinct rays 4 and 11 of an injective realization in $\mathbb{C}^3$ would force $[v_1] = [v_9]$; no injective realization exists (Sec. S3 of the supplementary material).

2. *1967 directed TITS* $\Gamma_1$*:* The 10-vertex graph $\Gamma_1$ of Lemma 1 in Ref.[4] (pp. 68, 70) implements the one-way implication $s(a_0) = 1 \implies s(a_9) = 1$. Completing its six uncompleted edges by fresh rays $23, 19, 18, 15, 14, 5$ yields the 16-ray, 9-context configuration $I_{16,9}$ in Fig. 2(d) (explicit label map in Sec. S3 of the supplementary material).

3. *1967 state-inseparable* $\Gamma_3$*:* In Ref.[4] (p. 70), $\Gamma_3$ forces $s(a) = s(b)$ in *every* two-valued state ($a, b$ are nonseparated by $\mathcal{S}(H)$). By contrast, our chromatically inseparable examples below have $\mathcal{S}(H)$ separating all vertices, with $c(u) = c(v)$ forced only on $\mathcal{S}_{\text{ext}}(H)$.

For distinct nonadjacent terminals $u, v \in V$, an ordinary TIFS[33] obeys $s(u) = 1 \implies s(v) = 0$ (equivalently $s(u) + s(v) \leq 1$, which is always bidirectional), and a directed ordinary TITS obeys $s(u) = 1 \implies s(v) = 1$. When $\mathcal{C}_d(H) \neq \varnothing$, an *unanchored chromatic TIFS* obeys $c(u) = 0 \implies c(v) \in \{1, \ldots, d-1\}$, and an *unanchored chromatic TITS* obeys $c(u) = 0 \implies c(v) = 0$.

**Proposition 14** (Palette symmetry and purely chromatic gadgets)**.** *For chromatically complete $H$ and nonadjacent $u, v \in V$, the unanchored chromatic TIFS is equivalent to $c(u) \neq c(v)$ for all $c \in \mathcal{C}_d(H)$ (an* implicit edge[11] *(Def. 3.1)), and the unanchored chromatic TITS is equivalent to $u \sim_c v$ (an* implicit identity*). Both chromatic relations are bidirectional. Ordinary TIFS and directed ordinary TITS entail their chromatic counterparts, and with only $c(u) = 0$ fixed, the allowed output set $\{c(v) : c \in \mathcal{C}_d(H),\ c(u) = 0\}$ is one of $\{0\}$, $\{1, \ldots, d-1\}$, or $\{0, \ldots, d-1\}$.*

*Proof.* Global palette permutations act transitively on $\{0, \ldots, d-1\}$ and, stabilizing 0, transitively on $\{1, \ldots, d-1\}$. Since every color class $s_a(x) = \mathbf{1}_{\{c(x)=a\}}$ is in $\mathcal{S}_{\text{ext}}(H) \subseteq \mathcal{S}(H)$, any ordinary TIFS or one-way TITS forces $c(u) \neq c(v)$ or $c(u) = c(v)$, respectively. □

We call a chromatic TIFS *purely chromatic* if some $s \in \mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$ satisfies $s(u) = s(v) = 1$, and a chromatic identity $u \sim_c v$ *two-way purely chromatic* if ordinary states in $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$ realize both $(s(u), s(v)) = (1, 0)$ and $(0, 1)$, so neither directed ordinary TITS holds. Since a state with $s(u) = s(v) = 1$ cannot be a color class when $c(u) \neq c(v)$ for all $c$, and a state with $s(u) \neq s(v)$ cannot be one when $u \sim_c v$, the qualifier $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$ is automatic: a chromatic TIFS is purely chromatic exactly when the ordinary TIFS fails, and a chromatic identity is two-way purely chromatic exactly when both directed ordinary TITS fail.

### B. Faithful Yu–Oh restrictions $H_{17,10}$ and $H_{21,13}$ and exhaustive inventory search

The base-coloring partition of $H_{\text{YO}}^{+}$ in Theorem 13 directly produces chromatically inseparable faithful restrictions when diagonal sectors are omitted. Retain the four base contexts $\{1, 2, 3\}$, $\{7, 8, 9\}$, $\{4, 5, 6\}$, $\{2, 8, 5\}$ of Eq. (15). Keeping the two diagonal sectors at 10 and 11 defines $H_{17,10}$ on 17 rays and 10 contexts [Fig. 1(a)]; keeping the three sectors at $10, 11, 13$ defines $H_{21,13}$ on 21 rays and 13 contexts [Fig. 1(b)]. Deleting $\{4, 5, 6\}$, $\{4, 11, 24\}$, $\{4, 10, 18\}$ from $H_{17,10}$ recovers the 13-ray, 7-context completed Specker bug $B_{13,7}$ [Fig. 2(b)].

**Theorem 15** (Chromatic inseparability in $H_{17,10}$ and $H_{21,13}$)**.** *Both $H_{17,10}$ and $H_{21,13}$ are linear of Berge girth five and inherit injective edge-faithful realizations in $\mathbb{R}^3$ from $H_{\text{YO}}^{+}$:*

**(a)** $H_{17,10}$ (17 rays, 10 contexts, 2 inseparable pairs)

**(b)** $H_{21,13}$ (21 rays, 13 contexts, 21 inseparable pairs)





FIG. 1. Faithful three-dimensional Yu–Oh restrictions in $\mathbb{R}^3$, drawn in the common vertex layout of completed Yu–Oh. (a) $H_{17,10}$, omitting diagonal sectors $h_0 = 12$ and $h_3 = 13$, with four normalized strong 3-colorings and chromatically inseparable pairs $10 \sim_c 24$ and $11 \sim_c 18$; (b) $H_{21,13}$, omitting only $h_0 = 12$, with two normalized strong 3-colorings and 21 chromatically inseparable pairs across nine chromatic signature classes.

1. *$H_{17,10}$ has* 18 *unital separating states,* 4 *normalized strong* 3*-colorings (Table S41), and* 10 *extendable states. Its only distinct chromatically inseparable pairs are* $(10, 24)$ *and* $(11, 18)$*. The ordinary implications* $10 \Rightarrow 24$ *and* $11 \Rightarrow 18$ *already hold, but their converses fail; these identities are not two-way purely chromatic. Relative to the anchor* $(c(2), c(8), c(5)) = (0, 1, 2)$*, its four colorings restrict on* $(9, 10, 11)$ *to* $(0, 2, 1)$*,* $(0, 2, 0)$*,* $(2, 1, 2)$*,* $(2, 0, 2)$*, yielding the anchored implications*

$$\begin{aligned} c(9) = 0 &\implies c(11) \in \{0, 1\}, \\ c(10) = 0 &\implies c(11) = 2. \end{aligned} \tag{18}$$

2. *$H_{21,13}$ has* 21 *unital separating states,* 2 *normalized strong* 3*-colorings, and* 6 *extendable states. Its nine chromatic signature classes* $(c_1(v), c_2(v)) \in \{0, 1, 2\}^2$ *are listed in Table S44, giving* $3\binom{4}{2} + 3\binom{2}{2} = 21$ *chromatically inseparable pairs. Exactly* 15 *are two-way purely chromatic identities (Table S70). The remaining six pairs obey the directed ordinary implications*

$$\begin{aligned} &11 \Rightarrow 18, \quad 11 \Rightarrow 21, \quad 13 \Rightarrow 15, \\ &13 \Rightarrow 19, \quad 10 \Rightarrow 20, \quad 10 \Rightarrow 24, \end{aligned}$$

*each with a false converse. This classification exhausts all* 21 *pairs; no distinct vertices are inseparable under all ordinary states.*

*Proof.* By Theorem 13(1), among the 8 base colorings in Eq. (16), sector 11 excludes $(1, 1, 0)$, $(0, 0, 1)$ and sector 10 excludes $(0, 0, 0)$, $(1, 1, 1)$. The 4 surviving $(b_1, b_2, b_3)$ extend uniquely to $H_{17,10}$ (Table S41): in each, $N(10) = \{1, 9, 4\}$ and $N(11) = \{3, 7, 4\}$ are 2-colored with $c(10) \neq c(4)$ and $c(11) \neq c(4)$, and $c(10) \neq c(11)$ by the restrictions to $(9, 10, 11)$ listed in (1), forcing $c(24) = c(10)$ in $\{4, 11, 24\}$ and $c(18) = c(11)$ in $\{4, 10, 18\}$. Adding sector 13 in $H_{21,13}$ additionally excludes $(0, 1, 1)$, $(1, 0, 0)$, leaving only $(1, 0, 1)$ and $(0, 1, 0)$, whose column signatures are listed in Table S44. Each retained diagonal adds 3 states to the 12 base states, giving $12 + 2 \times 3 = 18$ and $12 + 3 \times 3 = 21$ states (Sec. S2 of the supplementary material). Testing both directed truth patterns on these 21 states gives the exhaustive 15-versus-6 classification; Table S70 of the supplementary material supplies witnesses for all 15 two-way pairs, together with a reproducible enumeration procedure. □

The full chromatic signatures and both-direction truth witnesses are in the supplement. The witnesses are generated from the complete state families and checked against every context.

Physically, every noncontextual eigenvalue assignment on $H_{17,10}$ gives the nonorthogonal rays 10 and 24, and likewise 11 and 18, the same eigenvalue, although ordinary truth assignments distinguish them. In $H_{21,13}$, fifteen of the twenty-one forced identities are not even shadowed by a one-way truth implication: they exist only at the level of noncontextual eigenvalues.

**Proposition 16** (Complete inventory classification)**.** *Among all* $2^{16} - 1 = 65{,}535$ *nonempty context subsets* $E' \subseteq E(H_{\mathrm{YO}}^{+})$ *with inherited coordinates from Tables S1–S3, exactly* 539 *are connected and edge-faithful, and among these* precisely ten *are* 3*-colorable with a nontrivial chromatic inseparability class: the* $\binom{4}{2} = 6$ *two-sector restrictions (*17 *rays,* 10 *contexts,* 18 *states,* 4 *normalized colorings,* 2 *inseparable pairs) and the* $\binom{4}{3} = 4$ *three-sector restrictions (*21 *rays,* 13 *contexts,* 21 *states,* 2 *normalized colorings,* 21 *inseparable pairs). Thus* 17 *rays and* 10 *contexts are minimal for chromatic inseparability within the faithful restrictions of* $H_{\mathrm{YO}}^{+}$ *with inherited coordinates.*

*Proof.* Since $H_{\mathrm{YO}}^{+}$ is linear and edge-faithful, a subset $E'$ is edge-faithful if and only if no omitted context contains two vertices of $V' = \bigcup E'$. Exact enumeration by two independent coloring algorithms over all 539 connected faithful subsets certifies the ten solutions listed in Table S71. Disconnected restrictions separate distinct components by independent palette permutations. □

The ten retained-sector cases and binary search masks are enumerated in the supplement. The restriction to inherited coordinates matters: $I_{16,9}$ of Sec. IV C is $H_{17,10}$ without the context $\{4, 5, 6\}$; it has 16 rays and 9 contexts and keeps both identities, but it is edge-faithful only with refitted coordinates.

### C. Faithful three-dimensional TIFS, TITS, and purely chromatic gadgets

Figure 2 and Table II synthesize our faithful real 3-dimensional gadgets (coordinates and state/coloring tables in Secs. S1–S2 of the supplementary material):

- *Pentagon* $P_5$ *[Fig. 2(a)]:* Contexts $\{1, 6, 2\}$, $\{2, 7, 3\}$, $\{3, 8, 4\}$, $\{4, 9, 5\}$, $\{5, 10, 1\}$ have 5 normalized 3-colorings, 11 states, and 10 extendable states, separating all vertex pairs (showing that chromatic separation does not require $\mathcal{S}_{\mathrm{ext}} = \mathcal{S}$). Its degree-one vertices obey the purely chromatic ternary exclusion $\neg(c(6) = c(7) = c(9))$: assigning color 0 to $6, 7, 9$ forces $1, 2, 3, 4, 5$ into $\{1, 2\}$ with $c(1) = c(3)$ and $c(4) = c(5)$, contradicting $\{4, 9, 5\}$, whereas the nonextendable state $\{6, 7, 8, 9, 10\}$ (whose complement is the 5-cycle 12345) assigns 1 to all three terminals.

- *Specker bug* $B_{13,7}$ *and completed* $\Gamma_1$ $I_{16,9}$ *[Fig. 2(b),(d)]:* $B_{13,7}$ is an ordinary and chromatic TIFS ($c(10) \neq c(11)$, overlap 1/9). Adjoining $\{4, 11, 24\}$ and $\{4, 10, 18\}$ gives $I_{16,9}$ (deleting $\{4, 5, 6\}$ from $H_{17,10}$ with refitted faithful coordinates), where $c(10) \neq c(11)$ forces $c(4)$ to be the third color, so $c(24) = c(10)$ and $c(18) = c(11)$ via the one-way ordinary TITS $s(10) = 1 \implies s(24) = 1$ (overlap 25/27).

**Proposition 17** (Purely chromatic binary TIFS $D_{16,9}$ and two-way identity $T_{19,11}$)**.** *Let* $D_{16,9} = H_{17,10} \setminus \{\{1, 10, 19\}\}$ *(with orphan* 19 *removed) and* $T_{19,11} = D_{16,9} \cup \{\{14, 26, 27\}, \{18, 26, 28\}\}$*, realized faithfully in* $\mathbb{R}^3$ *[Fig. 2(c),(e)]:*

1. *Every* $c \in \mathcal{C}_3(D_{16,9})$ *satisfies* $c(14) \neq c(18)$ *(*7 *normalized colorings,* 14 *extendable states), whereas the nonextendable state* $\{1, 5, 7, 14, 18, 23, 24\} \in \mathcal{S}(D_{16,9})$ *assigns* $s(14) = s(18) = 1$ *(squared terminal overlap* 16/2025*).*





FIG. 2. Faithful real three-dimensional logical gadgets in $\mathbb{R}^3$. (a) Pentagon $P_5$ with purely chromatic ternary exclusion $\neg(c(6) = c(7) = c(9))$; (b) completed Specker bug $B_{13,7}$ (ordinary TIFS $c(10) \neq c(11)$); (c) purely chromatic binary TIFS $D_{16,9}$ ($c(14) \neq c(18)$); (d) completed historical $\Gamma_1$ TITS $I_{16,9}$ ($10 \sim_c 24, 11 \sim_c 18$); (e) two-way purely chromatic identity $T_{19,11}$ ($14 \sim_c 28, 18 \sim_c 27$); (f) $H_{21,13}$ showing three of its fifteen two-way purely chromatic identities: $2 \sim_c 14, 5 \sim_c 22, 8 \sim_c 23$. Red triangles forbid monochromatic triples; red dashed arrows denote $c(u) \neq c(v)$; blue dotted arrows denote $u \sim_c v$.

2. *Every* $c \in \mathcal{C}_3(T_{19,11})$ *satisfies* $14 \sim_c 28$ *and* $18 \sim_c 27$ *(7 normalized colorings, 25 states, 14 extendable), while for both pairs, nonextendable states* $\{1, 5, 7, 14, 18, 23, 24\}$ *and* $\{1, 6, 8, 10, 11, 27, 28\}$ *realize* $(1, 0)$ *and* $(0, 1)$*, violating both directed ordinary TITS implications (squared overlap* 2009/2025*; Table S70).*

*Proof.* (1) If $c(14) = c(18) = 0$, set $c(3) = 1, c(11) = 2$ by $\{3, 11, 14\}$. Then $\{4, 11, 24\}$ and $\{4, 10, 18\}$ force $c(4) = 1$ and $c(10) = 2$. In $\{7, 8, 9\}$, $7 \perp 11$ and $9 \perp 10$ avoid color 2, forcing $c(8) = 2$. Then $\{1, 2, 3\}$ and $\{4, 5, 6\}$ make 2 and 5 avoid color 1, so $\{2, 8, 5\}$ forces $c(2) = c(5) = 0$, a contradiction. (2) In $T_{19,11}$, $c(14) \neq c(18)$ forces their common neighbor 26 ($r_{26} \parallel r_{14} \times r_{18}$) to take the third color, whence $c(27) = c(18)$ and $c(28) = c(14)$. □

### D. Apex extensions and faithful four-dimensional linear gadgets

**Proposition 18** (Apex extension)**.** *Embedding any* 3*-uniform hypergraph* $H$ *realized in* $\mathbb{F}^3$ *into* $\mathbb{F}^d$ *(*$d \geq 3$*) and appending* $d - 3$ *mutually orthogonal coordinate apices to every context preserves injectivity, edge-faithfulness,* $\chi(H^{(d)}) = \chi(H) + d - 3$*, normalized coloring count, and old chromatic inseparability classes (with apices as singletons), while adding* $d - 3$ *apex-only states (all extendable when* $\chi(H) = 3$*) to the zero-extended states of* $H$*.*

*Proof.* In every coloring of $H^{(d)}$, the $d - 3$ apices occupy $d - 3$ distinct colors across all contexts, leaving a 3-coloring of $H$; similarly, any state either sets one apex to 1 (forcing all old vertices to 0) or sets all apices to 0 and restricts to $\mathcal{S}(H)$. □

TABLE II. Faithful geometric gadgets. $N_c = |\mathcal{C}_d|/d!$ fixes one context palette. Terminal relations describe colors unless explicitly identified as ordinary truth implications.

| Gadget | $d$ | $\lvert V\rvert/\lvert E\rvert$ | $N_c$ | $\lvert\mathcal{S}\rvert$ | $\lvert\mathcal{S}_{\text{ext}}\rvert$ | Relation and mechanism |
|---|---|---|---|---|---|---|
| $P_5$ | 3 | 10/5 | 5 | 11 | 10 | Ternary exclusion on $6, 7, 9$; purely chromatic. |
| $B_{13,7}$ | 3 | 13/7 | 6 | 14 | 12 | $c(10) \neq c(11)$; ordinary TIFS. |
| $D_{16,9}$ | 3 | 16/9 | 7 | 21 | 14 | $c(14) \neq c(18)$; purely chromatic TIFS. |
| $I_{16,9}$ | 3 | 16/9 | 6 | 22 | 12 | $10 \sim_c 24$, $11 \sim_c 18$; directed ordinary TITS. |
| $T_{19,11}$ | 3 | 19/11 | 7 | 25 | 14 | $14 \sim_c 28$, $18 \sim_c 27$; both truth directions fail. |
| $H_{21,13}$ | 3 | 21/13 | 2 | 21 | 6 | 21 identical pairs: 15 two-way purely chromatic; six directed ordinary TITS. |
| $H_{17,10}$ | 3 | 17/10 | 4 | 18 | 10 | Two color-identical pairs from directed ordinary TITS. |
| $W^{(4)}_{11,5}$ | 4 | 11/5 | 5 | 12 | 11 | Apex ternary exclusion. |
| $W^{(4)}_{16,6}$ | 4 | 16/6 | 80 | 72 | 56 | Linear ternary exclusion on $6, 8, 11$. |
| $B^{(4)}_{14,7}$ | 4 | 14/7 | 6 | 15 | 13 | Apex ordinary TIFS. |
| $L^{(4)}_{18,7}$ | 4 | 18/7 | 144 | 68 | 68 | Linear ordinary TIFS $c(11) \neq c(15)$. |
| $D^{(4)}_{17,9}$ | 4 | 17/9 | 7 | 22 | 15 | Apex purely chromatic TIFS. |
| $L^{(4)}_{22,9}$ | 4 | 22/9 | 216 | 158 | 92 | Linear identity $11 \sim_c 20$ from directed ordinary TITS. |

Applying Proposition 18 at $d = 4$ to $P_5$, $B_{13,7}$, and $D_{16,9}$ yields the non-linear ($g = 2$) four-dimensional gadgets $W^{(4)}_{11,5}$, $B^{(4)}_{14,7}$, and $D^{(4)}_{17,9}$ in Fig. 3(a),(c),(e). To achieve *linearity* ($|e \cap e'| \leq 1$) in $\mathbb{R}^4$, we construct three faithful linear gadgets ($g = 3$) in Fig. 3(b),(d),(f) (Table II):

1. *Linear purely chromatic ternary gadget* $W^{(4)}_{16,6}$ *[Fig. 3(b)]:* Contexts $\{2, 0, 6, 1\}$, $\{2, 7, 13, 3\}$, $\{3, 0, 8, 4\}$, $\{4, 9, 14, 5\}$, $\{5, 10, 15, 1\}$, $\{0, 11, 12, 5\}$ satisfy $\neg(c(6) = c(8) = c(11))$ (if $c(6) = c(8) = c(11) = 0$ and $c(0) = 3$, then $1, 2, 3, 4, 5$ must 2-color a 5-cycle), while $\{6, 7, 8, 9, 10, 11\} \in \mathcal{S}(W^{(4)}_{16,6})$ sets all three terminals to 1.

2. *Linear ordinary TIFS* $L^{(4)}_{18,7}$ *[Fig. 3(d)]:* Contexts $\{1, 2, 4, 3\}$, $\{5, 8, 7, 6\}$, $\{4, 9, 10, 8\}$, $\{11, 1, 12, 5\}$, $\{11, 14, 13, 2\}$, $\{15, 3, 16, 7\}$, $\{15, 17, 18, 6\}$ enforce $s(11) = 1 \implies s(15) = 0$ and $c(11) \neq c(15)$ (if $c(11) = c(15) = a$, then $1, 2, 3$ and $5, 6, 7$ avoid $a$, forcing $c(4) = c(8) = a$ in $\{4, 9, 10, 8\}$). Using $r_{13} = (1, 0, 1, 1), r_{14} = (2, 0, -1, -1)$ avoids an unlisted orthogonality $13 \perp 15$; all 68 states are extendable (terminal overlap 1/42).

3. *Linear directed TITS and color identity* $L^{(4)}_{22,9}$ *[Fig. 3(f)]:* Adjoining $\{11, 21, 22, 19\}$ and $\{13, 20, 15, 19\}$ to $L^{(4)}_{18,7}$ (with $r_{13} = (1, 0, 4, 4), r_{14} = (8, 0, -1, -1)$ so $r_{13} \perp r_{15}$) makes $\{11, 13, 15, 19\}$ pairwise differently colored, forcing $11 \sim_c 20$ and $s(11) = 1 \implies s(20) = 1$ (while $\{1, 6, 9, 14, 16, 20, 21\}$ has $s(11) = 0, s(20) = 1$; terminal overlap 41/42).

## V. THE 37-RAY COMPLEX HESSE HYPERGRAPH AND RATIONAL FOUR-DIMENSIONAL COMPLETIONS

### A. The 37-ray complex Hesse hypergraph $H_{37}$ and its 76-event orthomodular lattice

With $\zeta = \exp(2\pi i/3)$ ($\zeta^2 + \zeta + 1 = 0$) and $a, k \in \mathbb{Z}_3$, define 37 rays in $\mathbb{C}^3$ (core rays $z, M_{ak}, P_k, Q_k$ and unique cross-product completion rays $A_k, B_{ak}, C_{ak}$, numbered $1, \ldots, 37$





FIG. 3. Faithful real four-dimensional gadgets in $\mathbb{R}^4$. Panels (a),(c),(e) are common-apex lifts ($g(H) = 2$); panels (b),(d),(f) are linear ($g(H) = 3$). (a),(b) Purely chromatic ternary exclusions; (c),(d) ordinary TIFS; (e) purely chromatic binary TIFS $c(14) \neq c(18)$; (f) linear color identity $11 \sim_c 20$ from directed ordinary TITS $s(11) = 1 \implies s(20) = 1$.

via $z \leftrightarrow 1, M_{ak} \leftrightarrow 2{+}3a{+}k, P_k \leftrightarrow 11{+}k, Q_k \leftrightarrow 14{+}k, A_k \leftrightarrow 17{+}k, B_{ak} \leftrightarrow 20{+}3a{+}k, C_{ak} \leftrightarrow 29 + 3a + k$):

$$\begin{aligned} z &= (1,0,0), & M_{ak} &= (1, \zeta^{a+k}, \zeta^{a+2k}), \\ P_k &= (0,1,-\zeta^k), & Q_k &= (-\zeta^k, 0, 1), \\ A_k &= (0,1,\zeta^k), & B_{ak} &= (-2, \zeta^{a+k}, \zeta^{a+2k}), \\ C_{ak} &= (1, -2\zeta^{2a+k}, \zeta^{2k}), \end{aligned} \tag{19}$$

and the 24 complete orthogonal triad contexts (Fig. 4):

$$\begin{aligned} \mathcal{M}_a &= \{M_{a0}, M_{a1}, M_{a2}\}, & \mathcal{A}_k &= \{z, P_k, A_k\}, \\ \mathcal{B}_{ak} &= \{M_{ak}, P_k, B_{ak}\}, & \mathcal{C}_{ak} &= \{M_{a,a+k}, Q_k, C_{ak}\}. \end{aligned} \tag{20}$$

**Theorem 19** (Properties of $H_{37}$)**.** *The hypergraph $H_{37}$ defined by Eqs. (19)–(20) satisfies:*

1. *It is linear with Berge girth five,* 72 *edges, and $\omega(\Gamma(H_{37})) = 3$; its triangles are precisely its* 24 *contexts.*

2. *Equation (19) defines an injective edge-faithful representation in $\mathbb{C}^3$, whereas $H_{37}$ admits* no *injective orthogonal representation in $\mathbb{R}^3$.*

3. *Its chromatic numbers are $\chi(H_{37}) = 4 > 3 = \chi_f(\Gamma(H_{37})) = \omega(\Gamma(H_{37}))$.*

4. *Its* 24 *Boolean blocks paste into a* 76*-event orthomodular lattice $L(H_{37}) = \{0,1\} \cup \{a_i, a_i^{\perp} : 1 \le i \le 37\}$. Its full state space has $|\mathcal{S}(H_{37})| = 153$ states and is order determining on $L(H_{37})$, with exact minimum state-family sizes* 6 *for unital vertex separation,* 7 *for separation of all* 76 *events, and* 18 *for order determination.*





FIG. 4. Complete 3-uniform hypergraph diagram of $H_{37}$ (37 rays, 24 orthogonal triads, 72 edges in $\mathbb{C}^3$). Bold circles denote $e_1, e_2, e_3$ $(\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2)$; the other 21 smooth traces join core pairs through their unique completion rays $17, \ldots, 37$. Concentric rings record incident context colors.

*Proof.* (1)–(2) Zero patterns, magnitudes, and $\zeta$-phases distinguish all 37 rays. Algebraic inner products confirm that $\langle v_i, v_j \rangle = 0$ holds strictly for the $24\binom{3}{2} = 72$ context pairs: $\langle M_{ak}, M_{c\ell} \rangle = 1 + \zeta^{c-a}(\zeta^{\ell-k} + \zeta^{2(\ell-k)}) = 0 \iff (c = a, \ell \neq k)$; $P_j \perp M_{ak} \iff j = k$; $Q_j \perp M_{ak} \iff k = a + j$; $\langle M_{c\ell}, B_{ak} \rangle = -2 + \zeta^{a-c}(\zeta^{k-\ell} + \zeta^{2(k-\ell)}) = 0 \iff (c, \ell) = (a, k)$; $\langle M_{c\ell}, C_{ak} \rangle = 1 - 2\zeta^{2a+k-c-\ell} + \zeta^{2k-c-2\ell} = 0 \iff (c = a, \ell = a + k)$; and unequal term magnitudes exclude orthogonalities among $A, B, C$. Uniqueness of cross-product completions in $\mathbb{C}^3$ excludes Berge 3- and 4-cycles, while $(\mathcal{A}_0, \mathcal{B}_{00}, \mathcal{M}_0, \mathcal{B}_{01}, \mathcal{A}_1)$ is a Berge 5-cycle.

To exclude any injective representation in $\mathbb{R}^3$, suppose unit $z \in \mathbb{R}^3$ and real $p_k, m_{ak} \in \mathbb{R}^3$ realize $\mathcal{A}_k, \mathcal{B}_{ak}, \mathcal{M}_a$. If $m_{ak} \cdot z = 0$, then $\{z, p_k, m_{ak}\}$ is an orthogonal basis of $\mathbb{R}^3$, and for $\ell \neq k$, $p_\ell \perp z$ and $m_{a\ell} \perp m_{ak}$ with $p_\ell \perp m_{a\ell}$ force $[p_\ell] = [m_{ak}]$ or $[m_{a\ell}] = [z]$. Hence $m_{ak} \cdot z \neq 0$. Rescaling so $m_{ak} \cdot z = 1$ and choosing unit $u_k \in z^\perp \cap p_k^\perp$ gives $m_{ak} = z + t_{ak} u_k$ $(t_{ak} \in \mathbb{R} \backslash \{0\})$. Orthogonality of $\mathcal{M}_a$ yields $1 + t_{ak} t_{a\ell}(u_k \cdot u_\ell) = 0$ $(k \neq \ell)$. Setting $q_k = t_{ak}/t_{0k}$ gives $q_0 q_1 = q_0 q_2 = q_1 q_2 = 1$, whose only real solutions are $(q_0, q_1, q_2) = \pm(1, 1, 1)$. Thus at most two distinct bases $\mathcal{M}_a$ exist in $\mathbb{R}^3$, contradicting injectivity on the three bases $\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$. The argument uses only the 13 rays $z, P_k, M_{ak}$ and the orthogonalities among them, so already the 13-vertex subgraph of $\Gamma(H_{37})$ induced by these rays has no injective orthogonal representation in $\mathbb{R}^3$.

(3) In any 3-coloring with $c(z) = 0$, $P_0, P_1, P_2 \in \{1, 2\}$ cannot be monochromatic (as

$\mathcal{M}_a$ uses all three colors and $M_{ak} \sim P_k$). Up to symmetry $(c(P_0), c(P_1), c(P_2)) = (2, 1, 1)$, forcing $c(M_{a0}) = 1$ and $(c(M_{a1}), c(M_{a2})) = (x_a, 2 - x_a)$ with $x_a \in \{0, 2\}$. Since $N(Q_k) = \{M_{0k}, M_{1,k+1}, M_{2,k+2}\}$ avoids a color, $x_1 = 2 - x_2$, $x_0 = 2 - x_1$, $x_2 = 2 - x_0$, impossible on a 3-cycle. Thus $\chi(H_{37}) = 4$ (attained by $c(z) = 0, c(M_{ak}) = k, c(P_k) = c(Q_k) = 3$). The 6-state double cover $\mathcal{R}_6 = \{s_2, s_{30}, s_{73}, s_{99}, s_{109}, s_{139}\}$ covers every vertex twice, giving $\chi_f(\Gamma(H_{37})) = 3$.

(4) Each state is fixed by $t = (t_0, t_1, t_2) \in \mathbb{Z}_3^3$ with $s(M_{ak}) = \mathbf{1}_{k=t_a}$, $s(z) = \varepsilon \in \{0, 1\}$, and independent bits on $U(t) = \mathbb{Z}_3 \setminus \{t_0, t_1, t_2\}$ (for $\varepsilon = 0$) and $W(t) = \mathbb{Z}_3 \setminus \{t_a - a\}$, yielding $(1 + 2^{|U(t)|})2^{|W(t)|}$ states per $t$:

| Class of $t \in \mathbb{Z}_3^3$ | $\lvert\#t\rvert$ | $(\lvert U(t)\rvert, \lvert W(t)\rvert)$ | $(1 + 2^{\lvert U\rvert})2^{\lvert W\rvert}$ |
|---|---|---|---|
| $(c, c, c)$ | 3 | $(2, 0)$ | $5 \times 1 = 5$ |
| $(c, c+1, c+2)$ | 3 | $(0, 2)$ | $2 \times 4 = 8$ |
| $(c, c+2, c+1)$ | 3 | $(0, 0)$ | $2 \times 1 = 2$ |
| Two equal entries | 18 | $(1, 1)$ | $3 \times 2 = 6$ |

(21)

and hence $|\mathcal{S}(H_{37})| = 3 \cdot 5 + 3 \cdot 8 + 3 \cdot 2 + 18 \cdot 6 = 153$. Since $2^5 - 1 = 31 < 37$ and $2^6 = 64 < 76$, the explicit families $\Omega_6$ and $\Omega_7$ given in the supplement prove the minima 6 and 7. For order determination ($a_i \le a_j^\perp \iff i \sim j$), the 15 nonorthogonal pairs in $D = \{P_0, P_1, P_2, Q_0, Q_1, Q_2\}$ require 15 disjoint joint-one witnesses in $T \subset \mathcal{S}(H_{37})$ ($|T| = 27$), and the 9 pairs $\{z, C_{ak}\}$ require at least 2 states outside $T$. Exhaustive bit-set verification over all $4{,}096 \times 132 = 540{,}672$ candidate 17-state families shows each meets at most 75 of the 81 minimal witness sets $W_{xy}$, whereas the 18-state family $\Omega_{18}$ in the supplement meets all 81 (Sec. S3 of the supplementary material). □

Theorem 19(2) is not the first finite separation of complex from real three-dimensional orthogonality: Harding and Salinas Schmeis[34] (Thm. 2.6) construct, from mutually unbiased bases, a two-gadget configuration that is realizable in $\mathbb{C}^3$ but not in $\mathbb{R}^3$ (48 rays, 32 bases, 189 context states, and 180 geometric states). $H_{37}$ is smaller, linear of Berge girth five, and edge-faithful in $\mathbb{C}^3$, and its real obstruction already resides in the 13-ray subgraph on $z, P_k, M_{ak}$.

By Proposition 18, appending $d - 3$ common apices to $H_{37}$ in $\mathbb{C}^d$ ($d \ge 3$) gives faithful $d$-uniform hypergraphs with $|V| = d + 34$, $|E| = 24$, $\omega = d$, $\chi = d + 1$, and $|\mathcal{S}| = d + 150$.

An order-determining family represents the specified 76-event order, orthocomplements, and Boolean operations within listed contexts. It does not preserve arbitrary meets as set intersections: nonorthogonal atoms have meet zero although some states make both true. It is therefore not a Boolean-sublattice embedding of the full lattice, nor a claim of closure under every Hilbert operation. In particular, the minimum six-state separating family and the six-state uniform double cover solve different problems.

### B. Hyperplane obstruction, two-plane completion lemma, and canonical linear $H_{65}$ in $\mathbb{R}^4$

Appending a single apex $v_{26} = (0, 0, 0, 1)^T \in \mathbb{R}^4$ to $H_{\mathrm{YO}}^+$ gives the 26-ray, 16-context non-linear suspension $H_{\mathrm{YO}}^{(4)}$ (Fig. 5) with $\omega = 4, \chi = 5$, and 25 unital separating states (Proposition 18), but $|e \cap e'| = 2$ ($g = 2$). Restricted to the 13 Yu–Oh rays and the apex, its orthogonality graph is the Yu–Oh graph joined to one common vertex, the example that Cabello, Kleinmann, and Budroni[15] (Thm. 1) used to show that $\chi_f > d$ is necessary but not sufficient for state-independent contextuality; $H_{65}$ below is a linear, apex-free example of the same phenomenon.

**Proposition 20** (Hyperplane obstruction to linearity in $\mathbb{F}^4$)**.** *In any injective realization in $\mathbb{P}(\mathbb{F}^4)$, if two contexts $e, e'$ each have three rays in a common 3-dimensional hyperplane $W = z^\perp$, both must contain $[z]$; hence if $|e_W \cap e'_W| \ge 1$, then $|e \cap e'| \ge 2$, violating linearity.*





FIG. 5. The completed 25-ray, 16-context Yu–Oh hypergraph $H_{\mathrm{YO}}^{+}$ in $\mathbb{R}^3$ ($\omega = 3, \chi = 4$) and its non-linear 26-ray, 16-context apex suspension $H_{\mathrm{YO}}^{(4)}$ in $\mathbb{R}^4$ ($\omega = 4, \chi = 5, g = 2$) obtained by adjoining the common orthogonal apex $v_{26} = (0, 0, 0, 1)^T$ to every context.

*Proof.* Three mutually orthogonal rays in $W$ span $W$, forcing the fourth orthogonal ray of both $e$ and $e'$ to equal $[z] = W^{\perp}$. □

**Lemma 21** (Simultaneous two-plane completion in $\mathbb{R}^4$)**.** *Let $V_{\mathrm{core}} \subset \mathbb{Q}^4 \setminus \{0\}$ be a finite set of non-collinear rational vectors with 4-element orthogonal base contexts $E_{\mathrm{base}}$ ($|C \cap C'| \leq 1$) and let all remaining orthogonal pairs be the bridge pairs $E_{\mathrm{bridge}} = \{e_1, \ldots, e_M\}$ ($e_m = \{u_m, v_m\}$, $L_m = \mathrm{span}(v_{u_m}, v_{v_m})$, $P_m = L_m^{\perp}$) such that every core vertex lies in a base context or is a bridge endpoint.*

1. *The pairs $e_m$ can be simultaneously completed to 4-bases $E_m = \{u_m, v_m, w_{m,1}, w_{m,2}\}$ by fresh integer vectors $w_{m,1}, w_{m,2} \in P_m \cap \mathbb{Z}^4$ forming a linear edge-faithful hypergraph $H$ if and only if*

$$\begin{aligned} v_z &\notin L_m \quad (z \in V_{\mathrm{core}} \setminus e_m), \\ P_\ell &\neq L_m \quad (\ell \neq m). \end{aligned} \tag{22}$$

2. *$\omega(\Gamma(H)) = 4$ and $\chi(H) = \max\{4, \chi(G_{\mathrm{core}})\}$; here $G_{\mathrm{core}}$ is automatically $K_5$-free, since no five nonzero vectors of $\mathbb{R}^4$ are mutually orthogonal.*

3. *If all $G_{\mathrm{core}}$-independent base supports $S_1, \ldots, S_N$ form a unital separating family on the core, and the zero-profiles $Z(e_m) = \{\alpha : S_\alpha \cap e_m = \varnothing\}$ are nonempty and pairwise distinct, the $\leq 2N$ canonical extensions $s_{\alpha,\varepsilon}$ ($\varepsilon \in \{0, 1\}$, setting $(s_{\alpha,\varepsilon}(w_{m,1}), s_{\alpha,\varepsilon}(w_{m,2})) = (1 - \varepsilon, \varepsilon)$ when $\alpha \in Z(e_m)$ and $(0, 0)$ otherwise) are unital and separating on $V(H)$.*

*Proof.* If $v_z \in L_m$ then $P_m \subset v_z^{\perp}$, and if $P_\ell = L_m$ then $P_\ell \perp P_m$, forcing extra orthogonalities. Conversely, given Eq. (22), an integer orthogonal basis $\{\eta_A, \eta_B\} \subset P_m \cap \mathbb{Z}^4$ defines the infinite family $w_{m,1}(a, b) = a\eta_A + b\eta_B$, $w_{m,2}(a, b) = -\|\eta_B\|^2 b\eta_A + \|\eta_A\|^2 a\eta_B$, avoiding inductively the finitely many 1-dimensional intersections $z^{\perp} \cap P_m$ and $\mathbb{R}z \cap P_m$, where $z$ ranges over the core rays *and* the fresh rays placed at earlier steps, together with $P_m \cap L_r$ ($r > m$). The last condition keeps every fresh ray outside each later $L_r$, so that $z^{\perp} \cap P_r$ remains 1-dimensional for fresh $z$ as well. Linearity, coloring extension, and distinctness of canonical vertex profiles follow from Proposition 7 and distinctness of $Z(e_m)$. □





FIG. 6. Canonical linear 4-uniform hypergraph $H_{65}$ in $\mathbb{R}^4$ (65 rays, 29 contexts, 174 edges, $g = 3, \omega = 4, \chi = 5$). Thick colored curves denote the five core contexts $C_1, \ldots, C_5$; the 24 thinner lines denote the completed bridge contexts $E_1, \ldots, E_{24}$ carrying degree-one completion pairs $\{18, 19\}, \ldots, \{64, 65\}$.

To construct a *linear* 4-uniform hypergraph in $\mathbb{R}^4$, we take the 17-ray core $V_{\text{core}} = \{1, \ldots, 17\} \subset \{-1, 0, 1\}^4$, which is orthogonally equivalent to the 17-ray Peres restriction of Haynes *et al.*[19] (pp. 8–9) via the orthogonal matrix

$$T = \frac{1}{2}\begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & -1 & -1 & -1 \end{pmatrix}, \qquad T^T T = I_4. \tag{23}$$

The core consists of the coordinate basis $C_1$, an even-parity Hadamard basis $C_2$, three odd-parity hub rays $9, 10, 11$, and the three binary bridge pairs $B_1 = \{12, 13\}$, $B_2 = \{14, 15\}$, $B_3 = \{16, 17\}$. The remaining base contexts are $C_3 = \{9, 10, 12, 13\}$, $C_4 = \{9, 11, 14, 15\}$ and $C_5 = \{10, 11, 16, 17\}$. Complete coordinates and all compatible supports are in Tables S17–S18 and S81 of the supplementary material. Completing its 24 uncovered bridge edges $e_m \in (C_1 \cup C_2) \times (B_1 \cup B_2 \cup B_3)$ by fresh integer pairs $\{2m+16, 2m+17\} \subset \{18, \ldots, 65\}$ via Lemma 21 yields $H_{65} = (V_{65}, E_{29})$ (Fig. 6; Tables S17–S18 of the supplementary material):

$$\begin{aligned} E_{29} &= \{C_1, \ldots, C_5\} \cup \{E_m : 1 \le m \le 24\}, \\ E_m &= \{u_m, v_m, 2m + 16, 2m + 17\}. \end{aligned} \tag{24}$$

**Theorem 22** (Properties of $H_{65}$)**.** *The 65-ray, 29-context hypergraph $H_{65}$ satisfies:*

1. *It is linear ($g(H_{65}) = 3$) and edge-faithful in $\mathbb{R}^4$ (174 edges) with $\omega(\Gamma(H_{65})) = 4$, and nonconformal (13 uncovered maximal triangles: 12 bridge triangles plus $\{9, 10, 11\}$).*

2. *It has $\alpha(G_{\text{core}}) = 4$, $\chi_f(\Gamma(H_{65})) = 17/4 = 4.25 > 4$, and $\chi(H_{65}) = 5$.*

TABLE III. Bridge neighbors in the two core bases. The complete 24 state supports are supplementary.

| $b$ | $N(b)\cap C_1$ | $N(b)\cap C_2$ |
|---|---|---|
| 12 | $\{2,3\}$ | $\{5,8\}$ |
| 13 | $\{1,4\}$ | $\{6,7\}$ |
| 14 | $\{2,4\}$ | $\{6,8\}$ |
| 15 | $\{1,3\}$ | $\{5,7\}$ |
| 16 | $\{3,4\}$ | $\{7,8\}$ |
| 17 | $\{1,2\}$ | $\{5,6\}$ |

3. *Its core has* 24 *compatible state supports* $S_\alpha = \{i_1, i_2, h, b\}$ *(Table S81), each missing* 14 *bridges and yielding* $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$ *states and* 48 *canonical separating states.*

4. *Every* $s \in \mathcal{S}(H_{65})$ *obeys* $s(9) + s(10) + s(11) = 1$*, so adjoining the fourth hub ray* $q = (1,1,1,-1)^T$ *forces* $\widetilde{s}(q) = 0$*; moreover,* $A_9 = A_{16} \dot\cup A_{17}$ *on all of* $\mathcal{S}(H_{65})$*, forcing the false atom-order implication* $s(16) \le s(9)$ *and collapsing the* 1*-ray* $[v_9] = [v_{16} - v_{17}]$ *with the* 2*-plane* $\mathrm{span}(v_{16}, v_{17})$*.*

*Proof.* (1)–(2) No $K_4$ in $G_{\mathrm{core}}$ contains a bridge edge $\{u_m, v_m\}$, since $u_m \in C_a$ and $v_m \in B_k$ have exactly one common neighbor in $G_{\mathrm{core}}$, namely the second vertex of $N(v_m) \cap C_a$; this verifies $P_\ell \neq L_m$, and support-size / magnitude checks verify $v_z \notin L_m$ in Lemma 21. For any independent $I \subseteq V_{\mathrm{core}}$, $|I \cap C_1| \le 1$ and $|I \cap C_2| \le 1$. If $I \cap \{9,10,11\} \neq \varnothing$, $|I \cap (C_3 \cup C_4 \cup C_5)| \le 2$ so $|I| \le 4$; if $I \cap \{9,10,11\} = \varnothing$ and $I$ picks $(b_1, b_2, b_3) \in B_1 \times B_2 \times B_3$, even-parity triples cover $C_2$ and odd-parity triples cover $C_1$ (Table III), so again $|I| \le 3{+}1 = 4$. Thus $\alpha(G_{\mathrm{core}}) = 4$, $\chi_f \ge 17/4$ (matched by weights $1/8$ on the 24 core states and $1/16$ on the 20 remaining independent 4-sets), and $\chi(H_{65}) = 5$. (3)–(4) By parity, no state can avoid $\{9,10,11\}$, giving $s(9) + s(10) + s(11) = 1$, $3 \times 2 \times 2 \times 2 = 24$ core supports (Table S81), $24 - 10 = 14$ missed bridges per support, and $A_9 = A_{16} \dot\cup A_{17}$. □

## VI. EVENT ORDER, QUANTUM PROBABILITIES, AND NONEXTENDABLE WEIGHT

### A. Extendable state polytopes and nonextendable weight on $H_{17,10}$

For any context hypergraph $H$, the classical state polytope $\mathcal{P}(H) = \mathrm{conv}\,\mathcal{S}(H)$ and the extendable state polytope $\mathcal{P}_{\mathrm{ext}}(H) = \mathrm{conv}\,\mathcal{S}_{\mathrm{ext}}(H)$ satisfy $\mathcal{P}_{\mathrm{ext}}(H) \subseteq \mathcal{P}(H)$, with $p_u = p_v$ for all $p \in \mathcal{P}_{\mathrm{ext}}(H)$ whenever $u \sim_c v$ by Corollary 4. For $p \in \mathcal{P}(H)$, define its *minimum nonextendable weight*

$$\begin{aligned} \eta_H(p) = \min_{\mu} \quad & \sum_{s \notin \mathcal{S}_{\mathrm{ext}}} \mu_s, \\ \text{subject to} \quad & p = \sum_{s \in \mathcal{S}} \mu_s s, \\ & \mu_s \ge 0, \qquad \sum_s \mu_s = 1. \end{aligned} \tag{25}$$

**Proposition 23** (Nonextendable weight lower bound)**.** *For any chromatically inseparable pair* $u \sim_c v$ *in* $H$ *and any* $p \in \mathcal{P}(H)$*,* $\eta_H(p) \ge |p_u - p_v|$*.*

*Proof.* Every $s \in \mathcal{S}_{\mathrm{ext}}(H)$ satisfies $s(u) - s(v) = 0$, while every $s \notin \mathcal{S}_{\mathrm{ext}}(H)$ satisfies $|s(u) - s(v)| \le 1$, so $|p_u - p_v| \le \sum_{s \notin \mathcal{S}_{\mathrm{ext}}(H)} \mu_s |s(u) - s(v)| \le \sum_{s \notin \mathcal{S}_{\mathrm{ext}}(H)} \mu_s$. □

**Theorem 24** (Quantum preparations in $\mathcal{P}(H_{17,10}) \setminus \mathcal{P}_{\text{ext}}(H_{17,10})$)**.** *In the real FOR of $H_{17,10}$ (Table S2 of the supplementary material):*

1. *The pure quantum preparation along $r_{24} = (1, -1, 2)^T$,*

$$\rho_\star = \frac{1}{6}\begin{pmatrix} 1 & -1 & 2 \\ -1 & 1 & -2 \\ 2 & -2 & 4 \end{pmatrix}, \qquad p_i = \mathrm{Tr}(\rho_\star P_i) = \frac{(r_{24} \cdot r_i)^2}{6\|r_i\|^2}, \tag{26}$$

*belongs to $\mathcal{P}(H_{17,10})$ via the exact five-state convex combination (state indices in Table S40 of the supplementary material; see also Tables S46–S47)*

$$p = \frac{1}{12}s_4 + \frac{5}{36}s_9 + \frac{1}{36}s_{10} + \frac{7}{12}s_{14} + \frac{1}{6}s_{17}, \tag{27}$$

*yet satisfies $p_{24} = 1$ and $p_{10} = 8/9$ for the chromatically inseparable pair $(10, 24)$, so*

$$p \notin \mathcal{P}_{\text{ext}}(H_{17,10}) \qquad \text{and} \qquad \eta_{H_{17,10}}(p) = \frac{1}{9}. \tag{28}$$

2. *The pure preparation along the coordinate ray $z_3 = r_5 = (0, 0, 1)^T$, with Born vector $p'_i = (r_5 \cdot r_i)^2/\|r_i\|^2$, has $p'_{24} = p'_{18} = 2/3$ and $p'_{10} = p'_{11} = 1/3$, belongs to $\mathcal{P}(H_{17,10})$ via the four-state convex combination $p' = \frac{1}{3}s_3 + \frac{1}{6}s_4 + \frac{1}{3}s_{14} + \frac{1}{6}s_{15}$, and satisfies $\eta_{H_{17,10}}(p') = 1/3$.*

3. *For every density operator $\rho$ and both inseparable pairs $(u, v) \in \{(10, 24), (11, 18)\}$, $|p_u(\rho) - p_v(\rho)| \leq \|P_u - P_v\| = 1/3$. Hence Proposition 23 certifies at most $1/3$ on $H_{17,10}$, and the preparation in (2) attains this value.*

*Proof.* (1) Direct evaluation of Eq. (27) on all 17 vertices matches $p_i$ in Eq. (26) (Table S46), with $r_{10} = (1, -1, 1)^T$ giving $p_{10} = 4^2/(6 \cdot 3) = 8/9$. Since $10 \sim_c 24$, Proposition 23 gives $\eta_{H_{17,10}}(p) \geq |1-8/9| = 1/9$, which is attained in Eq. (27) because among $\{s_4, s_9, s_{10}, s_{14}, s_{17}\}$ precisely $s_4$ and $s_{10}$ are nonextendable, with total weight $1/12+1/36 = 1/9$. (2) Direct evaluation of the four-state mixture on all 17 vertices reproduces $p'$ (Tables S46–S47). Proposition 23 gives $\eta_{H_{17,10}}(p') \geq |2/3 - 1/3| = 1/3$, attained because $s_3, s_{14}$ are extendable and $s_4, s_{15}$ are not, with total weight $1/6 + 1/6 = 1/3$. (3) For unit vectors with squared overlap $\kappa$, $P_u - P_v$ has eigenvalues $\pm\sqrt{1-\kappa}$ and 0. Both pairs have $\kappa = (r_u \cdot r_v)^2/(\|r_u\|^2\|r_v\|^2) = 16/18 = 8/9$, so $|\,\mathrm{Tr}\,\rho(P_u - P_v)| \leq \|P_u - P_v\| = 1/3$. □

The exact support-and-weight certificates and the complete Born probability vectors are given in the supplement. These tables are regenerated from the checked state inventory, rather than manually transcribed.

These preparations exclude only models built from noncontextual eigenvalue assignments: their ordinary-state models are exhibited explicitly. A state is extendable exactly when some noncontextual eigenvalue assignment $c$ makes it the indicator of a single eigenvalue $\lambda_a$ in every context observable $A_e$ of Eq. (1). For a model built from noncontextual eigenvalue assignments, every chromatic identity $u \sim_c v$ entails the equality $p_u = p_v$; quantum mechanics violates it, and Proposition 23 converts the violation into a lower bound on $\eta$. The weight $\eta$ therefore measures how far the quantum predictions force any noncontextual truth-assignment model away from noncontextual eigenvalues: on $H_{17,10}$, at least one third of the weight in the preparation along $z_3$. Since $\mathcal{P}_{\text{ext}}(H_{17,10})$ lies in the hyperplane $p_{10} = p_{24}$, a Born vector in $\mathcal{P}(H_{17,10})$ generically lies outside $\mathcal{P}_{\text{ext}}(H_{17,10})$; the content of Theorem 24 is the size of the required weight. Conversely, averaging the $d$ color classes of any admissible coloring gives the maximally mixed Born vector $1/d$, so the restricted obstruction is preparation dependent. The $H_{17,10}$ ordinary implication $s(10) \leq s(24)$ also prevents order

determination for distinct ray atoms; vertex separation and order representation must not be conflated.

For a single system, nothing forces a classical model to use only extendable states. The supersinglet game of Sec. I does. A local hidden-variable model that reproduces its perfect correlations gives the $d$ parties the $d$ color classes of a strong $d$-coloring, so each party follows an extendable state. If $d-1$ parties measure the same basis and obtain all of its rays but $v$, the last particle is prepared along $v$, and in such a model its outcome probabilities lie in $\mathcal{P}_{\mathrm{ext}}(H)$. On $H_{17,10}$ the two preparations of Theorem 24 arise in this way from the bases $\{4,11,24\}$ and $\{2,8,5\}$. Both lie outside $\mathcal{P}_{\mathrm{ext}}(H_{17,10})$, so quantum mechanics contradicts every such local model, although $\chi(H_{17,10})=3$. More directly, if $u\sim_c v$, such a model never lets two different parties obtain $u$ and $v$. Quantum mechanics gives this event the probability $(1-\kappa)/[d(d-1)]$, where $\kappa$ is the squared overlap of the two rays, because the reduced state of two particles of the supersinglet is the normalized projector onto the antisymmetric subspace. For the pairs $(10,24)$ and $(11,18)$ of $H_{17,10}$ this probability is $1/54$.

### B. Order determination versus Born probabilities on $H_{37}$, $H_{\mathrm{YO}}^{(4)}$, and $H_{65}$

When $\chi(H)>d$ (Regime I), no noncontextual eigenvalue assignment exists, $\mathcal{P}_{\mathrm{ext}}(H)=\varnothing$, and the relevant probability question is whether Born assignments $p_i(\rho)=\mathrm{Tr}(\rho\Pi_i)$ lie in $\mathcal{P}(H)=\mathrm{conv}\,\mathcal{S}(H)$ (Table IV).

**Corollary 25** (Order determination does not imply classical probability models)**.** *The* 76*-event orthomodular lattice $L(H_{37})$ admits an order-determining state space $\mathcal{S}(H_{37})$ and a classical mixture for the maximally mixed state $I_3/3$, yet every $p\in\mathcal{P}(H_{37})$ satisfies the tight inequality*

$$\mathcal{C}_{37}(p)=\sum_{k=0}^{2}\big[p(P_k)+p(Q_k)\big]\le 2, \tag{29}$$

*whereas the complex FOR of $H_{37}$ satisfies*

$$\sum_{k=0}^{2}(\Pi_{P_k}+\Pi_{Q_k})=\mathrm{diag}(3/2,3/2,3)=\frac{3}{2}I_3+\frac{3}{2}|e_3\rangle\langle e_3|, \tag{30}$$

*yielding $\mathcal{C}_{37}(\rho)=\frac{3}{2}(1+\rho_{33})>2$ whenever $\rho_{33}>1/3$ (and $\mathcal{C}_{37}(|e_3\rangle\langle e_3|)=3$).*

*Proof.* In the state parametrization of Theorem 19(4), every $t\in\mathbb{Z}_3^3$ has $|U(t)|+|W(t)|\le 2$, establishing Eq. (29), while summing the normalized projectors of $P_k,Q_k$ in Eq. (19) using $1+\zeta+\zeta^2=0$ gives Eq. (30). Uniform weights $1/6$ on $\mathcal{R}_6$ reproduce $p_i(I_3/3)=1/3$. □

In dimension four, the non-linear suspension $H_{\mathrm{YO}}^{(4)}$ obeys $\mathcal{Y}(p)=p_{26}+\sum_{i=10}^{13}p_i\le 1$ on $\mathcal{P}(H_{\mathrm{YO}}^{(4)})$, whereas $\sum_{i=10}^{13}\Pi_i=\frac{4}{3}(I_4-\Pi_{26})$ gives $\mathcal{Y}(\rho)=1+(1-p_{26}(\rho))/3>1$ for every density operator $\rho\neq\Pi_{26}$[15]. For the canonical linear $H_{65}$, every $s\in\mathcal{S}(H_{65})$ has 4 active core vertices, so every $p\in\mathcal{P}(H_{65})$ obeys $\sum_{i=1}^{17}p_i=4$. However, with $q=(1,1,1,-1)^T$ and $Q=qq^T/4$, direct summation gives

$$\sum_{i=1}^{17}\Pi_i=4I_4+Q\quad\Longrightarrow\quad\sum_{i=1}^{17}p_i(\rho)=4+\mathrm{Tr}(\rho Q), \tag{31}$$

violating the classical bound 4 whenever $\mathrm{Tr}(\rho Q)>0$ (including $I_4/4$, where the sum is $17/4=\chi_f$, and $\rho=Q$, where it is 5). Nevertheless, on the orthogonal subspace $q^{\perp}$, the density operator $\rho_\star=(I_4-Q)/3$ has core Born probabilities $1/4$ on $C_1\cup C_2$, $1/3$ on $\{9,10,11\}$, and $1/6$ on $B_1\cup B_2\cup B_3$, and completion probabilities $r_{m,1}+r_{m,2}=7/12$. Choosing $\alpha\in\{1,\dots,24\}$ uniformly and $\lambda\in[0,1]$ uniformly and setting $\varepsilon_m=\mathbf{1}_{\{\lambda<\tau_m\}}$ with

TABLE IV. Event order and preparation-dependent probability models for the fixed realizations. Membership statements concern Born vectors $p(\rho)$; the restricted model is $\mathcal{P}_{\text{ext}}$.

| Configuration | Order determination | An ordinary-state model | A failed model or witness |
|---|---|---|---|
| $H_{17,10}$ | No: $s(10) \leq s(24)$ is a false ray inclusion. | $P_{24}$ and $P_{z_3}$ have ordinary mixtures. | Neither lies in $\mathcal{P}_{\text{ext}}$; $\eta = 1/9$ and $1/3$ (maximal pair certificate). |
| Completed Yu–Oh | No. | No Born preparation lies in $\mathcal{P}$. | Every preparation gives $\sum_{10}^{13} p_i = 4/3 > 1$. |
| $H_{37}$ | Yes on its specified 76-event lattice; 18 states minimal. | $I_3/3$ has a six-state mixture. | $\rho_{33} > 1/3$ violates the $P, Q$ bound. |
| $H_{\text{YO}}^{(4)}$ | No. | Pure apex $P_{26}$ has its singleton-state model. | Every $\rho \neq P_{26}$ violates the displayed bound. |
| $H_{65}$ | No: $s(16) \leq s(9)$ is a false ray inclusion. | $(I_4 - Q)/3$ has a model for all 65 rays. | $\text{Tr}(\rho Q) > 0$ violates the core sum bound. |

$\tau_m = \frac{12}{7} r_{m,2} \in \{4/9, 5/11, 6/11, 5/9\}$ on the 14 missed bridges of $S_\alpha$ gives $\mathbb{E}[s(w_{m,2})] = \frac{14}{24}\tau_m = r_{m,2}$, expressing $p(\rho_\star) \in \mathcal{P}(H_{65})$ as a convex mixture of at most 120 states. Rotating $V_{\text{core}}$ by $D_0 = I_4, D_1 = \text{diag}(1,1,-1,-1), D_2 = \text{diag}(1,-1,1,-1), D_3 = \text{diag}(1,-1,-1,1)$ gives a fixed enlarged family with

$$\sum_{a,i} \Pi_i^{(a)} = 17 I_4, \qquad \sum_{a,i} p_i^{(a)} \leq 16 \text{ classically.} \tag{32}$$

Each copy admits at most four true rays under geometric pair exclusions; repeated projectors are counted with multiplicity. The 24 distinct rays form the Peres set[35]. This is an enlargement, not a state-independent test of the original 65 rays.

## VII. KS ANCESTRY, SPECIFIED PARENTS, AND ABSTRACT VERSUS GEOMETRIC STATES

### A. The KS ancestry hypothesis and abstract transverse extensions

Motivated by the recurring parent structures of our examples, we formulate the *KS ancestry hypothesis*: informally, every noncolorable ($\chi(H) > d$) configuration has a larger KS ancestor. For realized ray configurations, its existence form follows from an elementary union argument, even without the chromatic premise; the substantive open question is whether the retained rays can be made essential to the contradiction (Sec. VIII):

**Proposition 26** (Same-dimensional finite KS ancestry)**.** *Every finite ray set $R \subset \mathbb{P}(\mathbb{F}^d)$ ($\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, $d \geq 3$) is a proper subset of a finite KS ray set $P \subset \mathbb{P}(\mathbb{F}^d)$ whose orthogonality graph and full-basis hypergraph induce those on $R$, with every ray of $P$ in a complete context. Consequently, a finite graph $G$ has an injective edge-faithful realization in $\mathbb{F}^d$ if and only if $G$ is an induced subgraph of a finite KS orthogonality graph in $\mathbb{F}^d$.*

*Proof.* Choose a finite full-basis KS set $K \subset \mathbb{P}(\mathbb{F}^d)$[4,36], orient $K$ by $U \in \text{U}(d)$ (or $\text{O}(d)$) so $UK \not\subseteq R$, and complete any uncovered rays of $R \cup UK$ to bases to obtain $P$. Any state on $P$ would restrict to a state on $UK$, a contradiction, and keeping the rays of $R$ fixed preserves all induced orthogonalities and full bases on $R$. □

By Theorem 11, $H_8$ has no injective realization, and $H_9, G_{32}$ have no ray assignment; none can embed into any 3-dimensional geometric KS parent. For abstract hypergraphs, Lemma 3 yields a direct chromatic-to-stateless construction:

**Proposition 27** (Transverse chromatic-to-stateless extension)**.** *For any finite $d$-uniform hypergraph $H = (V, E)$ ($d \geq 3$), let $T(H)$ take $d-1$ disjoint copies $H^{(1)}, \ldots, H^{(d-1)}$ and adjoin for each $v \in V$ a fresh vertex $w_v$ and the transverse context $T_v = \{v^{(1)}, \ldots, v^{(d-1)}, w_v\}$. Then $T(H)$ has $d|V|$ vertices, $(d-1)|E| + |V|$ contexts, is linear if $H$ is linear, and its two-valued states correspond bijectively to labeled strong $d$-colorings of $H$; in particular,*

$$\mathcal{S}(T(H)) = \varnothing \quad \Longleftrightarrow \quad \chi(H) > d. \tag{33}$$

*Proof.* A state $s \in \mathcal{S}(T(H))$ restricts to $d-1$ states on $H^{(k)}$ that are pairwise disjoint by $T_v$ (forcing $s(w_v) = 1 - \sum_{k=1}^{d-1} s(v^{(k)})$), which by Lemma 3 is equivalent to a strong $d$-coloring of $H$. □

In dimension $d = 3$, any edge $u \sim v$ of $H$ creates a Berge 4-loop $(u^{(1)}, v^{(1)}, v^{(2)}, u^{(2)})$ in $T(H)$, excluding an injective realization in $\mathbb{C}^3$ by the common orthogonal-complement argument; thus $T(H)$ is an abstract construction for a single system. It describes, however, the supersinglet game of Ref.[8] (Sec. I): the copies $H^{(k)}$ are the bases measured by $d-1$ of the parties, the transverse context $T_v$ states that exactly one party obtains $v$, and the fresh vertex $w_v$ stands for the last party, whose truth assignment is fixed by Lemma 3. Equation (33) is thus the combinatorial content of the statement that this game has no local hidden-variable model exactly when $\chi(H) > d$. For $H_9$, $T(H_9)$ (45 vertices, 33 contexts) is context-critical for statelessness on all 33 contexts; for $H^+_{\mathrm{YO}}$, $T(H^+_{\mathrm{YO}})$ (75 vertices, 57 contexts) is critical on all 32 horizontal contexts and 12 transverse contexts $v \in \{14, \ldots, 25\}$, but remains stateless upon deleting $T_v$ for $v \in \{1, \ldots, 13\}$ (Sec. S3 of the supplementary material).

### B. Explicit inclusions in known KS parents and Cabello completions

Beyond unrestricted existence, the following explicit unions use known KS components; the unions are not asserted to be previously named, smallest, or critical parents (explicit coordinate dictionaries and certificates in Secs. S1–S3 of the supplementary material):

1. *Bub's 49-ray context-critical real parent of $H^+_{\mathrm{YO}}$:*

   **Proposition 28** (Context-critical real parent of $H^+_{\mathrm{YO}}$)**.** *Adjoining to $H^+_{\mathrm{YO}}$ the eight rays $(0, 1, \pm 2)$, $(0, 2, \pm 1)$, $(1, \pm 2, 0)$, $(2, \pm 1, 0)$ yields 33 rays with 20 bases and 16 uncovered orthogonal pairs that eliminate all 24 states of $H^+_{\mathrm{YO}}$. Completing those 16 pairs yields the 49-ray, 36-context full-context Schütte–Bub KS configuration[20,37], containing $H^+_{\mathrm{YO}}$ as an induced restriction, in which* every *one of the 36 contexts is critical for $\mathcal{S}_{\mathrm{ctx}} = \varnothing$ (Table S83 of the supplementary material). If the orthogonalities of a deleted context are retained as geometric exclusions, only the 20 bases of the 33-ray stage remain critical; deleting any of the 16 completion contexts leaves $\mathcal{S}_{\mathrm{geom}} = \varnothing$.*

2. *MUB-inventory parents of $H_{37}$:* In the 165-ray, 130-context MUB inventory[18] formed by three 69-ray, 50-context KS lineages $L_0, L_1, L_2$ (where each $L_a$ completes the 36 uncovered pairs of Cabello's 33-ray, 14-basis KS set[38]; Pavičić[39] points out that this 33-ray set already occurs as a subset of a 69-ray, 50-basis set in Ref.[40]), $H_{37}$ embeds as an induced restriction via the seed-index matrix $j(a,k) = \left(\begin{smallmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 9 & 8 \end{smallmatrix}\right)$ mapping $z \leftrightarrow a_{1,1}$, $M_{ak} \leftrightarrow u_{j(a,k)}$, $P_k \leftrightarrow c_{1,k+1}$, $Q_k \leftrightarrow c_{2,k+1}$, $A_k \leftrightarrow b_{1,k+1}$, $B_{ak} \leftrightarrow d_{1,j(a,k)}$, $C_{ak} \leftrightarrow d_{2,j(a,a+k)}$. It lies in the 57-ray, 40-context intermediate spoke hypergraph (297 states) and in each 87-ray, 64-context KS union $H_{37} \cup L_a$.

3. *Four-dimensional Peres and Cabello parents of $H^{(4)}_{\mathrm{YO}}$ and $H_{65}$:* Taking the union with Peres's 24-ray set $P_{24}$[35] gives a 40-ray, 36-context KS parent of $H^{(4)}_{\mathrm{YO}}$ ($|H^{(4)}_{\mathrm{YO}} \cap P_{24}| = 10$; note that a single-apex lift of a 3-dimensional KS set always retains the unique apex-one state) and a 72-ray, 48-context KS parent of $H_{65}$ ($|H_{65} \cap P_{24}| = 17$). More economically:

**Proposition 29** (Three-ray KS extensions of $H_{65}$)**.** *Adjoining* $a = (1,0,0,1), b = (0,1,0,1), c = (0,0,1,1)$ *to* $H_{65}$ *yields a* 68*-ray,* 35*-basis KS parent containing* $H_{65}$ *as an induced restriction and Cabello's* 18*-ray,* 9*-context parity proof* $C_{18,9}$[21] *(*$C_3, C_4, C_5$ *plus* $D_1 = \{2,3,12,a\}, D_2 = \{1,3,15,b\}, D_3 = \{1,2,17,c\}, D_4 = \{6,7,13,a\}, D_5 = \{6,8,14,b\}, D_6 = \{7,8,16,c\}$*). Adjoining* $(0,1,-1,0)$, $(1,0,-1,0)$, $(1,-1,0,0)$ *instead yields a* 68*-ray,* 35*-basis KS parent containing a ray-critical* 20*-ray,* 11*-context parity proof* $C_{20,11}$ *(*$11 = 4s(4)+4s(5)+2\sum_{i\neq 4,5} s(i)$*). All* 20 *rays of* $C_{20,11}$ *belong to Peres's* 24*-ray set, and* $C_{20,11}$ *is one of its* 96 *critical* 20*-ray KS subsets, which are mutually isomorphic and include Kernaghan's set*[21,22]*.*

*Proof.* In $C_{18,9}$, summing $s$ over $C_3, C_4, C_5$ gives $3 = 2(s(9)+s(10)+s(11))+\sum_{k=12}^{17} s(k)$, so $s(12)+\cdots+s(17)$ is odd, whereas $D_1 \oplus D_4, D_2 \oplus D_5, D_3 \oplus D_6$ give $s(12)+s(13) \equiv s(2)+s(3)+s(6)+s(7)$, $s(14)+s(15) \equiv s(1)+s(3)+s(6)+s(8)$, $s(16)+s(17) \equiv s(1)+s(2)+s(7)+s(8)$ (mod 2), summing to 0 (mod 2). In $C_{20,11}$ (comprising $C_1, \ldots, C_5$ and six contexts on $\{1,\ldots,8,12,\ldots,17\}$ with $(0,1,-1,0)$, $(1,0,-1,0)$, $(1,-1,0,0)$), vertices 4, 5 have degree 4 and the other 18 vertices have degree 2, so $11 = 4s(4)+4s(5)+2\sum_{i\neq 4,5} s(i)$, an odd-equals-even contradiction; explicit states after deleting any single ray are certified in Sec. S3 of the supplementary material. □

Both 68-ray presentations have 35 bases covering 192 of 195 geometric orthogonal pairs and context intersections of size up to two. They are not edge-faithful in the listed bases. The 72-ray Peres union covers all pairs; no size minimum or criticality of either full parent is asserted.

4. *Faithful completion* $C_{36,18}$ *and abstract versus geometric states on* $C_{45,21}$*:*

**Corollary 30** (Faithful Cabello completion and vertex deletion)**.** *The* 9 *uncovered orthogonal pairs of* $C_{18,9}$ *satisfy Eq.* (22)*, so Lemma 21 yields a linear edge-faithful* 36*-ray,* 18*-context KS hypergraph* $C_{36,18}$ *in* $\mathbb{R}^4$ *with* $\omega = 4, \chi = 5$*, and* $\mathcal{S} = \varnothing$*. Deleting one vertex (say* $c$*) and its two contexts from* $C_{18,9}$ *leaves* 17 *rays,* 7 *bases,* 14 *uncovered pairs, and* 12 *core states whose abstract completion* $C_{45,21}$ *has* $\chi = 5$ *and* $|\mathcal{S}_{\mathrm{ctx}}(C_{45,21})| = 3{,}584$ *unital separating context states (*24 *canonical). However,* $v_{16} \in \mathrm{span}(v_1, v_2)$ *and* $v_{17} \in \mathrm{span}(v_7, v_8)$ *violate Eq.* (22)*, forcing four extra orthogonalities that reduce the geometric state space to* $|\mathcal{S}_{\mathrm{geom}}(C_{45,21})| = 2{,}048$ *with* $s(9) = s(16) = s(17) = 0$*, destroying unitality and separation (Table V).*

*Proof.* For $C_{18,9}$, the 9 uncovered pairs are the 6 core bridges $\{1,13\}$, $\{2,14\}$, $\{3,16\}$, $\{6,17\}$, $\{7,15\}$, $\{8,12\}$ from $H_{65}$ plus $\{11,a\}$, $\{10,b\}$, $\{9,c\}$. They satisfy Eq. (22) and are completed in Tables S28–S29 of the supplementary material. Deleting $c, D_3, D_6$ leaves 12 core supports: 10 miss 8 of the 14 uncovered pairs and 2 miss 9 (Table S84), giving $10 \cdot 2^8 + 2 \cdot 2^9 = 3{,}584$ context states. Since $v_{16} = v_1 + v_2$ and $v_{17} = \frac{1}{2}(v_7 - v_8)$, completing $\{1,2\}$ by $\{21,22\}$ and $\{7,8\}$ by $\{35,36\}$ forces $v_{16} \perp v_{21}, v_{22}$ and $v_{17} \perp v_{35}, v_{36}$ in the local labels of Tables S31–S32. If $s(16) = 1$, neither 1 nor 2 is selected in any admissible core support, so $s(21)+s(22) = 1$, contradicting geometric exclusion; the same argument gives $s(17) = 0$. The retained hub bases then force $s(9) = 0$. Exactly 8 core supports satisfy $s(9) = s(16) = s(17) = 0$, each missing 8 uncovered pairs. All their fresh-pair choices obey the four extra exclusions, yielding $8 \cdot 2^8 = 2{,}048$ geometric states. □

## VIII. DEPENDENCIES, COUNTEREXAMPLES, AND REMAINING QUESTIONS

The common dependency structure is incidence → states and colorings, followed by independent tests of geometry, event order, and quantum probabilities. A chromatic argument can be made on an abstract hypergraph; its operational interpretation requires

TABLE V. Context and geometric state spaces in the four-dimensional examples. Linearity of the listed incidence structure does not imply edge-faithfulness.

| Configuration | $\lvert V\rvert/\lvert E\rvert$ | $\lvert\mathcal{S}_{\mathrm{ctx}}\rvert$ | $\lvert\mathcal{S}_{\mathrm{geom}}\rvert$ | Scope |
|---|---|---|---|---|
| $C_{18,9}$ | 18/9 | 0 | 0 | Nine uncovered geometric pairs; parity KS proof. |
| $C_{36,18}$ | 36/18 | 0 | 0 | Faithful linear completion. |
| $C_{45,21}$ | 45/21 | 3,584 | 2,048 | Four additional pairs destroy geometric unitality and separation. |
| $H_{65}$ | 65/29 | 393,216 | 393,216 | Edge-faithful; all context states respect the geometry. |

a ray realization. Conversely, a realization does not select an ontic state family or a classical probability mixture. The exact-palette assumption, that the outcomes admit a noncontextual assignment of $d$ eigenvalues, adds a precise completion test at the level of whole truth assignments: a truth assignment is compatible with noncontextual eigenvalues exactly when the outcomes it leaves false can carry the remaining $d-1$ eigenvalues.

As a worked example, consider how many hidden states a classical model needs for the 13 Yu–Oh rays of Sec. III C. A hidden state is the complete classical description of the system: it fixes, or at least gives the probabilities of, the outcome of every measurement. A preparation produces a hidden state, possibly at random from several, and a quantum state then corresponds to a probability distribution over hidden states. In a generalized urn model[13] the hidden state is the ball that is drawn; in a deterministic noncontextual model it is a two-valued state. One fact suffices: if two rays $u$ and $v$ are orthogonal, a measurement in a basis containing both tells the preparations along $u$ and along $v$ apart with certainty. Hence no hidden state can arise from both preparations, since in that hidden state the measurement would have to yield $u$ with certainty and also $v$ with certainty.

Suppose now that a classical model uses only three hidden states $A$, $B$, $C$, as many as the dimension.

1. The rays $z_1 = (1,0,0)$, $z_2 = (0,1,0)$, $z_3 = (0,0,1)$ are mutually orthogonal, so they need three different hidden states, say $A$, $B$, $C$.

2. The rays $y_1^{\pm} = (0,1,\pm1)$ are orthogonal to $z_1$ and to each other, so they take $B$ and $C$ in some order. Likewise $y_2^{\pm} = (1,0,\pm1)$ take $A$ and $C$, and $y_3^{\pm} = (1,\pm1,0)$ take $A$ and $B$.

3. Each diagonal ray is orthogonal to exactly one ray of each pair: $h_0 = (1,1,1)$ to $y_1^-, y_2^-, y_3^-$; $h_1 = (1,-1,-1)$ to $y_1^-, y_2^+, y_3^+$; $h_2 = (1,-1,1)$ to $y_1^+, y_2^-, y_3^+$; and $h_3 = (1,1,-1)$ to $y_1^+, y_2^+, y_3^-$. Each diagonal ray needs a hidden state that none of its three neighbors uses.

4. A diagonal ray can use $A$ only if its $y_2$-neighbor has $C$ and its $y_3$-neighbor has $B$. This fixes two of the three signs of its neighbors, and exactly one diagonal ray has the required signs, because the four diagonal rays have the four sign patterns with an even number of plus signs. The same holds for $B$ and for $C$. So at most three diagonal rays find a free hidden state; the fourth is orthogonal to rays in $A$, $B$, and $C$.

Any classical model therefore needs at least four hidden states merely to tell these 13 preparations apart, whereas a qutrit realizes them in three dimensions. A model with exactly three hidden states, each tied to one eigenvalue, is a noncontextual eigenvalue assignment: in hidden state $A$, every context yields the outcome that carries $\lambda_0$, and likewise for $B$ and $C$. Steps 1–4 thus also show directly that the Yu–Oh rays admit no noncontextual eigenvalue assignment; Theorem 13 refines this count for completed Yu–Oh.

Table VI also explains why the probability and gadget examples carry different force. A one-way ordinary TITS already gives a symmetric color identity; a two-way purely chromatic identity requires counterstates in both truth directions. A ternary color exclusion may

TABLE VI. Logical boundaries of the integrated results. Each row prevents transfer of a conclusion between different assignment or representation problems.

| Tempting unrestricted inference | Counterexample or necessary qualification |
|---|---|
| $\chi > d$ excludes states or ray realizations | Completed Yu–Oh has faithful real rays and 24 unital separating states. |
| $\chi > d$ makes $I_d/d$ contextual[8] | One system needs $\chi_f > d$[16]: $H_{37}$ has $\chi = 4$, $\chi_f = 3$, and a six-state mixture for $I_3/3$. |
| Ordinary states separate ⇒ colors separate | Faithful $H_{17,10}$ and $H_{21,13}$ have forced color identities. |
| Colors separate ⇒ every state extends | The pentagon separates all vertex pairs but has one nonextendable state. |
| Vertex separation ⇒ event-order determination | $H_{17,10}$ has 10 ⇒ 24; $H_{65}$ has 16 ⇒ 9 and ray–plane collapse. |
| Order determination ⇒ all Born mixtures | The specified event order of $H_{37}$ is represented, while $\rho_{33} > 1/3$ violates the state inequality. |
| $\chi_f > d$ establishes fixed-measurement state independence | Ref.[15] (Thm. 1) (Yu–Oh plus apex); linear and apex-free: $H_{65}$ admits the Born model of $(I_4 - Q)/3$ on all 65 rays. |
| Abstract stateless extension ⇒ geometric KS parent | Closing a color-identical pair changes orthogonality; transverse four-loops exclude injective triadic geometry. |
| Generic KS ancestry explains the original chromatic gap | Unrestricted extension holds even for colorable sets and need not use the old rays essentially. |

survive even when every pair can be equal. A strict inclusion $\mathcal{P}_{\text{ext}} \subsetneq \mathcal{P}$ therefore need not be detectable through pair equalities alone.

Open problems concern global minimum faithful examples beyond the specified inventory, economical rational completions with fewer degree-one rays, residual-coloring certificates in dimensions above three, and geometric ancestry in which the retained rays are essential to the contradiction. The current statements establish existence and the recorded finite classifications, without asserting these stronger minima or essential-ancestry properties. All complete enumerations and the detailed verification scope remain in the supplement and electronic records.

## IX. CONCLUSION

We started from a simple physical requirement on intertwined complete experiments: that each is the measurement of a maximal, nondegenerate observable with one common spectrum of $d$ eigenvalues, and that the eigenvalue carried by an outcome is an element of physical reality in the sense of Einstein, Podolsky, and Rosen[3], the same in every experiment in which the outcome occurs. This requirement of noncontextual eigenvalues is exactly strong $d$-colorability of the context hypergraph. It asks for more than Kochen–Specker noncontextuality: not one classical truth assignment, but $d$ of them with disjoint supports covering every outcome. Its failure, $\chi(H) > d$, is therefore a weaker but more widespread form of contextuality. Completed Yu–Oh has 24 truth assignments separating all of its rays, yet its outcomes cannot carry three eigenvalues noncontextually. These truth assignments also give a classical realization, which fails in the same way, so $\chi(H) > d$ restricts the spectra of maximal observables rather than truth values. When $H$ is realized in $\mathbb{C}^d$, it also means that a classical model needs more than $d$ hidden states merely to tell apart preparations along orthogonal rays.

By unifying these common-spectrum chromatic obstructions ($\chi(H) > d$) with exact-palette chromatic separability constraints ($\chi(H) = d$, $\mathcal{S}_{\text{ext}}(H) \subsetneq \mathcal{S}(H)$), we have distinguished the requirements across noncontextual eigenvalue assignments, geometric ray realizations, orthomodular event representations, and quantum probability models. At the foundation,

$\chi(H) > d$ excludes noncontextual eigenvalues for uniform maximal context observables, whereas the exclusion of an injective realization for $H_8$ and of every ray assignment for $H_9, G_{32}$ requires direct Hermitian projection forcing, correcting the earlier spectral argument for $G_{32}$[6] and establishing four-quadrant independence even at Berge girth five. Within chromatically complete hypergraphs ($\chi(H) = d$), where noncontextual eigenvalue assignments exist, our completion criterion ($\chi(\Gamma(H)[V \setminus \operatorname{supp} s]) \leq d - 1$) characterizes which truth assignments arise from noncontextual eigenvalue assignments and explains the specified examples of chromatic identities ($u \sim_c v$) and purely chromatic TIFS/TITS/ternary constraints ($H_{17,10}, H_{21,13}, D_{16,9}, T_{19,11}, W^{(4)}_{16,6}$) and yields measurable nonextendable weights 1/9 and 1/3 for quantum ray preparations on $H_{17,10}$, the latter being the largest value a single chromatic identity can certify there. In higher complexity, the Hesse hypergraph $H_{37}$ (6/7/18 OML minima) and our rational four-dimensional completion $H_{65}$ (393,216 states) demonstrate that neither exact order determination nor a fractional chromatic gap decides classicality for all Born preparations, while the elementary KS ancestry result, the 68-ray KS parents of $H_{65}$, and the completions $C_{36,18}$ and $C_{45,21}$ of $C_{18,9}$ delineate unrestricted embedding from specified critical parents and geometric pair exclusions.

The common thread is that a classical account of quantum outcomes can succeed at the level of truth values and still fail at the level of eigenvalues. For the preparation along $z_3$, the quantum predictions on $H_{17,10}$ admit noncontextual truth-assignment models, but every such model must place at least one third of its weight on truth assignments that no noncontextual eigenvalue assignment produces.

## SUPPLEMENTARY MATERIAL

See the supplementary material for the explicit vector and context inventories, the complete two-valued state and coloring enumerations, the odd-cycle and separation-profile certificates, and the supplementary proofs and verification reductions.

## ACKNOWLEDGMENTS

I thank Mohammad Hadi Shekarriz for valuable comments and suggestions. This research was funded in part by the Austrian Science Fund (FWF), Grant DOI 10.55776/PIN5424624. The author acknowledges FWF and TU Wien Bibliothek for support through the Open Access Funding Programme. Generative AI tools were used for editorial work, including language editing and the LaTeX layout of figures and tables, and for independent checks of finite computations. The author reviewed all such output and takes full responsibility for the content of this article.

## AUTHOR DECLARATIONS

### Conflict of Interest

The author has no conflicts to disclose.

## DATA AVAILABILITY

The data that support the findings of this study are available within the article and its supplementary material. The verification programs and machine-readable certificates are available from the author upon reasonable request.

[1] J. von Neumann, Annals of Mathematics **32**, 191 (1931).

[2]P. R. Halmos, *Finite-Dimensional Vector Spaces*, Undergraduate Texts in Mathematics (Springer, New York, 1974).
[3]A. Einstein, B. Podolsky, and N. Rosen, Physical Review **47**, 777 (1935).
[4]S. Kochen and E. P. Specker, Journal of Mathematics and Mechanics **17**, 59 (1967).
[5]K. Svozil, Quantum Reports **2**, 278 (2020), arXiv:1808.00813.
[6]M. H. Shekarriz and K. Svozil, Journal of Mathematical Physics **63**, 032104 (2022), arXiv:2105.08520.
[7]K. Svozil, Entropy **27**, 387 (2025), arXiv:2501.15261.
[8]A. Cabello, "State-independent quantum contextuality and maximum nonlocality," (2012), version 1 (2011): "Simple unified proof of state-independent contextuality", arXiv:1112.5149 [quant-ph].
[9]R. J. Greechie, Journal of Combinatorial Theory. Series A **10**, 119 (1971).
[10]S. Yu and C. H. Oh, Physical Review Letters **108**, 030402 (2012), arXiv:1109.4396.
[11]J. A. Martin H., "Graph-chromatic implicit relations," (2009), arXiv:0901.1255 [math.CO].
[12]A. Dvurečenskij, S. Pulmannová, and K. Svozil, Helvetica Physica Acta **68**, 407 (1995), arXiv:1806.04271.
[13]K. Svozil, International Journal of Theoretical Physics **44**, 745 (2005), arXiv:quant-ph/0209136.
[14]S. Prakash, "Quantum memory advantage from contextuality," (2026), arXiv:2607.00507 [quant-ph].
[15]A. Cabello, M. Kleinmann, and C. Budroni, Physical Review Letters **114**, 250402 (2015), arXiv:1501.03432.
[16]R. Ramanathan and P. Horodecki, Physical Review Letters **112**, 040404 (2014), arXiv:1212.5933.
[17]G. Scarpa and S. Severini, IEEE Transactions on Information Theory **58**, 2524 (2012), arXiv:1106.0712.
[18]M. Navara and K. Svozil, "Construction of Kochen-Specker sets from mutually unbiased bases," (2025), arXiv:2509.08636 [quant-ph].
[19]G. Haynes, C. Park, A. Schaeffer, J. Webster, and L. H. Mitchell, The Electronic Journal of Combinatorics **17**, R55 (2010).
[20]J. Bub, Foundations of Physics **26**, 787 (1996).
[21]A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
[22]M. Kernaghan, Journal of Physics A: Mathematical and General **27**, L829 (1994).
[23]A. Silva and A. Ibiapina, "On orthogonal vector edge coloring," (2019), arXiv:1909.01918 [cs.DM].
[24]G. Brinkmann, M. Preissmann, and D. Sasaki, Discrete Mathematics & Theoretical Computer Science **17**, 369 (2015).
[25]G. Schrag, Discrete Mathematics **70**, 185 (1988).
[26]K. Svozil and J. Tkadlec, Journal of Mathematical Physics **37**, 5380 (1996).
[27]L. Hogben, K. F. Palmowski, D. E. Roberson, and S. Severini, Electronic Journal of Linear Algebra **32**, 98 (2017).
[28]M. Waegell and P. K. Aravind, Foundations of Physics **41**, 1786 (2011), arXiv:1103.6058 [quant-ph].
[29]R. Isaacs, The American Mathematical Monthly **82**, 221 (1975).
[30]D. Blanuša, Glasnik Matematičko-Fizički i Astronomski, Serija II **1**, 31 (1946).
[31]M. Pavičić, Entropy **21**, 1107 (2019).
[32]S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
[33]A. Cabello, J. R. Portillo, A. Solís, and K. Svozil, Physical Review A **98**, 012106 (2018), arXiv:1805.00796.
[34]J. Harding and R. Salinas Schmeis, International Journal of Theoretical Physics **64**, 211 (2025), arXiv:2505.13871 [math-ph].
[35]A. Peres, Journal of Physics A: Mathematical and General **24**, L175 (1991).
[36]A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **339**, 425 (2005).
[37]M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
[38]A. Cabello, Physical Review Letters **135**, 190203 (2025), arXiv:2508.07335 [quant-ph].
[39]M. Pavičić, "On simplest Kochen-Specker sets," (2025), arXiv:2512.10483 [quant-ph].
[40]M. Pavičić, Quantum **7**, 953 (2023).

# Supplementary Material for "Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments"

Karl Svozil
*Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, 1040 Vienna, Austria*

(Dated: November 5, 2026)

This supplementary material provides the complete vector and context enumerations, two-valued state tables, normalized strong coloring tables, odd-cycle nonextendability certificates, chromatic signature and separation profile tables, parent-embedding dictionaries, and supplementary proofs and finite verification certificates supporting the main text. Section S1 lists the explicit unnormalized coordinates and context tables for the completed Yu–Oh hypergraph $H_{\mathrm{YO}}^{+}$ and its faithful restrictions $H_{17,10}$ and $H_{21,13}$, the three- and four-dimensional logical gadgets, the 37-ray complex Hesse hypergraph $H_{37}$, the 26-ray suspension $H_{\mathrm{YO}}^{(4)}$ and 65-ray canonical linear completion $H_{65}$, the Cabello completions $C_{36,18}$ and $C_{45,21}$, and the specified finite KS parents (49-ray Schütte–Bub, 87-ray MUB-inventory union, 40- and 72-ray Peres unions, and 68-ray three-ray extensions). Section S2 gives the complete two-valued state tables, extendable-state classifications, normalized coloring tables, odd-cycle certificates, and separation profiles across both regimes. Section S3 presents the supplementary proofs and exact reductions, including Isaacs dot-product snarks and quantitative $\Delta_3$ robustness bounds for $G_{32}$, the 540,672-candidate reduction for the minimum 18-state order-determining family on $L(H_{37})$, the canonical completion selection rule for $H_{65}$, the forced-zero and ray-criticality certificates for the four-dimensional Cabello completions, transverse criticality deletion colorings, and historical Kochen–Specker gadget certificates.

## READING GUIDE AND VERIFICATION SCOPE

This supplement collects complete vector and context inventories (S1), state and coloring enumerations (S2), and extended proofs and certificates (S3). Labels are local to each configuration. "Exact palette" means $\chi(H) = d$, with no additional density or geometric assumption: the outcomes admit a noncontextual assignment of $d$ eigenvalues, that is, every complete experiment can be measured by a maximal observable with one common spectrum of $d$ eigenvalues, with each outcome carrying the same eigenvalue in every context. Colorings are such eigenvalue assignments, and extendable states are the truth assignments "the outcome with eigenvalue $\lambda_a$ occurs" that they produce.

The five-state and four-state $H_{17,10}$ mixtures and both-direction gadget witnesses use the complete state families. Every selected support satisfies every context equation, and rational arithmetic verifies every Born marginal and the attained nonextendable weights $1/9$ and $1/3$. The ordinary implication $10 \Rightarrow 24$ on $H_{17,10}$ gives a false inclusion between distinct rays and hence a failure of order determination.

The companion electronic data include all coordinates, state enumerations (including the compressed 393,216-state $H_{65}$ inventory), verification programs, and the exhaustive certificates for the eighteen-state Hesse minimum and the 65,535-subset Yu–Oh classification. Checking an eighteen-state family establishes an upper bound; the matching lower bound requires the exhaustive certificate in Sec. S3.3. The directories `part-a/data`, `part-a/verification`, `part-b/data`, and `part-b/verification` contain the underlying data and verifiers. Reports and source-integrity hashes are in `audit/`; Sec. S3.7 describes the joint verification entry point and its scope.

## S1. EXPLICIT UNNORMALIZED VECTORS AND CONTEXTS

All numerical vertex labels in this section are local to the named configuration and match the figures and theorems of the main text.

### S1.1. Completed Yu–Oh and the Petersen-derived contexts

The three-dimensional vectors below give all projective rays used in Sec. III C; signs are chosen to make the first nonzero integer coordinate positive. The 26-ray suspension appends zero and adds $e_4$. The abstract $G_{32}, H_9, H_8$ context lists in Eq. (10) and Theorem 11 specify their incidences completely. The geometric obstructions exclude a corresponding injective three-dimensional vector enumeration for these abstract examples.

TABLE S1: All 25 unnormalized completed Yu–Oh vectors.

| $i$ | $r_i$ | $i$ | $r_i$ |
|---|---|---|---|
| 1 | $(0,1,1)$ | 14 | $(2,1,1)$ |
| 2 | $(1,0,0)$ | 15 | $(1,2,-1)$ |
| 3 | $(0,1,-1)$ | 16 | $(1,-2,1)$ |
| 4 | $(1,1,0)$ | 17 | $(1,1,-2)$ |
| 5 | $(0,0,1)$ | 18 | $(1,-1,-2)$ |
| 6 | $(1,-1,0)$ | 19 | $(2,1,-1)$ |
| 7 | $(1,0,1)$ | 20 | $(2,-1,1)$ |
| 8 | $(0,1,0)$ | 21 | $(1,-2,-1)$ |
| 9 | $(1,0,-1)$ | 22 | $(1,1,2)$ |
| 10 | $(1,-1,1)$ | 23 | $(1,2,1)$ |
| 11 | $(1,-1,-1)$ | 24 | $(1,-1,2)$ |
| 12 | $(1,1,1)$ | 25 | $(2,-1,-1)$ |
| 13 | $(1,1,-1)$ | | |

### S1.2. Faithful Yu–Oh restrictions $H_{17,10}$ and $H_{21,13}$ and logical gadgets in dimensions three and four

Tables S2 and S3 record the restriction membership of the 25 parent rays and 16 parent contexts of $H^{+}_{\mathrm{YO}}$ in $H_{17,10}$ and $H_{21,13}$. Tables S4–S14 provide the explicit unnormalized integer coordinates and context lists for the faithful real 3-dimensional gadgets $P_5$, $B_{13,7}$, $D_{16,9}$, $I_{16,9}$, and $T_{19,11}$ and the faithful real 4-dimensional gadgets $W^{(4)}_{11,5}$, $W^{(4)}_{16,6}$, $B^{(4)}_{14,7}$, $L^{(4)}_{18,7}$, $D^{(4)}_{17,9}$, and $L^{(4)}_{22,9}$. Note that for the standalone gadgets $P_5$, $D_{16,9}$, $I_{16,9}$, and $T_{19,11}$, the vertex numbering is local to each gadget panel; in $L^{(4)}_{18,7}$ [Fig. 3(d)], $r_{13} = (1,0,1,1)$ and $r_{14} = (2,0,-1,-1)$ are chosen so that $r_{13} \not\perp r_{15}$, whereas in $L^{(4)}_{22,9}$ [Fig. 3(f)], $r_{13} = (1,0,4,4)$ and $r_{14} = (8,0,-1,-1)$ are used so that $r_{13} \perp r_{15}$ in the added context $\{13,20,15,19\}$.

TABLE S2: All 25 unnormalized parent vectors; membership of the two geometric examples.

| $i$ | $r_i$ | $H_{17,10}$ | $H_{21,13}$ | $i$ | $r_i$ | $H_{17,10}$ | $H_{21,13}$ |
|---|---|---|---|---|---|---|---|
| 1 | $(0,1,1)$ | yes | yes | 14 | $(2,1,1)$ | yes | yes |
| 2 | $(1,0,0)$ | yes | yes | 15 | $(1,2,-1)$ | yes | yes |
| 3 | $(0,1,-1)$ | yes | yes | 16 | $(1,-2,1)$ | – | – |
| 4 | $(1,1,0)$ | yes | yes | 17 | $(1,1,-2)$ | – | – |
| 5 | $(0,0,1)$ | yes | yes | 18 | $(1,-1,-2)$ | yes | yes |
| 6 | $(1,-1,0)$ | yes | yes | 19 | $(2,1,-1)$ | yes | yes |
| 7 | $(1,0,1)$ | yes | yes | 20 | $(2,-1,1)$ | – | yes |
| 8 | $(0,1,0)$ | yes | yes | 21 | $(1,-2,-1)$ | – | yes |
| 9 | $(1,0,-1)$ | yes | yes | 22 | $(1,1,2)$ | – | yes |
| 10 | $(1,-1,1)$ | yes | yes | 23 | $(1,2,1)$ | yes | yes |
| 11 | $(1,-1,-1)$ | yes | yes | 24 | $(1,-1,2)$ | yes | yes |
| 12 | $(1,1,1)$ | – | – | 25 | $(2,-1,-1)$ | – | – |
| 13 | $(1,1,-1)$ | – | yes | | | | |

TABLE S3: The 16 parent contexts and the two selected restrictions.

| $j$ | context | $H_{17,10}$ | $H_{21,13}$ |
|---|---|---|---|
| 1 | $\{1,2,3\}$ | yes | yes |
| 2 | $\{7,8,9\}$ | yes | yes |
| 3 | $\{4,5,6\}$ | yes | yes |
| 4 | $\{2,8,5\}$ | yes | yes |
| 5 | $\{3,12,25\}$ | – | – |
| 6 | $\{9,12,16\}$ | – | – |
| 7 | $\{6,12,17\}$ | – | – |
| 8 | $\{3,11,14\}$ | yes | yes |
| 9 | $\{7,11,15\}$ | yes | yes |
| 10 | $\{4,11,24\}$ | yes | yes |
| 11 | $\{9,10,23\}$ | yes | yes |
| 12 | $\{1,10,19\}$ | yes | yes |
| 13 | $\{4,10,18\}$ | yes | yes |
| 14 | $\{1,13,20\}$ | – | yes |
| 15 | $\{7,13,21\}$ | – | yes |
| 16 | $\{6,13,22\}$ | – | yes |

### S1.3. 3D panel (a): $P_5$

TABLE S4: Unnormalized vectors and all contexts of $P_5$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(1,0,0)$ | 1 | $\{1,6,2\}$ |
| 2 | $(0,1,1)$ | 2 | $\{2,7,3\}$ |
| 3 | $(1,1,-1)$ | 3 | $\{3,8,4\}$ |
| 4 | $(1,-2,-1)$ | 4 | $\{4,9,5\}$ |
| 5 | $(0,1,-2)$ | 5 | $\{5,10,1\}$ |
| 6 | $(0,1,-1)$ | | |
| 7 | $(2,-1,1)$ | | |
| 8 | $(1,0,1)$ | | |
| 9 | $(5,2,1)$ | | |
| 10 | $(0,2,1)$ | | |

### S1.4. 3D panel (b): $B_{13,7}$

TABLE S5: Unnormalized vectors and all contexts of $B_{13,7}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(0,1,1)$ | 1 | $\{1,2,3\}$ |
| 2 | $(1,0,0)$ | 2 | $\{7,8,9\}$ |
| 3 | $(0,1,-1)$ | 3 | $\{2,8,5\}$ |
| 5 | $(0,0,1)$ | 4 | $\{3,11,14\}$ |
| 7 | $(1,0,1)$ | 5 | $\{7,11,15\}$ |
| 8 | $(0,1,0)$ | 6 | $\{9,10,23\}$ |
| 9 | $(1,0,-1)$ | 7 | $\{1,10,19\}$ |
| 10 | $(1,-1,1)$ | | |
| 11 | $(1,-1,-1)$ | | |
| 14 | $(2,1,1)$ | | |
| 15 | $(1,2,-1)$ | | |
| 19 | $(2,1,-1)$ | | |
| 23 | $(1,2,1)$ | | |

### S1.5. 3D panel (c): $D_{16,9}$

TABLE S6: Unnormalized vectors and all contexts of $D_{16,9}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(0,1,2)$ | 1 | $\{1,2,3\}$ |
| 2 | $(1,0,0)$ | 2 | $\{7,8,9\}$ |
| 3 | $(0,2,-1)$ | 3 | $\{4,5,6\}$ |
| 4 | $(1,2,0)$ | 4 | $\{2,8,5\}$ |
| 5 | $(0,0,1)$ | 5 | $\{3,11,14\}$ |
| 6 | $(2,-1,0)$ | 6 | $\{7,11,15\}$ |
| 7 | $(1,0,1)$ | 7 | $\{4,11,24\}$ |
| 8 | $(0,1,0)$ | 8 | $\{9,10,23\}$ |
| 9 | $(1,0,-1)$ | 9 | $\{4,10,18\}$ |
| 10 | $(2,-1,2)$ | | |
| 11 | $(2,-1,-2)$ | | |
| 14 | $(5,2,4)$ | | |
| 15 | $(1,4,-1)$ | | |
| 18 | $(4,-2,-5)$ | | |
| 23 | $(1,4,1)$ | | |
| 24 | $(4,-2,5)$ | | |

### S1.6. 3D panel (d): $I_{16,9}$

TABLE S7: Unnormalized vectors and all contexts of $I_{16,9}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(0,1,1)$ | 1 | $\{1,2,3\}$ |
| 2 | $(1,0,0)$ | 2 | $\{7,8,9\}$ |
| 3 | $(0,1,-1)$ | 3 | $\{2,8,5\}$ |
| 4 | $(4,5,3)$ | 4 | $\{3,11,14\}$ |
| 5 | $(0,0,1)$ | 5 | $\{7,11,15\}$ |
| 7 | $(1,0,2)$ | 6 | $\{4,11,24\}$ |
| 8 | $(0,1,0)$ | 7 | $\{9,10,23\}$ |
| 9 | $(2,0,-1)$ | 8 | $\{1,10,19\}$ |
| 10 | $(1,-2,2)$ | 9 | $\{4,10,18\}$ |
| 11 | $(2,-1,-1)$ | | |
| 14 | $(1,1,1)$ | | |
| 15 | $(2,5,-1)$ | | |
| 18 | $(16,-5,-13)$ | | |
| 19 | $(4,1,-1)$ | | |
| 23 | $(2,5,4)$ | | |
| 24 | $(1,-5,7)$ | | |

### S1.7. 3D panel (e): $T_{19,11}$

TABLE S8: Unnormalized vectors and all contexts of $T_{19,11}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(0,1,2)$ | 1 | $\{1,2,3\}$ |
| 2 | $(1,0,0)$ | 2 | $\{7,8,9\}$ |
| 3 | $(0,2,-1)$ | 3 | $\{4,5,6\}$ |
| 4 | $(1,2,0)$ | 4 | $\{2,8,5\}$ |
| 5 | $(0,0,1)$ | 5 | $\{3,11,14\}$ |
| 6 | $(2,-1,0)$ | 6 | $\{7,11,15\}$ |
| 7 | $(1,0,1)$ | 7 | $\{4,11,24\}$ |
| 8 | $(0,1,0)$ | 8 | $\{9,10,23\}$ |
| 9 | $(1,0,-1)$ | 9 | $\{4,10,18\}$ |
| 10 | $(2,-1,2)$ | 10 | $\{14,26,27\}$ |
| 11 | $(2,-1,-2)$ | 11 | $\{18,26,28\}$ |
| 14 | $(5,2,4)$ | | |
| 15 | $(1,4,-1)$ | | |
| 18 | $(4,-2,-5)$ | | |
| 23 | $(1,4,1)$ | | |
| 24 | $(4,-2,5)$ | | |
| 26 | $(2,-41,18)$ | | |
| 27 | $(200,-82,-209)$ | | |
| 28 | $(241,82,160)$ | | |

### S1.8. 4D panel (a): $W^{(4)}_{11,5}$

TABLE S9: Unnormalized vectors and all contexts of $W^{(4)}_{11,5}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 0 | $(0,0,0,1)$ | 1 | $\{0,1,6,2\}$ |
| 1 | $(1,0,0,0)$ | 2 | $\{0,2,7,3\}$ |
| 2 | $(0,1,1,0)$ | 3 | $\{0,3,8,4\}$ |
| 3 | $(1,1,-1,0)$ | 4 | $\{0,4,9,5\}$ |
| 4 | $(1,-2,-1,0)$ | 5 | $\{0,5,10,1\}$ |
| 5 | $(0,1,-2,0)$ | | |
| 6 | $(0,1,-1,0)$ | | |
| 7 | $(2,-1,1,0)$ | | |
| 8 | $(1,0,1,0)$ | | |
| 9 | $(5,2,1,0)$ | | |
| 10 | $(0,2,1,0)$ | | |

### S1.9. 4D panel (b): $W^{(4)}_{16,6}$

TABLE S10: Unnormalized vectors and all contexts of $W^{(4)}_{16,6}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 0 | $(0,0,0,1)$ | 1 | $\{2,0,6,1\}$ |
| 1 | $(1,0,0,0)$ | 2 | $\{2,7,13,3\}$ |
| 2 | $(0,1,1,0)$ | 3 | $\{3,0,8,4\}$ |
| 3 | $(1,1,-1,0)$ | 4 | $\{4,9,14,5\}$ |
| 4 | $(1,-2,-1,0)$ | 5 | $\{5,10,15,1\}$ |
| 5 | $(0,1,-2,0)$ | 6 | $\{0,11,12,5\}$ |
| 6 | $(0,1,-1,0)$ | | |
| 7 | $(2,-1,1,1)$ | | |
| 8 | $(1,0,1,0)$ | | |
| 9 | $(5,2,1,1)$ | | |
| 10 | $(0,4,2,1)$ | | |
| 11 | $(1,2,1,0)$ | | |
| 12 | $(5,-2,-1,0)$ | | |
| 13 | $(2,-1,1,-6)$ | | |
| 14 | $(5,2,1,-30)$ | | |
| 15 | $(0,2,1,-10)$ | | |

### S1.10. 4D panel (c): $B^{(4)}_{14,7}$

TABLE S11: Unnormalized vectors and all contexts of $B^{(4)}_{14,7}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 0 | $(0,0,0,1)$ | 1 | $\{0,1,2,3\}$ |
| 1 | $(0,1,1,0)$ | 2 | $\{0,7,8,9\}$ |
| 2 | $(1,0,0,0)$ | 3 | $\{0,2,8,5\}$ |
| 3 | $(0,1,-1,0)$ | 4 | $\{0,3,11,14\}$ |
| 5 | $(0,0,1,0)$ | 5 | $\{0,7,11,15\}$ |
| 7 | $(1,0,1,0)$ | 6 | $\{0,1,10,19\}$ |
| 8 | $(0,1,0,0)$ | 7 | $\{0,9,10,23\}$ |
| 9 | $(1,0,-1,0)$ | | |
| 10 | $(1,-1,1,0)$ | | |
| 11 | $(1,-1,-1,0)$ | | |
| 14 | $(2,1,1,0)$ | | |
| 15 | $(1,2,-1,0)$ | | |
| 19 | $(2,1,-1,0)$ | | |
| 23 | $(1,2,1,0)$ | | |

### S1.11. 4D panel (d): $L^{(4)}_{18,7}$

TABLE S12: Unnormalized vectors and all contexts of $L^{(4)}_{18,7}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(1,0,0,0)$ | 1 | $\{1,2,4,3\}$ |
| 2 | $(0,1,0,0)$ | 2 | $\{5,8,7,6\}$ |
| 3 | $(0,0,1,0)$ | 3 | $\{4,9,10,8\}$ |
| 4 | $(0,0,0,1)$ | 4 | $\{11,1,12,5\}$ |
| 5 | $(0,1,-1,-1)$ | 5 | $\{11,14,13,2\}$ |
| 6 | $(4,-5,-11,6)$ | 6 | $\{15,3,16,7\}$ |
| 7 | $(1,-4,0,-4)$ | 7 | $\{15,17,18,6\}$ |
| 8 | $(4,1,1,0)$ | | |
| 9 | $(1,-5,1,0)$ | | |
| 10 | $(2,-1,-7,0)$ | | |
| 11 | $(0,0,1,-1)$ | | |
| 12 | $(0,2,1,1)$ | | |
| 13 | $(1,0,1,1)$ | | |
| 14 | $(2,0,-1,-1)$ | | |
| 15 | $(4,2,0,-1)$ | | |
| 16 | $(4,-5,0,6)$ | | |
| 17 | $(5,-9,7,2)$ | | |
| 18 | $(2,113,77,234)$ | | |

### S1.12. 4D panel (e): $D^{(4)}_{17,9}$

TABLE S13: Unnormalized vectors and all contexts of $D^{(4)}_{17,9}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 0 | $(0,0,0,1)$ | 1 | $\{0,1,2,3\}$ |
| 1 | $(0,1,2,0)$ | 2 | $\{0,7,8,9\}$ |
| 2 | $(1,0,0,0)$ | 3 | $\{0,4,5,6\}$ |
| 3 | $(0,2,-1,0)$ | 4 | $\{0,2,8,5\}$ |
| 4 | $(1,2,0,0)$ | 5 | $\{0,3,11,14\}$ |
| 5 | $(0,0,1,0)$ | 6 | $\{0,7,11,15\}$ |
| 6 | $(2,-1,0,0)$ | 7 | $\{0,9,10,23\}$ |
| 7 | $(1,0,1,0)$ | 8 | $\{0,4,11,24\}$ |
| 8 | $(0,1,0,0)$ | 9 | $\{0,4,10,18\}$ |
| 9 | $(1,0,-1,0)$ | | |
| 10 | $(2,-1,2,0)$ | | |
| 11 | $(2,-1,-2,0)$ | | |
| 14 | $(5,2,4,0)$ | | |
| 15 | $(1,4,-1,0)$ | | |
| 18 | $(4,-2,-5,0)$ | | |
| 23 | $(1,4,1,0)$ | | |
| 24 | $(4,-2,5,0)$ | | |

### S1.13. 4D panel (f): $L^{(4)}_{22,9}$

TABLE S14: Unnormalized vectors and all contexts of $L^{(4)}_{22,9}$.

| $i$ | $r_i$ | $j$ | Context |
|---|---|---|---|
| 1 | $(1,0,0,0)$ | 1 | $\{1,2,4,3\}$ |
| 2 | $(0,1,0,0)$ | 2 | $\{5,8,7,6\}$ |
| 3 | $(0,0,1,0)$ | 3 | $\{4,9,10,8\}$ |
| 4 | $(0,0,0,1)$ | 4 | $\{11,1,12,5\}$ |
| 5 | $(0,1,-1,-1)$ | 5 | $\{11,14,13,2\}$ |
| 6 | $(4,-5,-11,6)$ | 6 | $\{15,3,16,7\}$ |
| 7 | $(1,-4,0,-4)$ | 7 | $\{15,17,18,6\}$ |
| 8 | $(4,1,1,0)$ | 8 | $\{11,21,22,19\}$ |
| 9 | $(1,-5,1,0)$ | 9 | $\{13,20,15,19\}$ |
| 10 | $(2,-1,-7,0)$ | | |
| 11 | $(0,0,1,-1)$ | | |
| 12 | $(0,2,1,1)$ | | |
| 13 | $(1,0,4,4)$ | | |
| 14 | $(8,0,-1,-1)$ | | |
| 15 | $(4,2,0,-1)$ | | |
| 16 | $(4,-5,0,6)$ | | |
| 17 | $(5,-9,7,2)$ | | |
| 18 | $(2,113,77,234)$ | | |
| 19 | $(16,-33,-2,-2)$ | | |
| 20 | $(4,2,-21,20)$ | | |
| 21 | $(22,12,-11,-11)$ | | |
| 22 | $(129,44,153,153)$ | | |

### S1.14. Hesse $H_{37}$: vectors, contexts, and non-collinearity

This appendix uses exactly the vertex numbers and context numbers of Fig. 4. All indices $a, k$ in the family notation are in $\mathbb{Z}_3$. The vectors are unnormalized and the inner product is Hermitian. Table S15 lists every vector, squared norm, color, and incident context. Table S16 lists all 24 contexts with their actual vectors. Together they specify the faithful orthogonal representation completely.

#### *S1.14.1. All 37 vectors, their colors, and incident contexts*

TABLE S15: Complete enumeration of the 37 vectors of $H_{37}$. The color $c(i) \in \{0,1,2,3\}$ is the explicit four-coloring in the main text. Context indices refer to Table S16.

| $i$ | Ray | $v_i$ | $\|v_i\|^2$ | $c(i)$ | Incident contexts |
|---|---|---|---|---|---|
| 1 | $z$ | $(1,0,0)$ | 1 | 0 | $\{4,5,6\}$ |
| 2 | $M_{00}$ | $(1,1,1)$ | 3 | 0 | $\{1,7,16\}$ |
| 3 | $M_{01}$ | $(1,\zeta,\zeta^2)$ | 3 | 1 | $\{1,8,17\}$ |
| 4 | $M_{02}$ | $(1,\zeta^2,\zeta)$ | 3 | 2 | $\{1,9,18\}$ |
| 5 | $M_{10}$ | $(1,\zeta,\zeta)$ | 3 | 0 | $\{2,10,21\}$ |
| 6 | $M_{11}$ | $(1,\zeta^2,1)$ | 3 | 1 | $\{2,11,19\}$ |
| 7 | $M_{12}$ | $(1,1,\zeta^2)$ | 3 | 2 | $\{2,12,20\}$ |
| 8 | $M_{20}$ | $(1,\zeta^2,\zeta^2)$ | 3 | 0 | $\{3,13,23\}$ |
| 9 | $M_{21}$ | $(1,1,\zeta)$ | 3 | 1 | $\{3,14,24\}$ |
| 10 | $M_{22}$ | $(1,\zeta,1)$ | 3 | 2 | $\{3,15,22\}$ |
| 11 | $P_0$ | $(0,1,-1)$ | 2 | 3 | $\{4,7,10,13\}$ |
| 12 | $P_1$ | $(0,1,-\zeta)$ | 2 | 3 | $\{5,8,11,14\}$ |
| 13 | $P_2$ | $(0,1,-\zeta^2)$ | 2 | 3 | $\{6,9,12,15\}$ |
| 14 | $Q_0$ | $(-1,0,1)$ | 2 | 3 | $\{16,19,22\}$ |
| 15 | $Q_1$ | $(-\zeta,0,1)$ | 2 | 3 | $\{17,20,23\}$ |
| 16 | $Q_2$ | $(-\zeta^2,0,1)$ | 2 | 3 | $\{18,21,24\}$ |
| 17 | $A_0$ | $(0,1,1)$ | 2 | 1 | $\{4\}$ |

*Continued on the next page.*

TABLE S15: Complete enumeration of the 37 vectors of $H_{37}$. The color $c(i) \in \{0,1,2,3\}$ is the explicit four-coloring in the main text. Context indices refer to Table S16. (continued).

| $i$ | Ray | $v_i$ | $\|v_i\|^2$ | $c(i)$ | Incident contexts |
|---|---|---|---|---|---|
| 18 | $A_1$ | $(0,1,\zeta)$ | 2 | 1 | {5} |
| 19 | $A_2$ | $(0,1,\zeta^2)$ | 2 | 1 | {6} |
| 20 | $B_{00}$ | $(-2,1,1)$ | 6 | 1 | {7} |
| 21 | $B_{01}$ | $(-2,\zeta,\zeta^2)$ | 6 | 2 | {8} |
| 22 | $B_{02}$ | $(-2,\zeta^2,\zeta)$ | 6 | 0 | {9} |
| 23 | $B_{10}$ | $(-2,\zeta,\zeta)$ | 6 | 1 | {10} |
| 24 | $B_{11}$ | $(-2,\zeta^2,1)$ | 6 | 2 | {11} |
| 25 | $B_{12}$ | $(-2,1,\zeta^2)$ | 6 | 0 | {12} |
| 26 | $B_{20}$ | $(-2,\zeta^2,\zeta^2)$ | 6 | 1 | {13} |
| 27 | $B_{21}$ | $(-2,1,\zeta)$ | 6 | 2 | {14} |
| 28 | $B_{22}$ | $(-2,\zeta,1)$ | 6 | 0 | {15} |
| 29 | $C_{00}$ | $(1,-2,1)$ | 6 | 1 | {16} |
| 30 | $C_{01}$ | $(1,-2\zeta,\zeta^2)$ | 6 | 2 | {17} |
| 31 | $C_{02}$ | $(1,-2\zeta^2,\zeta)$ | 6 | 0 | {18} |
| 32 | $C_{10}$ | $(1,-2\zeta^2,1)$ | 6 | 2 | {19} |
| 33 | $C_{11}$ | $(1,-2,\zeta^2)$ | 6 | 0 | {20} |
| 34 | $C_{12}$ | $(1,-2\zeta,\zeta)$ | 6 | 1 | {21} |
| 35 | $C_{20}$ | $(1,-2\zeta,1)$ | 6 | 0 | {22} |
| 36 | $C_{21}$ | $(1,-2\zeta^2,\zeta^2)$ | 6 | 1 | {23} |
| 37 | $C_{22}$ | $(1,-2,\zeta)$ | 6 | 2 | {24} |

### *S1.14.2. All 24 contexts and their orthogonal vectors*

TABLE S16: All 24 orthogonal triads, numbered in the same order as the context colors in Fig. 4. Vectors are given in increasing order of their vertex numbers. Each triad contributes three orthogonality edges; the 72 resulting pairs are all the orthogonal pairs.

| Context | Family | Vertices | Unnormalized vectors |
|---|---|---|---|
| $e_1$ | $\mathcal{M}_0$ | $\{2,3,4\}$ | $\{(1,1,1),\ (1,\zeta,\zeta^2),\ (1,\zeta^2,\zeta)\}$ |
| $e_2$ | $\mathcal{M}_1$ | $\{5,6,7\}$ | $\{(1,\zeta,\zeta),\ (1,\zeta^2,1),\ (1,1,\zeta^2)\}$ |
| $e_3$ | $\mathcal{M}_2$ | $\{8,9,10\}$ | $\{(1,\zeta^2,\zeta^2),\ (1,1,\zeta),\ (1,\zeta,1)\}$ |
| $e_4$ | $\mathcal{A}_0$ | $\{1,11,17\}$ | $\{(1,0,0),\ (0,1,-1),\ (0,1,1)\}$ |
| $e_5$ | $\mathcal{A}_1$ | $\{1,12,18\}$ | $\{(1,0,0),\ (0,1,-\zeta),\ (0,1,\zeta)\}$ |
| $e_6$ | $\mathcal{A}_2$ | $\{1,13,19\}$ | $\{(1,0,0),\ (0,1,-\zeta^2),\ (0,1,\zeta^2)\}$ |
| $e_7$ | $\mathcal{B}_{00}$ | $\{2,11,20\}$ | $\{(1,1,1),\ (0,1,-1),\ (-2,1,1)\}$ |
| $e_8$ | $\mathcal{B}_{01}$ | $\{3,12,21\}$ | $\{(1,\zeta,\zeta^2),\ (0,1,-\zeta),\ (-2,\zeta,\zeta^2)\}$ |
| $e_9$ | $\mathcal{B}_{02}$ | $\{4,13,22\}$ | $\{(1,\zeta^2,\zeta),\ (0,1,-\zeta^2),\ (-2,\zeta^2,\zeta)\}$ |
| $e_{10}$ | $\mathcal{B}_{10}$ | $\{5,11,23\}$ | $\{(1,\zeta,\zeta),\ (0,1,-1),\ (-2,\zeta,\zeta)\}$ |
| $e_{11}$ | $\mathcal{B}_{11}$ | $\{6,12,24\}$ | $\{(1,\zeta^2,1),\ (0,1,-\zeta),\ (-2,\zeta^2,1)\}$ |
| $e_{12}$ | $\mathcal{B}_{12}$ | $\{7,13,25\}$ | $\{(1,1,\zeta^2),\ (0,1,-\zeta^2),\ (-2,1,\zeta^2)\}$ |
| $e_{13}$ | $\mathcal{B}_{20}$ | $\{8,11,26\}$ | $\{(1,\zeta^2,\zeta^2),\ (0,1,-1),\ (-2,\zeta^2,\zeta^2)\}$ |
| $e_{14}$ | $\mathcal{B}_{21}$ | $\{9,12,27\}$ | $\{(1,1,\zeta),\ (0,1,-\zeta),\ (-2,1,\zeta)\}$ |
| $e_{15}$ | $\mathcal{B}_{22}$ | $\{10,13,28\}$ | $\{(1,\zeta,1),\ (0,1,-\zeta^2),\ (-2,\zeta,1)\}$ |
| $e_{16}$ | $\mathcal{C}_{00}$ | $\{2,14,29\}$ | $\{(1,1,1),\ (-1,0,1),\ (1,-2,1)\}$ |

*Continued on the next page.*

TABLE S16: All 24 orthogonal triads, numbered in the same order as the context colors in Fig. 4. Vectors are given in increasing order of their vertex numbers. Each triad contributes three orthogonality edges; the 72 resulting pairs are all the orthogonal pairs. (continued).

| Context | Family | Vertices | Unnormalized vectors |
|---|---|---|---|
| $e_{17}$ | $\mathcal{C}_{01}$ | $\{3,15,30\}$ | $\{(1,\zeta,\zeta^2),\ (-\zeta,0,1),\ (1,-2\zeta,\zeta^2)\}$ |
| $e_{18}$ | $\mathcal{C}_{02}$ | $\{4,16,31\}$ | $\{(1,\zeta^2,\zeta),\ (-\zeta^2,0,1),\ (1,-2\zeta^2,\zeta)\}$ |
| $e_{19}$ | $\mathcal{C}_{10}$ | $\{6,14,32\}$ | $\{(1,\zeta^2,1),\ (-1,0,1),\ (1,-2\zeta^2,1)\}$ |
| $e_{20}$ | $\mathcal{C}_{11}$ | $\{7,15,33\}$ | $\{(1,1,\zeta^2),\ (-\zeta,0,1),\ (1,-2,\zeta^2)\}$ |
| $e_{21}$ | $\mathcal{C}_{12}$ | $\{5,16,34\}$ | $\{(1,\zeta,\zeta),\ (-\zeta^2,0,1),\ (1,-2\zeta,\zeta)\}$ |
| $e_{22}$ | $\mathcal{C}_{20}$ | $\{10,14,35\}$ | $\{(1,\zeta,1),\ (-1,0,1),\ (1,-2\zeta,1)\}$ |
| $e_{23}$ | $\mathcal{C}_{21}$ | $\{8,15,36\}$ | $\{(1,\zeta^2,\zeta^2),\ (-\zeta,0,1),\ (1,-2\zeta^2,\zeta^2)\}$ |
| $e_{24}$ | $\mathcal{C}_{22}$ | $\{9,16,37\}$ | $\{(1,1,\zeta),\ (-\zeta^2,0,1),\ (1,-2,\zeta)\}$ |

#### *S1.14.3. Norm shells and a non-collinearity certificate*

The four squared-norm shells have sizes

$$\#\{i : \|v_i\|^2 = 1\} = 1, \quad \#\{i : \|v_i\|^2 = 2\} = 9, \quad \#\{i : \|v_i\|^2 = 3\} = 9, \quad \#\{i : \|v_i\|^2 = 6\} = 18. \tag{S1}$$

Exact evaluation over the $\binom{37}{2} = 666$ unordered pairs gives

$$\max_{i<j} \frac{|\langle v_i, v_j\rangle|^2}{\|v_i\|^2\|v_j\|^2} = \frac{3}{4} < 1. \tag{S2}$$

For example, the maximum is attained by $v_{11} = P_0 = (0,1,-1)$ and $v_{29} = C_{00} = (1,-2,1)$:

$$\frac{|\langle v_{11}, v_{29}\rangle|^2}{\|v_{11}\|^2\|v_{29}\|^2} = \frac{(-3)^2}{2\cdot 6} = \frac{3}{4}. \tag{S3}$$

Thus no two printed vectors represent the same ray. The maximum and all 72 zero inner products are checked exactly by the accompanying verifier; the family-wise argument in the main text explains why there are no further orthogonalities.

### S1.15. Four-dimensional $H^{(4)}_{\mathrm{YO}}$ and $H_{65}$: vectors and contexts

For direct human and computational verification, this appendix enumerates all unnormalized integer vectors in $\mathbb{Z}^4$ and all four-element contexts for the two four-dimensional examples $H_{65}$ and $H^{(4)}_{\mathrm{YO}}$:

1. The *canonical linear hypergraph* $H_{65} = (V_{65}, E_{29})$ (Theorem 22, Appendix S1.15.1), which has exactly 65 **pairwise non-collinear integer vectors** (17 core vectors $v_1, \ldots, v_{17}$ and 48 degree-1 completion vectors $v_{18}, \ldots, v_{65}$), 29 **four-element contexts** (5 core contexts $C_1, \ldots, C_5$ and 24 bridge-completion contexts $E_1, \ldots, E_{24}$), and 174 **orthogonality edges** ($29 \times \binom{4}{2} = 174$).

2. The *non-linear warm-up apex suspension* $H^{(4)}_{\mathrm{YO}} = (V_{26}, E^{(4)}_{16})$ (Proposition 20, Appendix S1.15.2), which has exactly 26 **pairwise non-collinear integer vectors** (13 Yu–Oh rays $v_1, \ldots, v_{13}$, 12 completion rays $v_{14}, \ldots, v_{25}$ in $\mathbb{R}^3 \times \{0\}$, and 1 apex ray $v_{26}$), 16 **four-element contexts**, and 73 **orthogonality edges** ($16 \times \binom{3}{2} = 48$ edges in $\mathbb{R}^3 \times \{0\}$ plus 25 apex edges).

*Canonical primitive normal form and human-verifiable non-collinearity.* Every vector $v_i = (x_{i,1}, x_{i,2}, x_{i,3}, x_{i,4})^T \in \mathbb{Z}^4$ listed below in both $H_{65}$ and $H_{\mathrm{YO}}^{(4)}$ is written in *canonical primitive projective form*:

$$\begin{gathered} \gcd\big(|x_{i,1}|, |x_{i,2}|, |x_{i,3}|, |x_{i,4}|\big) = 1, \\ \text{first nonzero entry of } v_i > 0. \end{gathered} \tag{S4}$$

Because every projective ray $[v] \in \mathbb{P}(\mathbb{Q}^4)$ possesses a *unique* representative in $\mathbb{Z}^4$ satisfying Eq. (S4), two vectors $v_i, v_j$ in canonical primitive form are collinear (proportional over $\mathbb{R}$ or $\mathbb{C}$) *if and only if they are identical* 4*-tuples* ($v_i = v_j$) and, in particular, have the same squared Euclidean norm $\|v_i\|^2 = \|v_j\|^2$.

#### *S1.15.1. The canonical linear* 65*-vertex,* 29*-context hypergraph* $H_{65}$

Table S17 enumerates all 65 unnormalized integer vectors $v_1, \ldots, v_{65} \in \mathbb{Z}^4$ of $H_{65}$, together with their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0, 1, 2, 3, 4\}$ in the strong 5-coloring extending Eq. (S28), and the contexts in $E_{29}$ to which each vertex belongs. Table S18 enumerates all 29 contexts $C_1, \ldots, C_5, E_1, \ldots, E_{24} \in E_{29}$, displaying side by side both the vertex indices $\{i, j, k, \ell\}$ and the four mutually orthogonal integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

To verify by hand that all 65 vectors in Table S17 are pairwise non-collinear across all $\binom{65}{2} = 2{,}080$ pairs, observe that all 65 tuples satisfy Eq. (S4) and partition by squared norm $\|v_i\|^2$ into seven small, disjoint norm shells ($4 + 6 + 7 + 12 + 12 + 12 + 12 = 65$):

- $\|v_i\|^2 = 1$ (4 vectors): indices $\{1, 2, 3, 4\}$, the 4 standard coordinate basis vectors;
- $\|v_i\|^2 = 2$ (6 vectors): indices $\{12, \ldots, 17\}$, one vector for each of the $\binom{4}{2} = 6$ coordinate-pair supports;
- $\|v_i\|^2 = 4$ (7 vectors): indices $\{5, \ldots, 11\}$, seven distinct $\pm 1$ vectors with $x_{i,1} = +1$;
- $\|v_i\|^2 = 11$ (12 vectors): indices $\{18, 20, 22, 25, 26, 28, 30, 33, 34, 36, 39, 41\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct sign/entry patterns of $\{0, \pm 1, \pm 1, \pm 3\}$;
- $\|v_i\|^2 = 18$ (12 vectors): indices $\{43, 45, 46, 49, 51, 53, 54, 56, 59, 61, 62, 65\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 2, \pm 2, \pm 3\}$;
- $\|v_i\|^2 = 22$ (12 vectors): indices $\{19, 21, 23, 24, 27, 29, 31, 32, 35, 37, 38, 40\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct patterns of $\{0, \pm 2, \pm 3, \pm 3\}$;
- $\|v_i\|^2 = 36$ (12 vectors): indices $\{42, 44, 47, 48, 50, 52, 55, 57, 58, 60, 63, 64\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 1, \pm 3, \pm 5\}$.

Because no two 4-tuples within any of the seven norm shells coincide, no two of the 65 vectors are collinear; quantitatively, the maximum squared cosine over all $\binom{65}{2} = 2{,}080$ pairs is $\max_{1 \le i < j \le 65} (v_i \cdot v_j)^2 / (\|v_i\|^2 \|v_j\|^2) = 19^2/(11 \times 36) = 361/396 \approx 0.9116 < 1$, attained by $(v_{41}, v_{47})$, whose squared norms are 11 and 36 and whose inner product is 19.

TABLE S17: Complete enumeration of the 65 unnormalized integer vectors $v_i \in \mathbb{Z}^4$ of $H_{65}$, their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0,1,2,3,4\}$, and the contexts in $E_{29}$ containing vertex $i$.

| $i$ | $v_i \in \mathbb{Z}^4$ | $\|v_i\|^2$ | $c(i)$ | Contexts containing $i$ | $i$ | $v_i \in \mathbb{Z}^4$ | $\|v_i\|^2$ | $c(i)$ | Contexts containing $i$ |
|---|---|---|---|---|---|---|---|---|---|
| 1 | $(1,0,0,0)$ | 1 | 0 | $C_1, E_3, E_7, E_{11}$ | 34 | $(1,-1,0,3)$ | 11 | 1 | $E_9 = \{3,16,34,35\}$ |
| 2 | $(0,1,0,0)$ | 1 | 1 | $C_1, E_1, E_5, E_{12}$ | 35 | $(3,-3,0,-2)$ | 22 | 3 | $E_9 = \{3,16,34,35\}$ |
| 3 | $(0,0,1,0)$ | 1 | 2 | $C_1, E_2, E_8, E_9$ | 36 | $(1,-1,3,0)$ | 11 | 1 | $E_{10} = \{4,16,36,37\}$ |
| 4 | $(0,0,0,1)$ | 1 | 3 | $C_1, E_4, E_6, E_{10}$ | 37 | $(3,-3,-2,0)$ | 22 | 2 | $E_{10} = \{4,16,36,37\}$ |
| 5 | $(1,1,1,1)$ | 4 | 0 | $C_2, E_{13}, E_{19}, E_{23}$ | 38 | $(0,2,3,3)$ | 22 | 1 | $E_{11} = \{1,17,38,39\}$ |
| 6 | $(1,1,-1,-1)$ | 4 | 1 | $C_2, E_{15}, E_{17}, E_{24}$ | 39 | $(0,3,-1,-1)$ | 11 | 3 | $E_{11} = \{1,17,38,39\}$ |
| 7 | $(1,-1,1,-1)$ | 4 | 2 | $C_2, E_{16}, E_{20}, E_{21}$ | 40 | $(2,0,-3,-3)$ | 22 | 0 | $E_{12} = \{2,17,40,41\}$ |
| 8 | $(1,-1,-1,1)$ | 4 | 4 | $C_2, E_{14}, E_{18}, E_{22}$ | 41 | $(3,0,1,1)$ | 11 | 3 | $E_{12} = \{2,17,40,41\}$ |
| 9 | $(1,1,-1,1)$ | 4 | 0 | $C_3, C_4$ | 42 | $(1,-5,3,1)$ | 36 | 1 | $E_{13} = \{5,12,42,43\}$ |
| 10 | $(1,-1,1,1)$ | 4 | 1 | $C_3, C_5$ | 43 | $(2,-1,-3,2)$ | 18 | 2 | $E_{13} = \{5,12,42,43\}$ |
| 11 | $(1,-1,-1,-1)$ | 4 | 3 | $C_4, C_5$ | 44 | $(1,-3,5,1)$ | 36 | 0 | $E_{14} = \{8,12,44,45\}$ |
| 12 | $(1,0,0,-1)$ | 2 | 3 | $C_3, E_1, E_2, E_{13}, E_{14}$ | 45 | $(2,3,1,2)$ | 18 | 1 | $E_{14} = \{8,12,44,45\}$ |
| 13 | $(0,1,1,0)$ | 2 | 4 | $C_3, E_3, E_4, E_{15}, E_{16}$ | 46 | $(1,-2,2,-3)$ | 18 | 0 | $E_{15} = \{6,13,46,47\}$ |
| 14 | $(1,0,1,0)$ | 2 | 2 | $C_4, E_5, E_6, E_{17}, E_{18}$ | 47 | $(5,-1,1,3)$ | 36 | 2 | $E_{15} = \{6,13,46,47\}$ |
| 15 | $(0,1,0,-1)$ | 2 | 1 | $C_4, E_7, E_8, E_{19}, E_{20}$ | 48 | $(3,-1,1,5)$ | 36 | 0 | $E_{16} = \{7,13,48,49\}$ |
| 16 | $(1,1,0,0)$ | 2 | 0 | $C_5, E_9, E_{10}, E_{21}, E_{22}$ | 49 | $(3,2,-2,-1)$ | 18 | 1 | $E_{16} = \{7,13,48,49\}$ |
| 17 | $(0,0,1,-1)$ | 2 | 2 | $C_5, E_{11}, E_{12}, E_{23}, E_{24}$ | 50 | $(1,3,-1,5)$ | 36 | 0 | $E_{17} = \{6,14,50,51\}$ |
| 18 | $(1,0,-3,1)$ | 11 | 0 | $E_1 = \{2,12,18,19\}$ | 51 | $(2,-3,-2,1)$ | 18 | 3 | $E_{17} = \{6,14,50,51\}$ |
| 19 | $(3,0,2,3)$ | 22 | 2 | $E_1 = \{2,12,18,19\}$ | 52 | $(1,5,-1,3)$ | 36 | 0 | $E_{18} = \{8,14,52,53\}$ |
| 20 | $(1,3,0,1)$ | 11 | 0 | $E_2 = \{3,12,20,21\}$ | 53 | $(2,1,-2,-3)$ | 18 | 1 | $E_{18} = \{8,14,52,53\}$ |
| 21 | $(3,-2,0,3)$ | 22 | 1 | $E_2 = \{3,12,20,21\}$ | 54 | $(3,-2,1,-2)$ | 18 | 2 | $E_{19} = \{5,15,54,55\}$ |
| 22 | $(0,1,-1,3)$ | 11 | 1 | $E_3 = \{1,13,22,23\}$ | 55 | $(3,1,-5,1)$ | 36 | 3 | $E_{19} = \{5,15,54,55\}$ |
| 23 | $(0,3,-3,-2)$ | 22 | 2 | $E_3 = \{1,13,22,23\}$ | 56 | $(1,2,3,2)$ | 18 | 0 | $E_{20} = \{7,15,56,57\}$ |
| 24 | $(2,-3,3,0)$ | 22 | 0 | $E_4 = \{4,13,24,25\}$ | 57 | $(5,1,-3,1)$ | 36 | 3 | $E_{20} = \{7,15,56,57\}$ |
| 25 | $(3,1,-1,0)$ | 11 | 1 | $E_4 = \{4,13,24,25\}$ | 58 | $(1,-1,-5,-3)$ | 36 | 1 | $E_{21} = \{7,16,58,59\}$ |
| 26 | $(1,0,-1,-3)$ | 11 | 0 | $E_5 = \{2,14,26,27\}$ | 59 | $(2,-2,-1,3)$ | 18 | 3 | $E_{21} = \{7,16,58,59\}$ |
| 27 | $(3,0,-3,2)$ | 22 | 3 | $E_5 = \{2,14,26,27\}$ | 60 | $(1,-1,-3,-5)$ | 36 | 1 | $E_{22} = \{8,16,60,61\}$ |
| 28 | $(1,-3,-1,0)$ | 11 | 0 | $E_6 = \{4,14,28,29\}$ | 61 | $(2,-2,3,-1)$ | 18 | 2 | $E_{22} = \{8,16,60,61\}$ |
| 29 | $(3,2,-3,0)$ | 22 | 1 | $E_6 = \{4,14,28,29\}$ | 62 | $(1,3,-2,-2)$ | 18 | 1 | $E_{23} = \{5,17,62,63\}$ |
| 30 | $(0,1,3,1)$ | 11 | 2 | $E_7 = \{1,15,30,31\}$ | 63 | $(5,-3,-1,-1)$ | 36 | 3 | $E_{23} = \{5,17,62,63\}$ |
| 31 | $(0,3,-2,3)$ | 22 | 3 | $E_7 = \{1,15,30,31\}$ | 64 | $(3,-5,-1,-1)$ | 36 | 0 | $E_{24} = \{6,17,64,65\}$ |
| 32 | $(2,3,0,3)$ | 22 | 0 | $E_8 = \{3,15,32,33\}$ | 65 | $(3,1,2,2)$ | 18 | 3 | $E_{24} = \{6,17,64,65\}$ |
| 33 | $(3,-1,0,-1)$ | 11 | 3 | $E_8 = \{3,15,32,33\}$ | | | | | |

TABLE S18: Complete enumeration of the 29 four-element contexts of $H_{65} = (V_{65}, E_{29})$, listing for each context both its vertex indices $\{i,j,k,\ell\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

| Context | Vertex set | $v_i$ | $v_j$ | $v_k$ | $v_\ell$ |
|---|---|---|---|---|---|
| *Five core 4-contexts $C_1, \ldots, C_5$ on $V_{\text{core}} = \{1,\ldots,17\}$:* | | | | | |
| $C_1$ | $\{1,2,3,4\}$ | $v_1 = (1,0,0,0)$ | $v_2 = (0,1,0,0)$ | $v_3 = (0,0,1,0)$ | $v_4 = (0,0,0,1)$ |
| $C_2$ | $\{5,6,7,8\}$ | $v_5 = (1,1,1,1)$ | $v_6 = (1,1,-1,-1)$ | $v_7 = (1,-1,1,-1)$ | $v_8 = (1,-1,-1,1)$ |
| $C_3$ | $\{9,10,12,13\}$ | $v_9 = (1,1,-1,1)$ | $v_{10} = (1,-1,1,1)$ | $v_{12} = (1,0,0,-1)$ | $v_{13} = (0,1,1,0)$ |
| $C_4$ | $\{9,11,14,15\}$ | $v_9 = (1,1,-1,1)$ | $v_{11} = (1,-1,-1,-1)$ | $v_{14} = (1,0,1,0)$ | $v_{15} = (0,1,0,-1)$ |
| $C_5$ | $\{10,11,16,17\}$ | $v_{10} = (1,-1,1,1)$ | $v_{11} = (1,-1,-1,-1)$ | $v_{16} = (1,1,0,0)$ | $v_{17} = (0,0,1,-1)$ |
| *Twelve coordinate-to-bridge completion contexts $E_1, \ldots, E_{12}$ (between $C_1$ and $B_1 \cup B_2 \cup B_3$):* | | | | | |
| $E_1$ | $\{2,12,18,19\}$ | $v_2 = (0,1,0,0)$ | $v_{12} = (1,0,0,-1)$ | $v_{18} = (1,0,-3,1)$ | $v_{19} = (3,0,2,3)$ |
| $E_2$ | $\{3,12,20,21\}$ | $v_3 = (0,0,1,0)$ | $v_{12} = (1,0,0,-1)$ | $v_{20} = (1,3,0,1)$ | $v_{21} = (3,-2,0,3)$ |
| $E_3$ | $\{1,13,22,23\}$ | $v_1 = (1,0,0,0)$ | $v_{13} = (0,1,1,0)$ | $v_{22} = (0,1,-1,3)$ | $v_{23} = (0,3,-3,-2)$ |
| $E_4$ | $\{4,13,24,25\}$ | $v_4 = (0,0,0,1)$ | $v_{13} = (0,1,1,0)$ | $v_{24} = (2,-3,3,0)$ | $v_{25} = (3,1,-1,0)$ |
| $E_5$ | $\{2,14,26,27\}$ | $v_2 = (0,1,0,0)$ | $v_{14} = (1,0,1,0)$ | $v_{26} = (1,0,-1,-3)$ | $v_{27} = (3,0,-3,2)$ |
| $E_6$ | $\{4,14,28,29\}$ | $v_4 = (0,0,0,1)$ | $v_{14} = (1,0,1,0)$ | $v_{28} = (1,-3,-1,0)$ | $v_{29} = (3,2,-3,0)$ |
| $E_7$ | $\{1,15,30,31\}$ | $v_1 = (1,0,0,0)$ | $v_{15} = (0,1,0,-1)$ | $v_{30} = (0,1,3,1)$ | $v_{31} = (0,3,-2,3)$ |
| $E_8$ | $\{3,15,32,33\}$ | $v_3 = (0,0,1,0)$ | $v_{15} = (0,1,0,-1)$ | $v_{32} = (2,3,0,3)$ | $v_{33} = (3,-1,0,-1)$ |
| $E_9$ | $\{3,16,34,35\}$ | $v_3 = (0,0,1,0)$ | $v_{16} = (1,1,0,0)$ | $v_{34} = (1,-1,0,3)$ | $v_{35} = (3,-3,0,-2)$ |
| $E_{10}$ | $\{4,16,36,37\}$ | $v_4 = (0,0,0,1)$ | $v_{16} = (1,1,0,0)$ | $v_{36} = (1,-1,3,0)$ | $v_{37} = (3,-3,-2,0)$ |
| $E_{11}$ | $\{1,17,38,39\}$ | $v_1 = (1,0,0,0)$ | $v_{17} = (0,0,1,-1)$ | $v_{38} = (0,2,3,3)$ | $v_{39} = (0,3,-1,-1)$ |
| $E_{12}$ | $\{2,17,40,41\}$ | $v_2 = (0,1,0,0)$ | $v_{17} = (0,0,1,-1)$ | $v_{40} = (2,0,-3,-3)$ | $v_{41} = (3,0,1,1)$ |
| *Twelve Hadamard-to-bridge completion contexts $E_{13}, \ldots, E_{24}$ (between $C_2$ and $B_1 \cup B_2 \cup B_3$):* | | | | | |
| $E_{13}$ | $\{5,12,42,43\}$ | $v_5 = (1,1,1,1)$ | $v_{12} = (1,0,0,-1)$ | $v_{42} = (1,-5,3,1)$ | $v_{43} = (2,-1,-3,2)$ |
| $E_{14}$ | $\{8,12,44,45\}$ | $v_8 = (1,-1,-1,1)$ | $v_{12} = (1,0,0,-1)$ | $v_{44} = (1,-3,5,1)$ | $v_{45} = (2,3,1,2)$ |

*Continued on the next page.*

TABLE S18: All 29 contexts of $H_{65}$ (continued).

| Context | Vertex set | $v_i$ | $v_j$ | $v_k$ | $v_\ell$ |
|---|---|---|---|---|---|
| $E_{15}$ | $\{6,13,46,47\}$ | $v_6=(1,1,-1,-1)$ | $v_{13}=(0,1,1,0)$ | $v_{46}=(1,-2,2,-3)$ | $v_{47}=(5,-1,1,3)$ |
| $E_{16}$ | $\{7,13,48,49\}$ | $v_7=(1,-1,1,-1)$ | $v_{13}=(0,1,1,0)$ | $v_{48}=(3,-1,1,5)$ | $v_{49}=(3,2,-2,-1)$ |
| $E_{17}$ | $\{6,14,50,51\}$ | $v_6=(1,1,-1,-1)$ | $v_{14}=(1,0,1,0)$ | $v_{50}=(1,3,-1,5)$ | $v_{51}=(2,-3,-2,1)$ |
| $E_{18}$ | $\{8,14,52,53\}$ | $v_8=(1,-1,-1,1)$ | $v_{14}=(1,0,1,0)$ | $v_{52}=(1,5,-1,3)$ | $v_{53}=(2,1,-2,-3)$ |
| $E_{19}$ | $\{5,15,54,55\}$ | $v_5=(1,1,1,1)$ | $v_{15}=(0,1,0,-1)$ | $v_{54}=(3,-2,1,-2)$ | $v_{55}=(3,1,-5,1)$ |
| $E_{20}$ | $\{7,15,56,57\}$ | $v_7=(1,-1,1,-1)$ | $v_{15}=(0,1,0,-1)$ | $v_{56}=(1,2,3,2)$ | $v_{57}=(5,1,-3,1)$ |
| $E_{21}$ | $\{7,16,58,59\}$ | $v_7=(1,-1,1,-1)$ | $v_{16}=(1,1,0,0)$ | $v_{58}=(1,-1,-5,-3)$ | $v_{59}=(2,-2,-1,3)$ |
| $E_{22}$ | $\{8,16,60,61\}$ | $v_8=(1,-1,-1,1)$ | $v_{16}=(1,1,0,0)$ | $v_{60}=(1,-1,-3,-5)$ | $v_{61}=(2,-2,3,-1)$ |
| $E_{23}$ | $\{5,17,62,63\}$ | $v_5=(1,1,1,1)$ | $v_{17}=(0,0,1,-1)$ | $v_{62}=(1,3,-2,-2)$ | $v_{63}=(5,-3,-1,-1)$ |
| $E_{24}$ | $\{6,17,64,65\}$ | $v_6=(1,1,-1,-1)$ | $v_{17}=(0,0,1,-1)$ | $v_{64}=(3,-5,-1,-1)$ | $v_{65}=(3,1,2,2)$ |

#### *S1.15.2. The non-linear warm-up 26-vertex, 16-context apex suspension $H^{(4)}_{\mathrm{YO}}$*

For completeness, the 26 unnormalized integer vectors $v_1,\dots,v_{26}\in\mathbb{Z}^4$ of the non-linear warm-up hypergraph $H^{(4)}_{\mathrm{YO}}=(V_{26},E^{(4)}_{16})$ in Proposition 20 (written in canonical primitive form satisfying Eq. (S4)) are:

$$
\begin{aligned}
v_1&=(0,1,1,0), & v_2&=(1,0,0,0),\\
v_3&=(0,1,-1,0), & v_4&=(1,1,0,0),\\
v_5&=(0,0,1,0), & v_6&=(1,-1,0,0),\\
v_7&=(1,0,1,0), & v_8&=(0,1,0,0),\\
v_9&=(1,0,-1,0), & v_{10}&=(1,-1,1,0),\\
v_{11}&=(1,-1,-1,0), & v_{12}&=(1,1,1,0),\\
v_{13}&=(1,1,-1,0), & v_{14}&=(2,1,1,0),\\
v_{15}&=(1,2,-1,0), & v_{16}&=(1,-2,1,0),\\
v_{17}&=(1,1,-2,0), & v_{18}&=(1,-1,-2,0),\\
v_{19}&=(2,1,-1,0), & v_{20}&=(2,-1,1,0),\\
v_{21}&=(1,-2,-1,0), & v_{22}&=(1,1,2,0),\\
v_{23}&=(1,2,1,0), & v_{24}&=(1,-1,2,0),\\
v_{25}&=(2,-1,-1,0), & v_{26}&=(0,0,0,1).
\end{aligned}
\tag{S5}
$$

These 26 canonical primitive vectors partition by squared norm $\|v_i\|^2$ into four disjoint norm shells $(4+6+4+12=26)$:

- $\|v_i\|^2=1$ (4 vectors): $\{v_2,v_5,v_8,v_{26}\}$, the four coordinate basis vectors of $\mathbb{R}^4$;
- $\|v_i\|^2=2$ (6 vectors): $\{v_1,v_3,v_4,v_6,v_7,v_9\}$, the six binary rays in $\mathbb{R}^3\times\{0\}$;
- $\|v_i\|^2=3$ (4 vectors): $\{v_{10},v_{11},v_{12},v_{13}\}$, the four triad rays $(1,\pm1,\pm1,0)^T$;
- $\|v_i\|^2=6$ (12 vectors): $\{v_{14},\dots,v_{25}\}$, all $3\times4=12$ canonical primitive permutations and sign patterns of $(\pm2,\pm1,\pm1,0)^T$.

Since all 26 canonical 4-tuples are distinct, all $\binom{26}{2}=325$ pairs of vectors are strictly non-collinear, with maximum squared cosine $\max_{1\le i<j\le26}(v_i\cdot v_j)^2/(\|v_i\|^2\|v_j\|^2)=4^2/(3\times6)=8/9\approx0.8889<1$. Table S19 lists all 16 four-element contexts of $E^{(4)}_{16}$ side by side with their four mutually orthogonal integer vectors.

TABLE S19: Complete enumeration of the 16 four-element contexts of the non-linear warm-up apex suspension $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, listing for each context both its vertex indices $\{i, j, k, 26\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_{26}\} \subset \mathbb{Z}^4$.

| Context | Vertex set | $v_i$ | $v_j$ | $v_k$ | $v_{26}$ |
|---|---|---|---|---|---|
| $e_1^{(4)}$ | $\{1,2,3,26\}$ | $v_1 = (0,1,1,0)$ | $v_2 = (1,0,0,0)$ | $v_3 = (0,1,-1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_2^{(4)}$ | $\{7,8,9,26\}$ | $v_7 = (1,0,1,0)$ | $v_8 = (0,1,0,0)$ | $v_9 = (1,0,-1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_3^{(4)}$ | $\{4,5,6,26\}$ | $v_4 = (1,1,0,0)$ | $v_5 = (0,0,1,0)$ | $v_6 = (1,-1,0,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_4^{(4)}$ | $\{2,5,8,26\}$ | $v_2 = (1,0,0,0)$ | $v_5 = (0,0,1,0)$ | $v_8 = (0,1,0,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_5^{(4)}$ | $\{3,12,25,26\}$ | $v_3 = (0,1,-1,0)$ | $v_{12} = (1,1,1,0)$ | $v_{25} = (2,-1,-1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_6^{(4)}$ | $\{9,12,16,26\}$ | $v_9 = (1,0,-1,0)$ | $v_{12} = (1,1,1,0)$ | $v_{16} = (1,-2,1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_7^{(4)}$ | $\{6,12,17,26\}$ | $v_6 = (1,-1,0,0)$ | $v_{12} = (1,1,1,0)$ | $v_{17} = (1,1,-2,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_8^{(4)}$ | $\{3,11,14,26\}$ | $v_3 = (0,1,-1,0)$ | $v_{11} = (1,-1,-1,0)$ | $v_{14} = (2,1,1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_9^{(4)}$ | $\{7,11,15,26\}$ | $v_7 = (1,0,1,0)$ | $v_{11} = (1,-1,-1,0)$ | $v_{15} = (1,2,-1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{10}^{(4)}$ | $\{4,11,24,26\}$ | $v_4 = (1,1,0,0)$ | $v_{11} = (1,-1,-1,0)$ | $v_{24} = (1,-1,2,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{11}^{(4)}$ | $\{9,10,23,26\}$ | $v_9 = (1,0,-1,0)$ | $v_{10} = (1,-1,1,0)$ | $v_{23} = (1,2,1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{12}^{(4)}$ | $\{1,10,19,26\}$ | $v_1 = (0,1,1,0)$ | $v_{10} = (1,-1,1,0)$ | $v_{19} = (2,1,-1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{13}^{(4)}$ | $\{4,10,18,26\}$ | $v_4 = (1,1,0,0)$ | $v_{10} = (1,-1,1,0)$ | $v_{18} = (1,-1,-2,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{14}^{(4)}$ | $\{1,13,20,26\}$ | $v_1 = (0,1,1,0)$ | $v_{13} = (1,1,-1,0)$ | $v_{20} = (2,-1,1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{15}^{(4)}$ | $\{7,13,21,26\}$ | $v_7 = (1,0,1,0)$ | $v_{13} = (1,1,-1,0)$ | $v_{21} = (1,-2,-1,0)$ | $v_{26} = (0,0,0,1)$ |
| $e_{16}^{(4)}$ | $\{6,13,22,26\}$ | $v_6 = (1,-1,0,0)$ | $v_{13} = (1,1,-1,0)$ | $v_{22} = (1,1,2,0)$ | $v_{26} = (0,0,0,1)$ |

## S1.16. Explicit parent configurations and inclusion maps

### *S1.16.1. The 49-ray real Yu–Oh parent and published Bub labels*

Table S20 lists all rays, including the original 25. The final column gives the exact labels in the published full 49-ray encoding[1] (p. 14, Sec. 5(xi)) of Bub's set[2]. Under this map the 36 bases in Table S21 equal all 36 published contexts; their 108 pairs equal all geometric orthogonalities. In this labeling 26–33 are Eq. (S31), and 34–49 complete the sixteen remaining orthogonal pairs. The first 25 vectors are those of Table S1.

TABLE S20: All unnormalized vectors of the real 49-ray parent, with the verified published Bub labels.

| $i$ | $v_i$ | Published label |
|---|---|---|
| 1 | $(0,1,1)$ | `G` |
| 2 | $(1,0,0)$ | `1` |
| 3 | $(0,1,-1)$ | `P` |
| 4 | $(1,1,0)$ | `H` |
| 5 | $(0,0,1)$ | `3` |
| 6 | $(1,-1,0)$ | `U` |
| 7 | $(1,0,1)$ | `Q` |
| 8 | $(0,1,0)$ | `2` |
| 9 | $(1,0,-1)$ | `e` |
| 10 | $(1,-1,1)$ | `D` |
| 11 | $(1,-1,-1)$ | `h` |
| 12 | $(1,1,1)$ | `S` |
| 13 | $(1,1,-1)$ | `a` |
| 14 | $(2,1,1)$ | `g` |
| 15 | $(1,2,-1)$ | `n` |
| 16 | $(1,-2,1)$ | `d` |
| 17 | $(1,1,-2)$ | `T` |
| 18 | $(1,-1,-2)$ | `F` |
| 19 | $(2,1,-1)$ | `E` |
| 20 | $(2,-1,1)$ | `I` |
| 21 | $(1,-2,-1)$ | `k` |
| 22 | $(1,1,2)$ | `b` |
| 23 | $(1,2,1)$ | `A` |

| $i$ | $v_i$ | Published label |
|---|---|---|
| 24 | $(1,-1,2)$ | N |
| 25 | $(2,-1,-1)$ | R |
| 26 | $(0,1,2)$ | i |
| 27 | $(0,2,-1)$ | O |
| 28 | $(0,1,-2)$ | 6 |
| 29 | $(0,2,1)$ | 7 |
| 30 | $(1,2,0)$ | 5 |
| 31 | $(2,-1,0)$ | 4 |
| 32 | $(1,-2,0)$ | V |
| 33 | $(2,1,0)$ | j |
| 34 | $(2,1,-5)$ | Z |
| 35 | $(5,-2,1)$ | m |
| 36 | $(1,2,5)$ | W |
| 37 | $(5,2,-1)$ | c |
| 38 | $(1,-2,-5)$ | f |
| 39 | $(5,-1,2)$ | L |
| 40 | $(5,1,2)$ | 9 |
| 41 | $(2,1,5)$ | 8 |
| 42 | $(2,-1,-5)$ | J |
| 43 | $(5,2,1)$ | X |
| 44 | $(1,-2,5)$ | l |
| 45 | $(5,-1,-2)$ | Y |
| 46 | $(5,-2,-1)$ | B |
| 47 | $(1,2,-5)$ | C |
| 48 | $(5,1,-2)$ | M |
| 49 | $(2,-1,5)$ | K |

TABLE S21: All 36 complete bases of the real parent.

| $e$ | Context | $e$ | Context | $e$ | Context |
|---|---|---|---|---|---|
| 1 | $\{1,2,3\}$ | 2 | $\{1,10,19\}$ | 3 | $\{1,13,20\}$ |
| 4 | $\{2,5,8\}$ | 5 | $\{2,26,27\}$ | 6 | $\{2,28,29\}$ |
| 7 | $\{3,11,14\}$ | 8 | $\{3,12,25\}$ | 9 | $\{4,5,6\}$ |
| 10 | $\{4,10,18\}$ | 11 | $\{4,11,24\}$ | 12 | $\{5,30,31\}$ |
| 13 | $\{5,32,33\}$ | 14 | $\{6,12,17\}$ | 15 | $\{6,13,22\}$ |
| 16 | $\{7,8,9\}$ | 17 | $\{7,11,15\}$ | 18 | $\{7,13,21\}$ |
| 19 | $\{9,10,23\}$ | 20 | $\{9,12,16\}$ | 21 | $\{14,32,34\}$ |
| 22 | $\{15,26,35\}$ | 23 | $\{15,31,36\}$ | 24 | $\{16,26,37\}$ |
| 25 | $\{16,33,38\}$ | 26 | $\{17,29,39\}$ | 27 | $\{18,27,40\}$ |
| 28 | $\{19,32,41\}$ | 29 | $\{20,30,42\}$ | 30 | $\{21,28,43\}$ |
| 31 | $\{21,33,44\}$ | 32 | $\{22,27,45\}$ | 33 | $\{23,28,46\}$ |
| 34 | $\{23,31,47\}$ | 35 | $\{24,29,48\}$ | 36 | $\{25,30,49\}$ |

#### *S1.16.2. The 87-ray Hesse parent*

Use $\zeta = \exp(2\pi i/3)$ and $\zeta^2 + \zeta + 1 = 0$. The first 37 rays retain exactly the Hesse numbering; projective representatives have their first nonzero coordinate one. The inventory labels use the generation convention of Sec. VII B. Tables S22 and S23 list all 87 rays and all 64 complete bases. The six additional orthogonal pairs outside the 64 complete bases are listed separately: this larger parent's full orthogonality graph need not equal its context two-section. The retained Hesse restriction has all its original orthogonalities covered by contexts.

TABLE S22: All 87 unnormalized rays of the first Hesse/lineage union.

| $i$ | $v_i$ | Inventory label |
|---|---|---|
| 1 | $(1,0,0)$ | $a_{11}$ |
| 2 | $(1,1,1)$ | $u_1$ |
| 3 | $(1,\zeta,-1-\zeta)$ | $u_2$ |
| 4 | $(1,-1-\zeta,\zeta)$ | $u_3$ |
| 5 | $(1,\zeta,\zeta)$ | $u_4$ |
| 6 | $(1,-1-\zeta,1)$ | $u_5$ |
| 7 | $(1,1,-1-\zeta)$ | $u_6$ |
| 8 | $(1,-1-\zeta,-1-\zeta)$ | $u_7$ |
| 9 | $(1,1,\zeta)$ | $u_9$ |
| 10 | $(1,\zeta,1)$ | $u_8$ |
| 11 | $(0,1,-1)$ | $c_{11}$ |
| 12 | $(0,1,-\zeta)$ | $c_{12}$ |
| 13 | $(0,1,1+\zeta)$ | $c_{13}$ |
| 14 | $(1,0,-1)$ | $c_{21}$ |
| 15 | $(1,0,1+\zeta)$ | $c_{22}$ |
| 16 | $(1,0,-\zeta)$ | $c_{23}$ |
| 17 | $(0,1,1)$ | $b_{11}$ |
| 18 | $(0,1,\zeta)$ | $b_{12}$ |
| 19 | $(0,1,-1-\zeta)$ | $b_{13}$ |
| 20 | $(1,\frac{-1}{2},\frac{-1}{2})$ | $d_{11}$ |
| 21 | $(1,-\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$ | $d_{12}$ |
| 22 | $(1,\frac{1}{2}+\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$ | $d_{13}$ |
| 23 | $(1,-\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$ | $d_{14}$ |
| 24 | $(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{-1}{2})$ | $d_{15}$ |
| 25 | $(1,\frac{-1}{2},\frac{1}{2}+\frac{1}{2}\zeta)$ | $d_{16}$ |
| 26 | $(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$ | $d_{17}$ |
| 27 | $(1,\frac{-1}{2},-\frac{1}{2}\zeta)$ | $d_{19}$ |
| 28 | $(1,-\frac{1}{2}\zeta,\frac{-1}{2})$ | $d_{18}$ |
| 29 | $(1,-2,1)$ | $d_{21}$ |
| 30 | $(1,-2\zeta,-1-\zeta)$ | $d_{22}$ |
| 31 | $(1,2+2\zeta,\zeta)$ | $d_{23}$ |
| 32 | $(1,2+2\zeta,1)$ | $d_{25}$ |
| 33 | $(1,-2,-1-\zeta)$ | $d_{26}$ |
| 34 | $(1,-2\zeta,\zeta)$ | $d_{24}$ |
| 35 | $(1,-2\zeta,1)$ | $d_{28}$ |
| 36 | $(1,2+2\zeta,-1-\zeta)$ | $d_{27}$ |
| 37 | $(1,-2,\zeta)$ | $d_{29}$ |
| 38 | $(0,1,0)$ | $a_{21}$ |
| 39 | $(0,0,1)$ | $a_{31}$ |
| 40 | $(1,\frac{-1}{2},\frac{1}{2})$ | $b_{1121}$ |
| 41 | $(1,-\frac{1}{2}\zeta,\frac{-1}{2}-\frac{1}{2}\zeta)$ | $b_{1122}$ |
| 42 | $(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{1}{2}\zeta)$ | $b_{1123}$ |
| 43 | $(1,\frac{1}{2},\frac{-1}{2})$ | $b_{1131}$ |
| 44 | $(1,\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$ | $b_{1132}$ |
| 45 | $(1,\frac{-1}{2}-\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$ | $b_{1133}$ |
| 46 | $(1,1,-1)$ | $b_{121}$ |
| 47 | $(1,\zeta,1+\zeta)$ | $b_{122}$ |
| 48 | $(1,-1-\zeta,-\zeta)$ | $b_{123}$ |
| 49 | $(1,-1,1)$ | $b_{131}$ |
| 50 | $(1,-\zeta,-1-\zeta)$ | $b_{132}$ |
| 51 | $(1,1+\zeta,\zeta)$ | $b_{133}$ |
| 52 | $(1,0,1)$ | $b_{21}$ |
| 53 | $(1,-2,-1)$ | $b_{2121}$ |
| 54 | $(1,-2\zeta,1+\zeta)$ | $b_{2122}$ |
| 55 | $(1,2+2\zeta,-\zeta)$ | $b_{2123}$ |
| 56 | $(1,0,-1-\zeta)$ | $b_{22}$ |
| 57 | $(1,2,-1)$ | $b_{2231}$ |
| 58 | $(1,2\zeta,1+\zeta)$ | $b_{2232}$ |
| 59 | $(1,-2-2\zeta,-\zeta)$ | $b_{2233}$ |
| 60 | $(1,0,\zeta)$ | $b_{23}$ |
| 61 | $(1,-1,-1)$ | $b_{231}$ |
| 62 | $(1,-\zeta,1+\zeta)$ | $b_{232}$ |
| 63 | $(1,1+\zeta,-\zeta)$ | $b_{233}$ |
| 64 | $(1,1,0)$ | $b_{31}$ |
| 65 | $(1,-1,-2)$ | $b_{3131}$ |

| $i$ | $v_i$ | Inventory label |
|---|---|---|
| 66 | $(1, -\zeta, 2+2\zeta)$ | $b_{3132}$ |
| 67 | $(1, 1+\zeta, -2\zeta)$ | $b_{3133}$ |
| 68 | $(1, \zeta, 0)$ | $b_{32}$ |
| 69 | $(1, -1, 2)$ | $b_{3231}$ |
| 70 | $(1, -\zeta, -2-2\zeta)$ | $b_{3232}$ |
| 71 | $(1, 1+\zeta, 2\zeta)$ | $b_{3233}$ |
| 72 | $(1, -1-\zeta, 0)$ | $b_{33}$ |
| 73 | $(1, -1, 0)$ | $c_{31}$ |
| 74 | $(1, -\zeta, 0)$ | $c_{32}$ |
| 75 | $(1, 1+\zeta, 0)$ | $c_{33}$ |
| 76 | $(1, 1, -2)$ | $d_{31}$ |
| 77 | $(1, \zeta, 2+2\zeta)$ | $d_{32}$ |
| 78 | $(1, -1-\zeta, -2\zeta)$ | $d_{33}$ |
| 79 | $(1, 2, 1)$ | $e_{11}$ |
| 80 | $(1, 2\zeta, -1-\zeta)$ | $e_{12}$ |
| 81 | $(1, -2-2\zeta, \zeta)$ | $e_{13}$ |
| 82 | $(1, 1, 2)$ | $e_{21}$ |
| 83 | $(1, \zeta, -2-2\zeta)$ | $e_{22}$ |
| 84 | $(1, -1-\zeta, 2\zeta)$ | $e_{23}$ |
| 85 | $(1, \frac{1}{2}, \frac{1}{2})$ | $e_{31}$ |
| 86 | $(1, \frac{1}{2}\zeta, \frac{-1}{2} - \frac{1}{2}\zeta)$ | $e_{32}$ |
| 87 | $(1, \frac{-1}{2} - \frac{1}{2}\zeta, \frac{1}{2}\zeta)$ | $e_{33}$ |

TABLE S23: All 64 complete bases of the first 87-ray Hesse parent.

| $e$ | Context | $e$ | Context | $e$ | Context |
|---|---|---|---|---|---|
| 1 | $\{1, 11, 17\}$ | 2 | $\{1, 12, 18\}$ | 3 | $\{1, 13, 19\}$ |
| 4 | $\{1, 38, 39\}$ | 5 | $\{2, 3, 4\}$ | 6 | $\{2, 11, 20\}$ |
| 7 | $\{2, 14, 29\}$ | 8 | $\{2, 73, 76\}$ | 9 | $\{3, 12, 21\}$ |
| 10 | $\{3, 15, 30\}$ | 11 | $\{3, 74, 77\}$ | 12 | $\{4, 13, 22\}$ |
| 13 | $\{4, 16, 31\}$ | 14 | $\{4, 75, 78\}$ | 15 | $\{5, 6, 7\}$ |
| 16 | $\{5, 11, 23\}$ | 17 | $\{5, 16, 34\}$ | 18 | $\{6, 12, 24\}$ |
| 19 | $\{6, 14, 32\}$ | 20 | $\{7, 13, 25\}$ | 21 | $\{7, 15, 33\}$ |
| 22 | $\{8, 9, 10\}$ | 23 | $\{8, 11, 26\}$ | 24 | $\{8, 15, 36\}$ |
| 25 | $\{9, 12, 27\}$ | 26 | $\{9, 16, 37\}$ | 27 | $\{10, 13, 28\}$ |
| 28 | $\{10, 14, 35\}$ | 29 | $\{11, 61, 85\}$ | 30 | $\{12, 62, 86\}$ |
| 31 | $\{13, 63, 87\}$ | 32 | $\{14, 38, 52\}$ | 33 | $\{14, 49, 79\}$ |
| 34 | $\{15, 38, 56\}$ | 35 | $\{15, 50, 80\}$ | 36 | $\{16, 38, 60\}$ |
| 37 | $\{16, 51, 81\}$ | 38 | $\{17, 40, 46\}$ | 39 | $\{17, 43, 49\}$ |
| 40 | $\{18, 41, 47\}$ | 41 | $\{18, 44, 50\}$ | 42 | $\{19, 42, 48\}$ |
| 43 | $\{19, 45, 51\}$ | 44 | $\{39, 64, 73\}$ | 45 | $\{39, 68, 74\}$ |
| 46 | $\{39, 72, 75\}$ | 47 | $\{46, 47, 48\}$ | 48 | $\{46, 52, 53\}$ |
| 49 | $\{46, 73, 82\}$ | 50 | $\{47, 54, 56\}$ | 51 | $\{47, 74, 83\}$ |
| 52 | $\{48, 55, 60\}$ | 53 | $\{48, 75, 84\}$ | 54 | $\{49, 50, 51\}$ |
| 55 | $\{49, 64, 65\}$ | 56 | $\{50, 66, 68\}$ | 57 | $\{51, 67, 72\}$ |
| 58 | $\{52, 57, 61\}$ | 59 | $\{56, 58, 62\}$ | 60 | $\{59, 60, 63\}$ |
| 61 | $\{61, 62, 63\}$ | 62 | $\{61, 64, 69\}$ | 63 | $\{62, 68, 70\}$ |
| 64 | $\{63, 71, 72\}$ | | | | |

Its six further orthogonal pairs are $\{5, 74\}$, $\{6, 75\}$, $\{7, 73\}$, $\{8, 75\}$, $\{9, 73\}$, $\{10, 74\}$.

#### *S1.16.3. Peres24 and the complete four-dimensional extensions*

The coordinates of $H_{65}$ and $H_{\mathrm{YO}}^{(4)}$ are those already listed above. Table S24 explicitly enumerates the known Peres rays. In each union, retain the original local labels and append the missing rays in the order displayed in Table S25. Tables S26 and S27 list every available complete basis. Full orthogonality-pair lists accompany the coordinates in the data; the 40-ray union has additional pairs outside its complete contexts.

TABLE S24: All 24 primitive unnormalized Peres rays, in the certificate order.

| $i$ | $p_i$ |
|---|---|
| 1 | $(1,0,0,0)$ |
| 2 | $(0,1,0,0)$ |
| 3 | $(0,0,1,0)$ |
| 4 | $(0,0,0,1)$ |
| 5 | $(1,-1,0,0)$ |
| 6 | $(1,1,0,0)$ |
| 7 | $(1,0,-1,0)$ |
| 8 | $(1,0,1,0)$ |
| 9 | $(1,0,0,-1)$ |
| 10 | $(1,0,0,1)$ |
| 11 | $(0,1,-1,0)$ |
| 12 | $(0,1,1,0)$ |
| 13 | $(0,1,0,-1)$ |
| 14 | $(0,1,0,1)$ |
| 15 | $(0,0,1,-1)$ |
| 16 | $(0,0,1,1)$ |
| 17 | $(1,-1,-1,-1)$ |
| 18 | $(1,-1,-1,1)$ |
| 19 | $(1,-1,1,-1)$ |
| 20 | $(1,-1,1,1)$ |
| 21 | $(1,1,-1,-1)$ |
| 22 | $(1,1,-1,1)$ |
| 23 | $(1,1,1,-1)$ |
| 24 | $(1,1,1,1)$ |

TABLE S25: Every ray added to the original four-dimensional examples to obtain the stated parents.

| Original | New label | Added vector |
|---|---|---|
| $H_{\mathrm{YO}}^{(4)}$ | 27 | $(1,0,0,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 28 | $(1,0,0,1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 29 | $(0,1,0,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 30 | $(0,1,0,1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 31 | $(0,0,1,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 32 | $(0,0,1,1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 33 | $(1,-1,-1,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 34 | $(1,-1,-1,1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 35 | $(1,-1,1,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 36 | $(1,-1,1,1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 37 | $(1,1,-1,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 38 | $(1,1,-1,1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 39 | $(1,1,1,-1)$ |
| $H_{\mathrm{YO}}^{(4)}$ | 40 | $(1,1,1,1)$ |
| $H_{65}$ | 66 | $(1,-1,0,0)$ |
| $H_{65}$ | 67 | $(1,0,-1,0)$ |
| $H_{65}$ | 68 | $(1,0,0,1)$ |
| $H_{65}$ | 69 | $(0,1,-1,0)$ |
| $H_{65}$ | 70 | $(0,1,0,1)$ |
| $H_{65}$ | 71 | $(0,0,1,1)$ |
| $H_{65}$ | 72 | $(1,1,1,-1)$ |

TABLE S26: All 36 complete bases of the 40-ray parent.

| $e$ | Context | $e$ | Context |
|---|---|---|---|
| 1 | $\{1,2,3,26\}$ | 2 | $\{1,3,27,28\}$ |
| 3 | $\{1,10,19,26\}$ | 4 | $\{1,13,20,26\}$ |
| 5 | $\{1,27,36,38\}$ | 6 | $\{1,28,35,37\}$ |
| 7 | $\{2,5,8,26\}$ | 8 | $\{2,5,29,30\}$ |
| 9 | $\{2,8,31,32\}$ | 10 | $\{3,11,14,26\}$ |
| 11 | $\{3,12,25,26\}$ | 12 | $\{3,27,34,40\}$ |
| 13 | $\{3,28,33,39\}$ | 14 | $\{4,5,6,26\}$ |
| 15 | $\{4,6,31,32\}$ | 16 | $\{4,10,18,26\}$ |
| 17 | $\{4,11,24,26\}$ | 18 | $\{4,31,33,36\}$ |
| 19 | $\{4,32,34,35\}$ | 20 | $\{5,8,27,28\}$ |
| 21 | $\{6,12,17,26\}$ | 22 | $\{6,13,22,26\}$ |
| 23 | $\{6,31,37,40\}$ | 24 | $\{6,32,38,39\}$ |
| 25 | $\{7,8,9,26\}$ | 26 | $\{7,9,29,30\}$ |
| 27 | $\{7,11,15,26\}$ | 28 | $\{7,13,21,26\}$ |
| 29 | $\{7,29,33,38\}$ | 30 | $\{7,30,34,37\}$ |
| 31 | $\{9,10,23,26\}$ | 32 | $\{9,12,16,26\}$ |
| 33 | $\{9,29,35,40\}$ | 34 | $\{9,30,36,39\}$ |
| 35 | $\{33,36,38,39\}$ | 36 | $\{34,35,37,40\}$ |

TABLE S27: All 48 complete bases of the 72-ray parent.

| $e$ | Context | $e$ | Context |
|---|---|---|---|
| 1 | $\{1,2,3,4\}$ | 2 | $\{1,2,17,71\}$ |
| 3 | $\{1,3,15,70\}$ | 4 | $\{1,4,13,69\}$ |
| 5 | $\{1,13,22,23\}$ | 6 | $\{1,15,30,31\}$ |
| 7 | $\{1,17,38,39\}$ | 8 | $\{2,3,12,68\}$ |
| 9 | $\{2,4,14,67\}$ | 10 | $\{2,12,18,19\}$ |
| 11 | $\{2,14,26,27\}$ | 12 | $\{2,17,40,41\}$ |
| 13 | $\{3,4,16,66\}$ | 14 | $\{3,12,20,21\}$ |
| 15 | $\{3,15,32,33\}$ | 16 | $\{3,16,34,35\}$ |
| 17 | $\{4,13,24,25\}$ | 18 | $\{4,14,28,29\}$ |
| 19 | $\{4,16,36,37\}$ | 20 | $\{5,6,7,8\}$ |
| 21 | $\{5,6,17,66\}$ | 22 | $\{5,7,15,67\}$ |
| 23 | $\{5,8,12,69\}$ | 24 | $\{5,12,42,43\}$ |
| 25 | $\{5,15,54,55\}$ | 26 | $\{5,17,62,63\}$ |
| 27 | $\{6,7,13,68\}$ | 28 | $\{6,8,14,70\}$ |
| 29 | $\{6,13,46,47\}$ | 30 | $\{6,14,50,51\}$ |
| 31 | $\{6,17,64,65\}$ | 32 | $\{7,8,16,71\}$ |
| 33 | $\{7,13,48,49\}$ | 34 | $\{7,15,56,57\}$ |
| 35 | $\{7,16,58,59\}$ | 36 | $\{8,12,44,45\}$ |
| 37 | $\{8,14,52,53\}$ | 38 | $\{8,16,60,61\}$ |
| 39 | $\{9,10,11,72\}$ | 40 | $\{9,10,12,13\}$ |
| 41 | $\{9,11,14,15\}$ | 42 | $\{9,66,71,72\}$ |
| 43 | $\{10,11,16,17\}$ | 44 | $\{10,67,70,72\}$ |
| 45 | $\{11,68,69,72\}$ | 46 | $\{12,13,68,69\}$ |
| 47 | $\{14,15,67,70\}$ | 48 | $\{16,17,66,71\}$ |

## S1.17. Cabello completions and the two 68-ray parents

### *S1.17.1. The eighteen-ray input and faithful 36-ray completion*

All labels in this subsection are local. In $C_{18,9}$ and $C_{36,18}$, $18 = a$, $19 = b$, and $20 = c$; labels 4 and 5 are absent. They do not denote the completion rays of $H_{65}$. The core is the first eighteen rows of Table S28; labels 21–38 are fresh. The contexts of $C_{18,9}$ are the first nine rows of Table S29.

TABLE S28: All 36 unnormalized vectors of the faithful Cabello completion.

| $i$ | $v_i$ |
|---|---|
| 1 | $(1,0,0,0)$ |
| 2 | $(0,1,0,0)$ |
| 3 | $(0,0,1,0)$ |
| 6 | $(1,1,-1,-1)$ |
| 7 | $(1,-1,1,-1)$ |
| 8 | $(1,-1,-1,1)$ |
| 9 | $(1,1,-1,1)$ |
| 10 | $(1,-1,1,1)$ |
| 11 | $(1,-1,-1,-1)$ |
| 12 | $(1,0,0,-1)$ |
| 13 | $(0,1,1,0)$ |
| 14 | $(1,0,1,0)$ |
| 15 | $(0,1,0,-1)$ |
| 16 | $(1,1,0,0)$ |
| 17 | $(0,0,1,-1)$ |
| 18 | $(1,0,0,1)$ |
| 19 | $(0,1,0,1)$ |
| 20 | $(0,0,1,1)$ |
| 21 | $(0,1,-1,3)$ |
| 22 | $(0,3,-3,-2)$ |
| 23 | $(1,0,-1,3)$ |
| 24 | $(3,0,-3,-2)$ |
| 25 | $(1,-1,0,3)$ |
| 26 | $(3,-3,0,-2)$ |
| 27 | $(3,1,2,2)$ |
| 28 | $(3,-5,-1,-1)$ |
| 29 | $(3,2,1,2)$ |
| 30 | $(3,-1,-5,-1)$ |
| 31 | $(2,3,1,2)$ |
| 32 | $(1,-3,5,1)$ |
| 33 | $(4,2,3,-3)$ |
| 34 | $(5,-7,-1,1)$ |
| 35 | $(4,3,2,-3)$ |
| 36 | $(5,-1,-7,1)$ |
| 37 | $(3,4,2,-3)$ |
| 38 | $(1,-5,7,-1)$ |

TABLE S29: All eighteen full bases of the faithful Cabello completion.

| $e$ | Context |
|---|---|
| 1 | $\{9,10,12,13\}$ |
| 2 | $\{9,11,14,15\}$ |
| 3 | $\{10,11,16,17\}$ |
| 4 | $\{2,3,12,18\}$ |
| 5 | $\{1,3,15,19\}$ |
| 6 | $\{1,2,17,20\}$ |
| 7 | $\{6,7,13,18\}$ |
| 8 | $\{6,8,14,19\}$ |
| 9 | $\{7,8,16,20\}$ |
| 10 | $\{1,13,21,22\}$ |
| 11 | $\{2,14,23,24\}$ |
| 12 | $\{3,16,25,26\}$ |
| 13 | $\{6,17,27,28\}$ |
| 14 | $\{7,15,29,30\}$ |
| 15 | $\{8,12,31,32\}$ |
| 16 | $\{9,20,33,34\}$ |
| 17 | $\{10,19,35,36\}$ |
| 18 | $\{11,18,37,38\}$ |

The orthogonal matrix $Q = M/\sqrt{2}$, with

$$M = \begin{pmatrix} 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \qquad M^T M = 2I_4, \tag{S6}$$

maps the published eighteen-ray set and all nine bases of Ref.[3] (Eqs. (1)–(9)) onto this realization. Table S30 gives the complete ray map; the input representatives are unnormalized and signs are immaterial.

TABLE S30: Exact ray map from the originally published Cabello coordinates.

| Published vector $u$ | Local label of $[Qu]$ |
|---|---|
| $(0,0,0,1)$ | 12 |
| $(0,0,1,0)$ | 13 |
| $(0,0,1,1)$ | 11 |
| $(0,1,-1,0)$ | 8 |
| $(0,1,0,-1)$ | 1 |
| $(0,1,0,0)$ | 18 |
| $(1,-1,-1,-1)$ | 19 |
| $(1,-1,-1,1)$ | 16 |
| $(1,-1,0,0)$ | 9 |
| $(1,-1,1,-1)$ | 17 |
| $(1,0,-1,0)$ | 2 |
| $(1,0,0,-1)$ | 7 |
| $(1,0,0,1)$ | 6 |
| $(1,0,1,0)$ | 3 |
| $(1,1,-1,1)$ | 15 |
| $(1,1,0,0)$ | 10 |
| $(1,1,1,-1)$ | 14 |
| $(1,1,1,1)$ | 20 |

#### *S1.17.2. An injective 45-ray completion after deleting $c$*

Delete local ray 20 and contexts 6 and 9 from the first nine contexts above. The seven remaining bases are followed by fourteen fresh-pair contexts in Table S32. The next table prints all 45 vectors; 4, 5, and 20 are absent and 21–48 are fresh. Its only uncovered orthogonal pairs are $\{16,21\}, \{16,22\}, \{17,35\}, \{17,36\}$.

TABLE S31: All 45 unnormalized vectors of the vertex-deleted completion.

| $i$ | $v_i$ |
|---|---|
| 1 | $(1,0,0,0)$ |
| 2 | $(0,1,0,0)$ |
| 3 | $(0,0,1,0)$ |
| 6 | $(1,1,-1,-1)$ |
| 7 | $(1,-1,1,-1)$ |
| 8 | $(1,-1,-1,1)$ |
| 9 | $(1,1,-1,1)$ |
| 10 | $(1,-1,1,1)$ |
| 11 | $(1,-1,-1,-1)$ |
| 12 | $(1,0,0,-1)$ |
| 13 | $(0,1,1,0)$ |
| 14 | $(1,0,1,0)$ |
| 15 | $(0,1,0,-1)$ |
| 16 | $(1,1,0,0)$ |
| 17 | $(0,0,1,-1)$ |
| 18 | $(1,0,0,1)$ |
| 19 | $(0,1,0,1)$ |
| 21 | $(0,0,1,2)$ |
| 22 | $(0,0,2,-1)$ |
| 23 | $(0,1,-1,3)$ |
| 24 | $(0,3,-3,-2)$ |
| 25 | $(0,1,2,2)$ |
| 26 | $(0,4,-1,-1)$ |
| 27 | $(1,0,-1,3)$ |
| 28 | $(3,0,-3,-2)$ |
| 29 | $(1,0,2,2)$ |
| 30 | $(4,0,-1,-1)$ |
| 31 | $(1,-1,0,3)$ |
| 32 | $(3,-3,0,-2)$ |
| 33 | $(5,3,4,4)$ |
| 34 | $(7,-9,-1,-1)$ |
| 35 | $(1,1,3,3)$ |
| 36 | $(3,3,-1,-1)$ |
| 37 | $(5,4,3,4)$ |
| 38 | $(7,-1,-9,-1)$ |
| 39 | $(2,-2,-1,3)$ |
| 40 | $(1,-1,-5,-3)$ |
| 41 | $(4,5,3,4)$ |
| 42 | $(1,-7,9,1)$ |
| 43 | $(4,-4,-1,-9)$ |
| 44 | $(5,-5,13,3)$ |
| 45 | $(6,5,4,-5)$ |
| 46 | $(9,-1,-11,1)$ |
| 47 | $(3,4,2,-3)$ |
| 48 | $(1,-5,7,-1)$ |

TABLE S32: All 21 full bases of the vertex-deleted completion.

| $e$ | Context |
|---|---|
| 1 | $\{9, 10, 12, 13\}$ |
| 2 | $\{9, 11, 14, 15\}$ |
| 3 | $\{10, 11, 16, 17\}$ |
| 4 | $\{2, 3, 12, 18\}$ |
| 5 | $\{1, 3, 15, 19\}$ |
| 6 | $\{6, 7, 13, 18\}$ |
| 7 | $\{6, 8, 14, 19\}$ |
| 8 | $\{1, 2, 21, 22\}$ |
| 9 | $\{1, 13, 23, 24\}$ |
| 10 | $\{1, 17, 25, 26\}$ |
| 11 | $\{2, 14, 27, 28\}$ |
| 12 | $\{2, 17, 29, 30\}$ |
| 13 | $\{3, 16, 31, 32\}$ |
| 14 | $\{6, 17, 33, 34\}$ |
| 15 | $\{7, 8, 35, 36\}$ |
| 16 | $\{7, 15, 37, 38\}$ |
| 17 | $\{7, 16, 39, 40\}$ |
| 18 | $\{8, 12, 41, 42\}$ |
| 19 | $\{8, 16, 43, 44\}$ |
| 20 | $\{10, 19, 45, 46\}$ |
| 21 | $\{11, 18, 47, 48\}$ |

#### *S1.17.3. All vectors and bases of the two 68-ray parents*

In each parent retain all 65 vectors of Appendix S1.15.1, with their original labels, and add the three vectors in the corresponding rows of Table S33; Tables S34 and S35 list all 35 bases of each parent. Thus the six rows specify every added ray; the two parent inventories use distinct local meanings for labels 66–68. These labels also differ from the 72-ray parent numbering in Appendix S1.16.

TABLE S33: The three added rays in each local 68-ray parent inventory.

| Parent variant | $i$ | $v_i$ |
|---|---|---|
| 18–9 subsystem | 66 | $(1, 0, 0, 1)$ |
| 18–9 subsystem | 67 | $(0, 1, 0, 1)$ |
| 18–9 subsystem | 68 | $(0, 0, 1, 1)$ |
| 20–11 subsystem | 66 | $(0, 1, -1, 0)$ |
| 20–11 subsystem | 67 | $(1, 0, -1, 0)$ |
| 20–11 subsystem | 68 | $(1, -1, 0, 0)$ |

TABLE S34: All 35 full bases of 68-ray parent variant 1.

| $e$ | Context | $e$ | Context |
|---|---|---|---|
| 1 | $\{1,2,3,4\}$ | 2 | $\{1,2,17,68\}$ |
| 3 | $\{1,3,15,67\}$ | 4 | $\{1,13,22,23\}$ |
| 5 | $\{1,15,30,31\}$ | 6 | $\{1,17,38,39\}$ |
| 7 | $\{2,3,12,66\}$ | 8 | $\{2,12,18,19\}$ |
| 9 | $\{2,14,26,27\}$ | 10 | $\{2,17,40,41\}$ |
| 11 | $\{3,12,20,21\}$ | 12 | $\{3,15,32,33\}$ |
| 13 | $\{3,16,34,35\}$ | 14 | $\{4,13,24,25\}$ |
| 15 | $\{4,14,28,29\}$ | 16 | $\{4,16,36,37\}$ |
| 17 | $\{5,6,7,8\}$ | 18 | $\{5,12,42,43\}$ |
| 19 | $\{5,15,54,55\}$ | 20 | $\{5,17,62,63\}$ |
| 21 | $\{6,7,13,66\}$ | 22 | $\{6,8,14,67\}$ |
| 23 | $\{6,13,46,47\}$ | 24 | $\{6,14,50,51\}$ |
| 25 | $\{6,17,64,65\}$ | 26 | $\{7,8,16,68\}$ |
| 27 | $\{7,13,48,49\}$ | 28 | $\{7,15,56,57\}$ |
| 29 | $\{7,16,58,59\}$ | 30 | $\{8,12,44,45\}$ |
| 31 | $\{8,14,52,53\}$ | 32 | $\{8,16,60,61\}$ |
| 33 | $\{9,10,12,13\}$ | 34 | $\{9,11,14,15\}$ |
| 35 | $\{10,11,16,17\}$ | | |

TABLE S35: All 35 full bases of 68-ray parent variant 2.

| $e$ | Context | $e$ | Context |
|---|---|---|---|
| 1 | $\{1,2,3,4\}$ | 2 | $\{1,4,13,66\}$ |
| 3 | $\{1,13,22,23\}$ | 4 | $\{1,15,30,31\}$ |
| 5 | $\{1,17,38,39\}$ | 6 | $\{2,4,14,67\}$ |
| 7 | $\{2,12,18,19\}$ | 8 | $\{2,14,26,27\}$ |
| 9 | $\{2,17,40,41\}$ | 10 | $\{3,4,16,68\}$ |
| 11 | $\{3,12,20,21\}$ | 12 | $\{3,15,32,33\}$ |
| 13 | $\{3,16,34,35\}$ | 14 | $\{4,13,24,25\}$ |
| 15 | $\{4,14,28,29\}$ | 16 | $\{4,16,36,37\}$ |
| 17 | $\{5,6,7,8\}$ | 18 | $\{5,6,17,68\}$ |
| 19 | $\{5,7,15,67\}$ | 20 | $\{5,8,12,66\}$ |
| 21 | $\{5,12,42,43\}$ | 22 | $\{5,15,54,55\}$ |
| 23 | $\{5,17,62,63\}$ | 24 | $\{6,13,46,47\}$ |
| 25 | $\{6,14,50,51\}$ | 26 | $\{6,17,64,65\}$ |
| 27 | $\{7,13,48,49\}$ | 28 | $\{7,15,56,57\}$ |
| 29 | $\{7,16,58,59\}$ | 30 | $\{8,12,44,45\}$ |
| 31 | $\{8,14,52,53\}$ | 32 | $\{8,16,60,61\}$ |
| 33 | $\{9,10,12,13\}$ | 34 | $\{9,11,14,15\}$ |
| 35 | $\{10,11,16,17\}$ | | |

For variant 1 the nine-context parity subsystem consists of exactly $C_3, C_4, C_5, D_1, \ldots, D_6$ with $a = 66, b = 67, c = 68$. For variant 2 the twenty-ray subsystem consists of all rows whose labels lie in $\{1, \ldots, 17, 66, 67, 68\}$. The machine-readable inventories repeat every coordinate, basis, and inclusion map.

## S2. TWO-VALUED STATES, EXTENDABILITY CERTIFICATES, COLORINGS, AND SEPARATION PROFILES

### S2.1. State, coloring, odd-cycle certificate, and profile tables for chromatically complete configurations ($\chi(H) = d$)

This subsection provides the complete state and coloring enumerations and odd-cycle nonextendability certificates for $H_{6,3}$ (Table S36), $P_5$ (Table S37), $H_8$ (Tables S38 and S39), $H_{17,10}$ (Tables S40 and S41), $H_{21,13}$ (Tables S42, S43, and S44), the vertex separation profiles and Born probability decomposition on $H_{17,10}$ and $H_{21,13}$ (Tables S45–S47), the complete state and normalized coloring tables for all faithful 3- and 4-dimensional logical gadgets (Tables S48–S69 and S70), and the ten chromatically inseparable faithful restrictions of $H_{\mathrm{YO}}^{+}$ (Table S71).

TABLE S36: All two-valued states of $H_{6,3}$. The last column gives an odd cycle in the complement of each nonextendable support.

| $j$ | support of $s_j$ | extendable | odd cycle |
|---|---|---|---|
| 1 | $\{1,4\}$ | yes | – |
| 2 | $\{2,4,6\}$ | no | $(3,1,5)$ |
| 3 | $\{2,5\}$ | yes | – |
| 4 | $\{3,6\}$ | yes | – |

TABLE S37: All two-valued states of $P_5$. The last column gives an odd cycle in the complement of each nonextendable support.

| $j$ | support of $s_j$ | extendable | odd cycle |
|---|---|---|---|
| 1 | $\{1,3,9\}$ | yes | – |
| 2 | $\{1,4,7\}$ | yes | – |
| 3 | $\{1,7,8,9\}$ | yes | – |
| 4 | $\{2,4,10\}$ | yes | – |
| 5 | $\{2,5,8\}$ | yes | – |
| 6 | $\{2,8,9,10\}$ | yes | – |
| 7 | $\{3,5,6\}$ | yes | – |
| 8 | $\{3,6,9,10\}$ | yes | – |
| 9 | $\{4,6,7,10\}$ | yes | – |
| 10 | $\{5,6,7,8\}$ | yes | – |
| 11 | $\{6,7,8,9,10\}$ | no | $(3,2,1,5,4)$ |

TABLE S38: All two-valued states of $H_8$. The last column gives an odd cycle in the complement of each nonextendable support.

| $j$ | support of $s_j$ | extendable | odd cycle |
|---|---|---|---|
| 1 | $\{1,4,6,9\}$ | yes | – |
| 2 | $\{1,5,8,10,14\}$ | no | $(9,7,6,12,11)$ |
| 3 | $\{1,5,9,13,14\}$ | yes | – |
| 4 | $\{2,4,6,8,11\}$ | yes | – |
| 5 | $\{2,4,7,11,14\}$ | no | $(5,3,1,12,6)$ |
| 6 | $\{2,5,8,10,12\}$ | yes | – |
| 7 | $\{2,5,8,11,13,14\}$ | yes | – |
| 8 | $\{2,5,9,12,13\}$ | no | $(4,3,1,11,10)$ |
| 9 | $\{3,6,8,11,13\}$ | no | $(7,5,4,10,9)$ |
| 10 | $\{3,7,10,12\}$ | yes | – |
| 11 | $\{3,7,11,13,14\}$ | yes | – |

TABLE S39. All strong three-colorings of $H_8$ up to color permutation. The anchor is $(1,2,3) = (0,1,2)$.

| $i$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $c_1$ | 0 | 1 | 2 | 0 | 1 | 0 | 2 | 1 | 0 | 1 | 2 | 1 | 2 | 2 |
| $c_2$ | 0 | 1 | 2 | 0 | 1 | 0 | 2 | 1 | 0 | 2 | 1 | 2 | 1 | 1 |
| $c_3$ | 0 | 1 | 2 | 1 | 0 | 1 | 2 | 1 | 0 | 2 | 1 | 2 | 0 | 0 |

TABLE S40: All two-valued states of $H_{17,10}$. The last column gives an odd cycle in the complement of each nonextendable support.

| $j$ | support of $s_j$ | extendable | odd cycle |
|---|---|---|---|
| 1 | $\{1,4,8,14,15,23\}$ | yes | – |
| 2 | $\{1,5,7,14,18,23,24\}$ | no | $(4,11,3,2,8,9,10)$ |
| 3 | $\{1,5,9,11,18\}$ | yes | – |
| 4 | $\{1,5,9,14,15,18,24\}$ | no | $(11,3,2,8,7)$ |
| 5 | $\{1,6,8,11,18,23\}$ | yes | – |
| 6 | $\{1,6,8,14,15,18,23,24\}$ | no | $(11,3,2,5,4)$ |
| 7 | $\{2,4,7,14,19,23\}$ | yes | – |
| 8 | $\{2,4,9,14,15,19\}$ | yes | – |
| 9 | $\{2,6,7,10,14,24\}$ | yes | – |
| 10 | $\{2,6,7,14,18,19,23,24\}$ | no | $(11,3,1,10,4)$ |
| 11 | $\{2,6,9,11,18,19\}$ | yes | – |
| 12 | $\{2,6,9,14,15,18,19,24\}$ | no | $(11,3,1,10,4)$ |
| 13 | $\{3,4,8,15,19,23\}$ | yes | – |
| 14 | $\{3,5,7,10,24\}$ | yes | – |
| 15 | $\{3,5,7,18,19,23,24\}$ | no | $(8,2,1,10,9)$ |
| 16 | $\{3,5,9,15,18,19,24\}$ | no | $(7,8,2,1,10,4,11)$ |
| 17 | $\{3,6,8,10,15,24\}$ | yes | – |
| 18 | $\{3,6,8,15,18,19,23,24\}$ | no | $(5,2,1,10,4)$ |

TABLE S41. All strong three-colorings of $H_{17,10}$ up to color permutation. The anchor is $(2,8,5) = (0,1,2)$.

| $i$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 14 | 15 | 18 | 19 | 23 | 24 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $c_1$ | 1 | 0 | 2 | 0 | 2 | 1 | 2 | 1 | 0 | 2 | 1 | 0 | 0 | 1 | 0 | 1 | 2 |
| $c_2$ | 1 | 0 | 2 | 1 | 2 | 0 | 2 | 1 | 0 | 2 | 0 | 1 | 1 | 0 | 0 | 1 | 2 |
| $c_3$ | 2 | 0 | 1 | 0 | 2 | 1 | 0 | 1 | 2 | 1 | 2 | 0 | 1 | 2 | 0 | 0 | 1 |
| $c_4$ | 2 | 0 | 1 | 1 | 2 | 0 | 0 | 1 | 2 | 0 | 2 | 0 | 1 | 2 | 1 | 1 | 0 |

TABLE S42: All two-valued states of $H_{21,13}$. The last column gives an odd cycle in the complement of each nonextendable support.

| $j$ | support of $s_j$ | extendable | odd cycle |
|---|---|---|---|
| 1 | $\{1,4,8,14,15,21,22,23\}$ | no | $(7,11,3,2,5,6,13)$ |
| 2 | $\{1,5,7,14,18,22,23,24\}$ | no | $(4,11,3,2,8,9,10)$ |
| 3 | $\{1,5,9,11,18,21,22\}$ | yes | – |
| 4 | $\{1,5,9,14,15,18,21,22,24\}$ | no | $(11,3,2,8,7)$ |
| 5 | $\{1,6,8,11,18,21,23\}$ | yes | – |
| 6 | $\{1,6,8,14,15,18,21,23,24\}$ | no | $(11,3,2,5,4)$ |
| 7 | $\{2,4,7,14,19,20,22,23\}$ | no | $(8,9,10,1,13,6,5)$ |
| 8 | $\{2,4,9,13,14,15,19\}$ | yes | – |
| 9 | $\{2,4,9,14,15,19,20,21,22\}$ | no | $(11,3,1,13,7)$ |
| 10 | $\{2,6,7,10,14,20,24\}$ | yes | – |
| 11 | $\{2,6,7,14,18,19,20,23,24\}$ | no | $(11,3,1,10,4)$ |
| 12 | $\{2,6,9,11,18,19,20,21\}$ | no | $(5,4,10,1,13,7,8)$ |
| 13 | $\{2,6,9,14,15,18,19,20,21,24\}$ | no | $(11,3,1,10,4)$ |
| 14 | $\{3,4,8,13,15,19,23\}$ | yes | – |
| 15 | $\{3,4,8,15,19,20,21,22,23\}$ | no | $(5,2,1,13,6)$ |
| 16 | $\{3,5,7,10,20,22,24\}$ | yes | – |
| 17 | $\{3,5,7,18,19,20,22,23,24\}$ | no | $(8,2,1,10,9)$ |
| 18 | $\{3,5,9,13,15,18,19,24\}$ | no | $(7,8,2,1,10,4,11)$ |
| 19 | $\{3,5,9,15,18,19,20,21,22,24\}$ | no | $(8,2,1,13,7)$ |
| 20 | $\{3,6,8,10,15,20,21,24\}$ | no | $(4,5,2,1,13,7,11)$ |
| 21 | $\{3,6,8,15,18,19,20,21,23,24\}$ | no | $(5,2,1,10,4)$ |

TABLE S43. All strong three-colorings of $H_{21,13}$ up to color permutation. The anchor is $(2,8,5) = (0,1,2)$.

| $i$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 13 | 14 | 15 | 18 | 19 | 20 | 21 | 22 | 23 | 24 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $c_1$ | 1 | 0 | 2 | 0 | 2 | 1 | 2 | 1 | 0 | 2 | 1 | 0 | 0 | 0 | 1 | 0 | 2 | 1 | 2 | 1 | 2 |
| $c_2$ | 2 | 0 | 1 | 1 | 2 | 0 | 0 | 1 | 2 | 0 | 2 | 1 | 0 | 1 | 2 | 1 | 0 | 2 | 2 | 1 | 0 |

TABLE S44. The nine chromatic signatures of $H_{21,13}$. Columns give $c_1$ and rows give $c_2$ for the two normalized colorings in Table S43. Each cell is one inseparability class: its vertices have the same color in every admissible coloring. Ordinary two-valued states distinguish every pair, including vertices in the same cell.

| $c_2 \backslash c_1$ | 0 | 1 | 2 |
|---|---|---|---|
| 2 | $\{9\}$ | $\{1,11,18,21\}$ | $\{5,22\}$ |
| 1 | $\{4,13,15,19\}$ | $\{8,23\}$ | $\{3\}$ |
| 0 | $\{2,14\}$ | $\{6\}$ | $\{7,10,20,24\}$ |

TABLE S46. Valid states and exact weights reproducing two $H_{17,10}$ preparations: the five-state mixture for $P_{24}$ (upper block, Eq. (27)) and the four-state mixture for $P_{z_3}$, the preparation along $z_3 = r_5$ (lower block). State numbering is that of Table S40. Every context equation and every Born marginal are checked with rational arithmetic.

| State | Weight | Support | Extendable? |
|---|---|---|---|
| Preparation $P_{24}$, nonextendable weight $\frac{1}{9}$ | | | |
| $s_4$ | $\frac{1}{12}$ | $\{1, 5, 9, 14, 15, 18, 24\}$ | no |
| $s_9$ | $\frac{5}{36}$ | $\{2, 6, 7, 10, 14, 24\}$ | yes |
| $s_{10}$ | $\frac{1}{36}$ | $\{2, 6, 7, 14, 18, 19, 23, 24\}$ | no |
| $s_{14}$ | $\frac{7}{12}$ | $\{3, 5, 7, 10, 24\}$ | yes |
| $s_{17}$ | $\frac{1}{6}$ | $\{3, 6, 8, 10, 15, 24\}$ | yes |
| Preparation $P_{z_3}$, nonextendable weight $\frac{1}{3}$ | | | |
| $s_3$ | $\frac{1}{3}$ | $\{1, 5, 9, 11, 18\}$ | yes |
| $s_4$ | $\frac{1}{6}$ | $\{1, 5, 9, 14, 15, 18, 24\}$ | no |
| $s_{14}$ | $\frac{1}{3}$ | $\{3, 5, 7, 10, 24\}$ | yes |
| $s_{15}$ | $\frac{1}{6}$ | $\{3, 5, 7, 18, 19, 23, 24\}$ | no |

TABLE S45: Complete state profiles in hexadecimal, and all Born probabilities for the 17-ray preparation in Eq. (26). The probabilities are also reproduced by Eq. (27).

| $i$ | $H_{17,10}$ profile | $H_{21,13}$ profile | $p_i$ on $H_{17,10}$ |
|---|---|---|---|
| 1 | 3F | 3F | 1/12 |
| 2 | FC0 | 1FC0 | 1/6 |
| 3 | 3F000 | 1FE000 | 3/4 |
| 4 | 10C1 | 61C1 | 0 |
| 5 | E00E | 7800E | 2/3 |
| 6 | 30F30 | 181E30 | 1/3 |
| 7 | 6342 | 18642 | 3/4 |
| 8 | 31031 | 186031 | 1/6 |
| 9 | 8C8C | 6198C | 1/12 |
| 10 | 12100 | 88200 | 8/9 |
| 11 | 414 | 814 | 0 |
| 13 | – | 22080 | – |
| 14 | BEB | 17EB | 1/4 |
| 15 | 398A9 | 1E71A9 | 1/4 |
| 18 | 2CE3E | 171C3E | 1/9 |
| 19 | 2DEC0 | 177DC0 | 1/36 |
| 20 | – | 1DDF40 | – |
| 21 | – | 1C593D | – |
| 22 | – | 5C14F | – |
| 23 | 25273 | 116473 | 1/36 |
| 24 | 3EB2A | 1F962A | 1 |

TABLE S47. Exact Born probabilities $p_i = (r_{24} \cdot r_i)^2/(6\|r_i\|^2)$ for $P_{24}$ and $p'_i = (r_5 \cdot r_i)^2/\|r_i\|^2$ for $P_{z_3}$, also reproduced by the two mixtures of Table S46.

| $i$ | $p_i$ | $p'_i$ | $i$ | $p_i$ | $p'_i$ |
|---|---|---|---|---|---|
| 1 | $\frac{1}{12}$ | $\frac{1}{2}$ | 10 | $\frac{8}{9}$ | $\frac{1}{3}$ |
| 2 | $\frac{1}{6}$ | 0 | 11 | 0 | $\frac{1}{3}$ |
| 3 | $\frac{3}{4}$ | $\frac{1}{2}$ | 14 | $\frac{1}{4}$ | $\frac{1}{6}$ |
| 4 | 0 | 0 | 15 | $\frac{1}{4}$ | $\frac{1}{6}$ |
| 5 | $\frac{2}{3}$ | 1 | 18 | $\frac{1}{9}$ | $\frac{2}{3}$ |
| 6 | $\frac{1}{3}$ | 0 | 19 | $\frac{1}{36}$ | $\frac{1}{6}$ |
| 7 | $\frac{3}{4}$ | $\frac{1}{2}$ | 23 | $\frac{1}{36}$ | $\frac{1}{6}$ |
| 8 | $\frac{1}{6}$ | 0 | 24 | 1 | $\frac{2}{3}$ |
| 9 | $\frac{1}{12}$ | $\frac{1}{2}$ | | | |

### S2.2. State conventions for the additional gadgets

The state tables below are local to each labeled configuration. Supports are in lexicographic order; E is yes precisely for color-class supports. The three-dimensional pentagon and $H_{21,13}$ use the complete earlier tables and are not repeated. The four-dimensional apex-only states are included explicitly. All ordinary state families listed here are unital and separating.

### S2.3. All states of $B_{13,7}$

TABLE S48: All two-valued states of $B_{13,7}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{1,5,7,14,23\}$ | yes | 8 | $\{2,9,11,19\}$ | yes |
| 2 | $\{1,5,9,11\}$ | yes | 9 | $\{2,9,14,15,19\}$ | yes |
| 3 | $\{1,5,9,14,15\}$ | no | 10 | $\{3,5,7,10\}$ | yes |
| 4 | $\{1,8,11,23\}$ | yes | 11 | $\{3,5,7,19,23\}$ | no |
| 5 | $\{1,8,14,15,23\}$ | yes | 12 | $\{3,5,9,15,19\}$ | yes |
| 6 | $\{2,7,10,14\}$ | yes | 13 | $\{3,8,10,15\}$ | yes |
| 7 | $\{2,7,14,19,23\}$ | yes | 14 | $\{3,8,15,19,23\}$ | yes |

### S2.4. All states of $D_{16,9}$

TABLE S49: All two-valued states of $D_{16,9}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{1,4,8,14,15,23\}$ | yes | 12 | $\{2,6,7,10,14,24\}$ | yes |
| 2 | $\{1,5,7,10,14,24\}$ | yes | 13 | $\{2,6,7,14,18,23,24\}$ | no |
| 3 | $\{1,5,7,14,18,23,24\}$ | no | 14 | $\{2,6,9,11,18\}$ | yes |
| 4 | $\{1,5,9,11,18\}$ | yes | 15 | $\{2,6,9,14,15,18,24\}$ | no |
| 5 | $\{1,5,9,14,15,18,24\}$ | no | 16 | $\{3,4,8,15,23\}$ | yes |
| 6 | $\{1,6,8,10,11\}$ | yes | 17 | $\{3,5,7,10,24\}$ | yes |
| 7 | $\{1,6,8,10,14,15,24\}$ | no | 18 | $\{3,5,7,18,23,24\}$ | yes |
| 8 | $\{1,6,8,11,18,23\}$ | yes | 19 | $\{3,5,9,15,18,24\}$ | yes |
| 9 | $\{1,6,8,14,15,18,23,24\}$ | no | 20 | $\{3,6,8,10,15,24\}$ | yes |
| 10 | $\{2,4,7,14,23\}$ | yes | 21 | $\{3,6,8,15,18,23,24\}$ | no |
| 11 | $\{2,4,9,14,15\}$ | yes | | | |

### S2.5. All states of $I_{16,9}$

TABLE S50: All two-valued states of $I_{16,9}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{1,4,5,7,14,23\}$ | yes | 12 | $\{2,7,14,18,19,23,24\}$ | no |
| 2 | $\{1,4,5,9,14,15\}$ | no | 13 | $\{2,9,11,18,19\}$ | yes |
| 3 | $\{1,4,8,14,15,23\}$ | yes | 14 | $\{2,9,14,15,18,19,24\}$ | no |
| 4 | $\{1,5,7,14,18,23,24\}$ | no | 15 | $\{3,4,5,7,19,23\}$ | no |
| 5 | $\{1,5,9,11,18\}$ | yes | 16 | $\{3,4,5,9,15,19\}$ | yes |
| 6 | $\{1,5,9,14,15,18,24\}$ | no | 17 | $\{3,4,8,15,19,23\}$ | yes |
| 7 | $\{1,8,11,18,23\}$ | yes | 18 | $\{3,5,7,10,24\}$ | yes |
| 8 | $\{1,8,14,15,18,23,24\}$ | no | 19 | $\{3,5,7,18,19,23,24\}$ | no |
| 9 | $\{2,4,7,14,19,23\}$ | yes | 20 | $\{3,5,9,15,18,19,24\}$ | no |
| 10 | $\{2,4,9,14,15,19\}$ | yes | 21 | $\{3,8,10,15,24\}$ | yes |
| 11 | $\{2,7,10,14,24\}$ | yes | 22 | $\{3,8,15,18,19,23,24\}$ | no |

### S2.6. All states of $T_{19,11}$

TABLE S51: All two-valued states of $T_{19,11}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{1,4,8,14,15,23,28\}$ | yes | 14 | $\{2,6,7,14,18,23,24\}$ | no |
| 2 | $\{1,5,7,10,14,24,28\}$ | yes | 15 | $\{2,6,9,11,18,27\}$ | yes |
| 3 | $\{1,5,7,14,18,23,24\}$ | no | 16 | $\{2,6,9,14,15,18,24\}$ | no |
| 4 | $\{1,5,9,11,18,27\}$ | yes | 17 | $\{3,4,8,15,23,26\}$ | yes |
| 5 | $\{1,5,9,14,15,18,24\}$ | no | 18 | $\{3,4,8,15,23,27,28\}$ | no |
| 6 | $\{1,6,8,10,11,26\}$ | yes | 19 | $\{3,5,7,10,24,26\}$ | yes |
| 7 | $\{1,6,8,10,11,27,28\}$ | no | 20 | $\{3,5,7,10,24,27,28\}$ | no |
| 8 | $\{1,6,8,10,14,15,24,28\}$ | no | 21 | $\{3,5,7,18,23,24,27\}$ | yes |
| 9 | $\{1,6,8,11,18,23,27\}$ | yes | 22 | $\{3,5,9,15,18,24,27\}$ | yes |
| 10 | $\{1,6,8,14,15,18,23,24\}$ | no | 23 | $\{3,6,8,10,15,24,26\}$ | yes |
| 11 | $\{2,4,7,14,23,28\}$ | yes | 24 | $\{3,6,8,10,15,24,27,28\}$ | no |
| 12 | $\{2,4,9,14,15,28\}$ | yes | 25 | $\{3,6,8,15,18,23,24,27\}$ | no |
| 13 | $\{2,6,7,10,14,24,28\}$ | yes | | | |

### S2.7. All states of $W^{(4)}_{11,5}$

TABLE S52: All two-valued states of $W^{(4)}_{11,5}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{0\}$ | yes | 7 | $\{2,8,9,10\}$ | yes |
| 2 | $\{1,3,9\}$ | yes | 8 | $\{3,5,6\}$ | yes |
| 3 | $\{1,4,7\}$ | yes | 9 | $\{3,6,9,10\}$ | yes |
| 4 | $\{1,7,8,9\}$ | yes | 10 | $\{4,6,7,10\}$ | yes |
| 5 | $\{2,4,10\}$ | yes | 11 | $\{5,6,7,8\}$ | yes |
| 6 | $\{2,5,8\}$ | yes | 12 | $\{6,7,8,9,10\}$ | no |

### S2.8. All states of $W^{(4)}_{16,6}$

TABLE S53: All two-valued states of $W^{(4)}_{16,6}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{0,7,9,10\}$ | yes | 37 | $\{2,8,12,14,15\}$ | yes |
| 2 | $\{0,7,9,15\}$ | yes | 38 | $\{3,5,6\}$ | yes |
| 3 | $\{0,7,10,14\}$ | yes | 39 | $\{3,6,9,10,11\}$ | yes |
| 4 | $\{0,7,14,15\}$ | yes | 40 | $\{3,6,9,10,12\}$ | yes |
| 5 | $\{0,9,10,13\}$ | yes | 41 | $\{3,6,9,11,15\}$ | yes |
| 6 | $\{0,9,13,15\}$ | yes | 42 | $\{3,6,9,12,15\}$ | yes |
| 7 | $\{0,10,13,14\}$ | yes | 43 | $\{3,6,10,11,14\}$ | yes |
| 8 | $\{0,13,14,15\}$ | yes | 44 | $\{3,6,10,12,14\}$ | yes |
| 9 | $\{1,3,9,11\}$ | yes | 45 | $\{3,6,11,14,15\}$ | yes |
| 10 | $\{1,3,9,12\}$ | yes | 46 | $\{3,6,12,14,15\}$ | yes |
| 11 | $\{1,3,11,14\}$ | yes | 47 | $\{4,6,7,10,11\}$ | yes |
| 12 | $\{1,3,12,14\}$ | yes | 48 | $\{4,6,7,10,12\}$ | yes |
| 13 | $\{1,4,7,11\}$ | yes | 49 | $\{4,6,7,11,15\}$ | yes |
| 14 | $\{1,4,7,12\}$ | yes | 50 | $\{4,6,7,12,15\}$ | yes |
| 15 | $\{1,4,11,13\}$ | yes | 51 | $\{4,6,10,11,13\}$ | yes |
| 16 | $\{1,4,12,13\}$ | yes | 52 | $\{4,6,10,12,13\}$ | yes |
| 17 | $\{1,7,8,9,11\}$ | yes | 53 | $\{4,6,11,13,15\}$ | yes |
| 18 | $\{1,7,8,9,12\}$ | yes | 54 | $\{4,6,12,13,15\}$ | yes |
| 19 | $\{1,7,8,11,14\}$ | yes | 55 | $\{5,6,7,8\}$ | yes |
| 20 | $\{1,7,8,12,14\}$ | yes | 56 | $\{5,6,8,13\}$ | yes |
| 21 | $\{1,8,9,11,13\}$ | yes | 57 | $\{6,7,8,9,10,11\}$ | no |
| 22 | $\{1,8,9,12,13\}$ | yes | 58 | $\{6,7,8,9,10,12\}$ | no |
| 23 | $\{1,8,11,13,14\}$ | yes | 59 | $\{6,7,8,9,11,15\}$ | no |
| 24 | $\{1,8,12,13,14\}$ | yes | 60 | $\{6,7,8,9,12,15\}$ | no |
| 25 | $\{2,4,10,11\}$ | yes | 61 | $\{6,7,8,10,11,14\}$ | no |
| 26 | $\{2,4,10,12\}$ | yes | 62 | $\{6,7,8,10,12,14\}$ | no |
| 27 | $\{2,4,11,15\}$ | yes | 63 | $\{6,7,8,11,14,15\}$ | no |
| 28 | $\{2,4,12,15\}$ | yes | 64 | $\{6,7,8,12,14,15\}$ | no |
| 29 | $\{2,5,8\}$ | yes | 65 | $\{6,8,9,10,11,13\}$ | no |
| 30 | $\{2,8,9,10,11\}$ | yes | 66 | $\{6,8,9,10,12,13\}$ | no |
| 31 | $\{2,8,9,10,12\}$ | yes | 67 | $\{6,8,9,11,13,15\}$ | no |
| 32 | $\{2,8,9,11,15\}$ | yes | 68 | $\{6,8,9,12,13,15\}$ | no |
| 33 | $\{2,8,9,12,15\}$ | yes | 69 | $\{6,8,10,11,13,14\}$ | no |
| 34 | $\{2,8,10,11,14\}$ | yes | 70 | $\{6,8,10,12,13,14\}$ | no |
| 35 | $\{2,8,10,12,14\}$ | yes | 71 | $\{6,8,11,13,14,15\}$ | no |
| 36 | $\{2,8,11,14,15\}$ | yes | 72 | $\{6,8,12,13,14,15\}$ | no |

### S2.9. All states of $B^{(4)}_{14,7}$

TABLE S54: All two-valued states of $B^{(4)}_{14,7}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{0\}$ | yes | 9 | $\{2,9,11,19\}$ | yes |
| 2 | $\{1,5,7,14,23\}$ | yes | 10 | $\{2,9,14,15,19\}$ | yes |
| 3 | $\{1,5,9,11\}$ | yes | 11 | $\{3,5,7,10\}$ | yes |
| 4 | $\{1,5,9,14,15\}$ | no | 12 | $\{3,5,7,19,23\}$ | no |
| 5 | $\{1,8,11,23\}$ | yes | 13 | $\{3,5,9,15,19\}$ | yes |
| 6 | $\{1,8,14,15,23\}$ | yes | 14 | $\{3,8,10,15\}$ | yes |
| 7 | $\{2,7,10,14\}$ | yes | 15 | $\{3,8,15,19,23\}$ | yes |
| 8 | $\{2,7,14,19,23\}$ | yes | | | |

### S2.10. All states of $L^{(4)}_{18,7}$

TABLE S55: All two-valued states of $L^{(4)}_{18,7}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{1,6,9,13,16\}$ | yes | 35 | $\{3,5,9,13,18\}$ | yes |
| 2 | $\{1,6,9,14,16\}$ | yes | 36 | $\{3,5,9,14,17\}$ | yes |
| 3 | $\{1,6,10,13,16\}$ | yes | 37 | $\{3,5,9,14,18\}$ | yes |
| 4 | $\{1,6,10,14,16\}$ | yes | 38 | $\{3,5,10,13,17\}$ | yes |
| 5 | $\{1,7,9,13,17\}$ | yes | 39 | $\{3,5,10,13,18\}$ | yes |
| 6 | $\{1,7,9,13,18\}$ | yes | 40 | $\{3,5,10,14,17\}$ | yes |
| 7 | $\{1,7,9,14,17\}$ | yes | 41 | $\{3,5,10,14,18\}$ | yes |
| 8 | $\{1,7,9,14,18\}$ | yes | 42 | $\{3,6,9,11\}$ | yes |
| 9 | $\{1,7,10,13,17\}$ | yes | 43 | $\{3,6,9,12,13\}$ | yes |
| 10 | $\{1,7,10,13,18\}$ | yes | 44 | $\{3,6,9,12,14\}$ | yes |
| 11 | $\{1,7,10,14,17\}$ | yes | 45 | $\{3,6,10,11\}$ | yes |
| 12 | $\{1,7,10,14,18\}$ | yes | 46 | $\{3,6,10,12,13\}$ | yes |
| 13 | $\{1,8,13,15\}$ | yes | 47 | $\{3,6,10,12,14\}$ | yes |
| 14 | $\{1,8,13,16,17\}$ | yes | 48 | $\{3,8,11,17\}$ | yes |
| 15 | $\{1,8,13,16,18\}$ | yes | 49 | $\{3,8,11,18\}$ | yes |
| 16 | $\{1,8,14,15\}$ | yes | 50 | $\{3,8,12,13,17\}$ | yes |
| 17 | $\{1,8,14,16,17\}$ | yes | 51 | $\{3,8,12,13,18\}$ | yes |
| 18 | $\{1,8,14,16,18\}$ | yes | 52 | $\{3,8,12,14,17\}$ | yes |
| 19 | $\{2,5,9,15\}$ | yes | 53 | $\{3,8,12,14,18\}$ | yes |
| 20 | $\{2,5,9,16,17\}$ | yes | 54 | $\{4,5,13,15\}$ | yes |
| 21 | $\{2,5,9,16,18\}$ | yes | 55 | $\{4,5,13,16,17\}$ | yes |
| 22 | $\{2,5,10,15\}$ | yes | 56 | $\{4,5,13,16,18\}$ | yes |
| 23 | $\{2,5,10,16,17\}$ | yes | 57 | $\{4,5,14,15\}$ | yes |
| 24 | $\{2,5,10,16,18\}$ | yes | 58 | $\{4,5,14,16,17\}$ | yes |
| 25 | $\{2,6,9,12,16\}$ | yes | 59 | $\{4,5,14,16,18\}$ | yes |
| 26 | $\{2,6,10,12,16\}$ | yes | 60 | $\{4,6,11,16\}$ | yes |
| 27 | $\{2,7,9,12,17\}$ | yes | 61 | $\{4,6,12,13,16\}$ | yes |
| 28 | $\{2,7,9,12,18\}$ | yes | 62 | $\{4,6,12,14,16\}$ | yes |
| 29 | $\{2,7,10,12,17\}$ | yes | 63 | $\{4,7,11,17\}$ | yes |
| 30 | $\{2,7,10,12,18\}$ | yes | 64 | $\{4,7,11,18\}$ | yes |
| 31 | $\{2,8,12,15\}$ | yes | 65 | $\{4,7,12,13,17\}$ | yes |
| 32 | $\{2,8,12,16,17\}$ | yes | 66 | $\{4,7,12,13,18\}$ | yes |
| 33 | $\{2,8,12,16,18\}$ | yes | 67 | $\{4,7,12,14,17\}$ | yes |
| 34 | $\{3,5,9,13,17\}$ | yes | 68 | $\{4,7,12,14,18\}$ | yes |

### S2.11. All states of $D^{(4)}_{17,9}$

TABLE S56: All two-valued states of $D^{(4)}_{17,9}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{0\}$ | yes | 12 | $\{2,4,9,14,15\}$ | yes |
| 2 | $\{1,4,8,14,15,23\}$ | yes | 13 | $\{2,6,7,10,14,24\}$ | yes |
| 3 | $\{1,5,7,10,14,24\}$ | yes | 14 | $\{2,6,7,14,18,23,24\}$ | no |
| 4 | $\{1,5,7,14,18,23,24\}$ | no | 15 | $\{2,6,9,11,18\}$ | yes |
| 5 | $\{1,5,9,11,18\}$ | yes | 16 | $\{2,6,9,14,15,18,24\}$ | no |
| 6 | $\{1,5,9,14,15,18,24\}$ | no | 17 | $\{3,4,8,15,23\}$ | yes |
| 7 | $\{1,6,8,10,11\}$ | yes | 18 | $\{3,5,7,10,24\}$ | yes |
| 8 | $\{1,6,8,10,14,15,24\}$ | no | 19 | $\{3,5,7,18,23,24\}$ | yes |
| 9 | $\{1,6,8,11,18,23\}$ | yes | 20 | $\{3,5,9,15,18,24\}$ | yes |
| 10 | $\{1,6,8,14,15,18,23,24\}$ | no | 21 | $\{3,6,8,10,15,24\}$ | yes |
| 11 | $\{2,4,7,14,23\}$ | yes | 22 | $\{3,6,8,15,18,23,24\}$ | no |

### S2.12. All states of $L^{(4)}_{22,9}$

TABLE S57: All two-valued states of $L^{(4)}_{22,9}$. E indicates extendability.

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 1 | $\{1,6,9,13,16,21\}$ | yes | 80 | $\{2,8,12,16,17,20,21\}$ | no |
| 2 | $\{1,6,9,13,16,22\}$ | yes | 81 | $\{2,8,12,16,17,20,22\}$ | no |
| 3 | $\{1,6,9,14,16,19\}$ | yes | 82 | $\{2,8,12,16,18,19\}$ | yes |
| 4 | $\{1,6,9,14,16,20,21\}$ | no | 83 | $\{2,8,12,16,18,20,21\}$ | no |
| 5 | $\{1,6,9,14,16,20,22\}$ | no | 84 | $\{2,8,12,16,18,20,22\}$ | no |
| 6 | $\{1,6,10,13,16,21\}$ | yes | 85 | $\{3,5,9,13,17,21\}$ | yes |
| 7 | $\{1,6,10,13,16,22\}$ | yes | 86 | $\{3,5,9,13,17,22\}$ | yes |
| 8 | $\{1,6,10,14,16,19\}$ | yes | 87 | $\{3,5,9,13,18,21\}$ | yes |
| 9 | $\{1,6,10,14,16,20,21\}$ | no | 88 | $\{3,5,9,13,18,22\}$ | yes |
| 10 | $\{1,6,10,14,16,20,22\}$ | no | 89 | $\{3,5,9,14,17,19\}$ | yes |
| 11 | $\{1,7,9,13,17,21\}$ | yes | 90 | $\{3,5,9,14,17,20,21\}$ | no |
| 12 | $\{1,7,9,13,17,22\}$ | yes | 91 | $\{3,5,9,14,17,20,22\}$ | no |
| 13 | $\{1,7,9,13,18,21\}$ | yes | 92 | $\{3,5,9,14,18,19\}$ | yes |
| 14 | $\{1,7,9,13,18,22\}$ | yes | 93 | $\{3,5,9,14,18,20,21\}$ | no |
| 15 | $\{1,7,9,14,17,19\}$ | yes | 94 | $\{3,5,9,14,18,20,22\}$ | no |
| 16 | $\{1,7,9,14,17,20,21\}$ | no | 95 | $\{3,5,10,13,17,21\}$ | yes |
| 17 | $\{1,7,9,14,17,20,22\}$ | no | 96 | $\{3,5,10,13,17,22\}$ | yes |
| 18 | $\{1,7,9,14,18,19\}$ | yes | 97 | $\{3,5,10,13,18,21\}$ | yes |
| 19 | $\{1,7,9,14,18,20,21\}$ | no | 98 | $\{3,5,10,13,18,22\}$ | yes |
| 20 | $\{1,7,9,14,18,20,22\}$ | no | 99 | $\{3,5,10,14,17,19\}$ | yes |
| 21 | $\{1,7,10,13,17,21\}$ | yes | 100 | $\{3,5,10,14,17,20,21\}$ | no |
| 22 | $\{1,7,10,13,17,22\}$ | yes | 101 | $\{3,5,10,14,17,20,22\}$ | no |
| 23 | $\{1,7,10,13,18,21\}$ | yes | 102 | $\{3,5,10,14,18,19\}$ | yes |
| 24 | $\{1,7,10,13,18,22\}$ | yes | 103 | $\{3,5,10,14,18,20,21\}$ | no |
| 25 | $\{1,7,10,14,17,19\}$ | yes | 104 | $\{3,5,10,14,18,20,22\}$ | no |
| 26 | $\{1,7,10,14,17,20,21\}$ | no | 105 | $\{3,6,9,11,20\}$ | yes |
| 27 | $\{1,7,10,14,17,20,22\}$ | no | 106 | $\{3,6,9,12,13,21\}$ | yes |
| 28 | $\{1,7,10,14,18,19\}$ | yes | 107 | $\{3,6,9,12,13,22\}$ | yes |
| 29 | $\{1,7,10,14,18,20,21\}$ | no | 108 | $\{3,6,9,12,14,19\}$ | yes |
| 30 | $\{1,7,10,14,18,20,22\}$ | no | 109 | $\{3,6,9,12,14,20,21\}$ | no |
| 31 | $\{1,8,13,16,17,21\}$ | yes | 110 | $\{3,6,9,12,14,20,22\}$ | no |
| 32 | $\{1,8,13,16,17,22\}$ | yes | 111 | $\{3,6,10,11,20\}$ | yes |
| 33 | $\{1,8,13,16,18,21\}$ | yes | 112 | $\{3,6,10,12,13,21\}$ | yes |
| 34 | $\{1,8,13,16,18,22\}$ | yes | 113 | $\{3,6,10,12,13,22\}$ | yes |
| 35 | $\{1,8,14,15,21\}$ | yes | 114 | $\{3,6,10,12,14,19\}$ | yes |
| 36 | $\{1,8,14,15,22\}$ | yes | 115 | $\{3,6,10,12,14,20,21\}$ | no |
| 37 | $\{1,8,14,16,17,19\}$ | yes | 116 | $\{3,6,10,12,14,20,22\}$ | no |
| 38 | $\{1,8,14,16,17,20,21\}$ | no | 117 | $\{3,8,11,17,20\}$ | yes |
| 39 | $\{1,8,14,16,17,20,22\}$ | no | 118 | $\{3,8,11,18,20\}$ | yes |
| 40 | $\{1,8,14,16,18,19\}$ | yes | 119 | $\{3,8,12,13,17,21\}$ | yes |
| 41 | $\{1,8,14,16,18,20,21\}$ | no | 120 | $\{3,8,12,13,17,22\}$ | yes |
| 42 | $\{1,8,14,16,18,20,22\}$ | no | 121 | $\{3,8,12,13,18,21\}$ | yes |
| 43 | $\{2,5,9,15,21\}$ | yes | 122 | $\{3,8,12,13,18,22\}$ | yes |
| 44 | $\{2,5,9,15,22\}$ | yes | 123 | $\{3,8,12,14,17,19\}$ | yes |

| $j$ | Support | E | $j$ | Support | E |
|---|---|---|---|---|---|
| 45 | $\{2,5,9,16,17,19\}$ | yes | 124 | $\{3,8,12,14,17,20,21\}$ | no |
| 46 | $\{2,5,9,16,17,20,21\}$ | no | 125 | $\{3,8,12,14,17,20,22\}$ | no |
| 47 | $\{2,5,9,16,17,20,22\}$ | no | 126 | $\{3,8,12,14,18,19\}$ | yes |
| 48 | $\{2,5,9,16,18,19\}$ | yes | 127 | $\{3,8,12,14,18,20,21\}$ | no |
| 49 | $\{2,5,9,16,18,20,21\}$ | no | 128 | $\{3,8,12,14,18,20,22\}$ | no |
| 50 | $\{2,5,9,16,18,20,22\}$ | no | 129 | $\{4,5,13,16,17,21\}$ | yes |
| 51 | $\{2,5,10,15,21\}$ | yes | 130 | $\{4,5,13,16,17,22\}$ | yes |
| 52 | $\{2,5,10,15,22\}$ | yes | 131 | $\{4,5,13,16,18,21\}$ | yes |
| 53 | $\{2,5,10,16,17,19\}$ | yes | 132 | $\{4,5,13,16,18,22\}$ | yes |
| 54 | $\{2,5,10,16,17,20,21\}$ | no | 133 | $\{4,5,14,15,21\}$ | yes |
| 55 | $\{2,5,10,16,17,20,22\}$ | no | 134 | $\{4,5,14,15,22\}$ | yes |
| 56 | $\{2,5,10,16,18,19\}$ | yes | 135 | $\{4,5,14,16,17,19\}$ | yes |
| 57 | $\{2,5,10,16,18,20,21\}$ | no | 136 | $\{4,5,14,16,17,20,21\}$ | no |
| 58 | $\{2,5,10,16,18,20,22\}$ | no | 137 | $\{4,5,14,16,17,20,22\}$ | no |
| 59 | $\{2,6,9,12,16,19\}$ | yes | 138 | $\{4,5,14,16,18,19\}$ | yes |
| 60 | $\{2,6,9,12,16,20,21\}$ | no | 139 | $\{4,5,14,16,18,20,21\}$ | no |
| 61 | $\{2,6,9,12,16,20,22\}$ | no | 140 | $\{4,5,14,16,18,20,22\}$ | no |
| 62 | $\{2,6,10,12,16,19\}$ | yes | 141 | $\{4,6,11,16,20\}$ | yes |
| 63 | $\{2,6,10,12,16,20,21\}$ | no | 142 | $\{4,6,12,13,16,21\}$ | yes |
| 64 | $\{2,6,10,12,16,20,22\}$ | no | 143 | $\{4,6,12,13,16,22\}$ | yes |
| 65 | $\{2,7,9,12,17,19\}$ | yes | 144 | $\{4,6,12,14,16,19\}$ | yes |
| 66 | $\{2,7,9,12,17,20,21\}$ | no | 145 | $\{4,6,12,14,16,20,21\}$ | no |
| 67 | $\{2,7,9,12,17,20,22\}$ | no | 146 | $\{4,6,12,14,16,20,22\}$ | no |
| 68 | $\{2,7,9,12,18,19\}$ | yes | 147 | $\{4,7,11,17,20\}$ | yes |
| 69 | $\{2,7,9,12,18,20,21\}$ | no | 148 | $\{4,7,11,18,20\}$ | yes |
| 70 | $\{2,7,9,12,18,20,22\}$ | no | 149 | $\{4,7,12,13,17,21\}$ | yes |
| 71 | $\{2,7,10,12,17,19\}$ | yes | 150 | $\{4,7,12,13,17,22\}$ | yes |
| 72 | $\{2,7,10,12,17,20,21\}$ | no | 151 | $\{4,7,12,13,18,21\}$ | yes |
| 73 | $\{2,7,10,12,17,20,22\}$ | no | 152 | $\{4,7,12,13,18,22\}$ | yes |
| 74 | $\{2,7,10,12,18,19\}$ | yes | 153 | $\{4,7,12,14,17,19\}$ | yes |
| 75 | $\{2,7,10,12,18,20,21\}$ | no | 154 | $\{4,7,12,14,17,20,21\}$ | no |
| 76 | $\{2,7,10,12,18,20,22\}$ | no | 155 | $\{4,7,12,14,17,20,22\}$ | no |
| 77 | $\{2,8,12,15,21\}$ | yes | 156 | $\{4,7,12,14,18,19\}$ | yes |
| 78 | $\{2,8,12,15,22\}$ | yes | 157 | $\{4,7,12,14,18,20,21\}$ | no |
| 79 | $\{2,8,12,16,17,19\}$ | yes | 158 | $\{4,7,12,14,18,20,22\}$ | no |

### S2.13. Complete normalized colorings of the gadget panels

Each color word gives one representative under global color permutation; the vertices and the anchor convention are specified separately for each table. The full labeled family is recovered by all palette permutations. Equal color labels across different colorings carry no extra constraint.

### S2.14. Colorings of $P_5$

The columns of each word correspond, in order, to vertices $1,2,3,4,5,6,7,8,9,10$. The anchor context $(1,6,2)$ has colors $(0,\dots,2)$.

TABLE S58: Every normalized coloring of $P_5$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 0201211201 | 4 | 0210210211 |
| 2 | 0202111102 | 5 | 0212110002 |
| 3 | 0210110222 | | |

### S2.15. Colorings of $B_{13,7}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 5, 7, 8, 9, 10, 11, 14, 15, 19, 23$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S59: Every normalized coloring of $B_{13,7}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 1022012010221 | 4 | 2012012020111 |
| 2 | 1022210201101 | 5 | 2012012120100 |
| 3 | 1022210210001 | 6 | 2012210102102 |

### S2.16. Colorings of $D_{16,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S60: Every normalized coloring of $D_{16,9}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 1020210121102202 | 5 | 2010210121201201 |
| 2 | 1020212101100222 | 6 | 2011200120201210 |
| 3 | 1020212102100112 | 7 | 2011202102021012 |
| 4 | 1021202102011012 | | |

### S2.17. Colorings of $I_{16,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 14, 15, 18, 19, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S61: Every normalized coloring of $I_{16,9}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 1020221021001012 | 4 | 2010201212012001 |
| 2 | 1021221020110012 | 5 | 2011201202012110 |
| 3 | 1022201201021210 | 6 | 2012221010210021 |

### S2.18. Colorings of $T_{19,11}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24, 26, 27, 28$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S62: Every normalized coloring of $T_{19,11}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 1020210121102202120 | 5 | 2010210121201201120 |
| 2 | 1020212101100222120 | 6 | 2011200120201210120 |
| 3 | 1020212102100112210 | 7 | 2011202102021012102 |
| 4 | 1021202102011012201 | | |

### S2.19. Colorings of $H_{21,13}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 18, 19, 20, 21, 22, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S63: Every normalized coloring of $H_{21,13}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 102021210210001021212 | 2 | 201120012021012102210 |

### S2.20. Colorings of $W^{(4)}_{11,5}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10$. The anchor context $(0, 1, 6, 2)$ has colors $(0, \ldots, 3)$.

TABLE S64: Every normalized coloring of $W^{(4)}_{11,5}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 01312322312 | 4 | 01321321322 |
| 2 | 01313222213 | 5 | 01323221113 |
| 3 | 01321221333 | | |

### S2.21. Colorings of $W^{(4)}_{16,6}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15$. The anchor context $(2, 0, 6, 1)$ has colors $(0, \ldots, 3)$.

TABLE S65: Every normalized coloring of $W^{(4)}_{16,6}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 1302302101123322 | 41 | 1302022331003131 |
| 2 | 1302302101132322 | 42 | 1302022331030131 |
| 3 | 1302302101223321 | 43 | 1302022331103130 |
| 4 | 1302302101232321 | 44 | 1302022331130130 |
| 5 | 1302302102123312 | 45 | 1302022333003111 |
| 6 | 1302302102132312 | 46 | 1302022333030111 |
| 7 | 1302302102223311 | 47 | 1302022333103110 |
| 8 | 1302302102232311 | 48 | 1302022333130110 |
| 9 | 1302302301123122 | 49 | 1302322100003311 |
| 10 | 1302302301132122 | 50 | 1302322100030311 |
| 11 | 1302302301223121 | 51 | 1302322100103310 |
| 12 | 1302302301232121 | 52 | 1302322100130310 |
| 13 | 1302302302123112 | 53 | 1302322101003301 |
| 14 | 1302302302132112 | 54 | 1302322101030301 |
| 15 | 1302302302223111 | 55 | 1302322101103300 |
| 16 | 1302302302232111 | 56 | 1302322101130300 |
| 17 | 1303202101123232 | 57 | 1302322300003111 |
| 18 | 1303202101132232 | 58 | 1302322300030111 |
| 19 | 1303202101223231 | 59 | 1302322300103110 |
| 20 | 1303202101232231 | 60 | 1302322300130110 |
| 21 | 1303202103123212 | 61 | 1302322301003101 |

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 22 | 1303202103132212 | 62 | 1302322301030101 |
| 23 | 1303202103223211 | 63 | 1302322301103100 |
| 24 | 1303202103232211 | 64 | 1302322301130100 |
| 25 | 1303202201123132 | 65 | 1303022121003231 |
| 26 | 1303202201132132 | 66 | 1303022121030231 |
| 27 | 1303202201223131 | 67 | 1303022121103230 |
| 28 | 1303202201232131 | 68 | 1303022121130230 |
| 29 | 1303202203123112 | 69 | 1303022123003211 |
| 30 | 1303202203132112 | 70 | 1303022123030211 |
| 31 | 1303202203223111 | 71 | 1303022123103210 |
| 32 | 1303202203232111 | 72 | 1303022123130210 |
| 33 | 1302022131003331 | 73 | 1303022221003131 |
| 34 | 1302022131030331 | 74 | 1303022221030131 |
| 35 | 1302022131103330 | 75 | 1303022221103130 |
| 36 | 1302022131130330 | 76 | 1303022221130130 |
| 37 | 1302022133003311 | 77 | 1303022223003111 |
| 38 | 1302022133030311 | 78 | 1303022223030111 |
| 39 | 1302022133103310 | 79 | 1303022223103110 |
| 40 | 1302022133130310 | 80 | 1303022223130110 |

### S2.22. Colorings of $B^{(4)}_{14,7}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 5, 7, 8, 9, 10, 11, 14, 15, 19, 23$. The anchor context $(0, 1, 2, 3)$ has colors $(0, \ldots, 3)$.

TABLE S66: Every normalized coloring of $B^{(4)}_{14,7}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 01233213212331 | 4 | 01231132321321 |
| 2 | 01233312312221 | 5 | 01231231212333 |
| 3 | 01233312321121 | 6 | 01231231312322 |

### S2.23. Colorings of $L^{(4)}_{18,7}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18$. The anchor context $(1, 2, 4, 3)$ has colors $(0, \ldots, 3)$.

TABLE S67: Every normalized coloring of $L^{(4)}_{18,7}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 013212030123031203 | 73 | 013210230132021023 |
| 2 | 013212030123031230 | 74 | 013210230132021032 |
| 3 | 013212030123301203 | 75 | 013210230132201023 |
| 4 | 013212030123301230 | 76 | 013210230132201032 |
| 5 | 013212031023031203 | 77 | 013210231032021023 |
| 6 | 013212031023031230 | 78 | 013210231032021032 |
| 7 | 013212031023301203 | 79 | 013210231032201023 |
| 8 | 013212031023301230 | 80 | 013210231032201032 |

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 9 | 013210230123031023 | 81 | 013213201332020112 |
| 10 | 013210230123031032 | 82 | 013213201332020121 |
| 11 | 013210230123301023 | 83 | 013213201332200112 |
| 12 | 013210230123301032 | 84 | 013213201332200121 |
| 13 | 013210231023031023 | 85 | 013213203132020112 |
| 14 | 013210231023031032 | 86 | 013213203132020121 |
| 15 | 013210231023301023 | 87 | 013213203132200112 |
| 16 | 013210231023301032 | 88 | 013213203132200121 |
| 17 | 013213201323030112 | 89 | 013213201332021002 |
| 18 | 013213201323030121 | 90 | 013213201332021020 |
| 19 | 013213201323300112 | 91 | 013213201332201002 |
| 20 | 013213201323300121 | 92 | 013213201332201020 |
| 21 | 013213203123030112 | 93 | 013213203132021002 |
| 22 | 013213203123030121 | 94 | 013213203132021020 |
| 23 | 013213203123300112 | 95 | 013213203132201002 |
| 24 | 013213203123300121 | 96 | 013213203132201020 |
| 25 | 013213201323031002 | 97 | 013221030131022103 |
| 26 | 013213201323031020 | 98 | 013221030131022130 |
| 27 | 013213201323301002 | 99 | 013221030131202103 |
| 28 | 013213201323301020 | 100 | 013221030131202130 |
| 29 | 013213203123031002 | 101 | 013221031031022103 |
| 30 | 013213203123031020 | 102 | 013221031031022130 |
| 31 | 013213203123301002 | 103 | 013221031031202103 |
| 32 | 013213203123301020 | 104 | 013221031031202130 |
| 33 | 013231201321030123 | 105 | 013223010331021202 |
| 34 | 013231201321030132 | 106 | 013223010331021220 |
| 35 | 013231201321300123 | 107 | 013223010331201202 |
| 36 | 013231201321300132 | 108 | 013223010331201220 |
| 37 | 013231203121030123 | 109 | 013223013031021202 |
| 38 | 013231203121030132 | 110 | 013223013031021220 |
| 39 | 013231203121300123 | 111 | 013223013031201202 |
| 40 | 013231203121300132 | 112 | 013223013031201220 |
| 41 | 013232101321030213 | 113 | 013223010331022101 |
| 42 | 013232101321030231 | 114 | 013223010331022110 |
| 43 | 013232101321300213 | 115 | 013223010331202101 |
| 44 | 013232101321300231 | 116 | 013223010331202110 |
| 45 | 013232103121030213 | 117 | 013223013031022101 |
| 46 | 013232103121030231 | 118 | 013223013031022110 |
| 47 | 013232103121300213 | 119 | 013223013031202101 |
| 48 | 013232103121300231 | 120 | 013223013031202110 |
| 49 | 013230210321031023 | 121 | 013220130131022013 |
| 50 | 013230210321031032 | 122 | 013220130131022031 |
| 51 | 013230210321301023 | 123 | 013220130131202013 |
| 52 | 013230210321301032 | 124 | 013220130131202031 |
| 53 | 013230213021031023 | 125 | 013220131031022013 |
| 54 | 013230213021031032 | 126 | 013220131031022031 |
| 55 | 013230213021301023 | 127 | 013220131031202013 |
| 56 | 013230213021301032 | 128 | 013220131031202031 |

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 57 | 013232010321031203 | 129 | 013223101331020212 |
| 58 | 013232010321031230 | 130 | 013223101331020221 |
| 59 | 013232010321301203 | 131 | 013223101331200212 |
| 60 | 013232010321301230 | 132 | 013223101331200221 |
| 61 | 013232013021031203 | 133 | 013223103131020212 |
| 62 | 013232013021031230 | 134 | 013223103131020221 |
| 63 | 013232013021301203 | 135 | 013223103131200212 |
| 64 | 013232013021301230 | 136 | 013223103131200221 |
| 65 | 013212030132021203 | 137 | 013223101331022001 |
| 66 | 013212030132021230 | 138 | 013223101331022010 |
| 67 | 013212030132201203 | 139 | 013223101331202001 |
| 68 | 013212030132201230 | 140 | 013223101331202010 |
| 69 | 013212031032021203 | 141 | 013223103131022001 |
| 70 | 013212031032021230 | 142 | 013223103131022010 |
| 71 | 013212031032201203 | 143 | 013223103131202001 |
| 72 | 013212031032201230 | 144 | 013223103131202010 |

### S2.24. Colorings of $D^{(4)}_{17,9}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24$. The anchor context $(0, 1, 2, 3)$ has colors $(0, \ldots, 3)$.

TABLE S68: Every normalized coloring of $D^{(4)}_{17,9}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 01232132313123123 | 5 | 01233122312123132 |
| 2 | 01232312131123323 | 6 | 01231323123211213 |
| 3 | 01232313121122333 | 7 | 01233121321213231 |
| 4 | 01232313123122113 | | |

### S2.25. Colorings of $L^{(4)}_{22,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22$. The anchor context $(1, 2, 4, 3)$ has colors $(0, \ldots, 3)$.

TABLE S69: Every normalized coloring of $L^{(4)}_{22,9}$.

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 1 | 0132120301230312033201 | 109 | 0132120310320212032301 |
| 2 | 0132120301230312033210 | 110 | 0132120310320212032310 |
| 3 | 0132120301230312303201 | 111 | 0132120310320212302301 |
| 4 | 0132120301230312303210 | 112 | 0132120310320212302310 |
| 5 | 0132120310230312033201 | 113 | 0132120301322012030312 |
| 6 | 0132120310230312033210 | 114 | 0132120301322012030321 |
| 7 | 0132120310230312303201 | 115 | 0132120301322012300312 |
| 8 | 0132120310230312303210 | 116 | 0132120301322012300321 |
| 9 | 0132120301233012030213 | 117 | 0132120310322012030312 |
| 10 | 0132120301233012030231 | 118 | 0132120310322012030321 |

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 11 | 013212030l233012300213 | 119 | 01321203l0322012300312 |
| 12 | 013212030l233012300231 | 120 | 01321203l0322012300321 |
| 13 | 013212031023301203021 3 | 121 | 013213201332200112l302 |
| 14 | 013212031023301203023 l | 122 | 013213201332200112l320 |
| 15 | 013212031023301230021 3 | 123 | 013213201332200121l302 |
| 16 | 013212031023301230023 l | 124 | 013213201332200121l320 |
| 17 | 013213201323300112l203 | 125 | 013213203132200112l302 |
| 18 | 013213201323300112l230 | 126 | 013213203132200112l320 |
| 19 | 013213201323300121l203 | 127 | 013213203132200121l302 |
| 20 | 013213201323300121l230 | 128 | 013213203132200121l320 |
| 21 | 013213203123300112l203 | 129 | 013210230l320210232301 |
| 22 | 013213203123300112l230 | 130 | 013210230l320210232310 |
| 23 | 013213203123300121l203 | 131 | 013210230l320210322301 |
| 24 | 013213203123300121l230 | 132 | 013210230l320210322310 |
| 25 | 013210230l230310233201 | 133 | 013210231032021023230l |
| 26 | 013210230l230310233210 | 134 | 013210231032021023231 0 |
| 27 | 013210230l230310323201 | 135 | 013210231032021032230l |
| 28 | 013210230l230310323210 | 136 | 013210231032021032231 0 |
| 29 | 013210231023031023320l | 137 | 013210230l322010230312 |
| 30 | 013210231023031023321 0 | 138 | 013210230l322010230321 |
| 31 | 013210231023031032320l | 139 | 013210230l322010320312 |
| 32 | 013210231023031032321 0 | 140 | 013210230l322010320321 |
| 33 | 013210230l233010230213 | 141 | 013210231032201023031 2 |
| 34 | 013210230l233010230231 | 142 | 013210231032201023032 1 |
| 35 | 013210230l233010320213 | 143 | 013210231032201032031 2 |
| 36 | 013210230l233010320231 | 144 | 013210231032201032032 1 |
| 37 | 013210231023301023021 3 | 145 | 013213201332021002230l |
| 38 | 013210231023301023023 l | 146 | 013213201332021002231 0 |
| 39 | 013210231023301032021 3 | 147 | 013213201332021020230l |
| 40 | 013210231023301032023 l | 148 | 013213201332021020231 0 |
| 41 | 013213201323031002320l | 149 | 013213203132021002230l |
| 42 | 013213201323031002321 0 | 150 | 013213203132021002231 0 |
| 43 | 013213201323031020320l | 151 | 013213203132021020230l |
| 44 | 013213201323031020321 0 | 152 | 013213203132021020231 0 |
| 45 | 013213203123031002320l | 153 | 013213201332201002031 2 |
| 46 | 013213203123031002321 0 | 154 | 013213201332201002032 1 |
| 47 | 013213203123031020320l | 155 | 013213201332201020031 2 |
| 48 | 013213203123031020321 0 | 156 | 013213201332201020032 1 |
| 49 | 013213201323301002021 3 | 157 | 013213203132201002031 2 |
| 50 | 013213201323301002023 l | 158 | 013213203132201002032 1 |
| 51 | 013213201323301020021 3 | 159 | 013213203132201020031 2 |
| 52 | 013213201323301020023 l | 160 | 013213203132201020032 1 |
| 53 | 013213203123301002021 3 | 161 | 013223010331021202230l |
| 54 | 013213203123301002023 l | 162 | 013223010331021202231 0 |
| 55 | 013213203123301020021 3 | 163 | 013223010331021220230l |
| 56 | 013213203123301020023 l | 164 | 013223010331021220231 0 |
| 57 | 013231201321300123120 3 | 165 | 013223013031021202230l |
| 58 | 013231201321300123123 0 | 166 | 013223013031021202231 0 |

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 59 | 013231201321300132 1203 | 167 | 013223013031021220 2301 |
| 60 | 013231201321300132 1230 | 168 | 013223013031021220 2310 |
| 61 | 013231203121300123 1203 | 169 | 013223010331201202 0312 |
| 62 | 013231203121300123 1230 | 170 | 013223010331201202 0321 |
| 63 | 013231203121300132 1203 | 171 | 013223010331201220 0312 |
| 64 | 013231203121300132 1230 | 172 | 013223010331201220 0321 |
| 65 | 013232101321300213 1203 | 173 | 013223013031201202 0312 |
| 66 | 013232101321300213 1230 | 174 | 013223013031201202 0321 |
| 67 | 013232101321300231 1203 | 175 | 013223013031201220 0312 |
| 68 | 013232101321300231 1230 | 176 | 013223013031201220 0321 |
| 69 | 013232103121300213 1203 | 177 | 013221030131022103 1302 |
| 70 | 013232103121300213 1230 | 178 | 013221030131022103 1320 |
| 71 | 013232103121300231 1203 | 179 | 013221030131022130 1302 |
| 72 | 013232103121300231 1230 | 180 | 013221030131022130 1320 |
| 73 | 013230210321031023 3201 | 181 | 013221031031022103 1302 |
| 74 | 013230210321031023 3210 | 182 | 013221031031022103 1320 |
| 75 | 013230210321031032 3201 | 183 | 013221031031022130 1302 |
| 76 | 013230210321031032 3210 | 184 | 013221031031022130 1320 |
| 77 | 013230213021031023 3201 | 185 | 013223010331022101 1302 |
| 78 | 013230213021031023 3210 | 186 | 013223010331022101 1320 |
| 79 | 013230213021031032 3201 | 187 | 013223010331022110 1302 |
| 80 | 013230213021031032 3210 | 188 | 013223010331022110 1320 |
| 81 | 013230210321301023 0213 | 189 | 013223013031022101 1302 |
| 82 | 013230210321301023 0231 | 190 | 013223013031022101 1320 |
| 83 | 013230210321301032 0213 | 191 | 013223013031022110 1302 |
| 84 | 013230210321301032 0231 | 192 | 013223013031022110 1320 |
| 85 | 013230213021301023 0213 | 193 | 013223101331200212 1302 |
| 86 | 013230213021301023 0231 | 194 | 013223101331200212 1320 |
| 87 | 013230213021301032 0213 | 195 | 013223101331200221 1302 |
| 88 | 013230213021301032 0231 | 196 | 013223101331200221 1320 |
| 89 | 013232010321031203 3201 | 197 | 013223103131200212 1302 |
| 90 | 013232010321031203 3210 | 198 | 013223103131200212 1320 |
| 91 | 013232010321031230 3201 | 199 | 013223103131200221 1302 |
| 92 | 013232010321031230 3210 | 200 | 013223103131200221 1320 |
| 93 | 013232013021031203 3201 | 201 | 013220130131022013 1302 |
| 94 | 013232013021031203 3210 | 202 | 013220130131022013 1320 |
| 95 | 013232013021031230 3201 | 203 | 013220130131022031 1302 |
| 96 | 013232013021031230 3210 | 204 | 013220130131022031 1320 |
| 97 | 013232010321301203 0213 | 205 | 013220131031022013 1302 |
| 98 | 013232010321301203 0231 | 206 | 013220131031022013 1320 |
| 99 | 013232010321301230 0213 | 207 | 013220131031022031 1302 |
| 100 | 013232010321301230 0231 | 208 | 013220131031022031 1320 |
| 101 | 013232013021301203 0213 | 209 | 013223101331022001 1302 |
| 102 | 013232013021301203 0231 | 210 | 013223101331022001 1320 |
| 103 | 013232013021301230 0213 | 211 | 013223101331022010 1302 |
| 104 | 013232013021301230 0231 | 212 | 013223101331022010 1320 |
| 105 | 013212030132021203 2301 | 213 | 013223103131022001 1302 |
| 106 | 013212030132021203 2310 | 214 | 013223103131022001 1320 |

| $j$ | Color word | $j$ | Color word |
|---|---|---|---|
| 107 | 0132120301320212302301 | 215 | 0132231031310220101302 |
| 108 | 0132120301320212302310 | 216 | 0132231031310220101320 |

*Exhaustive classification of chromatic identities.* For a chromatically inseparable pair $(u, v)$, form the two directed witness sets

$$W_{10}(u,v) = \{j : s_j(u) = 1,\ s_j(v) = 0\},$$
$$W_{01}(u,v) = \{j : s_j(u) = 0,\ s_j(v) = 1\}.$$

Both sets are nonempty exactly for a two-way purely chromatic identity. If only $W_{10}$ is empty, $u \Rightarrow v$ holds and its converse fails; if only $W_{01}$ is empty, the implication is reversed. If both are empty, the pair is inseparable under all ordinary states. Any member of either witness set is automatically nonextendable, since every color-class state has equal values on a chromatically inseparable pair.

The following Python standard-library procedure reconstructs $H_{21,13}$ directly from its 13 contexts in Table S3, enumerates all binary assignments, and reconstructs normalized strong colorings as partitions into three disjoint state supports. Lexicographic support order agrees with Table S42. It prints both complete directed witness sets for every chromatic identity, rather than only the representatives in Table S70.

```
from itertools import combinations
E = [(1,2,3),(7,8,9),(4,5,6),(2,8,5),
     (3,11,14),(7,11,15),(4,11,24),
     (9,10,23),(1,10,19),(4,10,18),
     (1,13,20),(7,13,21),(6,13,22)]
V = sorted(set().union(*map(set, E)))
M = [sum(1 << V.index(v) for v in e) for e in E]
S = sorted(tuple(v for i,v in enumerate(V) if b >> i & 1)
           for b in range(1 << len(V))
           if all((b & m).bit_count() == 1 for m in M))
S = list(map(frozenset, S))
C = []
for blocks in combinations(S, 3):
    if any(a & b for a,b in combinations(blocks, 2)):
        continue
    blocks = [next(s for s in blocks if a in s) for a in (2,8,5)]
    C.append({v:k for k,s in enumerate(blocks) for v in s})
assert (len(S), len(C)) == (21, 2)
counts = [0, 0, 0]
for u,v in combinations(V, 2):
    if not all(c[u] == c[v] for c in C):
        continue
    W10 = [j for j,s in enumerate(S, 1) if u in s and v not in s]
    W01 = [j for j,s in enumerate(S, 1) if v in s and u not in s]
    k = 0 if W10 and W01 else 1 if W10 or W01 else 2
    counts[k] += 1
    print((u,v), W10, W01)
assert counts == [15, 6, 0]
```

TABLE S70. Both directed truth-pattern witnesses for all 15 two-way purely chromatic pairs of $H_{21,13}$ and both displayed pairs of $T_{19,11}$. For $H_{21,13}$, $s_j$ is the state in Table S42; for $T_{19,11}$, entries give supports. Every witness is nonextendable and satisfies every context equation.

| Configuration | Pair $(u,v)$ | $(s(u),s(v))=(1,0)$ | $(s(u),s(v))=(0,1)$ |
|---|---|---|---|
| $T_{19,11}$ | $(14,28)$ | $\{1,5,7,14,18,23,24\}$ | $\{1,6,8,10,11,27,28\}$ |
| $T_{19,11}$ | $(18,27)$ | $\{1,5,7,14,18,23,24\}$ | $\{1,6,8,10,11,27,28\}$ |
| $H_{21,13}$ | $(1,11)$ | $s_1$ | $s_{12}$ |
| $H_{21,13}$ | $(1,18)$ | $s_1$ | $s_{11}$ |
| $H_{21,13}$ | $(1,21)$ | $s_2$ | $s_9$ |
| $H_{21,13}$ | $(2,14)$ | $s_{12}$ | $s_1$ |
| $H_{21,13}$ | $(4,13)$ | $s_1$ | $s_{18}$ |
| $H_{21,13}$ | $(4,15)$ | $s_7$ | $s_4$ |
| $H_{21,13}$ | $(4,19)$ | $s_1$ | $s_{11}$ |
| $H_{21,13}$ | $(5,22)$ | $s_{18}$ | $s_1$ |
| $H_{21,13}$ | $(7,10)$ | $s_2$ | $s_{20}$ |
| $H_{21,13}$ | $(7,20)$ | $s_2$ | $s_9$ |
| $H_{21,13}$ | $(7,24)$ | $s_7$ | $s_4$ |
| $H_{21,13}$ | $(8,23)$ | $s_{20}$ | $s_2$ |
| $H_{21,13}$ | $(15,19)$ | $s_1$ | $s_7$ |
| $H_{21,13}$ | $(18,21)$ | $s_2$ | $s_1$ |
| $H_{21,13}$ | $(20,24)$ | $s_7$ | $s_2$ |

TABLE S71: All ten connected, edge-faithful, colorable restrictions with a forced color identity in the specified inventory.

| retained diagonals | rays | contexts | colorings | inseparable pairs |
|---|---|---|---|---|
| $\{10,11\}$ | 17 | 10 | 4 | $(10,24)$, $(11,18)$ |
| $\{10,12\}$ | 17 | 10 | 4 | $(10,16)$, $(12,23)$ |
| $\{10,13\}$ | 17 | 10 | 4 | $(10,20)$, $(13,19)$ |
| $\{11,12\}$ | 17 | 10 | 4 | $(11,25)$, $(12,14)$ |
| $\{11,13\}$ | 17 | 10 | 4 | $(11,21)$, $(13,15)$ |
| $\{12,13\}$ | 17 | 10 | 4 | $(12,22)$, $(13,17)$ |
| $\{10,11,12\}$ | 21 | 13 | 2 | 21 pairs |
| $\{10,11,13\}$ | 21 | 13 | 2 | 21 pairs |
| $\{10,12,13\}$ | 21 | 13 | 2 | 21 pairs |
| $\{11,12,13\}$ | 21 | 13 | 2 | 21 pairs |

### S2.26. An explicit state certificate for the completed Yu–Oh hypergraph

The first four contexts of Eq. (15) force exactly one of vertices 2, 5, and 8 to have value one. In the triad containing that coordinate vertex, both other vertices have value zero. In each of the other two triads, one of the two remaining vertices has value one. There are therefore $3 \cdot 2^2 = 12$ possible supports $T_k$ on vertices $1,\dots,9$, listed in Table S72.

For each such support, exactly one of vertices $10,11,12,13$ has no orthogonal neighbor in $T_k$; call it $\eta_k$. The other three are forced to zero, while $s(\eta_k)$ may be either zero or one. This gives precisely 24 assignments on the original 13 vertices. Every completion vertex $w$ belongs to one context $\{i,j,w\}$ and has the uniquely determined value

$$s(w) = 1 - s(i) - s(j). \tag{S7}$$

The listed assignments have $s(i) + s(j) \le 1$ on every original orthogonal pair, so these

extensions are valid two-valued states. Conversely, every state of the completed hypergraph restricts to one of these assignments.

For a compact separation certificate, encode the profile of vertex $i$ as the nonnegative integer

$$m_i = \sum_{k=1}^{12} \sum_{b=0}^{1} s_{k,b}(i)\, 2^{2(k-1)+b}.$$

In hexadecimal notation these 25 integers are

| $i$ | 1 | 2 | 3 | 4 | 5 |
|---|---|---|---|---|---|
| $m_i$ | 0000FF | 00FF00 | FF0000 | 030F03 | 3C003C |
| $i$ | 6 | 7 | 8 | 9 | 10 |
| $m_i$ | C0F0C0 | 0C330C | C300C3 | 30CC30 | 882000 |
| $i$ | 11 | 12 | 13 | 14 | 15 |
| $m_i$ | 0080A0 | 00020A | 220800 | 007F5F | F34C53 |
| $i$ | 16 | 17 | 18 | 19 | 20 |
| $m_i$ | CF31C5 | 3F0D35 | 74D0FC | 77DF00 | DDF700 |
| $i$ | 21 | 22 | 23 | 24 | 25 |
| $m_i$ | D1C4F3 | 1D073F | 4713CF | FC705C | 00FDF5 |

(S8)

All 25 integers are nonzero and pairwise distinct, certifying unitality and separation. Notably, the Hamming weights (number of states in which $s(i) = 1$) partition the vertices into three classes: 8 for $i \in \{1, \ldots, 9\}$, 3 for $i \in \{10, \ldots, 13\}$, and 13 for the completion vertices $i \in \{14, \ldots, 25\}$ (with $8+3+13 = 24$ in each completion context), in contrast to the uniform state weight 2 across all 15 vertices of $G_{32}$ and consistent with the fractional chromatic–clique gap $\chi_f(\Gamma(H^+_{\mathrm{YO}})) = 35/11 > 3 = \omega(\Gamma(H^+_{\mathrm{YO}}))$.

TABLE S72. The 24 states are indexed by $(k, b)$, where $1 \le k \le 12$ and $b \in \{0, 1\}$. Their support on vertices $1, \ldots, 13$ is $T_k$ when $b = 0$ and $T_k \cup \{\eta_k\}$ when $b = 1$. Equation (S7) determines all remaining values.

| $k$ | $T_k$ | $\eta_k$ |
|---|---|---|
| 1 | $\{1, 4, 8\}$ | 12 |
| 2 | $\{1, 5, 7\}$ | 12 |
| 3 | $\{1, 5, 9\}$ | 11 |
| 4 | $\{1, 6, 8\}$ | 11 |
| 5 | $\{2, 4, 7\}$ | 12 |
| 6 | $\{2, 4, 9\}$ | 13 |
| 7 | $\{2, 6, 7\}$ | 10 |
| 8 | $\{2, 6, 9\}$ | 11 |
| 9 | $\{3, 4, 8\}$ | 13 |
| 10 | $\{3, 5, 7\}$ | 10 |
| 11 | $\{3, 5, 9\}$ | 13 |
| 12 | $\{3, 6, 8\}$ | 10 |

### S2.27. Hesse $H_{37}$: all states and separating profiles

The canonical states $s_1, \dots, s_{153}$ are listed in Table S74. A two-valued coloring is a state with exactly one value one per context. Vertex numbers are those of Appendix S1.14 and Fig. 4: vertices $1, \dots, 16$ form the core, and $17, \dots, 37$ are completions. We list every state, a minimum separating family, and every vertex profile.

#### *S2.27.1. State parametrization and all 27 branches*

Choose $t = (t_0, t_1, t_2) \in \mathbb{Z}_3^3$ and $\varepsilon \in \{0, 1\}$. Put

$$s(M_{ak}) = \mathbf{1}_{k=t_a}, \qquad s(z) = \varepsilon, \tag{S9}$$

$$U(t) = \mathbb{Z}_3 \setminus \{t_0, t_1, t_2\}, \qquad W(t) = \mathbb{Z}_3 \setminus \{t_a - a : a \in \mathbb{Z}_3\}. \tag{S10}$$

For $\varepsilon = 0$, choose the values of $P_k$ freely for $k \in U(t)$, with other $P$ values zero. For $\varepsilon = 1$, all $P$ values are zero. Independently, choose the values of $Q_k$ freely for $k \in W(t)$, with other $Q$ values zero. All completion values are forced:

$$s(A_k) = 1 - \varepsilon - s(P_k), \tag{S11}$$

$$s(B_{ak}) = 1 - s(M_{ak}) - s(P_k), \tag{S12}$$

$$s(C_{ak}) = 1 - s(M_{a,a+k}) - s(Q_k). \tag{S13}$$

Enumerate $t$ lexicographically with $0 < 1 < 2$, then $\varepsilon = 1, 0$. Within each branch, list the available $P$ bits in increasing $k$, followed by the available $Q$ bits in increasing $k$, and order the resulting bit strings with 1 before 0. This rule fixes all state indices used in the text, tables, and verification data. Table S73 gives the index ranges of all 27 branches.

TABLE S73: All 27 state branches. The sets $U(t), W(t)$ identify the free core bits. The two choices of $\varepsilon$ give $N_0 = 2^{|U|+|W|}$ and $N_1 = 2^{|W|}$ states; their index ranges refer to Table S74.

| $t$ | $U(t)$ | $W(t)$ | $N_0$ | $r$ for $\varepsilon = 0$ | $N_1$ | $r$ for $\varepsilon = 1$ |
|---|---|---|---|---|---|---|
| $(0,0,0)$ | $\{1,2\}$ | $\varnothing$ | 4 | 2–5 | 1 | 1 |
| $(0,0,1)$ | $\{2\}$ | $\{1\}$ | 4 | 8–11 | 2 | 6–7 |
| $(0,0,2)$ | $\{1\}$ | $\{1\}$ | 4 | 14–17 | 2 | 12–13 |
| $(0,1,0)$ | $\{2\}$ | $\{2\}$ | 4 | 20–23 | 2 | 18–19 |
| $(0,1,1)$ | $\{2\}$ | $\{1\}$ | 4 | 26–29 | 2 | 24–25 |
| $(0,1,2)$ | $\varnothing$ | $\{1,2\}$ | 4 | 34–37 | 4 | 30–33 |
| $(0,2,0)$ | $\{1\}$ | $\{2\}$ | 4 | 40–43 | 2 | 38–39 |
| $(0,2,1)$ | $\varnothing$ | $\varnothing$ | 1 | 45 | 1 | 44 |
| $(0,2,2)$ | $\{1\}$ | $\{2\}$ | 4 | 48–51 | 2 | 46–47 |
| $(1,0,0)$ | $\{2\}$ | $\{0\}$ | 4 | 54–57 | 2 | 52–53 |
| $(1,0,1)$ | $\{2\}$ | $\{0\}$ | 4 | 60–63 | 2 | 58–59 |
| $(1,0,2)$ | $\varnothing$ | $\varnothing$ | 1 | 65 | 1 | 64 |
| $(1,1,0)$ | $\{2\}$ | $\{2\}$ | 4 | 68–71 | 2 | 66–67 |
| $(1,1,1)$ | $\{0,2\}$ | $\varnothing$ | 4 | 73–76 | 1 | 72 |
| $(1,1,2)$ | $\{0\}$ | $\{2\}$ | 4 | 79–82 | 2 | 77–78 |
| $(1,2,0)$ | $\varnothing$ | $\{0,2\}$ | 4 | 87–90 | 4 | 83–86 |
| $(1,2,1)$ | $\{0\}$ | $\{0\}$ | 4 | 93–96 | 2 | 91–92 |
| $(1,2,2)$ | $\{0\}$ | $\{2\}$ | 4 | 99–102 | 2 | 97–98 |
| $(2,0,0)$ | $\{1\}$ | $\{0\}$ | 4 | 105–108 | 2 | 103–104 |
| $(2,0,1)$ | $\varnothing$ | $\{0,1\}$ | 4 | 113–116 | 4 | 109–112 |
| $(2,0,2)$ | $\{1\}$ | $\{1\}$ | 4 | 119–122 | 2 | 117–118 |
| $(2,1,0)$ | $\varnothing$ | $\varnothing$ | 1 | 124 | 1 | 123 |
| $(2,1,1)$ | $\{0\}$ | $\{1\}$ | 4 | 127–130 | 2 | 125–126 |
| $(2,1,2)$ | $\{0\}$ | $\{1\}$ | 4 | 133–136 | 2 | 131–132 |
| $(2,2,0)$ | $\{1\}$ | $\{0\}$ | 4 | 139–142 | 2 | 137–138 |
| $(2,2,1)$ | $\{0\}$ | $\{0\}$ | 4 | 145–148 | 2 | 143–144 |
| $(2,2,2)$ | $\{0,1\}$ | $\varnothing$ | 4 | 150–153 | 1 | 149 |
| Total | | | 99 | | 54 | |

### *S2.27.2. Canonical two-valued colorings: all 153 states*

Table S74 explicitly lists all 153 states. The last column is the complete binary vector in vertex order $1, \ldots, 37$, separated after the 16 core entries. Every row has context sum one in all 24 contexts. The support-size distribution is

$$\begin{array}{c|cccccc} m & 12 & 13 & 14 & 15 & 16 & 18 \\ \#\{s : |\operatorname{supp} s| = m\} & 6 & 18 & 27 & 24 & 51 & 27 \end{array} \tag{S14}$$

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$.

| $r$ | Core support | Completion support | $\lvert\operatorname{supp} s_r\rvert$ | 37-bit state vector |
|---|---|---|---|---|
| 1 | $\{1,2,5,8\}$ | $\{21,22,24,25,27,28,30,31,32,33,35,37\}$ | 16 | 1100100100000000 00001101101101111010 1 |
| 2 | $\{2,5,8,12,13\}$ | $\{17,30,31,32,33,35,37\}$ | 12 | 0100100100011000 10000000000001111010 1 |
| 3 | $\{2,5,8,12\}$ | $\{17,19,22,25,28,30,31,32,33,35,37\}$ | 15 | 0100100100010000 10100100100101111010 1 |
| 4 | $\{2,5,8,13\}$ | $\{17,18,21,24,27,30,31,32,33,35,37\}$ | 15 | 0100100100001000 11001001001011110101 |
| 5 | $\{2,5,8\}$ | $\{17,18,19,21,22,24,25,27,28,30,31,32,33,35,37\}$ | 18 | 0100100100000000 11101101101101110101 |
| 6 | $\{1,2,5,9,15\}$ | $\{21,22,24,25,26,28,31,32,35\}$ | 14 | 1100100010000010 00001101110100110010 0 |
| 7 | $\{1,2,5,9\}$ | $\{21,22,24,25,26,28,30,31,32,33,35,36\}$ | 16 | 1100100010000000 00001101110101111011 0 |
| 8 | $\{2,5,9,13,15\}$ | $\{17,18,21,24,26,31,32,35\}$ | 13 | 0100100010001010 11001001010000110010 0 |
| 9 | $\{2,5,9,13\}$ | $\{17,18,21,24,26,30,31,32,33,35,36\}$ | 15 | 0100100010001000 11001001010001111011 0 |
| 10 | $\{2,5,9,15\}$ | $\{17,18,19,21,22,24,25,26,28,31,32,35\}$ | 16 | 0100100010000010 11101101110100110010 0 |
| 11 | $\{2,5,9\}$ | $\{17,18,19,21,22,24,25,26,28,30,31,32,33,35,36\}$ | 18 | 0100100010000000 11101101110101111011 0 |
| 12 | $\{1,2,5,10,15\}$ | $\{21,22,24,25,26,27,31,32,37\}$ | 14 | 1100100001000010 00001101111000110000 1 |
| 13 | $\{1,2,5,10\}$ | $\{21,22,24,25,26,27,30,31,32,33,36,37\}$ | 16 | 1100100001000000 00001101111001111001 1 |
| 14 | $\{2,5,10,12,15\}$ | $\{17,19,22,25,26,31,32,37\}$ | 13 | 0100100001010010 10100100110000110000 1 |
| 15 | $\{2,5,10,12\}$ | $\{17,19,22,25,26,30,31,32,33,36,37\}$ | 15 | 0100100001010000 10100100110001111001 1 |
| 16 | $\{2,5,10,15\}$ | $\{17,18,19,21,22,24,25,26,27,31,32,37\}$ | 16 | 0100100001000010 11101101111000110000 1 |
| 17 | $\{2,5,10\}$ | $\{17,18,19,21,22,24,25,26,27,30,31,32,33,36,37\}$ | 18 | 0100100001000000 11101101111001111001 1 |
| 18 | $\{1,2,6,8,16\}$ | $\{21,22,23,25,27,28,30,33,35\}$ | 14 | 1100010100000001 00001110101101010010 0 |
| 19 | $\{1,2,6,8\}$ | $\{21,22,23,25,27,28,30,31,33,34,35,37\}$ | 16 | 1100010100000000 00001110101101011011 1 |
| 20 | $\{2,6,8,13,16\}$ | $\{17,18,21,23,27,30,33,35\}$ | 13 | 0100010100001001 10010100010010010100 |
| 21 | $\{2,6,8,13\}$ | $\{17,18,21,23,27,30,31,33,34,35,37\}$ | 15 | 0100010100001000 11001010001011011101 |
| 22 | $\{2,6,8,16\}$ | $\{17,18,19,21,22,23,25,27,28,30,33,35\}$ | 16 | 0100010100000001 11101110101101010100 |
| 23 | $\{2,6,8\}$ | $\{17,18,19,21,22,23,25,27,28,30,31,33,34,35,37\}$ | 18 | 0100010100000000 11101110101101011101 |
| 24 | $\{1,2,6,9,15\}$ | $\{21,22,23,25,26,28,31,34,35\}$ | 14 | 1100010010000010 00001110110100100110 0 |
| 25 | $\{1,2,6,9\}$ | $\{21,22,23,25,26,28,30,31,33,34,35,36\}$ | 16 | 1100010010000000 00001110110101101111 0 |
| 26 | $\{2,6,9,13,15\}$ | $\{17,18,21,23,26,31,34,35\}$ | 13 | 0100010010001010 11001010010000100110 0 |
| 27 | $\{2,6,9,13\}$ | $\{17,18,21,23,26,30,31,33,34,35,36\}$ | 15 | 0100010010001000 11001010010001101111 0 |
| 28 | $\{2,6,9,15\}$ | $\{17,18,19,21,22,23,25,26,28,31,34,35\}$ | 16 | 0100010010000010 11101110110100100110 0 |
| 29 | $\{2,6,9\}$ | $\{17,18,19,21,22,23,25,26,28,30,31,33,34,35,36\}$ | 18 | 0100010010000000 11101110101101101110 |
| 30 | $\{1,2,6,10,15,16\}$ | $\{21,22,23,25,26,27\}$ | 12 | 1100010001000011 00001110111000000000 |
| 31 | $\{1,2,6,10,15\}$ | $\{21,22,23,25,26,27,31,34,37\}$ | 14 | 1100010001000010 00001110111000100100 1 |
| 32 | $\{1,2,6,10,16\}$ | $\{21,22,23,25,26,27,30,33,36\}$ | 14 | 1100010001000001 00001110111001001001 0 |
| 33 | $\{1,2,6,10\}$ | $\{21,22,23,25,26,27,30,31,33,34,36,37\}$ | 16 | 1100010001000000 00001110111001101101 1 |
| 34 | $\{2,6,10,15,16\}$ | $\{17,18,19,21,22,23,25,26,27\}$ | 14 | 0100010001000011 11101110111000000000 |
| 35 | $\{2,6,10,15\}$ | $\{17,18,19,21,22,23,25,26,27,31,34,37\}$ | 16 | 0100010001000010 11101110111000100100 1 |
| 36 | $\{2,6,10,16\}$ | $\{17,18,19,21,22,23,25,26,27,30,33,36\}$ | 16 | 0100010001000001 11101110111001001001 0 |
| 37 | $\{2,6,10\}$ | $\{17,18,19,21,22,23,25,26,27,30,31,33,34,36,37\}$ | 18 | 0100010001000000 11101110111001101101 1 |
| 38 | $\{1,2,7,8,16\}$ | $\{21,22,23,24,27,28,30,32,35\}$ | 14 | 1100001100000001 00001111000110101001 0 |
| 39 | $\{1,2,7,8\}$ | $\{21,22,23,24,27,28,30,31,32,34,35,37\}$ | 16 | 1100001100000000 00001111000110111011 1 |
| 40 | $\{2,7,8,12,16\}$ | $\{17,19,22,23,28,30,32,35\}$ | 13 | 0100001100010001 10100110000101010010 0 |
| 41 | $\{2,7,8,12\}$ | $\{17,19,22,23,28,30,31,32,34,35,37\}$ | 15 | 0100001100010000 10100110000101110111 1 |
| 42 | $\{2,7,8,16\}$ | $\{17,18,19,21,22,23,24,27,28,30,32,35\}$ | 16 | 0100001100000001 11101111001101010010 0 |
| 43 | $\{2,7,8\}$ | $\{17,18,19,21,22,23,24,27,28,30,31,32,34,35,37\}$ | 18 | 0100001100000000 11101111001101110111 1 |
| 44 | $\{1,2,7,9\}$ | $\{21,22,23,24,26,28,30,31,32,34,35,36\}$ | 16 | 1100001010000000 00001111010101110111 0 |
| 45 | $\{2,7,9\}$ | $\{17,18,19,21,22,23,24,26,28,30,31,32,34,35,36\}$ | 18 | 0100001010000000 11101111010101110111 0 |
| 46 | $\{1,2,7,10,16\}$ | $\{21,22,23,24,26,27,30,32,36\}$ | 14 | 1100001001000001 00001111010010100010 |
| 47 | $\{1,2,7,10\}$ | $\{21,22,23,24,26,27,30,31,32,34,36,37\}$ | 16 | 1100001001000000 00001111010011101011 |
| 48 | $\{2,7,10,12,16\}$ | $\{17,19,22,23,26,30,32,36\}$ | 13 | 0100001001010001 10100110010001010010 |
| 49 | $\{2,7,10,12\}$ | $\{17,19,22,23,26,30,31,32,34,36,37\}$ | 15 | 0100001001010000 10100110010001111011 |
| 50 | $\{2,7,10,16\}$ | $\{17,18,19,21,22,23,24,26,27,30,32,36\}$ | 16 | 0100001001000001 11101111010010100010 |
| 51 | $\{2,7,10\}$ | $\{17,18,19,21,22,23,24,26,27,30,31,32,34,36,37\}$ | 18 | 0100001001000000 11101111010011101011 |

*Continued on the next page.*

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

| $r$ | Core support | Completion support | $\lvert\operatorname{supp} s_r\rvert$ | 37-bit state vector |
|---|---|---|---|---|
| 52 | $\{1,3,5,8,14\}$ | $\{20,22,24,25,27,28,31,33,37\}$ | 14 | 1010100100000100 0001010111001010001 |
| 53 | $\{1,3,5,8\}$ | $\{20,22,24,25,27,28,29,31,32,33,35,37\}$ | 16 | 1010100100000000 0001010111101110101 |
| 54 | $\{3,5,8,13,14\}$ | $\{17,18,20,24,27,31,33,37\}$ | 13 | 0010100100001100 1101000100100010001 |
| 55 | $\{3,5,8,13\}$ | $\{17,18,20,24,27,29,31,32,33,35,37\}$ | 15 | 0010100100001000 1101000100101011101 |
| 56 | $\{3,5,8,14\}$ | $\{17,18,19,20,22,24,25,27,28,31,33,37\}$ | 16 | 0010100100000100 1111010111001010001 |
| 57 | $\{3,5,8\}$ | $\{17,18,19,20,22,24,25,27,28,29,31,32,33,35,37\}$ | 18 | 0010100100000000 1111010111101110101 |
| 58 | $\{1,3,5,9,14\}$ | $\{20,22,24,25,26,28,31,33,36\}$ | 14 | 1010100010000100 0001010111010010010 |
| 59 | $\{1,3,5,9\}$ | $\{20,22,24,25,26,28,29,31,32,33,35,36\}$ | 16 | 1010100010000000 0001010111011101110 |
| 60 | $\{3,5,9,13,14\}$ | $\{17,18,20,24,26,31,33,36\}$ | 13 | 0010100010001100 1101000101000010010 |
| 61 | $\{3,5,9,13\}$ | $\{17,18,20,24,26,29,31,32,33,35,36\}$ | 15 | 0010100010001000 1101000101001011110 |
| 62 | $\{3,5,9,14\}$ | $\{17,18,19,20,22,24,25,26,28,31,33,36\}$ | 16 | 0010100010000100 1111010111010010010 |
| 63 | $\{3,5,9\}$ | $\{17,18,19,20,22,24,25,26,28,29,31,32,33,35,36\}$ | 18 | 0010100010000000 1111010111011110110 |
| 64 | $\{1,3,5,10\}$ | $\{20,22,24,25,26,27,29,31,32,33,36,37\}$ | 16 | 1010100001000000 0001010111101011110011 |
| 65 | $\{3,5,10\}$ | $\{17,18,19,20,22,24,25,26,27,29,31,32,33,36,37\}$ | 18 | 0010100001000000 1111010111101011110011 |
| 66 | $\{1,3,6,8,16\}$ | $\{20,22,23,25,27,28,29,33,35\}$ | 14 | 1010010100000001 0001011010111000010100 |
| 67 | $\{1,3,6,8\}$ | $\{20,22,23,25,27,28,29,31,33,34,35,37\}$ | 16 | 1010010100000000 0001011010111010011101 |
| 68 | $\{3,6,8,13,16\}$ | $\{17,18,20,23,27,29,33,35\}$ | 13 | 0010010100001001 1101001000101000010100 |
| 69 | $\{3,6,8,13\}$ | $\{17,18,20,23,27,29,31,33,34,35,37\}$ | 15 | 0010010100001000 1101001000101011101 |
| 70 | $\{3,6,8,16\}$ | $\{17,18,19,20,22,23,25,27,28,29,33,35\}$ | 16 | 0010010100000001 1111011010111000010100 |
| 71 | $\{3,6,8\}$ | $\{17,18,19,20,22,23,25,27,28,29,31,33,34,35,37\}$ | 18 | 0010010100000000 1111011010111010011101 |
| 72 | $\{1,3,6,9\}$ | $\{20,22,23,25,26,28,29,31,33,34,35,36\}$ | 16 | 1010010010000000 0001011010110101011110 |
| 73 | $\{3,6,9,11,13\}$ | $\{18,29,31,33,34,35,36\}$ | 12 | 0010010010101000 0100000000001010111110 |
| 74 | $\{3,6,9,11\}$ | $\{18,19,22,25,28,29,31,33,34,35,36\}$ | 15 | 0010010010100000 0110010010011010111110 |
| 75 | $\{3,6,9,13\}$ | $\{17,18,20,23,26,29,31,33,34,35,36\}$ | 15 | 0010010010001000 1101001001001010111110 |
| 76 | $\{3,6,9\}$ | $\{17,18,19,20,22,23,25,26,28,29,31,33,34,35,36\}$ | 18 | 0010010010000000 1111011011011010111110 |
| 77 | $\{1,3,6,10,16\}$ | $\{20,22,23,25,26,27,29,33,36\}$ | 14 | 1010010001000001 0001011011101000010010 |
| 78 | $\{1,3,6,10\}$ | $\{20,22,23,25,26,27,29,31,33,34,36,37\}$ | 16 | 1010010001000000 0001011011101010011011 |
| 79 | $\{3,6,10,11,16\}$ | $\{18,19,22,25,27,29,33,36\}$ | 13 | 0010010001100001 0110010010100010010 |
| 80 | $\{3,6,10,11\}$ | $\{18,19,22,25,27,29,31,33,34,36,37\}$ | 15 | 0010010001100000 0110010010101011011 |
| 81 | $\{3,6,10,16\}$ | $\{17,18,19,20,22,23,25,26,27,29,33,36\}$ | 16 | 0010010001000001 1111011011010010010 |
| 82 | $\{3,6,10\}$ | $\{17,18,19,20,22,23,25,26,27,29,31,33,34,36,37\}$ | 18 | 0010010001000000 1111011011010011011 |
| 83 | $\{1,3,7,8,14,16\}$ | $\{20,22,23,24,27,28\}$ | 12 | 1010001100000101 0001011100110000000000 |
| 84 | $\{1,3,7,8,14\}$ | $\{20,22,23,24,27,28,31,34,37\}$ | 14 | 1010001100000100 0001011100110010001001 |
| 85 | $\{1,3,7,8,16\}$ | $\{20,22,23,24,27,28,29,32,35\}$ | 14 | 1010001100000001 0001011100110010010100 |
| 86 | $\{1,3,7,8\}$ | $\{20,22,23,24,27,28,29,31,32,34,35,37\}$ | 16 | 1010001100000000 0001011100111011011101 |
| 87 | $\{3,7,8,14,16\}$ | $\{17,18,19,20,22,23,24,27,28\}$ | 14 | 0010001100000101 1111011100110000000000 |
| 88 | $\{3,7,8,14\}$ | $\{17,18,19,20,22,23,24,27,28,31,34,37\}$ | 16 | 0010001100000100 1111011100110010001001 |
| 89 | $\{3,7,8,16\}$ | $\{17,18,19,20,22,23,24,27,28,29,32,35\}$ | 16 | 0010001100000001 1111011100111001001100 |
| 90 | $\{3,7,8\}$ | $\{17,18,19,20,22,23,24,27,28,29,31,32,34,35,37\}$ | 18 | 0010001100000000 1111011100111011011101 |
| 91 | $\{1,3,7,9,14\}$ | $\{20,22,23,24,26,28,31,34,36\}$ | 14 | 1010001010000100 0001011101010010001010 |
| 92 | $\{1,3,7,9\}$ | $\{20,22,23,24,26,28,29,31,32,34,35,36\}$ | 16 | 1010001010000000 0001011101011011011110 |
| 93 | $\{3,7,9,11,14\}$ | $\{18,19,22,24,28,31,34,36\}$ | 13 | 0010001010100100 0110010100010010010 |
| 94 | $\{3,7,9,11\}$ | $\{18,19,22,24,28,29,31,32,34,35,36\}$ | 15 | 0010001010100000 0110010100011011011110 |
| 95 | $\{3,7,9,14\}$ | $\{17,18,19,20,22,23,24,26,28,31,34,36\}$ | 16 | 0010001010000100 1111011101010010010 |
| 96 | $\{3,7,9\}$ | $\{17,18,19,20,22,23,24,26,28,29,31,32,34,35,36\}$ | 18 | 0010001010000000 1111011101011011011110 |
| 97 | $\{1,3,7,10,16\}$ | $\{20,22,23,24,26,27,29,32,36\}$ | 14 | 1010001001000001 0001011101101001000010 |
| 98 | $\{1,3,7,10\}$ | $\{20,22,23,24,26,27,29,31,32,34,36,37\}$ | 16 | 1010001001000000 0001011101101011011011 |
| 99 | $\{3,7,10,11,16\}$ | $\{18,19,22,24,27,29,32,36\}$ | 13 | 0010001001100001 0110010100101001000010 |
| 100 | $\{3,7,10,11\}$ | $\{18,19,22,24,27,29,31,32,34,36,37\}$ | 15 | 0010001001100000 0110010100101011011011 |
| 101 | $\{3,7,10,16\}$ | $\{17,18,19,20,22,23,24,26,27,29,32,36\}$ | 16 | 0010001001000001 1111011101101001000010 |
| 102 | $\{3,7,10\}$ | $\{17,18,19,20,22,23,24,26,27,29,31,32,34,36,37\}$ | 18 | 0010001001000000 1111011101101011011011 |
| 103 | $\{1,4,5,8,14\}$ | $\{20,21,24,25,27,28,30,33,37\}$ | 14 | 1001100100000100 0001100111011010010001 |
| 104 | $\{1,4,5,8\}$ | $\{20,21,24,25,27,28,29,30,32,33,35,37\}$ | 16 | 1001100100000000 0001100111011110110101 |
| 105 | $\{4,5,8,12,14\}$ | $\{17,19,20,25,28,30,33,37\}$ | 13 | 0001100100101100 1011000010010100010001 |
| 106 | $\{4,5,8,12\}$ | $\{17,19,20,25,28,29,30,32,33,35,37\}$ | 15 | 0001100100010000 1011000010011101101101 |
| 107 | $\{4,5,8,14\}$ | $\{17,18,19,20,21,24,25,27,28,30,33,37\}$ | 16 | 0001100100000100 1111100111011010010001 |
| 108 | $\{4,5,8\}$ | $\{17,18,19,20,21,24,25,27,28,29,30,32,33,35,37\}$ | 18 | 0001100100000000 1111100111011110110101 |
| 109 | $\{1,4,5,9,14,15\}$ | $\{20,21,24,25,26,28\}$ | 12 | 1001100010000110 0001100111010000000000 |
| 110 | $\{1,4,5,9,14\}$ | $\{20,21,24,25,26,28,30,33,36\}$ | 14 | 1001100010000100 0001100111010100010010 |
| 111 | $\{1,4,5,9,15\}$ | $\{20,21,24,25,26,28,29,32,35\}$ | 14 | 1001100010000010 0001100111011001000100 |
| 112 | $\{1,4,5,9\}$ | $\{20,21,24,25,26,28,29,30,32,33,35,36\}$ | 16 | 1001100010000000 0001100111011110110110 |
| 113 | $\{4,5,9,14,15\}$ | $\{17,18,19,20,21,24,25,26,28\}$ | 14 | 0001100010000110 1111100111010000000000 |
| 114 | $\{4,5,9,14\}$ | $\{17,18,19,20,21,24,25,26,28,30,33,36\}$ | 16 | 0001100010000100 1111100111010100010010 |

*Continued on the next page.*

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

| $r$ | Core support | Completion support | $\lvert\operatorname{supp} s_r\rvert$ | 37-bit state vector |
|---|---|---|---|---|
| 115 | $\{4,5,9,15\}$ | $\{17,18,19,20,21,24,25,26,28,29,32,35\}$ | 16 | 0001100010000010 1111100111011001001 00 |
| 116 | $\{4,5,9\}$ | $\{17,18,19,20,21,24,25,26,28,29,30,32,33,35,36\}$ | 18 | 0001100010000000 1111100111011101101 10 |
| 117 | $\{1,4,5,10,15\}$ | $\{20,21,24,25,26,27,29,32,37\}$ | 14 | 1001100001000010 0001100111101001000 01 |
| 118 | $\{1,4,5,10\}$ | $\{20,21,24,25,26,27,29,30,32,33,36,37\}$ | 16 | 1001100001000000 0001100111101101100 11 |
| 119 | $\{4,5,10,12,15\}$ | $\{17,19,20,25,26,29,32,37\}$ | 13 | 0001100001010010 1011000011001001000 01 |
| 120 | $\{4,5,10,12\}$ | $\{17,19,20,25,26,29,30,32,33,36,37\}$ | 15 | 0001100001010000 1011000011001101100 11 |
| 121 | $\{4,5,10,15\}$ | $\{17,18,19,20,21,24,25,26,27,29,32,37\}$ | 16 | 0001100001000010 1111100111101001000 01 |
| 122 | $\{4,5,10\}$ | $\{17,18,19,20,21,24,25,26,27,29,30,32,33,36,37\}$ | 18 | 0001100001000000 1111100111101101100 11 |
| 123 | $\{1,4,6,8\}$ | $\{20,21,23,25,27,28,29,30,33,34,35,37\}$ | 16 | 1001010100000000 0001101010111001110 01 |
| 124 | $\{4,6,8\}$ | $\{17,18,19,20,21,23,25,27,28,29,30,33,34,35,37\}$ | 18 | 0001010100000000 1111101010111001110 01 |
| 125 | $\{1,4,6,9,15\}$ | $\{20,21,23,25,26,28,29,34,35\}$ | 14 | 1001010010000010 0001101011010000110 00 |
| 126 | $\{1,4,6,9\}$ | $\{20,21,23,25,26,28,29,30,33,34,35,36\}$ | 16 | 1001010010000000 0001101011011001111 10 |
| 127 | $\{4,6,9,11,15\}$ | $\{18,19,21,25,28,29,34,35\}$ | 13 | 0001010010100010 0110100010011000011 00 |
| 128 | $\{4,6,9,11\}$ | $\{18,19,21,25,28,29,30,33,34,35,36\}$ | 15 | 0001010010100000 0110100010011001111 10 |
| 129 | $\{4,6,9,15\}$ | $\{17,18,19,20,21,23,25,26,28,29,34,35\}$ | 16 | 0001010010000010 1111101011011000011 00 |
| 130 | $\{4,6,9\}$ | $\{17,18,19,20,21,23,25,26,28,29,30,33,34,35,36\}$ | 18 | 0001010010000000 1111101011011001111 10 |
| 131 | $\{1,4,6,10,15\}$ | $\{20,21,23,25,26,27,29,34,37\}$ | 14 | 1001010001000010 0001101011010000100 01 |
| 132 | $\{1,4,6,10\}$ | $\{20,21,23,25,26,27,29,30,33,34,36,37\}$ | 16 | 1001010001000000 0001101011010011011 00 |
| 133 | $\{4,6,10,11,15\}$ | $\{18,19,21,25,27,29,34,37\}$ | 13 | 0001010001100010 0110100010101000001 00 |
| 134 | $\{4,6,10,11\}$ | $\{18,19,21,25,27,29,30,33,34,36,37\}$ | 15 | 0001010001100000 0110100010101100111 01 |
| 135 | $\{4,6,10,15\}$ | $\{17,18,19,20,21,23,25,26,27,29,34,37\}$ | 16 | 0001010001000010 1111101011010000100 01 |
| 136 | $\{4,6,10\}$ | $\{17,18,19,20,21,23,25,26,27,29,30,33,34,36,37\}$ | 18 | 0001010001000000 1111101011011001111 01 |
| 137 | $\{1,4,7,8,14\}$ | $\{20,21,23,24,27,28,30,34,37\}$ | 14 | 1001001100000100 0001101100110100001 00 |
| 138 | $\{1,4,7,8\}$ | $\{20,21,23,24,27,28,29,30,32,34,35,37\}$ | 16 | 1001001100000000 0001101100111101101 01 |
| 139 | $\{4,7,8,12,14\}$ | $\{17,19,20,23,28,30,34,37\}$ | 13 | 0001001100010100 1011001000010001001 00 |
| 140 | $\{4,7,8,12\}$ | $\{17,19,20,23,28,29,30,32,34,35,37\}$ | 15 | 0001001100010000 1011001000011101101 01 |
| 141 | $\{4,7,8,14\}$ | $\{17,18,19,20,21,23,24,27,28,30,34,37\}$ | 16 | 0001001100000100 1111101100110100001 00 |
| 142 | $\{4,7,8\}$ | $\{17,18,19,20,21,23,24,27,28,29,30,32,34,35,37\}$ | 18 | 0001001100000000 1111101100111101101 01 |
| 143 | $\{1,4,7,9,14\}$ | $\{20,21,23,24,26,28,30,34,36\}$ | 14 | 1001001010000100 0001101101010100010 10 |
| 144 | $\{1,4,7,9\}$ | $\{20,21,23,24,26,28,29,30,32,34,35,36\}$ | 16 | 1001001010000000 0001101101011101110 10 |
| 145 | $\{4,7,9,11,14\}$ | $\{18,19,21,24,28,30,34,36\}$ | 13 | 0001001010100100 0110100010010001010 00 |
| 146 | $\{4,7,9,11\}$ | $\{18,19,21,24,28,29,30,32,34,35,36\}$ | 15 | 0001001010100000 0110100010001110101 10 |
| 147 | $\{4,7,9,14\}$ | $\{17,18,19,20,21,23,24,26,28,30,34,36\}$ | 16 | 0001001010000100 1111101101010100010 10 |
| 148 | $\{4,7,9\}$ | $\{17,18,19,20,21,23,24,26,28,29,30,32,34,35,36\}$ | 18 | 0001001010000000 1111101101011101110 10 |
| 149 | $\{1,4,7,10\}$ | $\{20,21,23,24,26,27,29,30,32,34,36,37\}$ | 16 | 1001001001000000 0001101101011010011 00 |
| 150 | $\{4,7,10,11,12\}$ | $\{19,29,30,32,34,36,37\}$ | 12 | 0001001001110000 0010000000011010011 00 |
| 151 | $\{4,7,10,11\}$ | $\{18,19,21,24,27,29,30,32,34,36,37\}$ | 15 | 0001001001100000 0110100100110101011 00 |
| 152 | $\{4,7,10,12\}$ | $\{17,19,20,23,26,29,30,32,34,36,37\}$ | 15 | 0001001001010000 1011001001001101011 00 |
| 153 | $\{4,7,10\}$ | $\{17,18,19,20,21,23,24,26,27,29,30,32,34,36,37\}$ | 18 | 0001001001000000 1111101101011010011 00 |

#### *S2.27.3. A minimum six-state unital, separating family*

Define $\Omega_6 = \{\sigma_1, \ldots, \sigma_6\}$ in the following order:

$$(\sigma_1, \ldots, \sigma_6) = (s_3, s_{31}, s_{75}, s_{89}, s_{113}, s_{151}). \tag{S15}$$

Table S75 prints their complete supports. Table S76 gives all 37 profiles $(\sigma_1(i), \ldots, \sigma_6(i))$ and their active state sets. Every profile is nonzero and no two coincide, so $\Omega_6$ is unital and separating. Since five states supply at most $2^5 - 1 = 31 < 37$ nonzero binary profiles, no smaller unital, separating family can exist.

TABLE S75: Complete supports and state vectors of the minimum unital, separating family $\Omega_6$. The full state indices are those of Table S74.

| State | Full index | Complete support | Weight | 37-bit vector |
|---|---|---|---|---|
| $\sigma_1$ | $s_3$ | $\{2,5,8,12,17,19,22,25,28,30,31,32,33,35,37\}$ | 15 | 0100100100010000 1010010010011110101 |
| $\sigma_2$ | $s_{31}$ | $\{1,2,6,10,15,21,22,23,25,26,27,31,34,37\}$ | 14 | 1100010001000010 0000111011100010010001 |
| $\sigma_3$ | $s_{75}$ | $\{3,6,9,13,17,18,20,23,26,29,31,33,34,35,36\}$ | 15 | 0010010010001000 1101001001001011110 |
| $\sigma_4$ | $s_{89}$ | $\{3,7,8,16,17,18,19,20,22,23,24,27,28,29,32,35\}$ | 16 | 0010001100000001 1111011100111001001 00 |
| $\sigma_5$ | $s_{113}$ | $\{4,5,9,14,15,17,18,19,20,21,24,25,26,28\}$ | 14 | 0001100010000110 1111100111010000000 0 |
| $\sigma_6$ | $s_{151}$ | $\{4,7,10,11,18,19,21,24,27,29,30,32,34,36,37\}$ | 15 | 0001001001100000 0110100100101101010 11 |

TABLE S76: All 37 vertex profiles on $\Omega_6$, in two panels. Here $X_i^{(6)} = \{j : \sigma_j(i) = 1\}$, $W_6 = |X_i^{(6)}|$, and $W_{153}$ is the weight across the full state space. The six-bit rows are nonzero and pairwise distinct.

| $i$ | Ray | $X_i^{(6)}$ | Profile | $W_6$ | $W_{153}$ | $i$ | Ray | $X_i^{(6)}$ | Profile | $W_6$ | $W_{153}$ |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | $z$ | $\{2\}$ | 010000 | 1 | 54 | 20 | $B_{00}$ | $\{3,4,5\}$ | 001110 | 3 | 86 |
| 2 | $M_{00}$ | $\{1,2\}$ | 110000 | 2 | 51 | 21 | $B_{01}$ | $\{2,5,6\}$ | 010011 | 3 | 86 |
| 3 | $M_{01}$ | $\{3,4\}$ | 001100 | 2 | 51 | 22 | $B_{02}$ | $\{1,2,4\}$ | 110100 | 3 | 86 |
| 4 | $M_{02}$ | $\{5,6\}$ | 000011 | 2 | 51 | 23 | $B_{10}$ | $\{2,3,4\}$ | 011100 | 3 | 86 |
| 5 | $M_{10}$ | $\{1,5\}$ | 100010 | 2 | 51 | 24 | $B_{11}$ | $\{4,5,6\}$ | 000111 | 3 | 86 |
| 6 | $M_{11}$ | $\{2,3\}$ | 011000 | 2 | 51 | 25 | $B_{12}$ | $\{1,2,5\}$ | 110010 | 3 | 86 |
| 7 | $M_{12}$ | $\{4,6\}$ | 000101 | 2 | 51 | 26 | $B_{20}$ | $\{2,3,5\}$ | 011010 | 3 | 86 |
| 8 | $M_{20}$ | $\{1,4\}$ | 100100 | 2 | 51 | 27 | $B_{21}$ | $\{2,4,6\}$ | 010101 | 3 | 86 |
| 9 | $M_{21}$ | $\{3,5\}$ | 001010 | 2 | 51 | 28 | $B_{22}$ | $\{1,4,5\}$ | 100110 | 3 | 86 |
| 10 | $M_{22}$ | $\{2,6\}$ | 010001 | 2 | 51 | 29 | $C_{00}$ | $\{3,4,6\}$ | 001101 | 3 | 76 |
| 11 | $P_0$ | $\{6\}$ | 000001 | 1 | 16 | 30 | $C_{01}$ | $\{1,6\}$ | 100001 | 2 | 76 |
| 12 | $P_1$ | $\{1\}$ | 100000 | 1 | 16 | 31 | $C_{02}$ | $\{1,2,3\}$ | 111000 | 3 | 76 |
| 13 | $P_2$ | $\{3\}$ | 001000 | 1 | 16 | 32 | $C_{10}$ | $\{1,4,6\}$ | 100101 | 3 | 76 |
| 14 | $Q_0$ | $\{5\}$ | 000010 | 1 | 26 | 33 | $C_{11}$ | $\{1,3\}$ | 101000 | 2 | 76 |
| 15 | $Q_1$ | $\{2,5\}$ | 010010 | 2 | 26 | 34 | $C_{12}$ | $\{2,3,6\}$ | 011001 | 3 | 76 |
| 16 | $Q_2$ | $\{4\}$ | 000100 | 1 | 26 | 35 | $C_{20}$ | $\{1,3,4\}$ | 101100 | 3 | 76 |
| 17 | $A_0$ | $\{1,3,4,5\}$ | 101110 | 4 | 83 | 36 | $C_{21}$ | $\{3,6\}$ | 001001 | 2 | 76 |
| 18 | $A_1$ | $\{3,4,5,6\}$ | 001111 | 4 | 83 | 37 | $C_{22}$ | $\{1,2,6\}$ | 110001 | 3 | 76 |
| 19 | $A_2$ | $\{1,4,5,6\}$ | 100111 | 4 | 83 | | | | | | |

#### *S2.27.4. All full 153-state vertex profiles*

Table S77 prints the transpose of the complete state table: the full 153-bit profile of every vertex, in three consecutive blocks of 51 states. Thus it also enumerates the complete active state set $X_i = \{s_r : s_r(i) = 1\}$ for every vertex: a one in position $r$ means exactly $s_r \in X_i$. The row weights are 54 for $z$, 51 for $M$, 16 for $P$, 26 for $Q$, 83 for $A$, 86 for $B$, and 76 for $C$.

TABLE S77: Complete 153-bit vertex profiles, with consecutive 51-bit blocks. Every row is nonzero and all 37 rows are distinct.

| $i$ | Ray | $W_{153}$ | $s_1 \ldots s_{51}$ | $s_{52} \ldots s_{102}$ | $s_{103} \ldots s_{153}$ |
|---|---|---|---|---|---|
| 1 | $z$ | 54 | 100001100001100001100001100001111000011000010110000 | 110000110000101100001000011000011110000110000110000 | 110000111100001100001011000011000011000011000010000 |
| 2 | $M_{00}$ | 51 | 111111111111111111111111111111111111111111111111111 | 000000000000000000000000000000000000000000000000000 | 000000000000000000000000000000000000000000000000000 |
| 3 | $M_{01}$ | 51 | 000000000000000000000000000000000000000000000000000 | 111111111111111111111111111111111111111111111111111 | 000000000000000000000000000000000000000000000000000 |
| 4 | $M_{02}$ | 51 | 000000000000000000000000000000000000000000000000000 | 000000000000000000000000000000000000000000000000000 | 111111111111111111111111111111111111111111111111111 |
| 5 | $M_{10}$ | 51 | 111111111111111110000000000000000000000000000000000 | 111111111111110000000000000000000000000000000000000 | 111111111111111111100000000000000000000000000000000 |
| 6 | $M_{11}$ | 51 | 000000000000000011111111111111111111000000000000000 | 000000000000011111111111111111000000000000000000000 | 000000000000000000111111111111110000000000000000000 |
| 7 | $M_{12}$ | 51 | 000000000000000000000000000000000000011111111111111 | 000000000000000000000000000000011111111111111111111 | 000000000000000000000000000000000111111111111111111 |
| 8 | $M_{20}$ | 51 | 111110000000000011111100000000000000011111100000000 | 111111000000001111110000000000111111100000000000000 | 111111000000000000011000000000000111111000000000000 |
| 9 | $M_{21}$ | 51 | 000001111110000000000011111100000000000000011000000 | 000000111111000000001111100000000000001111110000000 | 000000111111100000000111111000000000000011111100000 |
| 10 | $M_{22}$ | 51 | 000000000011111100000000000111111110000000111111000 | 000000000011000000000011111100000000000000111111000 | 000000000000011111000000011111100000000000011111000 |
| 11 | $P_0$ | 16 | 000000000000000000000000000000000000000000000000000 | 000000000000000000000110000110000000000011000011000 | 000000000000000000000011000011000000000011000011000 |
| 12 | $P_1$ | 16 | 011000000000011000000000000000000000000011000001100 | 000000000000000000000000000000000000000000000000000 | 001100000000000011000000000000000000000011000000001010 |
| 13 | $P_2$ | 16 | 010100011000000000011000011000000000000000000000000 | 001100001100000011000101000000000000000000000000000 | 000000000000000000000000000000000000000000000000000 |
| 14 | $Q_0$ | 26 | 000000000000000000000000000000000000000000000000000 | 101010101010000000000000000000011001100101010000000 | 101010110011000000000000000000000010101010101000000 |
| 15 | $Q_1$ | 26 | 000001010101010100000001010101100110000000000000000 | 000000000000000000000000000000000000000000000000000 | 000000101010101010100010101010101000000000000000000 |
| 16 | $Q_2$ | 26 | 000000000000000010101000000101010101010010010100000 | 000000000000010101000000101010101010000000101010000 | 000000000000000000000000000000000000000000000000000 |
| 17 | $A_0$ | 83 | 011110011110011100111001111000011110011110100111100 | 111110011101001111000110000110000111100001100001100 | 111110000111001111010000110000110011110000110001100 |
| 18 | $A_1$ | 83 | 000110011110000110011100111100001111000011010000110 | 111110011101001111011110011100000111100111100111101 | 000011000011110000110100111100111100001100111100101 |
| 19 | $A_2$ | 83 | 001010000110011100001100001100001111001111010011111 | 000011000011010000011001010011110000111100111100111 | 001111000011110011110100111100111100001100111101111 |
| 20 | $B_{00}$ | 86 | 000000000000000000000000000000000000000000000000000 | 111111111111111111111001110011111111111100111100111 | 111111111111111111111100111001111111111001110011111 |
| 21 | $B_{01}$ | 86 | 100111111111100111111111111111111111111100111111100111 | 000000000000000000000000000000000000000000000000000 | 110011111111111001111111111111111100111111110101111 |
| 22 | $B_{02}$ | 86 | 101011100111111111100111100111111111111111111111111111 | 110011110011111100111010111111111111111111111111111 | 000000000000000000000000000000000000000000000000000000 |
| 23 | $B_{10}$ | 86 | 000000000000000011111111111111111111111111111111111111 | 000000000000011111100111001111111111100111100111111 | 000000000000000000011100111001111111111001110011111 |
| 24 | $B_{11}$ | 86 | 100111111111100110000000000000000000011001111110011 | 111111111111110000000000000000011111111111111111111 | 110011111111111001100000000000000110011111111101011 |
| 25 | $B_{12}$ | 86 | 101011100111111111100111100111111111100000000000000 | 110011110011111100111010111111100000000000000000000 | 111111111111111111111111111111111111111100000000000000 |
| 26 | $B_{20}$ | 86 | 000001111111111110000001111111111111000000111111111 | 000000111111100000010011100110000000110011110011111 | 000000111111111111110011001111001100000011001110011 |
| 27 | $B_{21}$ | 86 | 100110000001100111111100000011111111111001100110011 | 111111000000111111110000011111111111111000000111111 | 110011000000001100111100000011111111100110000010101 |
| 28 | $B_{22}$ | 86 | 101011100110000001100111100110000000011111111100000 | 110011110011001100111010100000011111111111111000000 | 111111111111110000001111111100000011111111111100000 |
| 29 | $C_{00}$ | 76 | 000000000000000000000000000000000000000000000000000 | 010101010101111111111111111111111001100110101011111111 | 010101001100111111111111111111111111010101010111111 |
| 30 | $C_{01}$ | 76 | 111110101010101011111110101010011001111111111111111 | 000000000000000000000000000000000000000000000000000 | 111111010101010101011101010101011111111111111111111 |
| 31 | $C_{02}$ | 76 | 111111111111111101010111111101010101010101011010101 | 111111111111110101011111101010101010111111101010100 | 000000000000000000000000000000000000000000000000000 |
| 32 | $C_{10}$ | 76 | 111111111111111100000000000000000000111111111111111 | 010101010101100000000000000000011001101010111111111 | 010101001100111111100000000000000101010101011111111 |
| 33 | $C_{11}$ | 76 | 111110101010101011111101010100110011000000000000000 | 111111111111111111111111111111110000000000000000000 | 111111010101010101110101010101010000000000000000000 |
| 34 | $C_{12}$ | 76 | 000000000000000001010111111101010101010101011010101 | 000000000000000101011111101010101010111111101010100 | 000000000000000000001111111111111111111111111111111 |
| 35 | $C_{20}$ | 76 | 111111111110000001111111111100000000111111110000000 | 010101010101001111111111100000000110011010101000000 | 010101001100110000011111111100000001010101010100000 |
| 36 | $C_{21}$ | 76 | 000000101010101010000000101010011001100000011111111 | 000000111111110000001111111111100000000111111111111 | 000000010101010101010001010101010100000011111111111 |
| 37 | $C_{22}$ | 76 | 111110000001111110101010000000101010101010100010101 | 111111000000110101010000001010101010100000010101 | 111111000000011111111100000011111111111100000011111 |

### *S2.27.5. The six-state double cover*

The fractional-coloring certificate uses a second six-state family,

$$\mathcal{R}_6 = \{s_2, s_{30}, s_{73}, s_{99}, s_{109}, s_{139}\}. \tag{S16}$$

Table S78 specifies these states in the branch parametrization. Each of the 37 vertices occurs in exactly two of their supports. Thus weights 1/2 give a fractional coloring of total weight three, and uniform weights 1/6 reproduce the maximally mixed probabilities 1/3 for every ray. The family $\Omega_6$ in Eq. (S15) supplies the separating certificate; $\mathcal{R}_6$ supplies the double-cover certificate.

TABLE S78: Explicit six-state double cover $\mathcal{R}_6$. Omitted $P, Q$ values are zero; completion values follow from the state equations.

| $j$ | State | $t$ | $\varepsilon$ | Active $P$ subscripts | Active $Q$ subscripts |
|---|---|---|---|---|---|
| 1 | $s_2$ | $(0,0,0)$ | 0 | $\{1,2\}$ | $\varnothing$ |
| 2 | $s_{30}$ | $(0,1,2)$ | 1 | $\varnothing$ | $\{1,2\}$ |
| 3 | $s_{73}$ | $(1,1,1)$ | 0 | $\{0,2\}$ | $\varnothing$ |
| 4 | $s_{99}$ | $(1,2,2)$ | 0 | $\{0\}$ | $\{2\}$ |
| 5 | $s_{109}$ | $(2,0,1)$ | 1 | $\varnothing$ | $\{0,1\}$ |
| 6 | $s_{139}$ | $(2,2,0)$ | 0 | $\{1\}$ | $\{0\}$ |

#### *S2.27.6. Order determination and exact verification*

The companion file `sources/hesse/hesse37-canonical-evaluations.txt` records the 24 numbered contexts and all 153 binary state vectors. The parametrization in this appendix defines their canonical order independently of any external program.

Extend the canonical states to the 76 events of $L(H_{37})$ as in Sec. V A. If $X_x = \{r : s_r(x) = 1\}$, then the full profiles satisfy

$$X_x \subseteq X_y \quad \Longleftrightarrow \quad x \leq y. \tag{S17}$$

The verifier checks all $76^2 = 5{,}776$ ordered event pairs against the incidence order in Eq. (S27). Whenever $x \not\leq y$, the companion file `sources/hesse/hesse37-order-witnesses.csv` gives the least canonical index $r$ with $s_r(x) = 1$ and $s_r(y) = 0$. Its event labels `a1`,...,`a37` denote atoms and `c1`,...,`c37` their complements; `0` and `1` denote the bounds. Every one of the 594 distinct nonorthogonal atom pairs also has a joint-one state. Thus the full family is order determining; the six-state family $\Omega_6$ supplies the minimum unital, separating vertex representation in Eq. (S26). The distinct minima for event separation and order determination are certified in Appendix S3.3.

The verifier `hesse37-verify.py` uses exact rational arithmetic in $\mathbb{Q}(\zeta)$, with no numerical tolerance or external solver. It checks distinct rays, all 666 Hermitian inner products, the 24 bases, all graph triangles, linearity, the chromatic number and core criticality, all 153 states, the minimum separating family, both profile tables, the maximum squared ray overlap, the six-state double cover, and all event-order comparisons. It also compares the reconstructed state vectors with the archived enumeration. The accompanying machine-readable data give the same numbering and context ordering as the figure and these appendices. From the extracted joint project directory, run

```
python verification/verify-integrated.py
```

### S2.28. Four-dimensional $H_{\mathrm{YO}}^{(4)}$ and $H_{65}$: states and profiles

This appendix provides the complete enumeration of all two-valued states on the four-dimensional examples $H_{\mathrm{YO}}^{(4)}$ and $H_{65}$: the 25 two-valued states of the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)}$ (Appendix S2.28.1), and the complete branch-by-branch parametrization of all $24 \times 2^{14} = 393{,}216$ two-valued states—together with the 48 canonical unital and separating states $s_{\alpha,\varepsilon}$ and the 65 vertex state-profiles—of the canonical linear hypergraph $H_{65}$ (Appendix S2.28.2).

#### *S2.28.1. Complete enumeration of the 25 two-valued states on $H_{\mathrm{YO}}^{(4)}$*

Table S79 lists all 25 two-valued states $s_1, \ldots, s_{25} : V_{26} \to \{0,1\}$ on $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$. States $s_1, \ldots, s_{24}$ have $s_r(26) = 0$ and correspond to the 24 two-valued states of the completed

Yu–Oh hypergraph $H_{\mathrm{YO}}^{+}$ in $\mathbb{R}^3$ (grouped into 12 pairs of states sharing the same support on the three coordinate contexts $\{1,2,3\}$, $\{4,5,6\}$, $\{7,8,9\}$), while $s_{25}$ is the unique apex state with support $\{26\}$. Table S80 displays the transposed $26 \times 25$ vertex-by-state binary matrix, listing for each vertex $i \in \{1,\dots,26\}$ its state weight $\sum_{r=1}^{25} s_r(i) \in \{1,3,8,13\}$, the set of states $A_i = \{s_r : s_r(i) = 1\}$, and its 25-bit profile $(s_1(i),\dots,s_{25}(i))$.

Inspection of Table S80 highlights three structural properties of the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$:

1. *Unitality and atom separation:* Every row of Table S80 has positive weight (1, 3, 8, or 13) and all 26 binary rows are distinct ($A_i \neq A_j$ for all $1 \le i < j \le 26$), so $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is nonempty, unital, and separating on atoms.

2. *Exclusion of simultaneous* 1*s on the non-orthogonal triad atoms* $\{10,11,12,13\}$*:* Although the four Yu–Oh triad rays $v_{10}, v_{11}, v_{12}, v_{13}$ are pairwise non-orthogonal ($v_i \cdot v_j = \pm 1 \neq 0$ for $10 \le i < j \le 13$), their active state sets in Table S80 are pairwise disjoint:

$$\begin{aligned} A_{10} &= \{s_{13}, s_{19}, s_{23}\}, \quad A_{11} = \{s_5, s_7, s_{15}\}, \\ A_{12} &= \{s_1, s_3, s_9\}, \qquad A_{13} = \{s_{11}, s_{17}, s_{21}\}. \end{aligned} \tag{S18}$$

   Consequently, $\sum_{i=10}^{13} s(i) \le 1$ holds for every $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$, and no two-valued state assigns $(1,1)$ to any of the $\binom{4}{2} = 6$ non-orthogonal pairs $\{10,11\}$, $\{10,12\}$, $\{10,13\}$, $\{11,12\}$, $\{11,13\}$, $\{12,13\}$.

3. *Failure of order-determinacy:* Even though $A_i \neq A_j$ for all $i \neq j$, the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is not order-determining on atoms or compound propositions[4]. In particular, each triad atom $t \in \{10,11,12,13\}$ (*weight* 3) is strictly dominated across all 25 states ($A_t \subsetneq A_w$, i.e., $s(t) \le s(w)$ for all $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$) by three non-orthogonal completion atoms $w \in \{14,\dots,25\}$ (*weight* 13):

$$\begin{aligned} A_{10} &\subsetneq A_{16},\ A_{20},\ A_{24}, \quad A_{11} \subsetneq A_{18},\ A_{21},\ A_{25}, \\ A_{12} &\subsetneq A_{14},\ A_{22},\ A_{23}, \quad A_{13} \subsetneq A_{15},\ A_{17},\ A_{19}, \end{aligned} \tag{S19}$$

   together with compound-event inclusions such as $A_{12} \subsetneq A_3 \,\dot\cup\, A_{14}$, $A_{11} \subsetneq A_3 \,\dot\cup\, A_{25}$, and their unions with the apex set $A_{26} = \{s_{25}\}$.

TABLE S79: Complete enumeration of the 25 two-valued states $s_1,\dots,s_{25} : V_{26} \to \{0,1\}$ on the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, showing for each state its support on the 13-ray Yu–Oh core $\{1,\dots,13\}$, its forced support on $\{14,\dots,26\}$, its total support size $|S_r|$, and its 26-bit binary vector $(s_r(1\dots13)\, s_r(14\dots25)\, s_r(26))$.

| State | Core support | Completion/apex support | $\lvert S_r\rvert$ | 26-bit state vector |
|---|---|---|---|---|
| $s_1$ | $\{1,4,8,12\}$ | $\{14,15,21,22,23\}$ | 9 | 1001000100010 110000011100 0 |
| $s_2$ | $\{1,4,8\}$ | $\{14,15,16,17,21,22,23,25\}$ | 11 | 1001000100000 111100011101 0 |
| $s_3$ | $\{1,5,7,12\}$ | $\{14,18,22,23,24\}$ | 9 | 1000101000010 100010001110 0 |
| $s_4$ | $\{1,5,7\}$ | $\{14,16,17,18,22,23,24,25\}$ | 11 | 1000101000000 101110001111 0 |
| $s_5$ | $\{1,5,9,11\}$ | $\{17,18,21,22,25\}$ | 9 | 1000100010100 000110011001 0 |
| $s_6$ | $\{1,5,9\}$ | $\{14,15,17,18,21,22,24,25\}$ | 11 | 1000100010000 110110011011 0 |
| $s_7$ | $\{1,6,8,11\}$ | $\{16,18,21,23,25\}$ | 9 | 1000010100100 001010010101 0 |
| $s_8$ | $\{1,6,8\}$ | $\{14,15,16,18,21,23,24,25\}$ | 11 | 1000010100000 111010010111 0 |
| $s_9$ | $\{2,4,7,12\}$ | $\{14,19,20,22,23\}$ | 9 | 0101001000010 100001101100 0 |
| $s_{10}$ | $\{2,4,7\}$ | $\{14,16,17,19,20,22,23,25\}$ | 11 | 0101001000000 101101101101 0 |
| $s_{11}$ | $\{2,4,9,13\}$ | $\{14,15,17,19,25\}$ | 9 | 0101000010001 110101000001 0 |
| $s_{12}$ | $\{2,4,9\}$ | $\{14,15,17,19,20,21,22,25\}$ | 11 | 0101000010000 110101111001 0 |
| $s_{13}$ | $\{2,6,7,10\}$ | $\{14,16,20,24,25\}$ | 9 | 0100011001000 101000100011 0 |
| $s_{14}$ | $\{2,6,7\}$ | $\{14,16,18,19,20,23,24,25\}$ | 11 | 0100011000000 101011100111 0 |
| $s_{15}$ | $\{2,6,9,11\}$ | $\{18,19,20,21,25\}$ | 9 | 0100010010100 000011110001 0 |

*Continued on the next page.*

TABLE S79: All 25 states of $H_{\mathrm{YO}}^{(4)}$ (continued).

| State | Core support | Completion/apex support | $\lvert S_r\rvert$ | 26-bit state vector |
|---|---|---|---|---|
| $s_{16}$ | $\{2,6,9\}$ | $\{14,15,18,19,20,21,24,25\}$ | 11 | 0100010010000 110011110011 0 |
| $s_{17}$ | $\{3,4,8,13\}$ | $\{15,16,17,19,23\}$ | 9 | 0011000100001 011101000100 0 |
| $s_{18}$ | $\{3,4,8\}$ | $\{15,16,17,19,20,21,22,23\}$ | 11 | 0011000100000 011101111100 0 |
| $s_{19}$ | $\{3,5,7,10\}$ | $\{16,17,20,22,24\}$ | 9 | 0010101001000 001100101010 0 |
| $s_{20}$ | $\{3,5,7\}$ | $\{16,17,18,19,20,22,23,24\}$ | 11 | 0010101000000 001111101110 0 |
| $s_{21}$ | $\{3,5,9,13\}$ | $\{15,17,18,19,24\}$ | 9 | 0010100010001 010111000010 0 |
| $s_{22}$ | $\{3,5,9\}$ | $\{15,17,18,19,20,21,22,24\}$ | 11 | 0010100010000 010111111010 0 |
| $s_{23}$ | $\{3,6,8,10\}$ | $\{15,16,20,21,24\}$ | 9 | 0010010101000 011000110010 0 |
| $s_{24}$ | $\{3,6,8\}$ | $\{15,16,18,19,20,21,23,24\}$ | 11 | 0010010100000 011011110110 0 |
| $s_{25}$ | $\emptyset$ | $\{26\}$ | 1 | 0000000000000 000000000000 1 |

TABLE S80: Vertex state profiles across the 25 two-valued states $s_1,\dots,s_{25}$ of $H_{\mathrm{YO}}^{(4)}$, showing for each vertex $i \in \{1,\dots,26\}$ its state weight $\sum_{r=1}^{25} s_r(i)$, the active state indices $\{r : s_r(i)=1\}$, and the 25-bit profile $(s_{1\dots12}(i)\, s_{13\dots24}(i)\, s_{25}(i))$. All 26 rows are nonzero and pairwise distinct.

| Vertex $i$ | Weight $\sum_r s_r(i)$ | Active states $\{r \in \{1,\dots,25\} : s_r(i)=1\}$ | 25-bit profile $(s_{1\dots12}(i)\, s_{13\dots24}(i)\, s_{25}(i))$ |
|---|---|---|---|
| 1 | 8 | $\{1,2,3,4,5,6,7,8\}$ | 111111110000 000000000000 0 |
| 2 | 8 | $\{9,10,11,12,13,14,15,16\}$ | 000000001111 111100000000 0 |
| 3 | 8 | $\{17,18,19,20,21,22,23,24\}$ | 000000000000 0000111111111 0 |
| 4 | 8 | $\{1,2,9,10,11,12,17,18\}$ | 110000001111 000011000000 0 |
| 5 | 8 | $\{3,4,5,6,19,20,21,22\}$ | 001111000000 000000111100 0 |
| 6 | 8 | $\{7,8,13,14,15,16,23,24\}$ | 000000110000 111100000011 0 |
| 7 | 8 | $\{3,4,9,10,13,14,19,20\}$ | 001100001100 110000110000 0 |
| 8 | 8 | $\{1,2,7,8,17,18,23,24\}$ | 110000110000 000011000011 0 |
| 9 | 8 | $\{5,6,11,12,15,16,21,22\}$ | 000011000011 001100001100 0 |
| 10 | 3 | $\{13,19,23\}$ | 000000000000 100000100010 0 |
| 11 | 3 | $\{5,7,15\}$ | 000010100000 001000000000 0 |
| 12 | 3 | $\{1,3,9\}$ | 101000001000 000000000000 0 |
| 13 | 3 | $\{11,17,21\}$ | 000000000010 000010001000 0 |
| 14 | 13 | $\{1,2,3,4,6,8,9,10,11,12,13,14,16\}$ | 111101011111 110100000000 0 |
| 15 | 13 | $\{1,2,6,8,11,12,16,17,18,21,22,23,24\}$ | 110001010011 000111001111 0 |
| 16 | 13 | $\{2,4,7,8,10,13,14,17,18,19,20,23,24\}$ | 010100110100 110011110011 0 |
| 17 | 13 | $\{2,4,5,6,10,11,12,17,18,19,20,21,22\}$ | 010111000111 000011111100 0 |
| 18 | 13 | $\{3,4,5,6,7,8,14,15,16,20,21,22,24\}$ | 001111110000 011100011101 0 |
| 19 | 13 | $\{9,10,11,12,14,15,16,17,18,20,21,22,24\}$ | 000000001111 011111011101 0 |
| 20 | 13 | $\{9,10,12,13,14,15,16,18,19,20,22,23,24\}$ | 000000001101 111101110111 0 |
| 21 | 13 | $\{1,2,5,6,7,8,12,15,16,18,22,23,24\}$ | 110011110001 001101000111 0 |
| 22 | 13 | $\{1,2,3,4,5,6,9,10,12,18,19,20,22\}$ | 111111001101 000001110100 0 |
| 23 | 13 | $\{1,2,3,4,7,8,9,10,14,17,18,20,24\}$ | 111100111100 010011010001 0 |
| 24 | 13 | $\{3,4,6,8,13,14,16,19,20,21,22,23,24\}$ | 001101010000 110100111111 0 |
| 25 | 13 | $\{2,4,5,6,7,8,10,11,12,13,14,15,16\}$ | 010111110111 111100000000 0 |
| 26 | 1 | $\{25\}$ | 000000000000 000000000000 1 |

### *S2.28.2. Complete parametrization of all* 393,216 *two-valued states and the* 48 *canonical separating states on* $H_{65}$

Every two-valued state $s \in \mathcal{S}(H_{65})$ on the canonical linear hypergraph $H_{65} = (V_{65}, E_{29})$ is uniquely indexed by a pair $(\alpha, \boldsymbol{\varepsilon})$ consisting of:

1. a branch index $\alpha \in \{1,\dots,24\}$ selecting one of the 24 four-element core state supports $S_\alpha \subset V_{\mathrm{core}} = \{1,\dots,17\}$; and
2. a 14-bit binary vector $\boldsymbol{\varepsilon} = (\varepsilon_m)_{m\in\mathcal{F}_\alpha} \in \{0,1\}^{14}$ indexed by the 14 *free completion contexts*

$$\mathcal{F}_\alpha = \big\{m \in \{1,\dots,24\} : S_\alpha \cap \{u_m, v_m\} = \emptyset\big\},$$

whose complement is the set of 10 *hit completion contexts* $\mathcal{H}_\alpha = \{1,\dots,24\} \setminus \mathcal{F}_\alpha$.

Explicitly, the state $s_{\alpha,\boldsymbol{\varepsilon}} : V_{65} \to \{0,1\}$ takes values

$$\begin{aligned} s_{\alpha,\boldsymbol{\varepsilon}}(i) &= \mathbf{1}_{i\in S_\alpha} \qquad (1 \le i \le 17), \\ \big(s_{\alpha,\boldsymbol{\varepsilon}}(2m+16),\, s_{\alpha,\boldsymbol{\varepsilon}}(2m+17)\big) &= \begin{cases} (0,0), & m \in \mathcal{H}_\alpha, \\ (1-\varepsilon_m,\, \varepsilon_m), & m \in \mathcal{F}_\alpha. \end{cases} \end{aligned} \tag{S20}$$

Because $|\mathcal{F}_\alpha| = 14$ for every $\alpha \in \{1,\dots,24\}$, each branch $\alpha$ yields $2^{14} = 16{,}384$ states $s_{\alpha,\boldsymbol{\varepsilon}}$, giving $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$ two-valued states in total. Setting all 14 free bits equal to $\varepsilon \in \{0,1\}$ ($\varepsilon_m = \varepsilon$ for all $m \in \mathcal{F}_\alpha$) yields the 48 *canonical states* $s_{\alpha,0}, s_{\alpha,1}$ of Eq. (S29), which have support size $|S_\alpha| + 14 = 18$:

$$\begin{aligned} \operatorname{supp}(s_{\alpha,0}) &= S_\alpha \cup \big\{2m+16 : m \in \mathcal{F}_\alpha\big\}, \\ \operatorname{supp}(s_{\alpha,1}) &= S_\alpha \cup \big\{2m+17 : m \in \mathcal{F}_\alpha\big\}. \end{aligned} \tag{S21}$$

Table S81 enumerates all 24 branches $\alpha = 1,\dots,24$, specifying $S_\alpha$, the 10 hit context indices $\mathcal{H}_\alpha \subset \{1,\dots,24\}$, the 14 free context indices $\mathcal{F}_\alpha \subset \{1,\dots,24\}$, and the 24-bit free-context indicator string. Table S82 lists the resulting state profiles and weights for all 65 vertices of $H_{65}$ across both the 48 canonical states $\{s_{\alpha,\varepsilon}\}$ and the full space of 393,216 states: across the 48 canonical states $(s_{1,0}, s_{1,1}, \dots, s_{24,0}, s_{24,1})$, each '1' in row $i$ of Table S82 expands to '(1,1)' for core vertices $i \in \{1,\dots,17\}$, to '(1,0)' for first completion vertices $w_{m,1} = 2m+16$, and to '(0,1)' for second completion vertices $w_{m,2} = 2m+17$ (and each '0' expands to '(0,0)'), making all 65 canonical 48-bit vertex profiles nonzero and pairwise distinct by inspection.

TABLE S81: Complete branch-by-branch parametrization of the $24 \times 2^{14} = 393{,}216$ two-valued states $s_{\alpha,\boldsymbol{\varepsilon}}$ and the 48 canonical states $s_{\alpha,0}$,$s_{\alpha,1}$ on $H_{65}$ via Eqs. (S20)–(S21). For each $\alpha \in \{1,\dots,24\}$, the table lists the 4-element core support $S_\alpha$, the 10 hit completion context indices $m \in \mathcal{H}_\alpha$ where $(s(2m+16),s(2m+17)) = (0,0)$ is forced on $E_m$, the 14 free completion context indices $m \in \mathcal{F}_\alpha$ where $(s(2m+16),s(2m+17)) = (1-\varepsilon_m,\varepsilon_m)$ with $\varepsilon_m \in \{0,1\}$ on $E_m$, and the 24-bit indicator string $(\mathbf{1}_{1\in\mathcal{F}_\alpha} \dots \mathbf{1}_{12\in\mathcal{F}_\alpha}\, \mathbf{1}_{13\in\mathcal{F}_\alpha} \dots \mathbf{1}_{24\in\mathcal{F}_\alpha})$.

| Branch | Core support $S_\alpha$ | Hit contexts $\mathcal{H}_\alpha$ | Free contexts $\mathcal{F}_\alpha$ | Free indicator $m = 1\dots24$ |
|---|---|---|---|---|
| $S_1$ | {1,5,9,16} | {3,7,9,10,11,13,19,21,22,23} | {1,2,4,5,6,8,12,14,15,16,17,18,20,24} | 110111010001 011111010001 |
| $S_2$ | {1,5,10,14} | {3,5,6,7,11,13,17,18,19,23} | {1,2,4,8,9,10,12,14,15,16,20,21,22,24} | 110100011101 011100011101 |
| $S_3$ | {1,6,9,16} | {3,7,9,10,11,15,17,21,22,24} | {1,2,4,5,6,8,12,13,14,16,18,19,20,23} | 110111010001 110101110010 |
| $S_4$ | {1,6,11,12} | {1,2,3,7,11,13,14,15,17,24} | {4,5,6,8,9,10,12,16,18,19,20,21,22,23} | 000111011101 000101111110 |
| $S_5$ | {1,7,10,14} | {3,5,6,7,11,16,17,18,20,21} | {1,2,4,8,9,10,12,13,14,15,19,22,23,24} | 110100011101 111000100111 |
| $S_6$ | {1,7,11,12} | {1,2,3,7,11,13,14,16,20,21} | {4,5,6,8,9,10,12,15,17,18,19,22,23,24} | 000111011101 001011100111 |
| $S_7$ | {2,5,9,16} | {1,5,9,10,12,13,19,21,22,23} | {2,3,4,6,7,8,11,14,15,16,17,18,20,24} | 011101110010 011111010001 |
| $S_8$ | {2,5,11,13} | {1,3,4,5,12,13,15,16,19,23} | {2,6,7,8,9,10,11,14,17,18,20,21,22,24} | 010001111110 010011011101 |
| $S_9$ | {2,6,9,16} | {1,5,9,10,12,15,17,21,22,24} | {2,3,4,6,7,8,11,13,14,16,18,19,20,23} | 011101110010 110101110010 |
| $S_{10}$ | {2,6,10,15} | {1,5,7,8,12,15,17,19,20,24} | {2,3,4,6,9,10,11,13,14,16,18,21,22,23} | 011101001110 110101001110 |
| $S_{11}$ | {2,8,10,15} | {1,5,7,8,12,14,18,19,20,22} | {2,3,4,6,9,10,11,13,15,16,17,21,23,24} | 011101001110 101110001011 |
| $S_{12}$ | {2,8,11,13} | {1,3,4,5,12,14,15,16,18,22} | {2,6,7,8,9,10,11,13,17,19,20,21,23,24} | 010001111110 100010111011 |
| $S_{13}$ | {3,5,10,14} | {2,5,6,8,9,13,17,18,19,23} | {1,3,4,7,10,11,12,14,15,16,20,21,22,24} | 101100100111 011100011101 |
| $S_{14}$ | {3,5,11,13} | {2,3,4,8,9,13,15,16,19,23} | {1,5,6,7,10,11,12,14,17,18,20,21,22,24} | 100011100111 010011011101 |
| $S_{15}$ | {3,7,9,17} | {2,8,9,11,12,16,20,21,23,24} | {1,3,4,5,6,7,10,13,14,15,17,18,19,22} | 101111100100 111011100100 |
| $S_{16}$ | {3,7,10,14} | {2,5,6,8,9,16,17,18,20,21} | {1,3,4,7,10,11,12,13,14,15,19,22,23,24} | 101100100111 111000100111 |
| $S_{17}$ | {3,8,9,17} | {2,8,9,11,12,14,18,22,23,24} | {1,3,4,5,6,7,10,13,15,16,17,19,20,21} | 101111100100 101110111000 |
| $S_{18}$ | {3,8,11,13} | {2,3,4,8,9,14,15,16,18,22} | {1,5,6,7,10,11,12,13,17,19,20,21,23,24} | 100011100111 100010111011 |
| $S_{19}$ | {4,6,10,15} | {4,6,7,8,10,15,17,19,20,24} | {1,2,3,5,9,11,12,13,14,16,18,21,22,23} | 111010001011 110101001110 |
| $S_{20}$ | {4,6,11,12} | {1,2,4,6,10,13,14,15,17,24} | {3,5,7,8,9,11,12,16,18,19,20,21,22,23} | 001010111011 000101111110 |
| $S_{21}$ | {4,7,9,17} | {4,6,10,11,12,16,20,21,23,24} | {1,2,3,5,7,8,9,13,14,15,17,18,19,22} | 111010111000 111011100100 |
| $S_{22}$ | {4,7,11,12} | {1,2,4,6,10,13,14,16,20,21} | {3,5,7,8,9,11,12,15,17,18,19,22,23,24} | 001010111011 001011100111 |
| $S_{23}$ | {4,8,9,17} | {4,6,10,11,12,14,18,22,23,24} | {1,2,3,5,7,8,9,13,15,16,17,19,20,21} | 111010111000 101110111000 |
| $S_{24}$ | {4,8,10,15} | {4,6,7,8,10,14,18,19,20,22} | {1,2,3,5,9,11,12,13,15,16,17,21,23,24} | 111010001011 101110001011 |

TABLE S82: Vertex state profiles and weights for all 65 vertices of $H_{65}$, showing for each core vertex $i \in \{1,\dots,17\}$ and each completion pair $(w_{m,1}, w_{m,2}) = (2m+16, 2m+17)$ $(m = 1,\dots,24)$ the number of active branches $W_{24}$, the weight $W_{48} = \sum_{\alpha,\varepsilon} s_{\alpha,\varepsilon}(i)$ across the 48 canonical states, the weight $W_{393,216} = \sum_{s\in\mathcal{S}(H_{65})} s(i)$ across all 393,216 states, the active branch indices $\alpha \in \{1,\dots,24\}$ (equal to $Z(e_m)$ for $(w_{m,1}, w_{m,2})$), and the 24-bit branch profile across $S_1,\dots,S_{24}$ (in which each `1` expands to `11` for core vertices $i$, to `10` for $w_{m,1}$, and to `01` for $w_{m,2}$ across the 48 canonical states).

| Vertex / pair | $W_{24}$ | $W_{48}$ | $W_{393,216}$ | Active branches $\alpha$ | 24-bit branch profile |
|---|---|---|---|---|---|
| Core 1 | 6 | 12 | 98,304 | $\{1,2,3,4,5,6\}$ | 11111100 00000000 00000000 |
| Core 2 | 6 | 12 | 98,304 | $\{7,8,9,10,11,12\}$ | 00000011 11110000 00000000 |
| Core 3 | 6 | 12 | 98,304 | $\{13,14,15,16,17,18\}$ | 00000000 00001111 11000000 |
| Core 4 | 6 | 12 | 98,304 | $\{19,20,21,22,23,24\}$ | 00000000 00000000 00111111 |
| Core 5 | 6 | 12 | 98,304 | $\{1,2,7,8,13,14\}$ | 11000011 00001100 00000000 |
| Core 6 | 6 | 12 | 98,304 | $\{3,4,9,10,19,20\}$ | 00110000 11000000 00110000 |
| Core 7 | 6 | 12 | 98,304 | $\{5,6,15,16,21,22\}$ | 00001100 00000011 00001100 |
| Core 8 | 6 | 12 | 98,304 | $\{11,12,17,18,23,24\}$ | 00000000 00110000 11000011 |
| Core 9 | 8 | 16 | 131,072 | $\{1,3,7,9,15,17,21,23\}$ | 10100010 10000010 10001010 |
| Core 10 | 8 | 16 | 131,072 | $\{2,5,10,11,13,16,19,24\}$ | 01001000 01101001 00100001 |
| Core 11 | 8 | 16 | 131,072 | $\{4,6,8,12,14,18,20,22\}$ | 00010101 00010100 01010100 |
| Core 12 | 4 | 8 | 65,536 | $\{4,6,20,22\}$ | 00010100 00000000 00010100 |
| Core 13 | 4 | 8 | 65,536 | $\{8,12,14,18\}$ | 00000001 00010100 01000000 |
| Core 14 | 4 | 8 | 65,536 | $\{2,5,13,16\}$ | 01001000 00001001 00000000 |
| Core 15 | 4 | 8 | 65,536 | $\{10,11,19,24\}$ | 00000000 01100000 00100001 |
| Core 16 | 4 | 8 | 65,536 | $\{1,3,7,9\}$ | 10100010 10000000 00000000 |
| Core 17 | 4 | 8 | 65,536 | $\{15,17,21,23\}$ | 00000000 00000010 10001010 |
| $E_1$ : (18, 19) | 14 | 14 | 114,688 | $\{1,2,3,5,13,14,15,16,17,18,19,21,23,24\}$ | 11101000 00001111 11101011 |
| $E_2$ : (20, 21) | 14 | 14 | 114,688 | $\{1,2,3,5,7,8,9,10,11,12,19,21,23,24\}$ | 11101011 11110000 00101011 |
| $E_3$ : (22, 23) | 14 | 14 | 114,688 | $\{7,9,10,11,13,15,16,17,19,20,21,22,23,24\}$ | 00000010 11101011 10111111 |
| $E_4$ : (24, 25) | 14 | 14 | 114,688 | $\{1,2,3,4,5,6,7,9,10,11,13,15,16,17\}$ | 11111110 11101011 10000000 |
| $E_5$ : (26, 27) | 14 | 14 | 114,688 | $\{1,3,4,6,14,15,17,18,19,20,21,22,23,24\}$ | 10110100 00000110 11111111 |
| $E_6$ : (28, 29) | 14 | 14 | 114,688 | $\{1,3,4,6,7,8,9,10,11,12,14,15,17,18\}$ | 10110111 11110110 11000000 |
| $E_7$ : (30, 31) | 14 | 14 | 114,688 | $\{7,8,9,12,13,14,15,16,17,18,20,21,22,23\}$ | 00000011 10011111 11011110 |
| $E_8$ : (32, 33) | 14 | 14 | 114,688 | $\{1,2,3,4,5,6,7,8,9,12,20,21,22,23\}$ | 11111111 10010000 00011110 |
| $E_9$ : (34, 35) | 14 | 14 | 114,688 | $\{2,4,5,6,8,10,11,12,19,20,21,22,23,24\}$ | 01011101 01110000 00111111 |
| $E_{10}$ : (36, 37) | 14 | 14 | 114,688 | $\{2,4,5,6,8,10,11,12,13,14,15,16,17,18\}$ | 01011101 01111111 11000000 |
| $E_{11}$ : (38, 39) | 14 | 14 | 114,688 | $\{7,8,9,10,11,12,13,14,16,18,19,20,22,24\}$ | 00000011 11111101 01110101 |
| $E_{12}$ : (40, 41) | 14 | 14 | 114,688 | $\{1,2,3,4,5,6,13,14,16,18,19,20,22,24\}$ | 11111100 00001101 01110101 |
| $E_{13}$ : (42, 43) | 14 | 14 | 114,688 | $\{3,5,9,10,11,12,15,16,17,18,19,21,23,24\}$ | 00101000 11110011 11101011 |
| $E_{14}$ : (44, 45) | 14 | 14 | 114,688 | $\{1,2,3,5,7,8,9,10,13,14,15,16,19,21\}$ | 11101011 11001111 00101000 |
| $E_{15}$ : (46, 47) | 14 | 14 | 114,688 | $\{1,2,5,6,7,11,13,15,16,17,21,22,23,24\}$ | 11001110 00101011 10001111 |
| $E_{16}$ : (48, 49) | 14 | 14 | 114,688 | $\{1,2,3,4,7,9,10,11,13,17,19,20,23,24\}$ | 11110010 11101000 10110011 |
| $E_{17}$ : (50, 51) | 14 | 14 | 114,688 | $\{1,6,7,8,11,12,14,15,17,18,21,22,23,24\}$ | 10000111 00110110 11001111 |
| $E_{18}$ : (52, 53) | 14 | 14 | 114,688 | $\{1,3,4,6,7,8,9,10,14,15,19,20,21,22\}$ | 10110111 11000110 00111100 |
| $E_{19}$ : (54, 55) | 14 | 14 | 114,688 | $\{3,4,5,6,9,12,15,16,17,18,20,21,22,23\}$ | 00111100 10010011 11011110 |
| $E_{20}$ : (56, 57) | 14 | 14 | 114,688 | $\{1,2,3,4,7,8,9,12,13,14,17,18,20,23\}$ | 11110011 10011100 11010010 |
| $E_{21}$ : (58, 59) | 14 | 14 | 114,688 | $\{2,4,8,10,11,12,13,14,17,18,19,20,23,24\}$ | 01010001 01111100 11110011 |
| $E_{22}$ : (60, 61) | 14 | 14 | 114,688 | $\{2,4,5,6,8,10,13,14,15,16,19,20,21,22\}$ | 01011101 01001111 00111100 |
| $E_{23}$ : (62, 63) | 14 | 14 | 114,688 | $\{3,4,5,6,9,10,11,12,16,18,19,20,22,24\}$ | 00111100 11110001 01110101 |
| $E_{24}$ : (64, 65) | 14 | 14 | 114,688 | $\{1,2,5,6,7,8,11,12,13,14,16,18,22,24\}$ | 11001111 00111101 01000101 |

### S2.29. Parent-state and deletion certificates

The 49-ray, 87-ray, 40-ray, and 72-ray parents have no context states. The two smaller 68-ray parents are treated in Appendix S2.30.1. The first and last two have the independently verified complete bases printed in Appendix S1.16; the 87-ray parent contains the known 69-ray KS lineage. The 70-ray single-apex lift of the completed Cabello parent instead has the unique state supported at its apex.

For the real Yu–Oh parent, the intermediate 33 rays admit 192 assignments to the twenty bases alone and none after its sixteen additional pair exclusions. The sixteen missing pairs are uniquely completed by rays 34–49. The x-axis-plane extension by rays 26–29 alone has 72

basis assignments; all 24 original states extend. Including its eight additional pair exclusions leaves eight extensions, all with $s(2) = 1$. The z-axis-plane case is identical, with $s(5) = 1$. The two added bases on either axis each exclude twelve of the sixteen original axis-zero states when their further pair exclusions are included; eight original states are excluded by both, leaving none. Thus the axis forcing uses the complete orthogonality constraints, not the added bases alone.

TABLE S83. Context-state counts after deleting each single context of the real 49-ray parent. Every count is positive. Context numbering is Table S21. Deleting a completion context 21–36 leaves its fresh ray in no context; that ray is counted as an unconstrained free vertex, which doubles the count. If the orthogonalities of the deleted context are kept as geometric exclusions, deletions 21–36 leave no state, while deletions 1–20 leave 1, 2, or 4 states.

| Deleted context | States | Deleted context | States | Deleted context | States |
|---|---|---|---|---|---|
| 1 | 12 | 13 | 2 | 25 | 2 |
| 2 | 4 | 14 | 4 | 26 | 2 |
| 3 | 4 | 15 | 4 | 27 | 2 |
| 4 | 32 | 16 | 16 | 28 | 2 |
| 5 | 2 | 17 | 4 | 29 | 2 |
| 6 | 2 | 18 | 4 | 30 | 2 |
| 7 | 4 | 19 | 4 | 31 | 2 |
| 8 | 4 | 20 | 4 | 32 | 2 |
| 9 | 12 | 21 | 2 | 33 | 2 |
| 10 | 4 | 22 | 2 | 34 | 2 |
| 11 | 4 | 23 | 2 | 35 | 2 |
| 12 | 2 | 24 | 2 | 36 | 2 |

All complete states of $H_{65}$ are additionally supplied as `data/h65-all-states.csv.gz`. Each row gives its branch index, the fourteen-bit choice index within that branch, and its 65-bit state vector in vertex order. There are exactly $24 \times 2^{14} = 393{,}216$ data rows. This supplements the complete parametrization and the printed 48-state family; it does not assert that 48 is minimal. The 153 Hesse states, their event-order witnesses, and all original printed profiles remain in the accompanying Hesse data.

## S2.30. States of the Cabello restrictions and completions

### *S2.30.1. States of the vertex-deleted 45-ray completion*

The nine-context $C_{18,9}$, its eighteen-context faithful completion, and both 68-ray parents have no context states by their retained parity proofs. For the 45-ray construction, use exactly the local labels in Tables S31 and S32. Table S84 lists all twelve admissible core supports, and Table S85 lists the fourteen bridge pairs with their fresh labels and zero-profiles. For a support $S_\alpha$, let $F_\alpha$ be the bridge indices missed by $S_\alpha$. On every hit bridge both fresh values are zero; on each missed bridge independently choose exactly one of its two fresh vertices. This enumerates every context state once.

TABLE S84: All twelve core supports; $m = |F_\alpha|$. The final column specifies extendibility under all geometric pair exclusions.

| $\alpha$ | $S_\alpha$ | $m$ | Geometric extension |
|---|---|---|---|
| 1 | $\{1, 6, 11, 12\}$ | 8 | yes |
| 2 | $\{1, 10, 14, 18\}$ | 8 | yes |
| 3 | $\{3, 7, 10, 14\}$ | 8 | yes |
| 4 | $\{3, 8, 11, 13\}$ | 8 | yes |
| 5 | $\{3, 13, 14, 17\}$ | 8 | no |
| 6 | $\{2, 6, 10, 15\}$ | 8 | yes |
| 7 | $\{6, 12, 15, 16\}$ | 8 | no |
| 8 | $\{8, 10, 15, 18\}$ | 8 | yes |
| 9 | $\{9, 16, 18, 19\}$ | 9 | no |
| 10 | $\{9, 17, 18, 19\}$ | 9 | no |
| 11 | $\{2, 11, 13, 19\}$ | 8 | yes |
| 12 | $\{7, 11, 12, 19\}$ | 8 | yes |

TABLE S85: The fourteen bridge pairs, fresh labels, and zero-profiles; each $Z_m$ is the set of missed-bridge branch indices.

| $m$ | Bridge | Fresh pair | $Z_m$ |
|---|---|---|---|
| 1 | $\{1, 2\}$ | $\{21, 22\}$ | $\{3, 4, 5, 7, 8, 9, 10, 12\}$ |
| 2 | $\{1, 13\}$ | $\{23, 24\}$ | $\{3, 6, 7, 8, 9, 10, 12\}$ |
| 3 | $\{1, 17\}$ | $\{25, 26\}$ | $\{3, 4, 6, 7, 8, 9, 11, 12\}$ |
| 4 | $\{2, 14\}$ | $\{27, 28\}$ | $\{1, 4, 7, 8, 9, 10, 12\}$ |
| 5 | $\{2, 17\}$ | $\{29, 30\}$ | $\{1, 2, 3, 4, 7, 8, 9, 12\}$ |
| 6 | $\{3, 16\}$ | $\{31, 32\}$ | $\{1, 2, 6, 8, 10, 11, 12\}$ |
| 7 | $\{6, 17\}$ | $\{33, 34\}$ | $\{2, 3, 4, 8, 9, 11, 12\}$ |
| 8 | $\{7, 8\}$ | $\{35, 36\}$ | $\{1, 2, 5, 6, 7, 9, 10, 11\}$ |
| 9 | $\{7, 15\}$ | $\{37, 38\}$ | $\{1, 2, 4, 5, 9, 10, 11\}$ |
| 10 | $\{7, 16\}$ | $\{39, 40\}$ | $\{1, 2, 4, 5, 6, 8, 10, 11\}$ |
| 11 | $\{8, 12\}$ | $\{41, 42\}$ | $\{2, 3, 5, 6, 9, 10, 11\}$ |
| 12 | $\{8, 16\}$ | $\{43, 44\}$ | $\{1, 2, 3, 5, 6, 10, 11, 12\}$ |
| 13 | $\{10, 19\}$ | $\{45, 46\}$ | $\{1, 4, 5, 7\}$ |
| 14 | $\{11, 18\}$ | $\{47, 48\}$ | $\{3, 5, 6, 7\}$ |

Taking the first fresh vertex on every missed bridge gives $s_{\alpha,0}$, and taking the second gives $s_{\alpha,1}$. Table S86 explicitly enumerates all 24 canonical context states. Their profiles are nonzero and pairwise distinct on all 45 vertices. The twelve branch supports and fourteen distinct nonempty zero-profiles verify this by Lemma 21(3).

TABLE S86: All 24 canonical context-state supports of the 45-ray example. Some fail geometric pair exclusions.

| $\alpha$ | $\varepsilon$ | Support |
|---|---|---|
| 1 | 0 | $\{1,6,11,12,27,29,31,35,37,39,43,45\}$ |
| 1 | 1 | $\{1,6,11,12,28,30,32,36,38,40,44,46\}$ |
| 2 | 0 | $\{1,10,14,18,29,31,33,35,37,39,41,43\}$ |
| 2 | 1 | $\{1,10,14,18,30,32,34,36,38,40,42,44\}$ |
| 3 | 0 | $\{3,7,10,14,21,23,25,29,33,41,43,47\}$ |
| 3 | 1 | $\{3,7,10,14,22,24,26,30,34,42,44,48\}$ |
| 4 | 0 | $\{3,8,11,13,21,25,27,29,33,37,39,45\}$ |
| 4 | 1 | $\{3,8,11,13,22,26,28,30,34,38,40,46\}$ |
| 5 | 0 | $\{3,13,14,17,21,35,37,39,41,43,45,47\}$ |
| 5 | 1 | $\{3,13,14,17,22,36,38,40,42,44,46,48\}$ |
| 6 | 0 | $\{2,6,10,15,23,25,31,35,39,41,43,47\}$ |
| 6 | 1 | $\{2,6,10,15,24,26,32,36,40,42,44,48\}$ |
| 7 | 0 | $\{6,12,15,16,21,23,25,27,29,35,45,47\}$ |
| 7 | 1 | $\{6,12,15,16,22,24,26,28,30,36,46,48\}$ |
| 8 | 0 | $\{8,10,15,18,21,23,25,27,29,31,33,39\}$ |
| 8 | 1 | $\{8,10,15,18,22,24,26,28,30,32,34,40\}$ |
| 9 | 0 | $\{9,16,18,19,21,23,25,27,29,33,35,37,41\}$ |
| 9 | 1 | $\{9,16,18,19,22,24,26,28,30,34,36,38,42\}$ |
| 10 | 0 | $\{9,17,18,19,21,23,27,31,35,37,39,41,43\}$ |
| 10 | 1 | $\{9,17,18,19,22,24,28,32,36,38,40,42,44\}$ |
| 11 | 0 | $\{2,11,13,19,25,31,33,35,37,39,41,43\}$ |
| 11 | 1 | $\{2,11,13,19,26,32,34,36,38,40,42,44\}$ |
| 12 | 0 | $\{7,11,12,19,21,23,25,27,29,31,33,43\}$ |
| 12 | 1 | $\{7,11,12,19,22,24,26,28,30,32,34,44\}$ |

Ten branches miss eight pairs and two miss nine. Therefore

$$|\mathcal{S}_{\mathrm{ctx}}| = 10\,2^8 + 2\,2^9 = 3{,}584, \qquad |\mathcal{S}_{\mathrm{geom}}| = 8\,2^8 = 2048. \tag{S22}$$

The geometric branches are $1,2,3,4,6,8,11,12$; arbitrary fresh choices within each of these branches remain geometrically compatible. The other branches are excluded by the four forced pairs. Both full enumerations are supplied as compressed CSV data, in addition to this complete finite parametrization. Every geometric state vanishes on 9, 16, and 17, so this family is neither unital nor separating.

For an explicit counterexample, the context state supported at

$$\{6,12,15,16,21,23,25,27,29,35,45,47\}$$

selects both 16 and 21, although $v_{16} \cdot v_{21} = 0$.

#### *S2.30.2. Deleting one of the nine Cabello contexts*

Number the nine contexts as the first nine rows of Table S29. For every deletion there are 26 context states, six geometric states, and a proper four-coloring of the remaining context graph. The color strings in Table S87 use the ordered ray list $(1,2,3,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20)$. The geometric states vanish precisely on the omitted context.

TABLE S87: All single-context deletion counts and four-coloring witnesses for $C_{18,9}$.

| $e$ | Omitted context | $\lvert\mathcal{S}_{\mathrm{ctx}}\rvert$ | $\lvert\mathcal{S}_{\mathrm{geom}}\rvert$ | Four-color string |
|---|---|---|---|---|
| 1 | $\{9,10,12,13\}$ | 26 | 6 | 012102203020112333 |
| 2 | $\{9,11,14,15\}$ | 26 | 6 | 012021231012102333 |
| 3 | $\{10,11,16,17\}$ | 26 | 6 | 012021320012102333 |
| 4 | $\{2,3,12,18\}$ | 26 | 6 | 013103302120113322 |
| 5 | $\{1,3,15,19\}$ | 26 | 6 | 103130320102113232 |
| 6 | $\{1,2,17,20\}$ | 26 | 6 | 330130230103112222 |
| 7 | $\{6,7,13,18\}$ | 26 | 6 | 012221213030102333 |
| 8 | $\{6,8,14,19\}$ | 26 | 6 | 012202230013112333 |
| 9 | $\{7,8,16,20\}$ | 26 | 6 | 012021320012113332 |

For deletion of context 9, $\{7,8,16,20\}$, Table S88 explicitly enumerates all 26 supports; exactly the six marked rows satisfy every geometric pair exclusion. All nine full deletion-state lists accompany the data.

TABLE S88: All 26 states after deleting context 9; geometric compatibility is marked explicitly.

| $j$ | Support | Geometric |
|---|---|---|
| 1 | $\{1,6,11,12\}$ | yes |
| 2 | $\{1,7,8,11,12\}$ | no |
| 3 | $\{1,7,12,14,16\}$ | no |
| 4 | $\{1,8,9,16,18\}$ | no |
| 5 | $\{1,10,14,18\}$ | yes |
| 6 | $\{2,6,10,15\}$ | yes |
| 7 | $\{2,7,8,10,15\}$ | no |
| 8 | $\{2,8,13,15,16\}$ | no |
| 9 | $\{2,7,9,16,19\}$ | no |
| 10 | $\{2,11,13,19\}$ | yes |
| 11 | $\{3,6,9,17\}$ | no |
| 12 | $\{3,7,8,9,17\}$ | no |
| 13 | $\{3,13,14,17\}$ | yes |
| 14 | $\{6,12,15,17\}$ | no |
| 15 | $\{7,8,12,15,17\}$ | no |
| 16 | $\{9,17,18,19\}$ | yes |
| 17 | $\{3,6,9,16,20\}$ | no |
| 18 | $\{3,7,8,9,16,20\}$ | no |
| 19 | $\{3,7,10,14,20\}$ | no |
| 20 | $\{3,8,11,13,20\}$ | no |
| 21 | $\{3,13,14,16,20\}$ | no |
| 22 | $\{6,12,15,16,20\}$ | no |
| 23 | $\{7,11,12,19,20\}$ | no |
| 24 | $\{7,8,12,15,16,20\}$ | no |
| 25 | $\{9,16,18,19,20\}$ | no |
| 26 | $\{8,10,15,18,20\}$ | no |

#### *S2.30.3. All single-ray deletions and the alternative parity subsystem*

The next table records the two span failures and all forced-zero core rays after geometric pair exclusion for every possible single-ray deletion. An entry $\{u,v\} : z$ means $v_z \in \mathrm{span}(v_u, v_v)$. Every row has twelve compatible core states, 3,584 abstract completion states, and 2,048 geometric completion states, with exactly four forced extra pairs and no other extra orthogonality in the supplied realization. The complete core supports and 24 canonical states for every row are also supplied in the data.

TABLE S89: All eighteen vertex-deletion checks; the completed geometric families fail unitality and separation in every case.

| Deleted ray | First span failure | Second span failure | Forced-zero core rays |
|---|---|---|---|
| 1 | $\{15,19\}:2$ | $\{17,20\}:3$ | $\{2,3,13\}$ |
| 2 | $\{12,18\}:1$ | $\{17,20\}:3$ | $\{1,3,14\}$ |
| 3 | $\{12,18\}:1$ | $\{15,19\}:2$ | $\{1,2,16\}$ |
| 6 | $\{13,18\}:8$ | $\{14,19\}:7$ | $\{7,8,17\}$ |
| 7 | $\{13,18\}:8$ | $\{16,20\}:6$ | $\{6,8,15\}$ |
| 8 | $\{14,19\}:7$ | $\{16,20\}:6$ | $\{6,7,12\}$ |
| 9 | $\{12,13\}:11$ | $\{14,15\}:10$ | $\{10,11,20\}$ |
| 10 | $\{12,13\}:11$ | $\{16,17\}:9$ | $\{9,11,19\}$ |
| 11 | $\{14,15\}:10$ | $\{16,17\}:9$ | $\{9,10,18\}$ |
| 12 | $\{2,3\}:13$ | $\{9,10\}:18$ | $\{8,13,18\}$ |
| 13 | $\{6,7\}:12$ | $\{9,10\}:18$ | $\{1,12,18\}$ |
| 14 | $\{6,8\}:15$ | $\{9,11\}:19$ | $\{2,15,19\}$ |
| 15 | $\{1,3\}:14$ | $\{9,11\}:19$ | $\{7,14,19\}$ |
| 16 | $\{7,8\}:17$ | $\{10,11\}:20$ | $\{3,17,20\}$ |
| 17 | $\{1,2\}:16$ | $\{10,11\}:20$ | $\{6,16,20\}$ |
| 18 | $\{2,3\}:13$ | $\{6,7\}:12$ | $\{11,12,13\}$ |
| 19 | $\{1,3\}:14$ | $\{6,8\}:15$ | $\{10,14,15\}$ |
| 20 | $\{1,2\}:16$ | $\{7,8\}:17$ | $\{9,16,17\}$ |

For the alternative subsystem, take parent variant 2 of Table S35 and retain vertices 1–17 and 66–68. It is stateless. Each ray deletion instead leaves the positive geometric state count below; restriction of any such state shows that no eighteen-ray subset can be KS.

TABLE S90: Geometric state counts after every single-ray deletion of the twenty-ray parity subsystem.

| Deleted ray | States | Deleted ray | States |
|---|---|---|---|
| 1 | 8 | 2 | 8 |
| 3 | 8 | 4 | 28 |
| 5 | 28 | 6 | 8 |
| 7 | 8 | 8 | 8 |
| 9 | 12 | 10 | 12 |
| 11 | 12 | 12 | 10 |
| 13 | 10 | 14 | 10 |
| 15 | 10 | 16 | 10 |
| 17 | 10 | 66 | 8 |
| 67 | 8 | 68 | 8 |

## S3. SUPPLEMENTARY PROOFS, REDUCTIONS, AND EXACT VERIFICATION CERTIFICATES

### S3.1. Formulas used by the finite certificates

The following formulas restore the explicit conventions used by the later certificates. State indices are local to the corresponding complete table. For Hesse, the six-state vertex-separating family, its seven-state event-separating extension, and the eighteen-state order-determining family are respectively

$$\Omega_6 = \{s_3, s_{31}, s_{75}, s_{89}, s_{113}, s_{151}\}. \tag{S23}$$

$$\Omega_7 = \Omega_6 \cup \{s_{84}\} \tag{S24}$$

$$\begin{aligned}\Omega_{18} = \{&s_2, s_8, s_{20}, s_{30}, s_{40}, s_{45},\\ &s_{54}, s_{64}, s_{73}, s_{79}, s_{83}, s_{93},\\ &s_{105}, s_{109}, s_{119}, s_{123}, s_{127}, s_{150}\}.\end{aligned} \tag{S25}$$

The members of $\Omega_6$ are denoted $\sigma_1, \dots, \sigma_6$ in the displayed order. Their atom profiles and the nontrivial atom–coatom order are

$$\kappa(i) = (\sigma_1(i), \dots, \sigma_6(i)). \tag{S26}$$

$$a_i \le a_j^{\perp} \quad \Longleftrightarrow \quad \{i, j\} \in E(\Gamma(H_{37})). \tag{S27}$$

For $H_{65}$, the following core coloring extends to the completed hypergraph. A base-compatible independent support $S_\alpha$ is extended canonically using $\varepsilon \in \{0, 1\}$. In the plane formula, $\eta_A, \eta_B$ are an integer orthogonal basis and $n_A = \|\eta_A\|^2$, $n_B = \|\eta_B\|^2$.

$$\begin{aligned}(c(1), \dots, c(17)) = (&0, 1, 2, 3, 0, 1, 2, 4, 0, 1, 3,\\ &3, 4, 2, 1, 0, 2).\end{aligned} \tag{S28}$$

$$\begin{aligned}&\big(s_{\alpha,\varepsilon}(w_{m,1}),\ s_{\alpha,\varepsilon}(w_{m,2})\big)\\ &\quad = \begin{cases}(0, 0), & |S_\alpha \cap \{u_m, v_m\}| = 1,\\ (1-\varepsilon,\ \varepsilon), & S_\alpha \cap \{u_m, v_m\} = \emptyset.\end{cases}\end{aligned} \tag{S29}$$

$$\begin{aligned}w_{m,1}(a, b) &= a\,\eta_A + b\,\eta_B,\\ w_{m,2}(a, b) &= -n_B b\,\eta_A + n_A a\,\eta_B\end{aligned} \tag{S30}$$

The eight rays adjoined to completed Yu–Oh and the nine uncovered pairs of the Cabello subsystem are fixed by the following local labels. The last display records the completed historical $\Gamma_1$ TITS incidence.

$$\begin{aligned}&(0, 1, 2), (0, 2, -1), (0, 1, -2), (0, 2, 1),\\ &(1, 2, 0), (2, -1, 0), (1, -2, 0), (2, 1, 0).\end{aligned} \tag{S31}$$

$$\begin{aligned}&\{1, 13\}, \{2, 14\}, \{3, 16\}, \{6, 17\}, \{7, 15\},\\ &\{8, 12\}, \{9, c\}, \{10, b\}, \{11, a\}.\end{aligned} \tag{S32}$$

$$E(I_{16,9}) = E(B_{13,7}) \cup \{\{4, 11, 24\}, \{4, 10, 18\}\}. \tag{S33}$$

*a. Fractional value of the faithful Cabello completion.* The geometric orthogonality graph of $C_{18,9}$ has independence number four and exactly 54 independent four-sets, with each of its 18 rays belonging to twelve of them. Assigning weight $1/12$ to each set gives a fractional cover of total weight $9/2$, matching $18/4$. Fresh pair completion preserves this value by the main completion proposition, proving $\chi_f(C_{36,18}) = 9/2$ as recorded in the main comparison. The complete 54-set certificate is in the electronic audit data.

### S3.2. Snark obstructions and quantitative robustness bounds

#### S3.2.1. Cubic graphs, Blanuša snarks, and Isaacs dot products

For a finite simple cubic graph $X$, let $H(X)$ be the context hypergraph whose vertices are the edges of $X$ and whose contexts are the three edges incident with each vertex of $X$. Then $\Gamma(H(X)) = L(X)$ and $\chi(H(X)) = \chi'(X)$. An orthogonal ray assignment for $H(X)$ in $\mathbb{F}^3$ is an assignment of nonzero vectors to the edges of $X$ such that incident edges are orthogonal.

Silva and Ibiapina[5] (Question 6) asked whether an orthogonal vector edge coloring in $\mathbb{R}^3$ can exist when $\chi'(X) = 4$. Theorem 11 excludes the Petersen graph $P$ even over $\mathbb{C}$. Using the multipole forcing law of Theorem 11(1), we can extend this exclusion to the two 18-vertex Blanuša snarks[6] and the infinite family of Isaacs dot-product snarks[7].

Recall that for a cubic graph $G$ with two nonadjacent edges $e_1 = \{x_1, y_1\}$ and $e_2 = \{x_2, y_2\}$, the Isaacs dot product $G \cdot P$ is formed by removing $e_1, e_2$ from $G$, removing two adjacent vertices $u, v$ from the Petersen graph $P$ (leaving the 4-pole $P \setminus \{u, v\}$ whose dual hypergraph is precisely $H_8$), and attaching the two dangling edges $\{2, 8\}$ of $u$ to $\{x_1, y_1\}$ and the two dangling edges $\{13, 14\}$ of $v$ to $\{x_2, y_2\}$; the two distinct choices of nonadjacent edges in $G = P$ produce the two Blanuša snarks $B_1$ and $B_2$[6–8].

**Corollary 1** (Obstruction for Blanuša and dot-product snarks)**.** *If a cubic graph $G$ admits no orthogonal vector edge coloring in $\mathbb{C}^3$, then neither does any Isaacs dot product $G \cdot P$. In particular, both 18-vertex Blanuša snarks $B_1, B_2 = P \cdot P$ and all iterated dot products $X = P \cdot P \cdots P$ satisfy $\chi(H(X)) = \chi'(X) = 4 > 3 = \omega(L(X))$ and admit no orthogonal vector edge coloring in $\mathbb{C}^3$.*

*Proof.* In any orthogonal vector edge coloring of $G \cdot P$ in $\mathbb{C}^3$, the restriction to the embedded copy of $H_8 = H(P \setminus \{u, v\})$ satisfies $[v_2] = [v_8]$ and $[v_{13}] = [v_{14}]$ by Theorem 11(1). Assigning the common ray $[v_2] = [v_8]$ to $e_1 = \{x_1, y_1\}$ and the common ray $[v_{13}] = [v_{14}]$ to $e_2 = \{x_2, y_2\}$ preserves orthogonality at $x_1, y_1, x_2, y_2$ and therefore yields a valid orthogonal vector edge coloring of $G$ in $\mathbb{C}^3$. Since $P$ admits no such coloring by Theorem 11, induction excludes $B_1, B_2$ and all iterated dot products with $P$. □

Generalized Petersen graphs[9] and flower snarks provide further families for comparing real, complex, injective, and edge-faithful realizability.

#### S3.2.2. Explicit analytic lower bound and tetrahedral tight-frame upper bounds for $\Delta_3$

Let $A$ be the 30 edges of $\Gamma(G_{32})$ and define the minimum total squared orthogonality error in $\mathbb{C}^3$:

$$\Delta_3 = \min_{\substack{v_i \in \mathbb{C}^3 \\ \|v_i\|=1 \ (1 \le i \le 15)}} \sum_{\{i,j\} \in A} |\langle v_i, v_j \rangle|^2. \tag{S34}$$

By compactness of $(S^5)^{15}$ and Theorem 11, $\Delta_3 > 0$. We now establish both a certified rational analytic lower bound and explicit tetrahedral tight-frame upper bounds.

**Proposition 2** (Quantitative bounds on $\Delta_3$). *Every assignment of unit vectors $v_1, \dots, v_{15} \in \mathbb{C}^3$ satisfies*

$$\varepsilon \coloneqq \max_{\{i,j\}\in A} |\langle v_i, v_j\rangle| > \frac{1}{120}, \tag{S35}$$

*and the minimum total squared error in Eq. (S34) satisfies*

$$\begin{aligned}\frac{1}{14400} < \Delta_3 &\le \min_{a^2+b^2+c^2=1} F(a,b,c) \\ &\le \frac{9275914990340646}{10001032226635921} < 0.927496 < \frac{18}{19},\end{aligned} \tag{S36}$$

*where*

$$F(a,b,c) = 12a^2 + 12(ab+bc-ca)^2 + 6(b^2-a^2-c^2)^2. \tag{S37}$$

*Proof.* To prove the lower bound (S35), let $\varepsilon = \max_{\{i,j\}\in A} |\langle v_i, v_j\rangle|$. For any context $e = \{x, y, z\}$, the Gram matrix of $\{v_x, v_y, v_z\}$ has diagonal entries 1 and off-diagonal entries of modulus at most $\varepsilon$, so Gershgorin's circle theorem implies $\|I_3 - (v_x v_x^\dagger + v_y v_y^\dagger + v_z v_z^\dagger)\|_{\mathrm{op}} \le 2\varepsilon$. Hence for any unit vectors $u, w \in \mathbb{C}^3$,

$$\Big|1 - \big(|\langle u, x\rangle|^2 + |\langle u, y\rangle|^2 + |\langle u, z\rangle|^2\big)\Big| \le 2\varepsilon, \tag{S38}$$

and if $\{u, x\}, \{w, z\} \in A$, then

$$\big|\langle u, w\rangle - \langle u, y\rangle\langle y, w\rangle\big| \le 4\varepsilon. \tag{S39}$$

In $\{5, 6, 7\}$ and $\{9, 10, 11\}$, the vectors $w_7 = v_{12} - v_5\langle v_5, v_{12}\rangle$ and $w_9 = v_4 - v_{11}\langle v_{11}, v_4\rangle$ satisfy $\|w_7 - v_7\langle v_7, v_{12}\rangle\| \le 3\varepsilon$ and $\|w_9 - v_9\langle v_9, v_4\rangle\| \le 3\varepsilon$, so $|\langle w_7, w_9\rangle| \le 7\varepsilon$. Expanding,

$$\begin{aligned}\langle w_7, w_9\rangle = {} & \langle v_{12}, v_4\rangle - \langle v_{12}, v_{11}\rangle\langle v_{11}, v_4\rangle - \langle v_{12}, v_5\rangle\langle v_5, v_4\rangle \\ & + \langle v_{12}, v_5\rangle\langle v_5, v_{11}\rangle\langle v_{11}, v_4\rangle.\end{aligned}$$

The two middle terms are at most $\varepsilon$ in modulus, since $|\langle v_{12}, v_{11}\rangle| \le \varepsilon$ and $|\langle v_5, v_4\rangle| \le \varepsilon$. Eq. (S39) on $\{1, 2, 3\}$ replaces each of $\langle v_{12}, v_4\rangle$, $\langle v_{12}, v_5\rangle$, $\langle v_5, v_{11}\rangle$, and $\langle v_{11}, v_4\rangle$ by its product through $v_2$ with error at most $4\varepsilon$, and hence the triple product with error at most $12\varepsilon$, since all factors have modulus at most one. The main terms combine to

$$\big(1 + |\langle v_2, v_5\rangle|^2 |\langle v_2, v_{11}\rangle|^2\big)\langle v_{12}, v_2\rangle\langle v_2, v_4\rangle,$$

whose real prefactor is at least one, so

$$\big|\langle v_{12}, v_2\rangle\langle v_2, v_4\rangle\big| \le (7 + 2 + 4 + 12)\varepsilon = 25\varepsilon.$$

By reflection symmetry, we may assume $|\langle v_2, v_4\rangle| \le 5\sqrt{\varepsilon}$. Applying Eq. (S38) to $v_4$ in $\{1, 2, 3\}$ and to $v_2$ in $\{3, 4, 5\}$ gives $|\langle v_1, v_4\rangle|^2 \ge 1 - 27\varepsilon - \varepsilon^2$ and $|\langle v_2, v_5\rangle|^2 \ge 1 - 27\varepsilon - \varepsilon^2$, so $\min_\theta \|v_1 - e^{i\theta} v_4\| \le \sqrt{54\varepsilon + 2\varepsilon^2}$ and $\min_\theta \|v_2 - e^{i\theta} v_5\| \le \sqrt{54\varepsilon + 2\varepsilon^2}$. Since $|\langle v_4, v_{10}\rangle| \le \varepsilon$ and $|\langle v_5, v_7\rangle| \le \varepsilon$, Eq. (S38) applied to $v_1$ in $\{9, 10, 11\}$ and to $v_2$ in $\{7, 8, 9\}$ gives

$$|\langle v_i, v_9\rangle|^2 \ge 1 - 2\varepsilon - \varepsilon^2 - \big(\varepsilon + \sqrt{54\varepsilon + 2\varepsilon^2}\big)^2 \qquad (i = 1, 2).$$

Comparing the sum of these two inequalities with $|\langle v_1, v_9\rangle|^2 + |\langle v_2, v_9\rangle|^2 \le 1 + 2\varepsilon$ from $\{1, 2, 3\}$ gives

$$1 \le 6\varepsilon + 2\varepsilon^2 + 2\big(\varepsilon + \sqrt{54\varepsilon + 2\varepsilon^2}\big)^2.$$

The right-hand side is strictly increasing for $\varepsilon > 0$, and at $\varepsilon = 1/120$ we have $54\varepsilon + 2\varepsilon^2 = 3241/7200 < (27/40)^2$, so the right-hand side is strictly bounded above by

$$\frac{6}{120} + \frac{2}{120^2} + 2\left(\frac{1}{120} + \frac{27}{40}\right)^2 = \frac{1417}{1440} < 1,$$

certifying $\varepsilon > 1/120$ and $\Delta_3 \geq \varepsilon^2 > 1/14400$.

For the upper bounds, set $v_4 = e_1$, $v_8 = e_2$, $v_{12} = e_3$, and for any unit vector $(a, b, c) \in S^2$, define the remaining 12 unit vectors as representatives of the tetrahedral $A_4$-orbit of rays:

$$\begin{aligned} v_1 &= (-b, -c, -a)^T, & v_2 &= (c, -a, b)^T, \\ v_3 &= (-a, -b, c)^T, & v_5 &= (-a, b, c)^T, \\ v_6 &= (-b, c, -a)^T, & v_7 &= (c, a, b)^T, \\ v_9 &= (-c, -a, b)^T, & v_{10} &= (-a, b, -c)^T, \\ v_{11} &= (b, -c, -a)^T, & v_{13} &= (a, b, c)^T, \\ v_{14} &= (b, c, -a)^T, & v_{15} &= (-c, a, b)^T. \end{aligned} \tag{S40}$$

These 15 vectors form a tight frame $\sum_{i=1}^{15} v_i v_i^T = 5I_3$, and the sum of squared overlaps on the 30 edges of $\Gamma(G_{32})$ equals $F(a, b, c)$ in Eq. (S37). Evaluating Eq. (S40) at the normalized integer direction $(a, b, c) = (1, 3, -3)/\sqrt{19}$ yields rational squared overlaps with $\Delta_3 \leq 342/361 = 18/19 \approx 0.947368$ and $\max_{\{i,j\}\in A} |\langle v_i, v_j\rangle|^2 = 1/19$; evaluating at the refined normalized integer direction $(a, b, c) = (2332, 7101, -6644)/\sqrt{100005161}$ yields the certified rational bound in Eq. (S36). □

Numerical minimization of $F$ on $S^2$ (as well as unrestricted local searches over $(S^5)^{15}$) finds an attained candidate minimum near 0.927495724691 at $(a, b, c) \approx (0.23319713, 0.71005263, -0.66441280)$; certifying whether the unrestricted global minimum $\Delta_3$ coincides with $\min_{S^2} F$ remains an open optimization problem. Choosing instead the rational unit vector $(a, b, c) = \frac{1}{3}(1, 2, -2)$ in Eq. (S40) makes $ab + bc - ca = 0$, so four of the ten contexts of $G_{32}$ ($\{1, 2, 3\}, \{5, 6, 7\}, \{9, 10, 11\}, \{13, 14, 15\}$) become exact orthonormal bases of $\mathbb{R}^3$, all 15 rays are distinct, and $\Delta_3 \leq 38/27$ with $\max_{\{i,j\}\in A} |\langle v_i, v_j\rangle| = 1/3$.

### S3.3. Hesse $H_{37}$: minimum state families

All state indices in this appendix use the canonical enumeration of Appendix S2.27. The minima for unital vertex separation and event separation have the counting lower bounds $2^5 - 1 < 37$ and $2^6 < 76$. The families in Eqs. (S23) and (S24) attain those bounds: their profiles give respectively 37 distinct nonzero atom profiles and 76 distinct event profiles. The seven-state family still has 462 ordered event pairs for which inclusion of the profiles does not imply the lattice order. The family in Eq. (S25) has no such pair and hence gives the upper bound eighteen for order determination. This appendix supplies a computer-assisted proof of the matching lower bound.

#### *S3.3.1. Reduction to families of seventeen states*

Let $D = \{P_0, P_1, P_2, Q_0, Q_1, Q_2\}$. For each unordered pair $\{u, v\} \subset D$, order determination requires a state with $s(u) = s(v) = 1$: the pair is nonorthogonal, so $u \not\leq v^\perp$. Every state takes value one on at most two members of $D$. Consequently the fifteen pairs require fifteen different states. Table S91 gives their complete witness classes. These classes are disjoint, their union $T$ has 27 states, and a choice of one witness from each class has $1^3 2^{12} = 4{,}096$ possibilities.

TABLE S91: Complete joint-one witness classes for the fifteen pairs in $D$. Each entry lists all canonical indices $r$ for which $s_r(u) = s_r(v) = 1$.

| Pair $\{u, v\}$ | Witness indices $r$ |
|---|---|
| $\{P_0, P_1\}$ | 150 |
| $\{P_0, P_2\}$ | 73 |
| $\{P_0, Q_0\}$ | 93, 145 |
| $\{P_0, Q_1\}$ | 127, 133 |
| $\{P_0, Q_2\}$ | 79, 99 |
| $\{P_1, P_2\}$ | 2 |
| $\{P_1, Q_0\}$ | 105, 139 |
| $\{P_1, Q_1\}$ | 14, 119 |
| $\{P_1, Q_2\}$ | 40, 48 |
| $\{P_2, Q_0\}$ | 54, 60 |
| $\{P_2, Q_1\}$ | 8, 26 |
| $\{P_2, Q_2\}$ | 20, 68 |
| $\{Q_0, Q_1\}$ | 109, 113 |
| $\{Q_0, Q_2\}$ | 83, 87 |
| $\{Q_1, Q_2\}$ | 30, 34 |

The nine nonorthogonal pairs $\{z, C_{ak}\}$ require further joint-one witnesses. No state in $T$ witnesses any of them. Indeed, if a state activates a $P$ vertex then $s(z) = 0$. If it instead activates two $Q$ vertices, the state equations force the selected $M$ vertices into the remaining $Q$ column, and all nine $C$ values are zero. Any state outside $T$ witnesses at most six of the nine pairs: for $s(z) = 1$, the three selected $M$ vertices force at least three $C$ values to vanish. Thus at least two states outside $T$ are necessary.

It follows that an order-determining family of at most seventeen states would have exactly seventeen members: one from each of the fifteen classes in Table S91, and two outside $T$ that together witness all nine $\{z, C_{ak}\}$ pairs. Enumerating the $\binom{126}{2}$ unordered pairs outside $T$ leaves exactly 132 with this property. The reduction therefore leaves precisely

$$4{,}096 \cdot 132 = 540{,}672 \tag{S41}$$

candidate families. The nine pairs alone do not require three extra states; for example, $s_1$ and $s_{72}$ jointly witness all nine. The remaining order requirements must also be checked.

#### *S3.3.2. Exhaustive order certificate*

For each of the 5,407 ordered event pairs $x \not\leq y$, define its nonempty witness set

$$W_{xy} = \{r \in \{1, \ldots, 153\} : s_r(x) = 1,\ s_r(y) = 0\}. \tag{S42}$$

A family is order determining exactly when its index set meets every $W_{xy}$. Repeated sets and proper supersets can be discarded, because meeting an inclusion-minimal set also meets every set containing it. There are exactly 81 inclusion-minimal witness sets. The exhaustive check forms each of the 540,672 families in Eq. (S41) and counts the minimal witness sets that it meets. The largest count is 75, strictly below 81. Hence no family of seventeen states is order determining. Together with $\Omega_{18}$, this proves the minimum eighteen.

The accompanying Hesse extension checker, invoked by the joint entry point, reconstructs the contexts and all 153 states from their symbolic parametrization, checks their agreement with `hesse37-data.json`, and performs this finite search using integer bit sets and the Python standard library. No numerical tolerance, optimization solver, or unreported search cutoff is used. It also verifies the printed seven- and eighteen-state families and every entry of Table S91. From the extracted joint project directory, run the joint entry point, without Python's optimization option:

```
python verification/verify-integrated.py
```

The generated file `verification/_work/hesse/hesse37-extensions-certificate.json` records all 81 minimal witness sets, a representative event pair for each, all 132 possible extra pairs, the coverage-count distribution over the 540,672 candidates, and all

event profiles for $\Omega_6$, $\Omega_7$, and $\Omega_{18}$. The generated file `verification/_work/hesse/hesse37-order18-witnesses.csv` records a witness in $\Omega_{18}$ for every $x \not\leq y$. These two files use the same numbered event labels as the full-family certificate in Appendix S2.27.6.

The program additionally checks Berge girth five from the incidence graph, computes every meet and join in the 76-event order, verifies the orthomodular law, and verifies the projector sum in Eq. (30) exactly in $\mathbb{Q}(\zeta)$. The nonexistence of an injective real representation is proved analytically in Sec. V A.

### S3.4. Four-dimensional $H_{\mathrm{YO}}^{(4)}$ and $H_{65}$: completion and verification

The coordinates in Table S17 can be reproduced by the following finite selection rule. Normalize an integer vector by dividing by the greatest common divisor of its entries and choosing the sign so that its first nonzero entry is positive. Form the set of normalized nonzero vectors obtained from $\{-3,-2,\dots,3\}^4$, ordered by squared norm, then maximum absolute entry, then lexicographically. Process bridge edges in the order $e_{13},\dots,e_{24},e_1,\dots,e_{12}$. For each pair, choose $\eta_A$ to be the first vector in this ordered pool orthogonal to its endpoints, and $\eta_B$ the first such vector also orthogonal to $\eta_A$.

For all $(a,b) \in \{-7,-6,\dots,7\}^2 \setminus \{(0,0)\}$, form the pair in Eq. (S30) and normalize both vectors. Discard pairs that repeat an already placed ray or have an orthogonality to any already placed vertex other than the two endpoints. Among the remaining ordered pairs choose the lexicographically smallest score

$$\left( \max_{j,k} |(w_{m,j})_k|,\ \|w_{m,1}\|^2 + \|w_{m,2}\|^2,\ w_{m,1}, w_{m,2} \right). \tag{S43}$$

In Eq. (S43), the order on the last two entries is the usual lexicographic order on their integer coordinates. This rule yields exactly the 48 completion vectors in Table S17. It is a recipe for this specific example; the general existence assertion is established independently by Lemma 21.

The companion verifier parses the integrated coordinate and state tables, reproduces this completion rule and all 393,216 state extensions, and checks the maximal cliques, fractional-coloring certificate, atom-order failure, forced hub-completion zero, and warm-up data. It is included with the complete state file and invoked by the joint entry point in Sec. S3.7.

### S3.5. Exact Cabello completion and forced-zero certificates

For each of the nine uncovered pairs in Eq. (S32), rational row reduction verifies that its two endpoint vectors span no third core ray. If two endpoint planes were orthogonal, their four endpoints would form another full basis. Exact four-clique enumeration gives only the original nine bases, all of whose pairs are already covered. This excludes such an orthogonal-plane pair. The integer vectors in Table S28 directly verify the resulting faithful completion: 36 distinct projective rays, eighteen full bases, and 108 orthogonal pairs, with every pair covered by exactly one listed basis. The nine-context graph and the full eighteen-ray orthogonality graph both fail four-colorability and admit five-colorings, as checked by exact finite search in the companion certificate. Lemma 21(2) transfers the chromatic value to the completion.

For the vertex-deleted core with $c$ removed, rational row reduction finds exactly the span failures

$$v_{16} \in \mathrm{span}(v_1, v_2), \qquad v_{17} \in \mathrm{span}(v_7, v_8).$$

There is no orthogonal-plane-pair obstruction. The finite-avoidance choice in Lemma 21 can therefore avoid all extra incidences except those forced by these two spans; the explicit 45-ray inventory has exactly the four pairs printed in Sec. S1.17.2. The two fresh vectors

completing $\{1,2\}$ must both be orthogonal to 16, and those completing $\{7,8\}$ must both be orthogonal to 17.

The twelve rows of Table S84 give a short complete proof of the resulting obstruction. Whenever 16 is selected, neither 1 nor 2 is selected. The context completing $\{1,2\}$ then needs one fresh value equal to one, while geometric exclusion with 16 requires both to be zero. Hence every geometric extension has $s(16) = 0$. The same argument with $\{7,8\}$ gives $s(17) = 0$. These implications use the displayed core states in addition to the span incidences; the span incidences alone would not imply them. The retained basis $\{10, 11, 16, 17\}$ gives $s(10)+s(11) = 1$. If $s(9) = 1$, the other two hub bases give $s(10) = s(11) = 0$, a contradiction. Thus 9, 16, and 17 have identical zero profiles.

Conversely, each of the eight indicated core rows avoids 16 and 17, so all its fresh-pair choices satisfy the four extra exclusions. Each misses eight bridges. This proves the exact count $8\,2^8 = 2{,}048$, as well as failure of geometric unitality and separation. The two excluded core rays remain impossible for every fresh-pair completion of these fixed planes; additional accidental orthogonalities cannot restore them. Table S89 supplies the individual span checks for all eighteen deletions. Their full core tables, exact integer completions, and extension counts are included in `data/cabello18-review-checks.json`.

To audit the ancestry result, enumerate every four-clique after adjoining either three-ray choice to all 65 original rays. In both cases there are exactly 35 bases, 195 geometric pairs, and 192 context-covered pairs; the maximum basis intersection is two. The first union contains the displayed eighteen-ray parity proof. The second union contains the twenty-ray proof with multiplicities $4, 4, 2, \ldots, 2$ and eleven contexts. Each of its twenty ray deletions has a compatible assignment by Table S90. All twenty rays lie in Peres's 24-ray set. Enumerating the $\binom{24}{4} = 10{,}626$ twenty-ray subsets of that set, with all bases they contain, gives exactly 96 ray-critical KS subsets, each with eleven bases and all mutually isomorphic; the twenty-ray proof is one of them. Since Peres's set contains Kernaghan's critical twenty-ray set and 95 others[3], the twenty-ray proof is isomorphic to Kernaghan's[10]. All coordinates, full basis lists, and the preserved $H_{65}$ restriction are checked independently of the unrestricted ancestry existence proposition.

### S3.6. Transverse extension and criticality certificates

For $d = 3$, number the two copies of an $n$-vertex input as $i$ and $n + i$, and the fresh transverse vertex as $2n + i$. The transverse contexts are $\{i, n + i, 2n + i\}$ and the horizontal contexts are the two numbered copies of the original list. This explicitly specifies all 45 vertices and 33 contexts for $T(H_9)$, and all 75 vertices and 57 contexts for $T(H^+_{\mathrm{YO}})$.

For each deleted original context $e$, a strong three-coloring of $H \setminus \{e\}$ having two equal colors and a singleton color on $e$ gives a state after deletion of one horizontal copy of $e$. Use the repeated-color indicator on the deleted copy and the singleton-color indicator on the intact copy. Both have exactly one active vertex in each of their required contexts and their supports are disjoint; the remaining transverse values are forced. For deletion in the other copy, interchange the two indicators. Table S92 supplies an explicit coloring for every required context of both inputs. The color strings use original vertex order, and their labels 0,1,2 may be permuted.

TABLE S92: Explicit deletion colorings with pattern (1,1,2) and the number of states after deleting the corresponding first-copy horizontal context.

| Input | Deleted original context | Color string | Extension states |
|---|---|---|---|
| $H_9$ | $\{1,2,3\}$ | 212010201201122 | 20 |
| $H_9$ | $\{1,11,12\}$ | 201020120121221 | 20 |
| $H_9$ | $\{2,8,15\}$ | 021020120121220 | 12 |
| $H_9$ | $\{3,4,5\}$ | 012120102012212 | 20 |
| $H_9$ | $\{4,10,13\}$ | 012102120121000 | 12 |
| $H_9$ | $\{5,6,7\}$ | 012010120121220 | 20 |
| $H_9$ | $\{6,12,14\}$ | 012102102012202 | 12 |
| $H_9$ | $\{7,8,9\}$ | 012010200121222 | 20 |
| $H_9$ | $\{9,10,11\}$ | 012010201121222 | 20 |
| $H_{\mathrm{YO}}^{+}$ | $\{1,2,3\}$ | 200021210112020102011222 1 | 16 |
| $H_{\mathrm{YO}}^{+}$ | $\{1,10,19\}$ | 210021201212020001011202 1 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{1,13,20\}$ | 120210201012120011200220 1 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{2,5,8\}$ | 021021012120201122211001 2 | 24 |
| $H_{\mathrm{YO}}^{+}$ | $\{3,11,14\}$ | 012021102120200122210001 1 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{3,12,25\}$ | 021210102101222020210101 0 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{4,5,6\}$ | 021200210102121110220221 0 | 16 |
| $H_{\mathrm{YO}}^{+}$ | $\{4,10,18\}$ | 021210201202121010120201 0 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{4,11,24\}$ | 021201012120201120211000 2 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{6,12,17\}$ | 021210102100222110210101 2 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{6,13,22\}$ | 021012102120200112210001 2 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{7,8,9\}$ | 021012002120101112222001 2 | 16 |
| $H_{\mathrm{YO}}^{+}$ | $\{7,11,15\}$ | 021201012100221120211001 2 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{7,13,21\}$ | 021012102120100112220001 2 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{9,10,23\}$ | 021210201102121010220201 0 | 8 |
| $H_{\mathrm{YO}}^{+}$ | $\{9,12,16\}$ | 021210102102222010210101 0 | 8 |

Deleting one transverse context $T_v$ instead permits an ordered pair of original state supports $S_1, S_2$ exactly when $S_1 \cap S_2 \subseteq \{v\}$. In these non-three-colorable inputs disjoint pairs do not exist, so the intersection must equal $\{v\}$; the isolated fresh vertex has two choices. For $H_9$ this gives four states for each $v = 1, \ldots, 12$ and twelve states for each $v = 13, 14, 15$. Together with Table S92, this proves criticality of all 33 contexts. For completed Yu–Oh it gives zero states at $v = 1, \ldots, 13$ and eight states at $v = 14, \ldots, 25$. Thus horizontal criticality does not imply criticality of this full transverse construction. The corresponding $G_{32}$ extension has 45 vertices and 35 contexts and is stateless. Its distinct six state supports have pairwise intersection size one; for $H_9$ the distinct eight supports have intersection size one in 21 pairs and size three in seven pairs.

### S3.7. Data and verification

The companion electronic package contains coordinate and context inventories, complete state and coloring enumerations, verification programs, exact parent inventories, and finite certificates. Vertex labels and state orderings agree with the tables in this supplement. The verification programs check the common Yu–Oh numbering, the published 49-ray parent map, the parent vectors and contexts, the full $H_{65}$ state inventory, and cross-reference targets. The finite lower-bound certificate for the eighteen-state Hesse minimum is the exhaustive 540,672-candidate argument of Sec. S3.3.

Run the companion verification entry point from the project directory:

```
python verification/verify-integrated.py
```

The exact checks use rational or $\mathbb{Q}(\zeta)$ arithmetic, with $\zeta^2 + \zeta + 1 = 0$. Reports and hashes accompany the source.

The finite checks certify the instances and certificates they explicitly test. The analytic realization and existence results also rely on their proofs in the text.

## S3.8. Historical Kochen–Specker gadget certificates (1965 Fig. 1 and 1967 $\Gamma_1$)

### *S3.8.1. The 1965 triangles and the completed historical TITS*

Identifying the repeated occurrences of vertices 1 and 2 in Kochen–Specker's 1965 Fig. 1 gives

$$\{134, 325, 2\,9\,11, 11\,1\,10, 479, 5\,8\,10, 678\}. \tag{S44}$$

These are also the triples used in the displayed function definitions on p. 182 of Ref.[11]. Their four state supports are

$$\{1,5,6,9\}, \quad \{2,4,6,10\}, \quad \{3,6,9,10\}, \quad \{4,5,6,11\}.$$

The TIFS between 1 and 2 is nonvacuous. Directly, if both were true, contexts 134, 325, 2 9 11, and 11 1 10 would make 3,4,5,9,10,11 false; contexts 479 and 5 8 10 would then make both 7 and 8 true, contradicting 678.

Every state has 6 true and 7,8 false, so no three-coloring can exist: the three color-class states of any coloring would need to be disjoint. The graph is abstract rather than an injective ray example. In a purported injective realization in $\mathbb{C}^3$, vertices 1 and 9 are both orthogonal to 4 and 11. Distinct rays 4 and 11 span a plane, whose orthogonal complement is one-dimensional; thus rays 1 and 9 coincide. This proves the obstruction even over the complex field. No claim that the 1965 authors presented this graph as a Hilbert-space realization is intended.

For the 1967 $\Gamma_1$[12] (pp. 68,70), the correspondence between its original vertices and $I_{16,9}$ is

| $\Gamma_1$ | $a_0$ | $a_1$ | $a_2$ | $a_3$ | $a_4$ | $a_5$ | $a_6$ | $a_7$ | $a_8$ | $a_9$ |
|---|---|---|---|---|---|---|---|---|---|---|
| $I_{16,9}$ | 10 | 9 | 1 | 7 | 3 | 8 | 2 | 11 | 4 | 24 |

The already complete triangles are $\{a_1, a_3, a_5\}$, $\{a_2, a_4, a_6\}$, and $\{a_7, a_8, a_9\}$. The six remaining edges

$$a_0a_1, \ a_0a_2, \ a_0a_8, \ a_3a_7, \ a_4a_7, \ a_5a_6$$

are completed by fresh vertices $23, 19, 18, 15, 14, 5$, respectively. The resulting nine contexts are exactly Eq. (S33). This also identifies its directed TITS $a_0 \to a_9$ with $10 \to 24$.


[1] M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
[2] J. Bub, Foundations of Physics **26**, 787 (1996).
[3] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
[4] K. Svozil and J. Tkadlec, Journal of Mathematical Physics **37**, 5380 (1996).
[5] A. Silva and A. Ibiapina, "On orthogonal vector edge coloring," (2019), arXiv:1909.01918 [cs.DM].
[6] D. Blanuša, Glasnik Matematičko-Fizički i Astronomski, Serija II **1**, 31 (1946).
[7] R. Isaacs, The American Mathematical Monthly **82**, 221 (1975).
[8] G. Brinkmann, M. Preissmann, and D. Sasaki, Discrete Mathematics & Theoretical Computer Science **17**, 369 (2015).
[9] G. Schrag, Discrete Mathematics **70**, 185 (1988).
[10] M. Kernaghan, Journal of Physics A: Mathematical and General **27**, L829 (1994).
[11] S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
[12] S. Kochen and E. P. Specker, Journal of Mathematics and Mechanics **17**, 59 (1967).