Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments

Karl Svozil [1, *]

[1] *Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria*

(Dated: November 5, 2026)

Suppose intertwined complete experiments each measure a nondegenerate maximal observable with one common spectrum, and the eigenvalue carried by a shared outcome is an element of physical reality, the same in every experiment. Such noncontextual eigenvalue assignments exist exactly when the context hypergraph is colorable with as many colors as the dimension. Their failure is weaker but more widespread than Kochen–Specker contextuality: completed Yu–Oh keeps truth assignments separating all its rays, yet admits no noncontextual eigenvalue assignment. Where such assignments exist, they can force distinct rays to share an eigenvalue; for one quantum preparation, a third of any classical model must use truth assignments that no eigenvalue assignment produces. We also correct an earlier inference from coloring to geometry and separate eigenvalue assignments from realizability and Born probabilities.

I. INTRODUCTION

An experimental context with d rank-one outcomes admits a nondegenerate observable with d distinct eigenvalues. These eigenvalues distinguish all d outcomes and provide the maximal resolution of a complete projective experiment within that context. Such an observable is maximal in the sense that every self-adjoint observable commuting with it is a function of it, so the statistics of the latter can be obtained by processing its measurement outcomes. Noninjective functions merge outcomes and reduce this resolution. This is the finite-dimensional setting of von Neumann's functional representation of commuting observables [1, Satz 8, p. 221]; see also Halmos [2, §84, pp. 171–172].

When complete experiments are intertwined, they share outcomes represented by the same rank-one projectors. We ask whether the eigenvalues of the maximal observables measured in such intertwined experiments can be assigned to their outcomes noncontextually, in the spirit of the elements of physical reality of Einstein, Podolsky, and Rosen [3]: whether every outcome carries a definite eigenvalue that belongs to the outcome itself, rather than to the experiment in which it is recorded. Three assumptions make this precise.

(i) *Maximality.* Each complete experiment measures a maximal observable, so its d outcomes are never merged and carry d distinct eigenvalues. A degenerate observable would discard part of what the experiment resolves.

(ii) *Uniformity.* All maximal observables share one spectrum of d eigenvalues $\Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$. The eigenvalues are nothing but labels of the d outcomes of a complete experiment. Since self-adjoint operators with a common nondegenerate spectrum are unitarily equivalent, uniformity says that every complete experiment measures one and the same quantity in a different orientation, $A_e = U_e A U_e^\dagger$.

(iii) *Noncontextuality of eigenvalues.* An outcome shared by several complete experiments carries the same eigenvalue in each of them, just as Kochen and Specker require a shared outcome to carry the same truth value [4].

No constraint is placed on how the d eigenvalues are distributed over the outcomes of different contexts; any permutation within a context is allowed. A noncontextual eigenvalue assignment in this sense labels the outcomes, the most elementary record a complete experiment produces, and assumption (iii) treats these labels as elements of physical reality.

These assumptions concern eigenvalues rather than which outcome occurs, yet they have immediate consequences for truth assignments. A two-valued state selects one outcome in every context and retains only the distinction between that outcome and the others. A noncontextual eigenvalue assignment yields d such states at once, one for each eigenvalue λ_a: "the outcome with eigenvalue λ_a occurs in every experiment." These d truth assignments have pairwise disjoint supports covering every outcome. A noncontextual eigenvalue assignment is thus a hidden-variable model whose hidden variable is the eigenvalue itself; its d equally weighted values reproduce the maximally mixed state. Because this requirement is more demanding than Kochen–Specker noncontextuality, which asks for a single truth assignment, its failure—*chromatic contextuality*—is a weaker form of contextuality that already appears in configurations retaining many classical truth assignments. Even when a noncontextual eigenvalue assignment exists, its consistency across intertwined contexts can force relations between distinct outcomes, such as equal eigenvalues for nonorthogonal rays, that are absent from the full family of individual truth assignments. We study both the existence of noncontextual eigenvalue assignments and the event distinctions and

* karl.svozil@tuwien.ac.at

logical relations that survive when such assignments are available [5–7].

To express these questions, let a d-uniform context hypergraph $H = (V, E)$ encode the complete experiments. An ordinary two-valued state $s : V \to \{0, 1\}$ selects exactly one outcome in every context, so $\sum_{v \in e} s(v) = 1$ for every $e \in E$. The Kochen–Specker (KS) theorem [4] establishes the nonexistence of such a global truth assignment for suitable finite families of intertwined projective contexts in dimension $d \geq 3$.

A strong d-coloring $c : V \to \{0, \ldots, d-1\}$ assigns all d distinct labels bijectively within every context and independently of context at shared vertices. It is the formal counterpart of a noncontextual eigenvalue assignment: $c(i) = a$ means that outcome i carries the eigenvalue λ_a in every context containing it, so assumptions (i)–(iii) hold if and only if $\chi(H) \leq d$. For an injective ray realization $i \mapsto [v_i] \in \mathbb{P}(\mathbb{F}^d)$, with $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, rank-one projectors P_i, and a common nondegenerate real spectrum $\Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$, it defines the maximal context observables

$$A_e = \sum_{i \in e} \lambda_{c(i)} P_i. \tag{1}$$

Because the indicator functions $s_a(v) = \mathbf{1}_{\{c(v)=a\}}$ ($a \in \{0, \ldots, d-1\}$) of any strong d-coloring form d two-valued states whose supports partition V, one has the strict one-way implication

$$\mathcal{S}(H) = \varnothing \quad \Longrightarrow \quad \chi(H) > d, \tag{2}$$

whose converse fails even when $\mathcal{S}(H)$ is unital and separates every pair of vertices. The earlier introduction in Ref. [6] inferred absence of global two-valued assignments from $\chi > d$. Equation (2) is the valid implication; completed Yu–Oh refutes the converse. In terms of eigenvalues, if no noncontextual truth assignment exists, no noncontextual eigenvalue assignment exists either; but noncontextual eigenvalues can be impossible while truth assignments, even a unital separating family of them, survive. The distinct Appendix B inference about geometry is corrected below. We say that H has an *exact palette*, or is *chromatically complete*, when $\chi(H) = d$, with no additional density or geometric assumption. This asymmetry separates chromatic contextuality into two distinct regimes that are unified in this work:

1. *Regime I, common-spectrum chromatic obstructions ($\chi(H) > d$, possibly with $\mathcal{S}(H) \neq \varnothing$):* Here no strong d-coloring exists ($\mathcal{C}_d(H) = \varnothing$), so the outcomes admit no noncontextual assignment of d eigenvalues: every family of maximal observables with one common spectrum must either merge outcomes or let a shared outcome change its eigenvalue from context to context. This happens even in examples in which classical truth assignments survive. A critical foundational clarification—correcting an interpretation in Appendix B of the earlier work with Shekarriz [6]—is that the chromatic obstruction $\chi(H) > d$ is an operator spectral-value obstruction and *does not* by itself obstruct geometric ray realizability in $\mathbb{F}^d$. Indeed, for finite 3-uniform linear hypergraphs of Berge girth five with unital separating state spaces, strong 3-colorability and injective $\mathbb{C}^3$ realizability are logically independent (Proposition 10). Ruling out geometric ray assignments for Greechie's Petersen-derived G_{32} [8] requires a direct Hermitian projection-identity argument (Theorem 11) rather than $\chi(G_{32}) = 4$.

2. *Regime II, exact-palette chromatic separability and logical gadgets ($\chi(H) = d$ with $\mathcal{S}_{\text{ext}}(H) \subsetneq \mathcal{S}(H)$):* In this regime, $\mathcal{C}_d(H) \neq \varnothing$, so noncontextual eigenvalue assignments exist, but some two-valued states $s \in \mathcal{S}(H)$ cannot be completed to a strong d-coloring: they are truth assignments that no noncontextual eigenvalue assignment produces. We characterize the extendable states $\mathcal{S}_{\text{ext}}(H)$ by the residual chromatic condition $\chi(\Gamma(H)[V \setminus \operatorname{supp} s]) \leq d - 1$ (Theorem 2), which in dimension $d = 3$ yields an odd-cycle nonextendability certificate. In the specified subhypergraphs—such as the faithful real 17-ray and 21-ray restrictions $H_{17,10}$ and $H_{21,13}$ of completed Yu–Oh [7, 9] (Theorem 15) and our faithful 3- and 4-dimensional logical gadgets (Secs. IV C and IV D)—ordinary states separate all vertices while every admissible d-coloring forces nonadjacent rays to share a color ($u \sim_c v$, chromatic inseparability / implicit identity [10]) or obeys purely chromatic true-implies-false (TIFS) or ternary exclusions violated by nonextendable states in $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$.

These two regimes are governed by a shared structural mechanism. Lemma 3 ($\chi(H) = d$ if and only if H admits $d - 1$ pairwise disjoint states) is the global unanchored form of Theorem 2. Likewise, the eight normalized 3-colorings of the four base contexts of completed Yu–Oh H^+_{YO} partition into four antipodal pairs that each make one of the four diagonal neighborhoods rainbow (Theorem 13): retaining all four diagonal sectors forces $\chi(H^+_{\text{YO}}) = 4$ (Regime I), whereas retaining two or three sectors leaves four or two normalized 3-colorings and forces two or 21 chromatically inseparable pairs in $H_{17,10}$ and $H_{21,13}$ (Regime II), which our exhaustive 65,535-subset inventory classification (Proposition 16) proves minimal within H^+_{YO}.

We further connect these combinatorial and geometric layers to orthomodular event representations and quantum probability polytopes $\mathcal{P}_{\text{ext}}(H) \subseteq \mathcal{P}(H) = \operatorname{conv} \mathcal{S}(H)$, where $\mathcal{P}_{\text{ext}}(H)$ collects the probability vectors of hidden-variable models that choose a noncontextual eigenvalue assignment and one of its d eigenvalues. On $H_{17,10}$, an explicit quantum ray preparation lies in $\mathcal{P}(H_{17,10}) \setminus \mathcal{P}_{\text{ext}}(H_{17,10})$ and requires nonextendable state weight exactly $1/9$; the preparation along the coordinate ray z_3 requires exactly $1/3$, the largest weight that a single chromatic identity of $H_{17,10}$ can certify (Theorem 24).

In complex dimension three, the 37-ray Hesse restriction H_{37} [11] has no injective $\mathbb{R}^3$ realization, pastes into a 76-event orthomodular lattice with exact minimum state-family sizes 6, 7, and 18 for vertex separation, event separation, and order determination (Theorem 19), and demonstrates that exact event-order determination does not guarantee classical mixtures for all Born preparations (Corollary 25). In real dimension four, our two-plane completion lemma (Lemma 21) produces the canonical linear 65-ray hypergraph H_{65} completing the 17-ray core of Haynes *et al.* [12] with $\chi = 5$, $\chi_f = 17/4$, and 393,216 states that fail atom-order determination and collapse a ray with a plane (Theorem 22). Finally, we formulate and prove the existence form of the KS ancestry hypothesis (Proposition 26), give an abstract transverse chromatic-to-stateless construction (Proposition 27), establish exact inclusions into specified KS parents (Bub's 49-ray set [13], the 165-ray MUB inventory, Peres's 24-ray geometry, and 68-ray extensions containing the 18-ray parity proof $C_{18,9}$ of Cabello *et al.* [14] and a copy $C_{20,11}$ of Kernaghan's 20-ray parity proof [15]), and contrast abstract context states $\mathcal{S}_{\mathrm{ctx}}$ with geometric states $\mathcal{S}_{\mathrm{geom}}$ on the 36-ray completion and 45-ray vertex deletion of $C_{18,9}$. Complete vector, state, and coloring enumerations and finite verification certificates are provided in the Supplemental Material [16], which includes Refs. [17–19].

II. UNIFIED FRAMEWORK: STATES, COLORINGS, SEPARABILITY, AND COMPLETION

A. Context hypergraphs, orthogonal realizations, and spectral regimes

All hypergraphs and graphs considered here are finite. A d-uniform *context hypergraph* $H = (V, E)$ ($d \geq 2$) consists of a nonempty collection E of d-element subsets of V, called *contexts*, with $V = \bigcup_{e\in E} e$. Its *two-section* (or *context*) *graph* $\Gamma(H)$ has vertex set V and edge set $E(\Gamma(H)) = \bigcup_{e\in E} \binom{e}{2}$. A hypergraph H is *linear* if $|e \cap e'| \leq 1$ for all distinct $e, e' \in E$. A *Berge cycle* of length $\ell \geq 2$ is an alternating sequence of distinct vertices $x_1, \ldots, x_\ell$ and distinct contexts $e_1, \ldots, e_\ell$ with $\{x_j, x_{j+1}\} \subseteq e_j$ cyclically; the least such ℓ is the *Berge girth* $g(H)$ (or ∞ if acyclic). In Greechie's pasting of Boolean blocks [8, 20], linearity ($g(H) \geq 3$) with $g(H) \geq 4$ yields an *orthomodular poset*, and $g(H) \geq 5$ yields an *orthomodular lattice* (OML).

With the Hermitian inner product $\langle x, y\rangle = \sum_j \overline{x_j} y_j$ on $\mathbb{F}^d$ ($\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$), we distinguish three levels of geometric realization:

1. An *orthogonal ray assignment* $\rho : V \to \mathbb{P}(\mathbb{F}^d)$, $i \mapsto [v_i]$ ($v_i \neq 0$), requires $\langle v_i, v_j\rangle = 0$ whenever $i \sim j$ in $\Gamma(H)$; nonadjacent vertices may share a ray.

2. An *injective orthogonal representation* additionally requires $[v_i] \neq [v_j]$ for all $i \neq j$.

3. An injective representation is *edge-faithful* (abbreviated FOR) if $\langle v_i, v_j\rangle = 0 \iff i \sim j$ in $\Gamma(H)$ for all $i \neq j$ [21].

A *strong k-coloring* of H is a map $c : V \to \{0, \ldots, k-1\}$ injective on every context $e \in E$, i.e., a proper k-coloring of $\Gamma(H)$; the least such k is $\chi(H) = \chi(\Gamma(H))$. We call H *chromatically complete* (equivalently, H has an exact palette) when $\chi(H) = d$ and write $\mathcal{C}_d(H)$ for the set of strong d-colorings of H. For a realized d-uniform hypergraph with rank-one projectors $P_i = v_i v_i^\dagger / \langle v_i, v_i\rangle$, three spectral-labeling regimes arise: (i) *context-dependent common spectrum*, where bijections $\ell_e : e \to \Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$ define $A_e = \sum_{i\in e} \ell_e(i) P_i$ with shared projectors carrying context-dependent eigenvalues; (ii) *context-independent vertex labeling* via a strong k-coloring c and distinct real scalars $\lambda_0, \ldots, \lambda_{k-1}$ in Eq. (1), where spectra vary across contexts whenever $|c(V)| > d$ (unavoidable when $\chi(H) > d$); and (iii) *chromatic completeness* ($k = d$), which provides both context-independent eigenvalues and a single d-element spectrum Σ for all A_e. Only regime (iii) meets all three assumptions of the Introduction: regime (i) gives up noncontextual eigenvalues, and regime (ii) keeps them at the price of more than d eigenvalues, so that different contexts have different spectra.

B. Two-valued states, color extendability, and the separation hierarchy

A *two-valued context state* is a function $s : V \to \{0, 1\}$ satisfying

$$\sum_{v\in e} s(v) = 1 \qquad (e \in E). \tag{3}$$

We write $\mathcal{S}(H) = \mathcal{S}_{\mathrm{ctx}}(H)$ for the set of context states. For a ray realization ρ, the *geometric state space*

$$\mathcal{S}_{\mathrm{geom}}(H, \rho) = \{s \in \mathcal{S}(H) : s(i) + s(j) \leq 1 \text{ whenever } i \neq j,\ v_i \perp v_j\} \tag{4}$$

additionally enforces orthogonality exclusions on uncovered pairs [22]; $\mathcal{S}_{\mathrm{geom}}(H, \rho) = \mathcal{S}(H)$ when ρ is edge-faithful. A state family $\Omega \subseteq \mathcal{S}(H)$ is *unital* if every $v \in V$ has $s(v) = 1$ for some $s \in \Omega$, and *vertex separating* if distinct $u, v \in V$ have $s(u) \neq s(v)$ for some $s \in \Omega$, inducing the partition set representation $A_i = \{s \in \Omega : s(i) = 1\}$. On an event poset $(L, \leq)$, Ω is *event separating* if distinct $x, y \in L$ have distinct profiles on Ω, and *order determining* if $x \leq y \iff (\forall s \in \Omega,\ s(x) \leq s(y))$.

Definition 1 (Chromatic separability and extendable states)**.** For chromatically complete H ($\mathcal{C}_d(H) \neq \varnothing$), vertices $u, v \in V$ are *chromatically inseparable*, written $u \sim_c v$, if

$$c(u) = c(v) \qquad \text{for every } c \in \mathcal{C}_d(H), \tag{5}$$

an *implicit identity* in graph coloring [10, Def. 3.8]. We call H *chromatically separating* if $u \sim_c v \implies u = v$. A state $s \in \mathcal{S}(H)$ is *extendable*, written $s \in \mathcal{S}_{\text{ext}}(H)$, if $\operatorname{supp} s = \{v \in V : s(v) = 1\}$ is a color class of some $c \in \mathcal{C}_d(H)$ [5–7].

Physically, $u \sim_c v$ means that every noncontextual eigenvalue assignment gives u and v the same eigenvalue, and an extendable state is exactly a truth assignment of the form "the outcome with eigenvalue λ_a occurs" for some noncontextual eigenvalue assignment c and eigenvalue λ_a.

Theorem 2 (Completion criterion)**.** *For $s \in \mathcal{S}(H)$ with $S = \operatorname{supp} s$, the following are equivalent:*

1. *$s \in \mathcal{S}_{\text{ext}}(H)$;*
2. *$\chi(\Gamma(H)[V \setminus S]) \leq d-1$ (so S is a critical independent set [10, Def. 3.3]);*
3. *s belongs to d states in $\mathcal{S}(H)$ with pairwise disjoint supports partitioning V;*
4. *s belongs to $d-1$ states in $\mathcal{S}(H)$ with pairwise disjoint supports.*

For $d = 3$, $s \notin \mathcal{S}_{\text{ext}}(H)$ if and only if $\Gamma(H)[V \setminus S]$ contains an odd cycle.

Proof. A strong d-coloring with color class S leaves a $(d-1)$-coloring on $V \setminus S$. Conversely, any $(d-1)$-coloring of $\Gamma(H)[V \setminus S]$ combined with a dth color on S is a strong d-coloring because each context meets S once and forms a $(d-1)$-clique in $V \setminus S$; its color indicators yield (3). Conversely, d states with disjoint supports $S_1, \ldots, S_d$ partitioning V define the strong d-coloring $c(v) = k$ for $v \in S_k$, since each context meets each S_k exactly once; hence (3) $\implies$ (1). Trivially (3) $\implies$ (4), and given $d-1$ disjoint state supports $S_1, \ldots, S_{d-1}$, the complement $s_d(v) = 1 - \sum_{k=1}^{d-1} s_k(v)$ is binary and sums to $d-(d-1) = 1$ on every context, giving (3). For $d = 3$, $\chi(\Gamma(H)[V \setminus S]) \leq 2$ holds if and only if $\Gamma(H)[V \setminus S]$ is bipartite (odd-cycle free). □

Applying the equivalence (3) $\iff$ (4) of Theorem 2 globally without anchoring a state s yields the characterization of chromatic completeness:

Lemma 3 (Disjoint states and strong coloring)**.** *A finite d-uniform hypergraph H satisfies $\chi(H) = d$ if and only if $\mathcal{S}(H)$ contains $d-1$ states with pairwise disjoint supports.*

Corollary 4 (The separation hierarchy)**.** *For chromatically complete H and $u, v \in V$,*

$$u \sim_c v \iff s(u) = s(v) \text{ for all } s \in \mathcal{S}_{\text{ext}}(H). \quad (6)$$

Thus inseparability by $\mathcal{S}(H)$ implies chromatic inseparability ($\sim_c$), equivalently chromatic separation implies separation by $\mathcal{S}(H)$, whereas the converse fails even for faithful real ray configurations. Moreover, $\mathcal{S}_{\text{ext}}(H)$ is always unital whenever $\mathcal{C}_d(H) \neq \varnothing$.

Proof. The indicator states $s_a(v) = \mathbf{1}_{\{c(v)=a\}}$ over all $c \in \mathcal{C}_d(H)$ and $a \in \{0, \ldots, d-1\}$ are precisely $\mathcal{S}_{\text{ext}}(H) \subseteq \mathcal{S}(H)$. For each fixed coloring, the d supports partition V. The inclusion need not be strict: for a single d-element context every state is extendable. Strict inclusion is an additional property established for the specified examples. □

Proposition 5 (Canonical chromatic quotient and closure test)**.** *Let H be chromatically complete.*

1. *The relation $\sim_c$ is an equivalence relation with no two equivalent vertices in a common context. The quotient $\overline{H} = (V/\!\sim_c, \overline{E})$ is d-uniform and chromatically separating, and pullback along $V \to V/\!\sim_c$ induces a bijection $\mathcal{C}_d(\overline{H}) \to \mathcal{C}_d(H)$.*
2. *For $u \neq v$, adjoining one context $\{u, v, w_1, \ldots, w_{d-2}\}$ with fresh vertices w_i yields chromatic number $d+1$ if and only if $u \sim_c v$, and d otherwise. Consequently, the* separable chromatic number *[5] of any nontrivial graph satisfies $\chi_{\text{sep}} \in \{\chi, \chi+1\}$.*

Proof. (1) Adjacent vertices receive distinct colors, so each context projects injectively to $V/\!\sim_c$, and every $c \in \mathcal{C}_d(H)$ factors uniquely through $V/\!\sim_c$. (2) A coloring $c \in \mathcal{C}_d(H)$ extends to the fresh context if and only if $c(u) \neq c(v)$; if $u \sim_c v$, recoloring u with a $(d+1)$th color separates (u, v) while preserving all old context inequalities. □

The quotient preserves colorings, not an injective realization. Closing an inseparable pair is likewise abstract: faithful coordinates make the pair nonorthogonal, so the added context cannot retain those rays.

Put $\mathcal{P}_{\text{ext}} = \operatorname{conv} \mathcal{S}_{\text{ext}}$ and $\mathcal{P} = \operatorname{conv} \mathcal{S}$. Thus $\mathcal{P}$ is the set of probability vectors of noncontextual truth-assignment models, and $\mathcal{P}_{\text{ext}}$ is the subset produced by models that choose a noncontextual eigenvalue assignment c and an eigenvalue λ_a at random and report, in every context, the outcome with eigenvalue λ_a. A binary vector cannot be a convex combination of distinct binary vectors, since each positive-weight term must agree at every coordinate. Consequently,

$$\mathcal{P}_{\text{ext}} = \mathcal{P} \iff \mathcal{S}_{\text{ext}} = \mathcal{S}. \quad (7)$$

This is stronger than equality of vertex-separation kernels: the pentagon has separating colors yet a nonextendable state. Pair identities alone do not describe all completion constraints.

C. Fractional coloring and fresh-vertex context completion

Write $\alpha(G)$ for the independence number and $\chi_f(G)$ for the fractional chromatic number of $G = \Gamma(H)$.

Proposition 6 (Fractional threshold and uniform state mixture)**.** *For any finite d-uniform context hypergraph H,*

$$\chi_f(\Gamma(H)) = d \quad \Longleftrightarrow \quad \frac{1}{d}\mathbf{1} \in \mathcal{P}(H) = \operatorname{conv}\mathcal{S}(H). \quad (8)$$

Proof. A state mixture $\sum_{s\in\mathcal{S}(H)} \mu_s s = \frac{1}{d}\mathbf{1}$ assigns weight $d\mu_s$ to the independent set $\operatorname{supp} s$, covering each vertex with weight 1 at total weight d, so $\chi_f(\Gamma(H)) \le d$; since every context is a d-clique, $\chi_f(\Gamma(H)) \ge \omega(\Gamma(H)) \ge d$. Conversely, a fractional coloring $\{w_I\}$ of total weight d obeys $d \le \sum_{v\in e}\sum_{I\ni v} w_I = \sum_I w_I|I\cap e| \le \sum_I w_I = d$ for each $e \in E$, forcing $|I \cap e| = 1$ whenever $w_I > 0$, so $\{w_I/d\}$ is a state mixture equal to $\frac{1}{d}\mathbf{1}$. □

Proposition 7 (Fresh-vertex context completion)**.** *Let G be a finite core graph with designated d-clique base contexts ($d \ge 3$). Complete selected uncovered edges $\{u, v\}$ into d-cliques using $d-2$ degree-one fresh vertices per edge, covering all core vertices and edges. For the resulting hypergraph H and $G' = \Gamma(H)$,*

$$\begin{aligned} \chi(G') &= \max\{d, \chi(G)\}, \\ \chi_f(G') &= \max\{d, \chi_f(G)\}, \\ |\mathcal{S}(H)| &= \sum_S (d-2)^{m(S)}, \end{aligned} \quad (9)$$

where S ranges over independent core supports meeting each base context once and $m(S)$ counts completed edges missed by S.

Proof. Any k-coloring ($k \ge d$) or measurable fractional palette of size $k = \max\{d, \chi_f(G)\}$ assigns disjoint colors/sets of measure 1 to u and v, leaving $k - 2 \ge d - 2$ colors (or measure $k - 2 \ge d - 2$) for the $d - 2$ fresh vertices. A core support S hitting $\{u, v\}$ forces its fresh vertices to 0, whereas a missed edge allows $d - 2$ choices of a single fresh vertex set to 1. □

Table I summarizes the principal configurations studied below.

The logical situation has two distinct branches. If a hypergraph has no admissible two-valued context state, it cannot admit a strong d-coloring, because the color classes of such a coloring would supply d admissible states. The converse implication fails: a noncontextual assignment of d eigenvalues to the outcomes may be impossible while ordinary states, including a unital separating family, remain available.

When a noncontextual eigenvalue assignment does exist, its color classes provide a nonempty unital family of extendable states inside the full context-state space. A particular ordinary state belongs to this restricted family precisely when deleting its support leaves a $(d-1)$-colorable graph. Chromatic identities describe pairs having equal eigenvalues in every noncontextual eigenvalue assignment, and hence equal values in every state of that restricted family; they need not hold throughout the ordinary state space. These assignment questions have to be distinguished from the separate requirements of an injective geometric realization, a representation of event order, and a classical mixture reproducing a specified Born probability vector.

III. INDEPENDENCE OF COLORING AND COORDINATIZATION, PETERSEN OBSTRUCTIONS, AND COMPLETED YU–OH

A. Small abstract examples and four-quadrant independence

The non-linear hypergraph $H_{4,2}$ with contexts $\{1, 2, 3\}$ and $\{1, 2, 4\}$ has three states $\{1\}$, $\{2\}$, $\{3, 4\}$ (all extendable), forces $3 \sim_c 4$ and $s(3) = s(4)$, and has no injective $\mathbb{C}^3$ realization because two orthogonal rays determine a unique orthogonal-complement ray.

Proposition 8 (Smallest linear chromatically inseparable hypergraph)**.** *The linear 3-uniform hypergraph $H_{6,3}$ with contexts $\{1, 2, 3\}$, $\{3, 4, 5\}$, $\{5, 6, 1\}$ of Berge girth three has four unital separating states $\{1, 4\}$, $\{2, 5\}$, $\{3, 6\}$, $\{2, 4, 6\}$, a unique strong 3-coloring up to permutation with inseparable classes $\{1, 4\}$, $\{2, 5\}$, $\{3, 6\}$, and three extendable states (as $V \setminus \{2, 4, 6\} = \{1, 3, 5\}$ is a triangle). No linear 3-uniform hypergraph with $|V| < 6$ or $|E| < 3$ has a chromatically inseparable pair, and $H_{6,3}$ admits no injective realization in $\mathbb{C}^3$.*

Proof. The triangle $\{1, 3, 5\}$ in $\Gamma(H_{6,3})$ forces three distinct colors on $1, 3, 5$, whence $c(4) = c(1)$, $c(2) = c(5)$, and $c(6) = c(3)$. Similarly, in $\mathbb{C}^3$, $\{v_1, v_3, v_5\}$ forms an orthonormal basis, forcing $[v_4] = [v_1]$, $[v_2] = [v_5]$, and $[v_6] = [v_3]$. With $|E| \le 2$, independent permutations of unshared vertices separate all pairs; by inclusion–exclusion, three linear 3-contexts require $|V| \ge 6$. □

Proposition 9 (Clique number at Berge girth at least four)**.** *If $H = (V, E)$ is an n-uniform linear hypergraph ($n \ge 3$) of Berge girth $g(H) \ge 4$, every clique $K \subseteq V$ of size $|K| \ge 3$ in $\Gamma(H)$ is contained in a unique context $e \in E$. Hence $\omega(\Gamma(H)) = n$, and chromatic incompleteness ($\chi(H) > n$) is equivalent to the chromatic–clique gap $\chi(\Gamma(H)) > \omega(\Gamma(H))$.*

Proof. For any three vertices $u, v, w \in K$, pairwise adjacency gives contexts $e_1 \supseteq \{u, v\}$, $e_2 \supseteq \{v, w\}$, $e_3 \supseteq \{w, u\}$. Distinct e_1, e_2, e_3 would form a Berge 3-cycle, contradicting $g(H) \ge 4$. By linearity, the shared context e containing $\{u, v, w\}$ is unique for $\{u, v\}$ and thus contains all of K. □

Proposition 10 (Logical independence of coloring and coordinatization)**.** *For finite 3-uniform context hypergraphs, all four combinations of chromatic completeness ($\chi(H) = 3$) and injective coordinatizability in $\mathbb{C}^3$ occur even within the orthomodular-lattice admissible class of linear hypergraphs of Berge girth $g(H) \ge 5$ [8, 20] with unital separating state spaces:*

TABLE I. Principal configurations. FOR means injective and edge-faithful; g is Berge girth. Counts refer to all context states and all color-class states.

Configuration	d	$\lvert V\rvert/\lvert E\rvert$	g	χ	χ_f	$\lvert\mathcal{S}\rvert$	$\lvert\mathcal{S}_{\text{ext}}\rvert$	Geometry and principal distinction
One triad	3	3/1	∞	3	3	3	3	Real FOR; colors separate.
$H_{6,3}$	3	6/3	3	3	3	4	3	No injective complex realization.
P_5	3	10/5	5	3	3	11	10	Real FOR; colors separate; ternary exclusion.
H_8	3	14/8	5	3	3	11	7	Repeated-ray assignments; no complex injection.
H_9	3	15/9	5	4	3	8	0	No complex ray assignment; context-critical.
G_{32}	3	15/10	5	4	3	6	0	No complex ray assignment.
$H_{17,10}$	3	17/10	5	3	3	18	10	Real FOR; two color-identical pairs.
$H_{21,13}$	3	21/13	5	3	3	21	6	Real FOR; 21 color-identical pairs.
Completed Yu–Oh	3	25/16	5	4	35/11	24	0	Real FOR; context-critical.
H_{37}	3	37/24	5	4	3	153	0	Complex FOR; no real injection; OML minima 6/7/18.
$H_{\text{YO}}^{(4)}$	4	26/16	2	5	46/11	25	0	Real FOR; common-apex suspension, nonlinear.
$C_{36,18}$	4	36/18	3	5	9/2	0	0	Faithful linear KS completion.
H_{65}	4	65/29	3	5	17/4	393,216	0	Real FOR; false atom order and compound collapse.

1. $\chi = 3$, *injectively realizable in* $\mathbb{R}^3$*: a single context* $C_1 = \{\{1,2,3\}\}$ *(or* $P_5, H_{17,10}, H_{21,13}$*);*

2. $\chi = 3$, *not injectively realizable in* $\mathbb{C}^3$*: the* 8*-context Petersen restriction* H_8 *(Theorem 11);*

3. $\chi = 4$, *edge-faithfully realizable in* $\mathbb{R}^3$*: completed Yu–Oh* H_{YO}^{+} *(Theorem 13);*

4. $\chi = 4$, *admitting no orthogonal ray assignment in* $\mathbb{C}^3$*: Greechie's* G_{32} *and* H_9 *(Theorem 11).*

Injectivity is essential to quadrant (2): any strong n-coloring c yields a noninjective ray assignment $i \mapsto [e_{c(i)+1}] \in \mathbb{P}(\mathbb{F}^n)$. At Berge girth four (an orthomodular poset that is not a lattice), $E_{\square} = \{\{1,2,3\}, \{3,4,5\}, \{5,6,7\}, \{7,8,1\}\}$ also satisfies $\chi = 3$ while forcing $[v_3] = [v_7] \perp \text{span}(v_1, v_5)$ whenever $[v_1] \neq [v_5]$.

B. Greechie's G_{32}, multipole forcing, and correction of the JMP spectral argument

Greechie's G_{32} [8, Fig. 6] has 15 vertices and ten contexts

$$E(G_{32}) = \{123, 345, 567, 789, 9\,10\,11, \\ 11\,12\,1, 4\,10\,13, 6\,12\,14, \\ 8\,2\,15, 13\,14\,15\}. \tag{10}$$

Each vertex lies in two contexts, so G_{32} is dual to the Petersen graph P, $\Gamma(G_{32}) = L(P)$, $\text{Aut}(G_{32}) \cong S_5$, and $\chi(G_{32}) = \chi'(P) = 4$. Its six two-valued state supports satisfy

$$|\mathcal{S}(G_{32})| = 6, \qquad |S_\alpha| = 5, \qquad |S_\alpha \cap S_\beta| = 1 \ (\alpha \neq \beta). \tag{11}$$

These supports yield the $\binom{6}{2} = 15$ two-element vertex profiles $B_i = \{\alpha : i \in S_\alpha\}$. Their nonempty pairwise intersections exclude even two disjoint states, confirming $\chi(G_{32}) = 4$ by Lemma 3.

In Appendix B of Shekarriz and Svozil [6], the chromatic obstruction $\chi(G_{32}) = 4 > 3$ was invoked to argue against a 3-dimensional realization of G_{32}. As Proposition 10 demonstrates via H_{YO}^{+} ($\chi = 4$, yet faithfully realizable in $\mathbb{R}^3$), $\chi(H) > 3$ rules out only a context-independent 3-value spectral assignment—a noncontextual assignment of the eigenvalues of uniform maximal observables to their outcomes—*not* geometric rays. A genuine geometric exclusion requires the projection identity: for an orthonormal basis $\{x, y, z\} \subset \mathbb{C}^3$ with $u \perp x$ and $w \perp z$,

$$\langle u, w\rangle = \langle u, y\rangle\langle y, w\rangle. \tag{12}$$

Theorem 11 (Multipole forcing and minimal complex obstructions)**.** *Let* $H_8 = G_{32} \setminus \{\{8,2,15\}, \{13,14,15\}\}$ *(*14 *vertices,* 8 *contexts) and* $H_9 = G_{32} \setminus \{\{13,14,15\}\}$ *(*15 *vertices,* 9 *contexts).*

1. *In every orthogonal ray assignment of* H_8 *in* $\mathbb{C}^3$ *(allowing repeated rays), the terminal rays satisfy*

$$[v_2] = [v_8] \qquad \textit{and} \qquad [v_{13}] = [v_{14}], \tag{13}$$

and either $[v_1] = [v_4] = [v_9]$ *or* $[v_3] = [v_7] = [v_{12}]$*. Thus* H_8 *has no injective realization in* $\mathbb{C}^3$*.*

2. H_8 *is chromatically complete (*$\chi(H_8) = 3$*) with* 11 *unital separating states, of which* 7 *are extendable (the* 4 *nonextendable states leave odd cycles in* $\Gamma(H_8)$*; see Table S38). Its* 3 *normalized* 3*-colorings (fixing* $(c(1), c(2), c(3)) = (0, 1, 2)$*) are listed in Table S39, forcing the* 6 *chromatically inseparable pairs* $(1,9)$*,* $(2,8)$*,* $(3,7)$*,* $(4,6)$*,* $(10,12)$*,* $(13,14)$*; deleting any single context of* H_8 *removes all forced*

color identities (leaving $6, 20, 20, 6, 20, 20, 6, 6$ *normalized colorings on the remaining active vertices, up to global color permutation). Isolated vertices are dropped: retaining the isolated vertex created by deleting* $\{1,2,3\}$*,* $\{7,8,9\}$*,* $\{4,10,13\}$*, or* $\{6,12,14\}$ *multiplies the corresponding count by three, from* 6 *to* 18*.*

3. *Neither* H_9*,* G_{32}*, nor the* 12*-vertex,* 21*-edge subgraph* Γ_0 *of* $\Gamma(H_9)$ *on* $\{1,\dots,12\}$ *admits any orthogonal ray assignment in* $\mathbb{C}^3$*. Moreover,* H_9 *is context-minimal and* Γ_0 *is edge-minimal: deleting any context of* H_9 *or edge of* Γ_0 *yields a* 3*-colorable structure realizable in* $\mathbb{R}^3$ *by standard basis rays.*

Proof. (1) Let $v_1,\dots,v_{14} \in \mathbb{C}^3$ be unit vectors for H_8. In $\{v_1,v_2,v_3\}$, $v_{11}, v_{12} \perp v_1$ and $v_4, v_5 \perp v_3$, so Eq. (12) gives $\langle x,y\rangle = \langle x,v_2\rangle\langle v_2,y\rangle$ for $x \in \{v_{11},v_{12}\}, y \in \{v_4,v_5\}$. Expanding v_{12} in $\{v_5,v_6,v_7\}$ ($v_{12} \perp v_6$) and v_4 in $\{v_9,v_{10},v_{11}\}$ ($v_4 \perp v_{10}$) yields $v_7\langle v_7,v_{12}\rangle = v_{12} - v_5\langle v_5,v_{12}\rangle$ and $v_9\langle v_9,v_4\rangle = v_4 - v_{11}\langle v_{11},v_4\rangle$. Since $v_7 \perp v_9$, $v_{12} \perp v_{11}$, and $v_5 \perp v_4$, their inner product vanishes:

$$\begin{aligned} 0 &= \langle v_{12},v_4\rangle + \langle v_{12},v_5\rangle\langle v_5,v_{11}\rangle\langle v_{11},v_4\rangle \\ &= \langle v_{12},v_2\rangle\langle v_2,v_4\rangle\big(1+|\langle v_2,v_5\rangle|^2|\langle v_2,v_{11}\rangle|^2\big). \end{aligned} \tag{14}$$

Thus $\langle v_2,v_4\rangle = 0$ or $\langle v_{12},v_2\rangle = 0$. If $\langle v_2,v_4\rangle = 0$, set $(v_1,v_3,v_2) = (e_1,e_2,e_3)$; then $v_4 \perp v_2, v_3 \implies [v_4] = [e_1]$, $\{3,4,5\} \implies [v_5] = [e_3] = [v_2]$, $\{9,10,11\}$ with $v_{10}, v_{11} \perp e_1$ forces $[v_9] = [e_1]$, $\{5,6,7\}$ and $\{7,8,9\}$ force $[v_7] = [e_2], [v_6] = [e_1], [v_8] = [e_3] = [v_2]$, and $\{4,10,13\}$, $\{9,10,11\}$, $\{11,12,1\}$, $\{6,12,14\}$ in $\mathrm{span}(e_2,e_3)$ force $[v_{13}] = [v_{11}] = [v_{14}]$ and $[v_1] = [v_4] = [v_9]$. Reflection symmetry gives the second branch when $\langle v_{12},v_2\rangle = 0$.

(2) In H_8, fixing $(c(1),c(2),c(3)) = (0,1,2)$ leaves 16 choices along $345, 567, 789, 9\,10\,11$, of which 13 fail on $11\,12\,1, 4\,10\,13, 6\,12\,14$, leaving c_1, c_2, c_3 in Table S39. Equal columns identify the 6 pairs.

(3) Both H_9 (hence G_{32}) and Γ_0 add the orthogonality $v_2 \perp v_8$ to H_8, contradicting $[v_2] = [v_8]$. For minimality, deleting $\{8,2,15\}$ from H_9 leaves H_8 ($\chi = 3$); deleting $\{4,10,13\}$ or $\{6,12,14\}$ is isomorphic by D_6 symmetry to deleting $\{8,2,15\}$; and deleting any ring context (say $\{1,2,3\}$) allows the 3-coloring $(c(1),\dots,c(15)) = (2,1,2,0,1,0,2,0,1,2,0,1,1,2,2)$. On Γ_0, the following deletions cover all 21 edges up to D_6 symmetry: $\{1,2\}$ (colored by 001020120121), $\{2,3\}$ (011020120121), $\{1,3\}$ (010120102012), and $\{4,10\}$ (012010201021), each yielding a standard-basis $\mathbb{R}^3$ assignment $i \mapsto [e_{c(i)+1}]$. Quantitative robustness bounds $1/14400 < \Delta_3 < 0.927496$ are proved in Supplement Sec. S3. □

The complete normalized coloring lists for H_8 and $H_{17,10}$ are in the supplement.

Corollary 12 (Blanuša and Isaacs dot-product snarks)**.** *If a cubic graph* G *admits no orthogonal vector edge coloring in* $\mathbb{C}^3$*, neither does any Isaacs dot product* $G \cdot P$ *[23]. In particular, both* 18*-vertex Blanuša snarks* $B_1, B_2 = P \cdot P$ *[24] and all iterated dot products* $X = P \cdot P \cdots P$ *satisfy* $\chi(H(X)) = 4 > 3 = \omega(L(X))$ *and admit no orthogonal ray assignment in* $\mathbb{C}^3$*.*

Proof. In $G \cdot P$, the embedded copy of $P \setminus \{u,v\}$ has dual hypergraph H_8, so Theorem 11(1) forces $[v_2] = [v_8]$ and $[v_{13}] = [v_{14}]$ on the two pairs of cut edges, inducing a valid orthogonal ray assignment on G. □

C. Completed Yu–Oh H^+_{YO}, base-coloring partition, and context criticality

Completing each of the 12 uncovered orthogonal pairs of the 13-ray Yu–Oh set [9] ($r_1,\dots,r_{13} \in \mathbb{R}^3$, consisting of $z_1 = r_2, z_2 = r_8, z_3 = r_5$, $y_1^{\pm} = r_{1,3}, y_2^{\pm} = r_{7,9}, y_3^{\pm} = r_{4,6}$, and diagonals $h_0 = r_{12}, h_1 = r_{11}, h_2 = r_{10}, h_3 = r_{13}$) by cross-product rays $r_{14},\dots,r_{25}$ (Table S2) yields the 25-ray, 16-context linear hypergraph H^+_{YO} of Berge girth five [7, 25]:

$$\begin{aligned} E^+_{\mathrm{YO}} &= \{123,789,456,285\} \cup \bigcup_{h=10}^{13} \mathcal{E}_h, \\ \mathcal{E}_h &= \{\{x,h,w_{xh}\} : x \in N(h)\}, \end{aligned} \tag{15}$$

All 25 vectors, 16 contexts, and restriction-membership flags are tabulated in the supplement. Here the neighborhoods are $N(10) = \{1,9,4\}$, $N(11) = \{3,7,4\}$, $N(12) = \{3,9,6\}$ and $N(13) = \{1,7,6\}$, and each w_{xh} is the unique completing ray. It has a real FOR in $\mathbb{R}^3$ (48 orthogonal pairs) and 24 unital separating states (each of the 12 binary assignments on the four base contexts leaves uniquely one diagonal h_j with all three neighbors zero, allowing $s(h_j) \in \{0,1\}$ and hence $12 + 4 \times 3 = 24$ states). Completed Yu–Oh is therefore a faithfully realized real configuration in which noncontextual truth assignments survive, enough of them to separate every pair of rays, while by the following theorem no noncontextual assignment of three eigenvalues to its outcomes exists.

Theorem 13 (Base-coloring partition and context criticality of H^+_{YO})**.** *The completed Yu–Oh hypergraph* H^+_{YO} *satisfies* $\omega(\Gamma(H^+_{\mathrm{YO}})) = 3$*,* $\chi_f(\Gamma(H^+_{\mathrm{YO}})) = 35/11 > 3 = \chi_f(\Gamma(G_{32}))$*, and* $\chi(H^+_{\mathrm{YO}}) = 4$*, and is context-critical for* $\chi > 3$*:*

1. *Fixing* $(c(2),c(8),c(5)) = (0,1,2)$*, the four base contexts* $\{1,2,3\}$*,* $\{7,8,9\}$*,* $\{4,5,6\}$*,* $\{2,8,5\}$ *admit eight strong* 3*-colorings indexed by* $(b_1,b_2,b_3) \in \{0,1\}^3$*:*

$$\begin{aligned} (c(1),c(3)) &= (2-b_1, 1+b_1), \\ (c(7),c(9)) &= (2-2b_2, 2b_2), \\ (c(4),c(6)) &= (1-b_3, b_3). \end{aligned} \tag{16}$$

Each antipodal pair of (b_1,b_2,b_3) *makes the neighborhood* $N(h_j)$ *of* uniquely one *diagonal ray* $h_j \in$

$\{10, 11, 12, 13\}$ *rainbow* ($\{0, 1, 2\}$) *while the other three neighborhoods are* 2*-colored:*

Antipodal binary patterns	*Blocked diagonal*
$101, 010$	12
$110, 001$	11
$000, 111$	10
$011, 100$	13

(17)

2. *Consequently,* $\chi(H_{\mathrm{YO}}^{+}) = 4$*, whereas deleting any single context* $e \in E_{\mathrm{YO}}^{+}$ *yields* $\chi(H_{\mathrm{YO}}^{+} \setminus \{e\}) = 3$.

Proof. Direct check of Eqs. (16)–(17) shows that every base 3-coloring blocks one h_j, so $\chi(H_{\mathrm{YO}}^{+}) \geq 4$, matched by the four-coloring $(c(1), \ldots, c(25)) = (2, 0, 1, 1, 2, 0, 2, 1, 0, 3, 0, 2, 1, 2, 1, 1, 1, 0, 0, 0, 0, 2, 1, 2, 0)$, in which only $h_2 = 10$ receives the fourth color. Deleting a sector context $\{x, h_j, w\}$ frees h_j in its antipodal base colorings by setting $c(h_j) = c(x)$; deleting $\{2, 8, 5\}$ allows $c(2) = c(8) = c(5) = 0$; deleting $\{1, 2, 3\}$ allows $(c(1), c(2), c(3)) = (0, 0, 1)$, $(c(7), c(8), c(9)) = (2, 1, 0)$, $(c(4), c(5), c(6)) = (1, 2, 0)$, and $(c(10), c(11), c(12), c(13)) = (2, 0, 2, 1)$; the cases $\{7, 8, 9\}$ and $\{4, 5, 6\}$ follow by the threefold cyclic coordinate symmetry of the Yu–Oh set. For χ_f, weight $1/2$ on the 6 states of G_{32} gives $\chi_f(G_{32}) = 3$, while Proposition 7 preserves the 13-ray value $\chi_f = 35/11$ [26]. □

IV. EXACT-PALETTE CHROMATIC SEPARABILITY AND LOGICAL GADGETS IN DIMENSIONS THREE AND FOUR

From now on we assume that noncontextual assignments of d eigenvalues exist ($\chi(H) = d$) and ask what they imply for the relations between distinct outcomes. Such assignments may force outcomes to share or to avoid an eigenvalue although ordinary truth assignments permit otherwise; these forced relations are their logical content.

A. Kochen–Specker antecedents and palette-invariant implications

Three historical Kochen–Specker constructions clarify the hierarchy of logical gadgets:

1. *1965 abstract TIFS precursor:* Identifying repeated labels v_1 and v_2 in Kochen and Specker's 1965 Fig. 1 [27, p. 182] yields an 11-vertex, 7-context hypergraph $\{134, 325, 2\,9\,11, 11\,1\,10, 479, 5\,8\,10, 678\}$ implementing $s(1) = 1 \implies s(2) = 0$. It has only four states ($\{1, 5, 6, 9\}$, $\{2, 4, 6, 10\}$, $\{3, 6, 9, 10\}$, $\{4, 5, 6, 11\}$), all with $s(6) = 1$ and $s(7) = s(8) = 0$, so $\chi > 3$; moreover, 1 and 9 are both orthogonal to 4 and 11, forcing $[v_1] = [v_9]$ in $\mathbb{C}^3$ (Supplement Sec. S3).

2. *1967 directed TITS* Γ_1*:* The 10-vertex graph Γ_1 of Lemma 1 in Ref. [4, pp. 68, 70] implements the one-way implication $s(a_0) = 1 \implies s(a_9) = 1$. Completing its six uncompleted edges by fresh rays $23, 19, 18, 15, 14, 5$ yields the 16-ray, 9-context configuration $I_{16,9}$ in Fig. 2(d) (explicit label map in Supplement Sec. S3).

3. *1967 state-inseparable* Γ_3*:* In Ref. [4, p. 70], Γ_3 forces $s(a) = s(b)$ in *every* two-valued state (a, b are nonseparated by $\mathcal{S}(H)$). By contrast, our chromatically inseparable examples below have $\mathcal{S}(H)$ separating all vertices, with $c(u) = c(v)$ forced only on $\mathcal{S}_{\mathrm{ext}}(H)$.

For distinct nonadjacent terminals $u, v \in V$, an ordinary TIFS [28] obeys $s(u) = 1 \implies s(v) = 0$ (equivalently $s(u) + s(v) \leq 1$, which is always bidirectional), and a directed ordinary TITS obeys $s(u) = 1 \implies s(v) = 1$. When $\mathcal{C}_d(H) \neq \varnothing$, an *unanchored chromatic TIFS* obeys $c(u) = 0 \implies c(v) \in \{1, \ldots, d-1\}$, and an *unanchored chromatic TITS* obeys $c(u) = 0 \implies c(v) = 0$.

Proposition 14 (Palette symmetry and purely chromatic gadgets)**.** *For chromatically complete* H *and nonadjacent* $u, v \in V$*, the unanchored chromatic TIFS is equivalent to* $c(u) \neq c(v)$ *for all* $c \in \mathcal{C}_d(H)$ *(an* implicit edge *[10, Def. 3.1]), and the unanchored chromatic TITS is equivalent to* $u \sim_c v$ *(an* implicit identity*). Both chromatic relations are bidirectional. Ordinary TIFS and directed ordinary TITS entail their chromatic counterparts, and with only* $c(u) = 0$ *fixed, the allowed output set* $\{c(v) : c \in \mathcal{C}_d(H),\ c(u) = 0\}$ *is one of* $\{0\}$, $\{1, \ldots, d-1\}$, *or* $\{0, \ldots, d-1\}$.

Proof. Global palette permutations act transitively on $\{0, \ldots, d-1\}$ and, stabilizing 0, transitively on $\{1, \ldots, d-1\}$. Since every color class $s_a(x) = \mathbf{1}_{\{c(x)=a\}}$ is in $\mathcal{S}_{\mathrm{ext}}(H) \subseteq \mathcal{S}(H)$, any ordinary TIFS or one-way TITS forces $c(u) \neq c(v)$ or $c(u) = c(v)$, respectively. □

We call a chromatic TIFS *purely chromatic* if some $s \in \mathcal{S}(H) \setminus \mathcal{S}_{\mathrm{ext}}(H)$ satisfies $s(u) = s(v) = 1$, and a chromatic identity $u \sim_c v$ *two-way purely chromatic* if ordinary states in $\mathcal{S}(H) \setminus \mathcal{S}_{\mathrm{ext}}(H)$ realize both $(s(u), s(v)) = (1, 0)$ and $(0, 1)$, so neither directed ordinary TITS holds. Since a state with $s(u) = s(v) = 1$ cannot be a color class when $c(u) \neq c(v)$ for all c, and a state with $s(u) \neq s(v)$ cannot be one when $u \sim_c v$, the qualifier $\mathcal{S}(H) \setminus \mathcal{S}_{\mathrm{ext}}(H)$ is automatic: a chromatic TIFS is purely chromatic exactly when the ordinary TIFS fails, and a chromatic identity is two-way purely chromatic exactly when both directed ordinary TITS fail.

B. Faithful Yu–Oh restrictions $H_{17,10}$ and $H_{21,13}$ and exhaustive inventory search

The base-coloring partition of H_{YO}^{+} in Theorem 13 directly produces chromatically inseparable faithful restrictions when diagonal sectors are omitted. Retain the

four base contexts $\{1,2,3\}$, $\{7,8,9\}$, $\{4,5,6\}$, $\{2,8,5\}$ of Eq. (15). Keeping the two diagonal sectors at 10 and 11 defines $H_{17,10}$ on 17 rays and 10 contexts [Fig. 1(a)]; keeping the three sectors at $10, 11, 13$ defines $H_{21,13}$ on 21 rays and 13 contexts [Fig. 1(b)]. Deleting $\{4,5,6\}$, $\{4,11,24\}$, $\{4,10,18\}$ from $H_{17,10}$ recovers the 13-ray, 7-context completed Specker bug $B_{13,7}$ [Fig. 2(b)].

Theorem 15 (Chromatic inseparability in $H_{17,10}$ and $H_{21,13}$)**.** *Both $H_{17,10}$ and $H_{21,13}$ are linear of Berge girth five and inherit injective edge-faithful realizations in $\mathbb{R}^3$ from H_{YO}^{+}:*

1. *$H_{17,10}$ has 18 unital separating states, 4 normalized strong 3-colorings (Table S41), and 10 extendable states. Its only distinct chromatically inseparable pairs are $(10,24)$ and $(11,18)$. The ordinary implications $10 \Rightarrow 24$ and $11 \Rightarrow 18$ already hold, but their converses fail; these identities are not two-way purely chromatic. Relative to the anchor $(c(2),c(8),c(5)) = (0,1,2)$, its four colorings restrict on $(9,10,11)$ to $(0,2,1)$, $(0,2,0)$, $(2,1,2)$, $(2,0,2)$, yielding the anchored implications*

$$\begin{aligned} c(9) = 0 &\implies c(11) \in \{0,1\}, \\ c(10) = 0 &\implies c(11) = 2. \end{aligned} \tag{18}$$

2. *$H_{21,13}$ has 21 unital separating states, 2 normalized strong 3-colorings, and 6 extendable states. Its nine chromatic signature classes $(c_1(v), c_2(v)) \in \{0,1,2\}^2$ are listed in Table S44, giving $3\binom{4}{2}+3\binom{2}{2} = 21$ chromatically inseparable pairs. Exactly 15 are two-way purely chromatic identities (Table S70). The remaining six pairs obey the directed ordinary implications*

$$\begin{aligned} &11 \Rightarrow 18, \quad 11 \Rightarrow 21, \quad 13 \Rightarrow 15, \\ &13 \Rightarrow 19, \quad 10 \Rightarrow 20, \quad 10 \Rightarrow 24, \end{aligned}$$

each with a false converse. This classification exhausts all 21 pairs; no distinct vertices are inseparable under all ordinary states.

Proof. By Theorem 13(1), among the 8 base colorings in Eq. (16), sector 11 excludes $(1,1,0)$, $(0,0,1)$ and sector 10 excludes $(0,0,0)$, $(1,1,1)$. The 4 surviving (b_1,b_2,b_3) extend uniquely to $H_{17,10}$ (Table S41): in each, $N(10) = \{1,9,4\}$ and $N(11) = \{3,7,4\}$ are 2-colored with $c(10) \neq c(4)$ and $c(11) \neq c(4)$, forcing $c(24) = c(10)$ in $\{4,11,24\}$ and $c(18) = c(11)$ in $\{4,10,18\}$. Adding sector 13 in $H_{21,13}$ additionally excludes $(0,1,1)$, $(1,0,0)$, leaving only $(1,0,1)$ and $(0,1,0)$, whose column signatures are listed in Table S44. Each retained diagonal adds 3 states to the 12 base states, giving $12+2\times3 = 18$ and $12+3\times3 = 21$ states (Supplement Sec. S2). Testing both directed truth patterns on these 21 states gives the exhaustive 15-versus-6 classification; Supplemental Table S70 supplies witnesses for all 15 two-way pairs, together with a reproducible enumeration procedure. □

The full chromatic signatures and both-direction truth witnesses are in the supplement. The witnesses are generated from the complete state families and checked against every context.

Physically, every noncontextual eigenvalue assignment on $H_{17,10}$ gives the nonorthogonal rays 10 and 24, and likewise 11 and 18, the same eigenvalue, although ordinary truth assignments distinguish them. In $H_{21,13}$, fifteen of the twenty-one forced identities are not even shadowed by a one-way truth implication: they exist only at the level of noncontextual eigenvalues.

Proposition 16 (Complete inventory classification)**.** *Among all $2^{16}-1 = 65{,}535$ nonempty context subsets $E' \subseteq E(H_{\mathrm{YO}}^{+})$ with inherited coordinates from Tables S1–S3, exactly 539 are connected and edge-faithful, and among these* precisely ten *are 3-colorable with a nontrivial chromatic inseparability class: the $\binom{4}{2} = 6$ two-sector restrictions (17 rays, 10 contexts, 18 states, 4 normalized colorings, 2 inseparable pairs) and the $\binom{4}{3} = 4$ three-sector restrictions (21 rays, 13 contexts, 21 states, 2 normalized colorings, 21 inseparable pairs). Thus 17 rays and 10 contexts are minimal for chromatic inseparability within the faithful restrictions of H_{YO}^{+}.*

Proof. Since H_{YO}^{+} is linear and edge-faithful, a subset E' is edge-faithful if and only if no omitted context contains two vertices of $V' = \bigcup E'$. Exact enumeration by two independent coloring algorithms over all 539 connected faithful subsets certifies the ten solutions listed in Table S71. Disconnected restrictions separate distinct components by independent palette permutations. □

The ten retained-sector cases and binary search masks are enumerated in the supplement.

C. Faithful three-dimensional TIFS, TITS, and purely chromatic gadgets

Figure 2 and Table II synthesize our faithful real 3-dimensional gadgets (coordinates and state/coloring tables in Supplement Secs. S1–S2):

- *Pentagon P_5 [Fig. 2(a)]:* Contexts $\{1,6,2\}$, $\{2,7,3\}$, $\{3,8,4\}$, $\{4,9,5\}$, $\{5,10,1\}$ have 5 normalized 3-colorings, 11 states, and 10 extendable states, separating all vertex pairs (showing that chromatic separation does not require $\mathcal{S}_{\mathrm{ext}} = \mathcal{S}$). Its degree-one vertices obey the purely chromatic ternary exclusion $\neg(c(6) = c(7) = c(9))$: assigning color 0 to $6,7,9$ forces $1,2,3,4,5$ into $\{1,2\}$ with $c(1) = c(3)$ and $c(4) = c(5)$, contradicting $\{4,9,5\}$, whereas the nonextendable state $\{6,7,8,9,10\}$ (whose complement is the 5-cycle 12345) assigns 1 to all three terminals.

- *Specker bug $B_{13,7}$ and completed Γ_1 $I_{16,9}$ [Fig. 2(b),(d)]:* $B_{13,7}$ is an ordinary and chromatic TIFS ($c(10) \neq c(11)$, overlap 1/9).

(a) $H_{17,10}$ (17 rays, 10 contexts, 2 inseparable pairs)

(b) $H_{21,13}$ (21 rays, 13 contexts, 21 inseparable pairs)




FIG. 1. Faithful three-dimensional Yu–Oh restrictions in $\mathbb{R}^3$, drawn in the common vertex layout of completed Yu–Oh. (a) $H_{17,10}$, omitting diagonal sectors $h_0 = 12$ and $h_3 = 13$, with four normalized strong 3-colorings and chromatically inseparable pairs $10 \sim_c 24$ and $11 \sim_c 18$; (b) $H_{21,13}$, omitting only $h_0 = 12$, with two normalized strong 3-colorings and 21 chromatically inseparable pairs across nine chromatic signature classes.

Adjoining $\{4, 11, 24\}$ and $\{4, 10, 18\}$ gives $I_{16,9}$ (deleting $\{4, 5, 6\}$ from $H_{17,10}$ with refitted faithful coordinates), where $c(10) \neq c(11)$ forces $c(4)$ to be the third color, so $c(24) = c(10)$ and $c(18) = c(11)$ via the one-way ordinary TITS $s(10) = 1 \implies s(24) = 1$ (overlap 25/27).

Proposition 17 (Purely chromatic binary TIFS $D_{16,9}$ and two-way identity $T_{19,11}$)**.** *Let* $D_{16,9} = H_{17,10} \setminus \{\{1, 10, 19\}\}$ *(with orphan* 19 *removed) and* $T_{19,11} = D_{16,9} \cup \{\{14, 26, 27\}, \{18, 26, 28\}\}$*, realized faithfully in* $\mathbb{R}^3$ *[Fig. 2(c),(e)]:*

1. *Every* $c \in \mathcal{C}_3(D_{16,9})$ *satisfies* $c(14) \neq c(18)$ *(*7 *normalized colorings,* 14 *extendable states), whereas the nonextendable state* $\{1, 5, 7, 14, 18, 23, 24\} \in \mathcal{S}(D_{16,9})$ *assigns* $s(14) = s(18) = 1$ *(squared terminal overlap* 16/2025*).*

2. *Every* $c \in \mathcal{C}_3(T_{19,11})$ *satisfies* $14 \sim_c 28$ *and* $18 \sim_c 27$ *(*7 *normalized colorings,* 25 *states,* 14 *extendable), while for both pairs, nonextendable states* $\{1, 5, 7, 14, 18, 23, 24\}$ *and* $\{1, 6, 8, 10, 11, 27, 28\}$ *realize* $(1, 0)$ *and* $(0, 1)$*, violating both directed ordinary TITS implications (squared overlap* 2009/2025*; Table S70).*

Proof. (1) If $c(14) = c(18) = 0$, set $c(3) = 1, c(11) = 2$ by $\{3, 11, 14\}$. Then $\{4, 11, 24\}$ and $\{4, 10, 18\}$ force $c(4) = 1$ and $c(10) = 2$. In $\{7, 8, 9\}$, $7 \perp 11$ and $9 \perp 10$ avoid color 2, forcing $c(8) = 2$. Then $\{1, 2, 3\}$ and $\{4, 5, 6\}$ make 2 and 5 avoid color 1, so $\{2, 8, 5\}$ forces $c(2) = c(5) = 0$, a contradiction. (2) In $T_{19,11}$, $c(14) \neq c(18)$ forces their common neighbor 26 ($r_{26} \parallel r_{14} \times r_{18}$) to take the third color, whence $c(27) = c(18)$ and $c(28) = c(14)$. □

D. Apex extensions and faithful four-dimensional linear gadgets

Proposition 18 (Apex extension)**.** *Embedding any* 3*-uniform hypergraph* H *realized in* $\mathbb{F}^3$ *into* $\mathbb{F}^d$ $(d \geq 3)$ *and appending* $d - 3$ *mutually orthogonal coordinate apices to every context preserves injectivity, edge-faithfulness,* $\chi(H^{(d)}) = \chi(H) + d - 3$*, normalized coloring count, and old chromatic inseparability classes (with apices as singletons), while adding* $d - 3$ *apex-only states (all extendable when* $\chi(H) = 3$*) to the zero-extended states of* H*.*

Proof. In every coloring of $H^{(d)}$, the $d - 3$ apices occupy $d-3$ distinct colors across all contexts, leaving a 3-coloring of H; similarly, any state either sets one apex to 1 (forcing all old vertices to 0) or sets all apices to 0 and restricts to $\mathcal{S}(H)$. □

Applying Proposition 18 at $d = 4$ to P_5, $B_{13,7}$, and $D_{16,9}$ yields the non-linear ($g = 2$) four-dimensional gadgets $W^{(4)}_{11,5}$, $B^{(4)}_{14,7}$, and $D^{(4)}_{17,9}$ in Fig. 3(a),(c),(e). To achieve *linearity* ($|e \cap e'| \leq 1$) in $\mathbb{R}^4$, we construct three faithful linear gadgets ($g = 3$) in Fig. 3(b),(d),(f) (Table II):

1. *Linear purely chromatic ternary gadget* $W^{(4)}_{16,6}$ *[Fig. 3(b)]:* Contexts $\{2, 0, 6, 1\}$, $\{2, 7, 13, 3\}$,




FIG. 2. Faithful real three-dimensional logical gadgets in $\mathbb{R}^3$. (a) Pentagon P_5 with purely chromatic ternary exclusion $\neg(c(6) = c(7) = c(9))$; (b) completed Specker bug $B_{13,7}$ (ordinary TIFS $c(10) \neq c(11)$); (c) purely chromatic binary TIFS $D_{16,9}$ ($c(14) \neq c(18)$); (d) completed historical Γ_1 TITS $I_{16,9}$ ($10 \sim_c 24, 11 \sim_c 18$); (e) two-way purely chromatic identity $T_{19,11}$ ($14 \sim_c 28, 18 \sim_c 27$); (f) $H_{21,13}$ showing three of its fifteen two-way purely chromatic identities: $2 \sim_c 14, 5 \sim_c 22, 8 \sim_c 23$. Red triangles forbid monochromatic triples; red dashed arrows denote $c(u) \neq c(v)$; blue dotted arrows denote $u \sim_c v$.

$\{3, 0, 8, 4\}$, $\{4, 9, 14, 5\}$, $\{5, 10, 15, 1\}$, $\{0, 11, 12, 5\}$ satisfy $\neg(c(6) = c(8) = c(11))$ (if $c(6) = c(8) = c(11) = 0$ and $c(0) = 3$, then $1, 2, 3, 4, 5$ must 2-color a 5-cycle), while $\{6, 7, 8, 9, 10, 11\} \in \mathcal{S}(W^{(4)}_{16,6})$ sets all three terminals to 1.

2. *Linear ordinary TIFS $L^{(4)}_{18,7}$ [Fig. 3(d)]:* Contexts $\{1, 2, 4, 3\}$, $\{5, 8, 7, 6\}$, $\{4, 9, 10, 8\}$, $\{11, 1, 12, 5\}$, $\{11, 14, 13, 2\}$, $\{15, 3, 16, 7\}$, $\{15, 17, 18, 6\}$ enforce $s(11) = 1 \implies s(15) = 0$ and $c(11) \neq c(15)$ (if $c(11) = c(15) = a$, then $1, 2, 3$ and $5, 6, 7$ avoid a, forcing $c(4) = c(8) = a$ in $\{4, 9, 10, 8\}$). Using $r_{13} = (1, 0, 1, 1), r_{14} = (2, 0, -1, -1)$ avoids an unlisted orthogonality $13 \perp 15$; all 68 states are extendable (terminal overlap 1/42).

3. *Linear directed TITS and color identity $L^{(4)}_{22,9}$ [Fig. 3(f)]:* Adjoining $\{11, 21, 22, 19\}$ and $\{13, 20, 15, 19\}$ to $L^{(4)}_{18,7}$ (with $r_{13} = (1, 0, 4, 4), r_{14} = (8, 0, -1, -1)$ so $r_{13} \perp r_{15}$) makes $\{11, 13, 15, 19\}$ pairwise differently colored, forcing $11 \sim_c 20$ and $s(11) = 1 \implies s(20) = 1$ (while $\{1, 6, 9, 14, 16, 20, 21\}$ has $s(11) = 0, s(20) = 1$; terminal overlap 41/42).

V. THE 37-RAY COMPLEX HESSE HYPERGRAPH AND RATIONAL FOUR-DIMENSIONAL COMPLETIONS

A. The 37-ray complex Hesse hypergraph H_{37} and its 76-event orthomodular lattice

With $\zeta = \exp(2\pi i/3)$ ($\zeta^2 + \zeta + 1 = 0$) and $a, k \in \mathbb{Z}_3$, define 37 rays in $\mathbb{C}^3$ (core rays z, M_{ak}, P_k, Q_k and unique cross-product completion rays A_k, B_{ak}, C_{ak}, numbered $1, \ldots, 37$ via $z \leftrightarrow 1, M_{ak} \leftrightarrow 2+3a+k, P_k \leftrightarrow 11+k, Q_k \leftrightarrow 14+k, A_k \leftrightarrow 17+k, B_{ak} \leftrightarrow 20+3a+k, C_{ak} \leftrightarrow 29+3a+k$):

$$\begin{aligned} z &= (1, 0, 0), & M_{ak} &= (1, \zeta^{a+k}, \zeta^{a+2k}), \\ P_k &= (0, 1, -\zeta^k), & Q_k &= (-\zeta^k, 0, 1), \\ A_k &= (0, 1, \zeta^k), & B_{ak} &= (-2, \zeta^{a+k}, \zeta^{a+2k}), \\ C_{ak} &= (1, -2\zeta^{2a+k}, \zeta^{2k}), \end{aligned} \tag{19}$$

and the 24 complete orthogonal triad contexts (Fig. 4):

$$\begin{aligned} \mathcal{M}_a &= \{M_{a0}, M_{a1}, M_{a2}\}, & \mathcal{A}_k &= \{z, P_k, A_k\}, \\ \mathcal{B}_{ak} &= \{M_{ak}, P_k, B_{ak}\}, & \mathcal{C}_{ak} &= \{M_{a,a+k}, Q_k, C_{ak}\}. \end{aligned} \tag{20}$$

TABLE II. Faithful geometric gadgets. $N_c = |\mathcal{C}_d|/d!$ fixes one context palette. Terminal relations describe colors unless explicitly identified as ordinary truth implications.

Gadget	d	$\lvert V\rvert/\lvert E\rvert$	N_c	$\lvert\mathcal{S}\rvert$	$\lvert\mathcal{S}_{\text{ext}}\rvert$	Relation and mechanism
P_5	3	10/5	5	11	10	Ternary exclusion on $6, 7, 9$; purely chromatic.
$B_{13,7}$	3	13/7	6	14	12	$c(10) \neq c(11)$; ordinary TIFS.
$D_{16,9}$	3	16/9	7	21	14	$c(14) \neq c(18)$; purely chromatic TIFS.
$I_{16,9}$	3	16/9	6	22	12	$10 \sim_c 24$, $11 \sim_c 18$; directed ordinary TITS.
$T_{19,11}$	3	19/11	7	25	14	$14 \sim_c 28$, $18 \sim_c 27$; both truth directions fail.
$H_{21,13}$	3	21/13	2	21	6	21 identical pairs: 15 two-way purely chromatic; six directed ordinary TITS.
$H_{17,10}$	3	17/10	4	18	10	Two color-identical pairs from directed ordinary TITS.
$W^{(4)}_{11,5}$	4	11/5	5	12	11	Apex ternary exclusion.
$W^{(4)}_{16,6}$	4	16/6	80	72	56	Linear ternary exclusion on $6, 8, 11$.
$B^{(4)}_{14,7}$	4	14/7	6	15	13	Apex ordinary TIFS.
$L^{(4)}_{18,7}$	4	18/7	144	68	68	Linear ordinary TIFS $c(11) \neq c(15)$.
$D^{(4)}_{17,9}$	4	17/9	7	22	15	Apex purely chromatic TIFS.
$L^{(4)}_{22,9}$	4	22/9	216	158	92	Linear identity $11 \sim_c 20$ from directed ordinary TITS.

Theorem 19 (Properties of H_{37})**.** *The hypergraph H_{37} defined by Eqs. (19)–(20) satisfies:*

1. *It is linear with Berge girth five,* 72 *edges, and $\omega(\Gamma(H_{37})) = 3$; its triangles are precisely its* 24 *contexts.*
2. *Equation (19) defines an injective edge-faithful representation in $\mathbb{C}^3$, whereas H_{37} admits* no *injective orthogonal representation in $\mathbb{R}^3$.*
3. *Its chromatic numbers are $\chi(H_{37}) = 4 > 3 = \chi_f(\Gamma(H_{37})) = \omega(\Gamma(H_{37}))$.*
4. *Its* 24 *Boolean blocks paste into a* 76*-event orthomodular lattice $L(H_{37}) = \{0, 1\} \cup \{a_i, a_i^\perp : 1 \leq i \leq 37\}$. Its full state space has $|\mathcal{S}(H_{37})| = 153$ states and is order determining on $L(H_{37})$, with exact minimum state-family sizes* 6 *for unital vertex separation,* 7 *for separation of all* 76 *events, and* 18 *for order determination.*

Proof. (1)–(2) Zero patterns, magnitudes, and ζ-phases distinguish all 37 rays. Algebraic inner products confirm that $\langle v_i, v_j\rangle = 0$ holds strictly for the $24\binom{3}{2} = 72$ context pairs: $\langle M_{ak}, M_{c\ell}\rangle = 1 + \zeta^{c-a}(\zeta^{\ell-k} + \zeta^{2(\ell-k)}) = 0 \iff (c = a, \ell \neq k)$; $P_j \perp M_{ak} \iff j = k$; $Q_j \perp M_{ak} \iff k = a + j$; $\langle M_{c\ell}, B_{ak}\rangle = -2 + \zeta^{a-c}(\zeta^{k-\ell} + \zeta^{2(k-\ell)}) = 0 \iff (c, \ell) = (a, k)$; $\langle M_{c\ell}, C_{ak}\rangle = 1 - 2\zeta^{2a+k-c-\ell} + \zeta^{2k-c-2\ell} = 0 \iff (c = a, \ell = a + k)$; and unequal term magnitudes exclude orthogonalities among A, B, C. Uniqueness of cross-product completions in $\mathbb{C}^3$ excludes Berge 3- and 4-cycles, while $(\mathcal{A}_0, \mathcal{B}_{00}, \mathcal{M}_0, \mathcal{B}_{01}, \mathcal{A}_1)$ is a Berge 5-cycle.

To exclude any injective representation in $\mathbb{R}^3$, suppose unit $z \in \mathbb{R}^3$ and real $p_k, m_{ak} \in \mathbb{R}^3$ realize $\mathcal{A}_k, \mathcal{B}_{ak}, \mathcal{M}_a$. If $m_{ak} \cdot z = 0$, then $\{z, p_k, m_{ak}\}$ is an orthogonal basis of $\mathbb{R}^3$, and for $\ell \neq k$, $p_\ell \perp z$ and $m_{a\ell} \perp m_{ak}$ with $p_\ell \perp m_{a\ell}$ force $[p_\ell] = [m_{ak}]$ or $[m_{a\ell}] = [z]$. Hence $m_{ak} \cdot z \neq 0$. Rescaling so $m_{ak} \cdot z = 1$ and choosing unit $u_k \in z^\perp \cap p_k^\perp$ gives $m_{ak} = z + t_{ak}u_k$ ($t_{ak} \in \mathbb{R} \setminus \{0\}$). Orthogonality of $\mathcal{M}_a$ yields $1 + t_{ak}t_{a\ell}(u_k \cdot u_\ell) = 0$ ($k \neq \ell$). Setting $q_k = t_{ak}/t_{0k}$ gives $q_0q_1 = q_0q_2 = q_1q_2 = 1$, whose only real solutions are $(q_0, q_1, q_2) = \pm(1, 1, 1)$. Thus at most two distinct bases $\mathcal{M}_a$ exist in $\mathbb{R}^3$, contradicting injectivity on the three bases $\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$. The argument uses only the 13 rays z, P_k, M_{ak} and the orthogonalities among them, so already the 13-vertex subgraph of $\Gamma(H_{37})$ induced by these rays has no injective orthogonal representation in $\mathbb{R}^3$.

(3) In any 3-coloring with $c(z) = 0$, $P_0, P_1, P_2 \in \{1, 2\}$ cannot be monochromatic (as $\mathcal{M}_a$ uses all three colors and $M_{ak} \sim P_k$). Up to symmetry $(c(P_0), c(P_1), c(P_2)) = (2, 1, 1)$, forcing $c(M_{a0}) = 1$ and $(c(M_{a1}), c(M_{a2})) = (x_a, 2 - x_a)$ with $x_a \in \{0, 2\}$. Since $N(Q_k) = \{M_{0k}, M_{1,k+1}, M_{2,k+2}\}$ avoids a color, $x_1 = 2 - x_2$, $x_0 = 2 - x_1$, $x_2 = 2 - x_0$, impossible on a 3-cycle. Thus $\chi(H_{37}) = 4$ (attained by $c(z) = 0, c(M_{ak}) = k, c(P_k) = c(Q_k) = 3$). The 6-state double cover $\mathcal{R}_6 = \{s_2, s_{30}, s_{73}, s_{99}, s_{109}, s_{139}\}$ covers every vertex twice, giving $\chi_f(\Gamma(H_{37})) = 3$.


(4) Each state is fixed by $t = (t_0, t_1, t_2) \in \mathbb{Z}_3^3$ with $s(M_{ak}) = \mathbf{1}_{k=t_a}$, $s(z) = \varepsilon \in \{0, 1\}$, and independent bits on $U(t) = \mathbb{Z}_3 \setminus \{t_0, t_1, t_2\}$ (for $\varepsilon = 0$) and $W(t) = \mathbb{Z}_3 \setminus \{t_a - a\}$, yielding $(1 + 2^{|U(t)|})2^{|W(t)|}$ states per t:

Class of $t \in \mathbb{Z}_3^3$	$\lvert\#t\rvert$	$(\lvert U(t)\rvert, \lvert W(t)\rvert)$	$(1 + 2^{\lvert U\rvert})2^{\lvert W\rvert}$
(c, c, c)	3	$(2, 0)$	$5 \times 1 = 5$
$(c, c + 1, c + 2)$	3	$(0, 2)$	$2 \times 4 = 8$
$(c, c + 2, c + 1)$	3	$(0, 0)$	$2 \times 1 = 2$
Two equal entries	18	$(1, 1)$	$3 \times 2 = 6$

(21)

and hence $|\mathcal{S}(H_{37})| = 3 \cdot 5 + 3 \cdot 8 + 3 \cdot 2 + 18 \cdot 6 = 153$. Since

(a) $W^{(4)}_{11,5}$ (11/5) **(b)** $W^{(4)}_{16,6}$ (16/6, linear) **(c)** $B^{(4)}_{14,7}$ (14/7)



(d) $L^{(4)}_{18,7}$ (18/7, linear) **(e)** $D^{(4)}_{17,9}$ (17/9) **(f)** $L^{(4)}_{22,9}$ (22/9, linear)




FIG. 3. Faithful real four-dimensional gadgets in $\mathbb{R}^4$. Panels (a),(c),(e) are common-apex lifts ($g(H) = 2$); panels (b),(d),(f) are linear ($g(H) = 3$). (a),(b) Purely chromatic ternary exclusions; (c),(d) ordinary TIFS; (e) purely chromatic binary TIFS $c(14) \neq c(18)$; (f) linear color identity $11 \sim_c 20$ from directed ordinary TITS $s(11) = 1 \implies s(20) = 1$.

$2^5 - 1 = 31 < 37$ and $2^6 = 64 < 76$, the explicit families Ω_6 and Ω_7 given in the supplement prove the minima 6 and 7. For order determination ($a_i \leq a_j^\perp \iff i \sim j$), the 15 nonorthogonal pairs in $D = \{P_0, P_1, P_2, Q_0, Q_1, Q_2\}$ require 15 disjoint joint-one witnesses in $T \subset \mathcal{S}(H_{37})$ ($|T| = 27$), and the 9 pairs $\{z, C_{ak}\}$ require at least 2 states outside T. Exhaustive bit-set verification over all $4{,}096 \times 132 = 540{,}672$ candidate 17-state families shows each meets at most 75 of the 81 minimal witness sets W_{xy}, whereas the 18-state family Ω_{18} in the supplement meets all 81 (Supplement Sec. S3). □

Theorem 19(2) is not the first finite separation of complex from real three-dimensional orthogonality: Harding and Salinas Schmeis [29, Thm. 2.6] construct, from mutually unbiased bases, a two-gadget configuration that is realizable in $\mathbb{C}^3$ but not in $\mathbb{R}^3$ (48 rays, 32 bases, 189 context states, and 180 geometric states). H_{37} is smaller, linear of Berge girth five, and edge-faithful in $\mathbb{C}^3$, and its real obstruction already resides in the 13-ray subgraph on z, P_k, M_{ak}.

By Proposition 18, appending $d - 3$ common apices to H_{37} in $\mathbb{C}^d$ ($d \geq 3$) gives faithful d-uniform hypergraphs with $|V| = d + 34$, $|E| = 24$, $\omega = d$, $\chi = d + 1$, and $|\mathcal{S}| = d + 150$.

An order-determining family represents the specified 76-event order, orthocomplements, and Boolean operations within listed contexts. It does not preserve arbitrary meets as set intersections: nonorthogonal atoms have meet zero although some states make both true. It is therefore not a Boolean-sublattice embedding of the full lattice, nor a claim of closure under every Hilbert operation. In particular, the minimum six-state separating family and the six-state uniform double cover solve different problems.

B. Hyperplane obstruction, two-plane completion lemma, and canonical linear H_{65} in $\mathbb{R}^4$

Appending a single apex $v_{26} = (0, 0, 0, 1)^T \in \mathbb{R}^4$ to H^+_{YO} gives the 26-ray, 16-context non-linear suspension $H^{(4)}_{\mathrm{YO}}$ (Fig. 5) with $\omega = 4, \chi = 5$, and 25 unital separating states (Proposition 18), but $|e \cap e'| = 2$ ($g = 2$). Restricted to the 13 Yu–Oh rays and the apex, its orthogonality graph is the Yu–Oh graph joined to one common vertex, the example that Cabello, Kleinmann, and Budroni [26, Thm. 1] used to show that $\chi_f > d$ is necessary but not sufficient for state-independent contextuality; H_{65} below




FIG. 4. Complete 3-uniform hypergraph diagram of H_{37} (37 rays, 24 orthogonal triads, 72 edges in $\mathbb{C}^3$). Bold circles denote e_1, e_2, e_3 ($\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$); the other 21 smooth traces join core pairs through their unique completion rays $17, \ldots, 37$. Concentric rings record incident context colors.

is a linear, apex-free example of the same phenomenon.

Proposition 20 (Hyperplane obstruction to linearity in $\mathbb{F}^4$)**.** *In any injective realization in $\mathbb{P}(\mathbb{F}^4)$, if two contexts e, e' each have three rays in a common 3-dimensional hyperplane $W = z^\perp$, both must contain $[z]$; hence if $|e_W \cap e'_W| \geq 1$, then $|e \cap e'| \geq 2$, violating linearity.*

Proof. Three mutually orthogonal rays in W span W, forcing the fourth orthogonal ray of both e and e' to equal $[z] = W^\perp$. □

Lemma 21 (Simultaneous two-plane completion in $\mathbb{R}^4$)**.** *Let $V_{\text{core}} \subset \mathbb{Q}^4 \setminus \{0\}$ be a finite set of non-collinear rational vectors with 4-element orthogonal base contexts E_{base} ($|C \cap C'| \leq 1$) and let all remaining orthogonal pairs be the bridge pairs $E_{\text{bridge}} = \{e_1, \ldots, e_M\}$ ($e_m = \{u_m, v_m\}$, $L_m = \text{span}(v_{u_m}, v_{v_m})$, $P_m = L_m^\perp$) such that every core vertex lies in a base context or is a bridge endpoint.*

1. *The pairs e_m can be simultaneously completed to 4-bases $E_m = \{u_m, v_m, w_{m,1}, w_{m,2}\}$ by fresh integer vectors $w_{m,1}, w_{m,2} \in P_m \cap \mathbb{Z}^4$ forming a linear edge-*




FIG. 5. The completed 25-ray, 16-context Yu–Oh hypergraph H_{YO}^{+} in $\mathbb{R}^3$ ($\omega = 3, \chi = 4$) and its non-linear 26-ray, 16-context apex suspension $H_{\mathrm{YO}}^{(4)}$ in $\mathbb{R}^4$ ($\omega = 4, \chi = 5, g = 2$) obtained by adjoining the common orthogonal apex $v_{26} = (0, 0, 0, 1)^T$ to every context.

faithful hypergraph H if and only if

$$\begin{aligned} v_z &\notin L_m \quad (z \in V_{\mathrm{core}} \setminus e_m), \\ P_\ell &\neq L_m \quad (\ell \neq m). \end{aligned} \tag{22}$$

2. *$\omega(\Gamma(H)) = 4$ and $\chi(H) = \max\{4, \chi(G_{\mathrm{core}})\}$; here G_{core} is automatically K_5-free, since no five nonzero vectors of $\mathbb{R}^4$ are mutually orthogonal.*

3. *If all G_{core}-independent base supports $S_1, \ldots, S_N$ form a unital separating family on the core, and the zero-profiles $Z(e_m) = \{\alpha : S_\alpha \cap e_m = \varnothing\}$ are nonempty and pairwise distinct, the $\leq 2N$ canonical extensions $s_{\alpha,\varepsilon}$ ($\varepsilon \in \{0, 1\}$, setting $(s_{\alpha,\varepsilon}(w_{m,1}), s_{\alpha,\varepsilon}(w_{m,2})) = (1-\varepsilon, \varepsilon)$ when $\alpha \in Z(e_m)$ and $(0, 0)$ otherwise) are unital and separating on $V(H)$.*

Proof. If $v_z \in L_m$ then $P_m \subset v_z^\perp$, and if $P_\ell = L_m$ then $P_\ell \perp P_m$, forcing extra orthogonalities. Conversely, given Eq. (22), an integer orthogonal basis $\{\eta_A, \eta_B\} \subset P_m \cap \mathbb{Z}^4$ defines the infinite family $w_{m,1}(a, b) = a\eta_A + b\eta_B$, $w_{m,2}(a, b) = -\|\eta_B\|^2 b\eta_A + \|\eta_A\|^2 a\eta_B$, avoiding inductively the finitely many 1-dimensional intersections $z^\perp \cap P_m$ and $\mathbb{R}z \cap P_m$, where z ranges over the core rays *and* the fresh rays placed at earlier steps, together with $P_m \cap L_r$ ($r > m$). The last condition keeps every fresh ray outside each later L_r, so that $z^\perp \cap P_r$ remains 1-dimensional for fresh z as well. Linearity, coloring extension, and distinctness of canonical vertex profiles follow from Proposition 7 and distinctness of $Z(e_m)$. □

To construct a *linear* 4-uniform hypergraph in $\mathbb{R}^4$, we take the 17-ray core $V_{\mathrm{core}} = \{1, \ldots, 17\} \subset \{-1, 0, 1\}^4$, which is orthogonally equivalent to the 17-ray Peres restriction of Haynes *et al.* [12, pp. 8–9] via the orthogonal matrix

$$T = \frac{1}{2}\begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & -1 & -1 & -1 \end{pmatrix}, \qquad T^T T = I_4. \tag{23}$$

The core consists of the coordinate basis C_1, an even-parity Hadamard basis C_2, three odd-parity hub rays $9, 10, 11$, and the three binary bridge pairs $B_1 = \{12, 13\}$, $B_2 = \{14, 15\}$, $B_3 = \{16, 17\}$. The remaining base contexts are $C_3 = \{9, 10, 12, 13\}$, $C_4 = \{9, 11, 14, 15\}$ and $C_5 = \{10, 11, 16, 17\}$. Complete coordinates and all compatible supports are in Supplemental Tables S17–S18 and S81. Completing its 24 uncovered bridge edges $e_m \in (C_1 \cup C_2) \times (B_1 \cup B_2 \cup B_3)$ by fresh integer pairs $\{2m + 16, 2m + 17\} \subset \{18, \ldots, 65\}$ via Lemma 21 yields $H_{65} = (V_{65}, E_{29})$ (Fig. 6; Supplemental Tables S17–S18):

$$\begin{aligned} E_{29} &= \{C_1, \ldots, C_5\} \cup \{E_m : 1 \leq m \leq 24\}, \\ E_m &= \{u_m, v_m, 2m + 16, 2m + 17\}. \end{aligned} \tag{24}$$




FIG. 6. Canonical linear 4-uniform hypergraph H_{65} in $\mathbb{R}^4$ (65 rays, 29 contexts, 174 edges, $g = 3, \omega = 4, \chi = 5$). Thick colored curves denote the five core contexts $C_1, \ldots, C_5$; the 24 thinner lines denote the completed bridge contexts $E_1, \ldots, E_{24}$ carrying degree-one completion pairs $\{18, 19\}, \ldots, \{64, 65\}$.

TABLE III. Bridge neighbors in the two core bases. The complete 24 state supports are supplementary.

b	$N(b) \cap C_1$	$N(b) \cap C_2$
12	$\{2, 3\}$	$\{5, 8\}$
13	$\{1, 4\}$	$\{6, 7\}$
14	$\{2, 4\}$	$\{6, 8\}$
15	$\{1, 3\}$	$\{5, 7\}$
16	$\{3, 4\}$	$\{7, 8\}$
17	$\{1, 2\}$	$\{5, 6\}$

Theorem 22 (Properties of H_{65})**.** *The* 65*-ray,* 29*-context hypergraph* H_{65} *satisfies:*

1. *It is linear* ($g(H_{65}) = 3$) *and edge-faithful in* $\mathbb{R}^4$ (174 *edges) with* $\omega(\Gamma(H_{65})) = 4$*, and nonconformal* (13 *uncovered maximal triangles:* 12 *bridge triangles plus* $\{9, 10, 11\}$).
2. *It has* $\alpha(G_{\text{core}}) = 4$, $\chi_f(\Gamma(H_{65})) = 17/4 = 4.25 > 4$, *and* $\chi(H_{65}) = 5$.
3. *Its core has* 24 *compatible state supports* $S_\alpha = \{i_1, i_2, h, b\}$ *(Table S81), each missing* 14 *bridges and yielding* $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$ *states and* 48 *canonical separating states.*
4. *Every* $s \in \mathcal{S}(H_{65})$ *obeys* $s(9) + s(10) + s(11) = 1$*, so adjoining the fourth hub ray* $q = (1, 1, 1, -1)^T$ *forces* $\widetilde{s}(q) = 0$*; moreover,* $A_9 = A_{16} \dot{\cup} A_{17}$ *on all of* $\mathcal{S}(H_{65})$*, forcing the false atom-order implication* $s(16) \le s(9)$ *and collapsing the* 1*-ray* $[v_9] = [v_{16} - v_{17}]$ *with the* 2*-plane* $\text{span}(v_{16}, v_{17})$.

Proof. (1)–(2) No K_4 in G_{core} contains a bridge edge $\{u_m, v_m\}$, since $u_m \in C_a$ and $v_m \in B_k$ have exactly one common neighbor in G_{core}, namely the second vertex of $N(v_m) \cap C_a$; this verifies $P_\ell \neq L_m$, and support-size / magnitude checks verify $v_z \notin L_m$ in Lemma 21. For any independent $I \subseteq V_{\text{core}}$, $|I \cap C_1| \le 1$ and $|I \cap C_2| \le 1$. If $I \cap \{9, 10, 11\} \neq \varnothing$, $|I \cap (C_3 \cup C_4 \cup C_5)| \le 2$ so $|I| \le 4$; if $I \cap \{9, 10, 11\} = \varnothing$ and I picks $(b_1, b_2, b_3) \in B_1 \times B_2 \times B_3$, even-parity triples cover C_2 and odd-parity triples cover C_1 (Table III), so again $|I| \le 3 + 1 = 4$. Thus $\alpha(G_{\text{core}}) = 4$, $\chi_f \ge 17/4$ (matched by weights $1/8$ on the 24 core states and $1/16$ on the 20 remaining independent 4-sets), and $\chi(H_{65}) = 5$. (3)–(4) By parity, no state can avoid $\{9, 10, 11\}$, giving $s(9)+s(10)+s(11) = 1$, $3 \times 2 \times 2 \times 2 = 24$

core supports (Table S81), $24-10=14$ missed bridges per support, and $A_9 = A_{16} \dot{\cup} A_{17}$. □

VI. EVENT ORDER, QUANTUM PROBABILITIES, AND NONEXTENDABLE WEIGHT

A. Extendable state polytopes and nonextendable weight on $H_{17,10}$

For any context hypergraph H, the classical state polytope $\mathcal{P}(H) = \operatorname{conv} \mathcal{S}(H)$ and the extendable state polytope $\mathcal{P}_{\text{ext}}(H) = \operatorname{conv} \mathcal{S}_{\text{ext}}(H)$ satisfy $\mathcal{P}_{\text{ext}}(H) \subseteq \mathcal{P}(H)$, with $p_u = p_v$ for all $p \in \mathcal{P}_{\text{ext}}(H)$ whenever $u \sim_c v$ by Corollary 4. For $p \in \mathcal{P}(H)$, define its *minimum nonextendable weight*

$$\begin{aligned} \eta_H(p) = \min_{\mu} \quad & \sum_{s \notin \mathcal{S}_{\text{ext}}} \mu_s, \\ \text{subject to} \quad & p = \sum_{s \in \mathcal{S}} \mu_s s, \\ & \mu_s \geq 0, \qquad \sum_s \mu_s = 1. \end{aligned} \tag{25}$$

Proposition 23 (Nonextendable weight lower bound)**.** *For any chromatically inseparable pair $u \sim_c v$ in H and any $p \in \mathcal{P}(H)$, $\eta_H(p) \geq |p_u - p_v|$.*

Proof. Every $s \in \mathcal{S}_{\text{ext}}(H)$ satisfies $s(u) - s(v) = 0$, while every $s \notin \mathcal{S}_{\text{ext}}(H)$ satisfies $|s(u) - s(v)| \leq 1$, so $|p_u - p_v| \leq \sum_{s \notin \mathcal{S}_{\text{ext}}(H)} \mu_s |s(u) - s(v)| \leq \sum_{s \notin \mathcal{S}_{\text{ext}}(H)} \mu_s$. □

Theorem 24 (Quantum preparations in $\mathcal{P}(H_{17,10}) \setminus \mathcal{P}_{\text{ext}}(H_{17,10})$)**.** *In the real FOR of $H_{17,10}$ (Supplemental Table S2):*

1. *The pure quantum preparation along $r_{24} = (1,-1,2)^T$,*

$$\rho_\star = \frac{1}{6} \begin{pmatrix} 1 & -1 & 2 \\ -1 & 1 & -2 \\ 2 & -2 & 4 \end{pmatrix}, \qquad p_i = \operatorname{Tr}(\rho_\star P_i) = \frac{(r_{24} \cdot r_i)^2}{6 \|r_i\|^2}, \tag{26}$$

belongs to $\mathcal{P}(H_{17,10})$ via the exact five-state convex combination (state indices in Supplemental Table S40; see also Tables S46–S47)

$$p = \frac{1}{12} s_4 + \frac{5}{36} s_9 + \frac{1}{36} s_{10} + \frac{7}{12} s_{14} + \frac{1}{6} s_{17}, \tag{27}$$

yet satisfies $p_{24} = 1$ and $p_{10} = 8/9$ for the chromatically inseparable pair $(10, 24)$, so

$$p \notin \mathcal{P}_{\text{ext}}(H_{17,10}) \qquad \text{and} \qquad \eta_{H_{17,10}}(p) = \frac{1}{9}. \tag{28}$$

2. *The pure preparation along the coordinate ray $z_3 = r_5 = (0,0,1)^T$, with Born vector $p'_i = (r_5 \cdot r_i)^2 / \|r_i\|^2$, has $p'_{24} = p'_{18} = 2/3$ and $p'_{10} = p'_{11} = 1/3$, belongs to $\mathcal{P}(H_{17,10})$ via the four-state convex combination $p' = \frac{1}{3} s_3 + \frac{1}{6} s_4 + \frac{1}{3} s_{14} + \frac{1}{6} s_{15}$, and satisfies $\eta_{H_{17,10}}(p') = 1/3$.*

3. *For every density operator ρ and both inseparable pairs $(u, v) \in \{(10, 24), (11, 18)\}$, $|p_u(\rho) - p_v(\rho)| \leq \|P_u - P_v\| = 1/3$. Hence Proposition 23 certifies at most $1/3$ on $H_{17,10}$, and the preparation in (2) attains this value.*

Proof. (1) Direct evaluation of Eq. (27) on all 17 vertices matches p_i in Eq. (26) (Table S46), with $r_{10} = (1,-1,1)^T$ giving $p_{10} = 4^2/(6 \cdot 3) = 8/9$. Since $10 \sim_c 24$, Proposition 23 gives $\eta_{H_{17,10}}(p) \geq |1 - 8/9| = 1/9$, which is attained in Eq. (27) because among $\{s_4, s_9, s_{10}, s_{14}, s_{17}\}$ precisely s_4 and s_{10} are nonextendable, with total weight $1/12 + 1/36 = 1/9$. (2) Direct evaluation of the four-state mixture on all 17 vertices reproduces p' (Tables S46–S47). Proposition 23 gives $\eta_{H_{17,10}}(p') \geq |2/3 - 1/3| = 1/3$, attained because s_3, s_{14} are extendable and s_4, s_{15} are not, with total weight $1/6 + 1/6 = 1/3$. (3) For unit vectors with squared overlap κ, $P_u - P_v$ has eigenvalues $\pm\sqrt{1-\kappa}$ and 0. Both pairs have $\kappa = (r_u \cdot r_v)^2 / (\|r_u\|^2 \|r_v\|^2) = 16/18 = 8/9$, so $|\operatorname{Tr} \rho(P_u - P_v)| \leq \|P_u - P_v\| = 1/3$. □

The exact support-and-weight certificates and the complete Born probability vectors are given in the supplement. These tables are regenerated from the checked state inventory, rather than manually transcribed.

These preparations exclude only models built from noncontextual eigenvalue assignments: their ordinary-state models are exhibited explicitly. A state is extendable exactly when some noncontextual eigenvalue assignment c makes it the indicator of a single eigenvalue λ_a in every context observable A_e of Eq. (1). For a model built from noncontextual eigenvalue assignments, every chromatic identity $u \sim_c v$ entails the equality $p_u = p_v$; quantum mechanics violates it, and Proposition 23 converts the violation into a lower bound on η. The weight η therefore measures how far the quantum predictions force any noncontextual truth-assignment model away from noncontextual eigenvalues: on $H_{17,10}$, at least one third of the weight in the preparation along z_3. Since $\mathcal{P}_{\text{ext}}(H_{17,10})$ lies in the hyperplane $p_{10} = p_{24}$, a Born vector in $\mathcal{P}(H_{17,10})$ generically lies outside $\mathcal{P}_{\text{ext}}(H_{17,10})$; the content of Theorem 24 is the size of the required weight. Conversely, averaging the d color classes of any admissible coloring gives the maximally mixed Born vector $1/d$, so the restricted obstruction is preparation dependent. The $H_{17,10}$ ordinary implication $s(10) \leq s(24)$ also prevents order determination for distinct ray atoms; vertex separation and order representation must not be conflated.

B. Order determination versus Born probabilities on H_{37}, $H_{\rm YO}^{(4)}$, and H_{65}

When $\chi(H) > d$ (Regime I), no noncontextual eigenvalue assignment exists, $\mathcal{P}_{\rm ext}(H) = \varnothing$, and the relevant probability question is whether Born assignments $p_i(\rho) = \mathrm{Tr}(\rho\Pi_i)$ lie in $\mathcal{P}(H) = \mathrm{conv}\,\mathcal{S}(H)$ (Table IV).

Corollary 25 (Order determination does not imply classical probability models)**.** *The* 76*-event orthomodular lattice* $L(H_{37})$ *admits an order-determining state space* $\mathcal{S}(H_{37})$ *and a classical mixture for the maximally mixed state* $I_3/3$*, yet every* $p \in \mathcal{P}(H_{37})$ *satisfies the tight inequality*

$$\mathcal{C}_{37}(p) = \sum_{k=0}^{2} \big[p(P_k) + p(Q_k)\big] \le 2, \tag{29}$$

whereas the complex FOR of H_{37} *satisfies*

$$\sum_{k=0}^{2} (\Pi_{P_k} + \Pi_{Q_k}) = \mathrm{diag}(3/2, 3/2, 3) = \frac{3}{2} I_3 + \frac{3}{2} |e_3\rangle\langle e_3|, \tag{30}$$

yielding $\mathcal{C}_{37}(\rho) = \frac{3}{2}(1+\rho_{33}) > 2$ *whenever* $\rho_{33} > 1/3$ *(and* $\mathcal{C}_{37}(|e_3\rangle\langle e_3|) = 3$*).*

Proof. In the state parametrization of Theorem 19(4), every $t \in \mathbb{Z}_3^3$ has $|U(t)|+|W(t)| \le 2$, establishing Eq. (29), while summing the normalized projectors of P_k, Q_k in Eq. (19) using $1 + \zeta + \zeta^2 = 0$ gives Eq. (30). Uniform weights $1/6$ on $\mathcal{R}_6$ reproduce $p_i(I_3/3) = 1/3$. □

In dimension four, the non-linear suspension $H_{\rm YO}^{(4)}$ obeys $\mathcal{Y}(p) = p_{26} + \sum_{i=10}^{13} p_i \le 1$ on $\mathcal{P}(H_{\rm YO}^{(4)})$, whereas $\sum_{i=10}^{13} \Pi_i = \frac{4}{3}(I_4 - \Pi_{26})$ gives $\mathcal{Y}(\rho) = 1 + (1 - p_{26}(\rho))/3 > 1$ for every density operator $\rho \neq \Pi_{26}$ [26]. For the canonical linear H_{65}, every $s \in \mathcal{S}(H_{65})$ has 4 active core vertices, so every $p \in \mathcal{P}(H_{65})$ obeys $\sum_{i=1}^{17} p_i = 4$. However, with $q = (1, 1, 1, -1)^T$ and $Q = qq^T/4$, direct summation gives

$$\sum_{i=1}^{17} \Pi_i = 4I_4 + Q \quad \Longrightarrow \quad \sum_{i=1}^{17} p_i(\rho) = 4 + \mathrm{Tr}(\rho Q), \tag{31}$$

violating the classical bound 4 whenever $\mathrm{Tr}(\rho Q) > 0$ (including $I_4/4$, where the sum is $17/4 = \chi_f$, and $\rho = Q$, where it is 5). Nevertheless, on the orthogonal subspace $q^\perp$, the density operator $\rho_\star = (I_4 - Q)/3$ has core Born probabilities $1/4$ on $C_1 \cup C_2$, $1/3$ on $\{9, 10, 11\}$, and $1/6$ on $B_1 \cup B_2 \cup B_3$, and completion probabilities $r_{m,1} + r_{m,2} = 7/12$. Choosing $\alpha \in \{1, \ldots, 24\}$ uniformly and $\lambda \in [0, 1]$ uniformly and setting $\varepsilon_m = \mathbf{1}_{\{\lambda < \tau_m\}}$ with $\tau_m = \frac{12}{7} r_{m,2} \in \{4/9, 5/11, 6/11, 5/9\}$ on the 14 missed bridges of S_α gives $\mathbb{E}[s(w_{m,2})] = \frac{14}{24}\tau_m = r_{m,2}$, expressing $p(\rho_\star) \in \mathcal{P}(H_{65})$ as a convex mixture of at most 120 states. Rotating $V_{\rm core}$ by $D_0 = I_4, D_1 = \mathrm{diag}(1, 1, -1, -1), D_2 = \mathrm{diag}(1, -1, 1, -1), D_3 = \mathrm{diag}(1, -1, -1, 1)$ gives a fixed enlarged family with

$$\sum_{a,i} \Pi_i^{(a)} = 17 I_4, \qquad \sum_{a,i} p_i^{(a)} \le 16 \text{ classically.} \tag{32}$$

Each copy admits at most four true rays under geometric pair exclusions; repeated projectors are counted with multiplicity. The 24 distinct rays form the Peres set [30]. This is an enlargement, not a state-independent test of the original 65 rays.

VII. KS ANCESTRY, SPECIFIED PARENTS, AND ABSTRACT VERSUS GEOMETRIC STATES

A. The KS ancestry hypothesis and abstract transverse extensions

Motivated by the recurring parent structures of our examples, we formulate the *KS ancestry hypothesis*: informally, every noncolorable ($\chi(H) > d$) configuration has a larger KS ancestor. For realized ray configurations, its existence form follows from an elementary union argument, even without the chromatic premise; the substantive open question is whether the retained rays can be made essential to the contradiction (Sec. VIII):

Proposition 26 (Same-dimensional finite KS ancestry)**.** *Every finite ray set* $R \subset \mathbb{P}(\mathbb{F}^d)$ *(*$\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$*,* $d \ge 3$*) is a proper subset of a finite KS ray set* $P \subset \mathbb{P}(\mathbb{F}^d)$ *whose orthogonality graph and full-basis hypergraph induce those on* R*, with every ray of* P *in a complete context. Consequently, a finite graph* G *has an injective edge-faithful realization in* $\mathbb{F}^d$ *if and only if* G *is an induced subgraph of a finite KS orthogonality graph in* $\mathbb{F}^d$*.*

Proof. Choose a finite full-basis KS set $K \subset \mathbb{P}(\mathbb{F}^d)$ [4, 31], orient K by $U \in \mathrm{U}(d)$ (or $\mathrm{O}(d)$) so $UK \not\subseteq R$, and complete any uncovered rays of $R \cup UK$ to bases to obtain P. Any state on P would restrict to a state on UK, a contradiction, and keeping the rays of R fixed preserves all induced orthogonalities and full bases on R. □

By Theorem 11, H_8 has no injective realization, and H_9, G_{32} have no ray assignment; none can embed into any 3-dimensional geometric KS parent. For abstract hypergraphs, Lemma 3 yields a direct chromatic-to-stateless construction:

Proposition 27 (Transverse chromatic-to-stateless extension)**.** *For any finite* d*-uniform hypergraph* $H = (V, E)$ *(*$d \ge 3$*), let* $T(H)$ *take* $d-1$ *disjoint copies* $H^{(1)}, \ldots, H^{(d-1)}$ *and adjoin for each* $v \in V$ *a fresh vertex* w_v *and the transverse context* $T_v = \{v^{(1)}, \ldots, v^{(d-1)}, w_v\}$*. Then* $T(H)$ *has* $d|V|$ *vertices,* $(d-1)|E| + |V|$ *contexts, is linear if* H *is linear, and its two-valued states correspond bijectively to labeled strong* d*-colorings of* H*; in particular,*

$$\mathcal{S}(T(H)) = \varnothing \quad \Longleftrightarrow \quad \chi(H) > d. \tag{33}$$

TABLE IV. Event order and preparation-dependent probability models for the fixed realizations. Membership statements concern Born vectors $p(\rho)$; the restricted model is $\mathcal{P}_{\text{ext}}$.

Configuration	Order determination	An ordinary-state model	A failed model or witness
$H_{17,10}$	No: $s(10) \le s(24)$ is a false ray inclusion.	P_{24} and P_{z_3} have ordinary mixtures.	Neither lies in $\mathcal{P}_{\text{ext}}$; $\eta = 1/9$ and $1/3$ (maximal pair certificate).
Completed Yu–Oh	No.	No Born preparation lies in $\mathcal{P}$.	Every preparation gives $\sum_{10}^{13} p_i = 4/3 > 1$.
H_{37}	Yes on its specified 76-event lattice; 18 states minimal.	$I_3/3$ has a six-state mixture.	$\rho_{33} > 1/3$ violates the P, Q bound.
$H_{\text{YO}}^{(4)}$	No.	Pure apex P_{26} has its singleton-state model.	Every $\rho \neq P_{26}$ violates the displayed bound.
H_{65}	No: $s(16) \le s(9)$ is a false ray inclusion.	$(I_4 - Q)/3$ has a model for all 65 rays.	$\text{Tr}(\rho Q) > 0$ violates the core sum bound.

Proof. A state $s \in \mathcal{S}(T(H))$ restricts to $d-1$ states on $H^{(k)}$ that are pairwise disjoint by T_v (forcing $s(w_v) = 1 - \sum_{k=1}^{d-1} s(v^{(k)})$), which by Lemma 3 is equivalent to a strong d-coloring of H. □

In dimension $d = 3$, any edge $u \sim v$ of H creates a Berge 4-loop $(u^{(1)}, v^{(1)}, u^{(2)}, v^{(2)})$ in $T(H)$, excluding an injective realization in $\mathbb{C}^3$ by the common orthogonal-complement argument; thus $T(H)$ is an abstract construction. For H_9, $T(H_9)$ (45 vertices, 33 contexts) is context-critical for statelessness on all 33 contexts; for H_{YO}^+, $T(H_{\text{YO}}^+)$ (75 vertices, 57 contexts) is critical on all 32 horizontal contexts and 12 transverse contexts $v \in \{14, \ldots, 25\}$, but remains stateless upon deleting T_v for $v \in \{1, \ldots, 13\}$ (Supplement Sec. S3).

B. Explicit inclusions in known KS parents and Cabello completions

Beyond unrestricted existence, the following explicit unions use known KS components; the unions are not asserted to be previously named, smallest, or critical parents (explicit coordinate dictionaries and certificates in Supplement Secs. S1–S3):

1. *Bub's 49-ray context-critical real parent of H_{YO}^+:*

Proposition 28 (Context-critical real parent of H_{YO}^+)**.** *Adjoining to H_{YO}^+ the eight rays $(0, 1, \pm 2)$, $(0, 2, \pm 1)$, $(1, \pm 2, 0)$, $(2, \pm 1, 0)$ yields 33 rays with 20 bases and 16 uncovered orthogonal pairs that eliminate all 24 states of H_{YO}^+. Completing those 16 pairs yields the 49-ray, 36-context full-context Schütte–Bub KS configuration [13, 32], containing H_{YO}^+ as an induced restriction, in which every one of the 36 contexts is critical for $\mathcal{S}_{\text{ctx}} = \varnothing$ (Supplemental Table S83). If the orthogonalities of a deleted context are retained as geometric exclusions, only the 20 bases of the 33-ray stage remain critical; deleting any of the 16 completion contexts leaves $\mathcal{S}_{\text{geom}} = \varnothing$.*

2. *MUB-inventory parents of H_{37}:* In the 165-ray, 130-context MUB inventory [11] formed by three 69-ray, 50-context KS lineages L_0, L_1, L_2 (where each L_a completes the 36 uncovered pairs of Cabello's 33-ray, 14-basis KS set [33]; Pavičić [34] points out that this 33-ray set already occurs as a subset of a 69-ray, 50-basis set in Ref. [35]), H_{37} embeds as an induced restriction via the seed-index matrix $j(a,k) = \left(\begin{smallmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 9 & 8 \end{smallmatrix}\right)$ mapping $z \leftrightarrow a_{1,1}$, $M_{ak} \leftrightarrow u_{j(a,k)}$, $P_k \leftrightarrow c_{1,k+1}$, $Q_k \leftrightarrow c_{2,k+1}$, $A_k \leftrightarrow b_{1,k+1}$, $B_{ak} \leftrightarrow d_{1,j(a,k)}$, $C_{ak} \leftrightarrow d_{2,j(a,a+k)}$. It lies in the 57-ray, 40-context intermediate spoke hypergraph (297 states) and in each 87-ray, 64-context KS union $H_{37} \cup L_a$.

3. *Four-dimensional Peres and Cabello parents of $H_{\text{YO}}^{(4)}$ and H_{65}:* Taking the union with Peres's 24-ray set P_{24} [30] gives a 40-ray, 36-context KS parent of $H_{\text{YO}}^{(4)}$ ($|H_{\text{YO}}^{(4)} \cap P_{24}| = 10$; note that a single-apex lift of a 3-dimensional KS set always retains the unique apex-one state) and a 72-ray, 48-context KS parent of H_{65} ($|H_{65} \cap P_{24}| = 17$). More economically:

Proposition 29 (Three-ray KS extensions of H_{65})**.** *Adjoining $a = (1,0,0,1), b = (0,1,0,1), c = (0,0,1,1)$ to H_{65} yields a 68-ray, 35-basis KS parent containing H_{65} as an induced restriction and Cabello's 18-ray, 9-context parity proof $C_{18,9}$ [14] (C_3, C_4, C_5 plus $D_1 = \{2,3,12,a\}, D_2 = \{1,3,15,b\}, D_3 = \{1,2,17,c\}, D_4 = \{6,7,13,a\}, D_5 = \{6,8,14,b\}, D_6 = \{7,8,16,c\}$). Adjoining $(0,1,-1,0)$, $(1,0,-1,0)$, $(1,-1,0,0)$ instead yields a 68-ray, 35-basis KS parent containing a ray-critical 20-ray, 11-context parity proof $C_{20,11}$ ($11 = 4s(4) + 4s(5) + 2\sum_{i \neq 4,5} s(i)$). All 20 rays of $C_{20,11}$ belong to Peres's 24-ray set, and $C_{20,11}$ is one of its 96 critical 20-ray KS subsets, which are mutually isomorphic and include Kernaghan's set [14, 15].*

Proof. In $C_{18,9}$, summing s over C_3, C_4, C_5 gives $3 = 2(s(9) + s(10) + s(11)) + \sum_{k=12}^{17} s(k)$, so $s(12) + \cdots + s(17)$ is odd, whereas $D_1 \oplus D_4, D_2 \oplus D_5, D_3 \oplus D_6$ give $s(12) + s(13) \equiv s(2) + s(3) + s(6) + s(7)$, $s(14) + s(15) \equiv s(1) + s(3) + s(6) + s(8)$, $s(16) + s(17) \equiv s(1) + s(2) + s(7) + s(8)$ (mod 2), summing to 0 (mod 2). In $C_{20,11}$ (comprising

$C_1, \ldots, C_5$ and six contexts on $\{1, \ldots, 8, 12, \ldots, 17\}$ with $(0, 1, -1, 0)$, $(1, 0, -1, 0)$, $(1, -1, 0, 0)$), vertices $4, 5$ have degree 4 and the other 18 vertices have degree 2, so $11 = 4s(4) + 4s(5) + 2\sum_{i \neq 4,5} s(i)$, an odd-equals-even contradiction; explicit states after deleting any single ray are certified in Supplement Sec. S3. □

Both 68-ray presentations have 35 bases covering 192 of 195 geometric orthogonal pairs and context intersections of size up to two. They are not edge-faithful in the listed bases. The 72-ray Peres union covers all pairs; no size minimum or criticality of either full parent is asserted.

4. *Faithful completion $C_{36,18}$ and abstract versus geometric states on $C_{45,21}$:*

Corollary 30 (Faithful Cabello completion and vertex deletion)**.** *The 9 uncovered orthogonal pairs of $C_{18,9}$ satisfy Eq. (22), so Lemma 21 yields a linear edge-faithful 36-ray, 18-context KS hypergraph $C_{36,18}$ in $\mathbb{R}^4$ with $\omega = 4, \chi = 5$, and $\mathcal{S} = \varnothing$. Deleting one vertex (say c) and its two contexts from $C_{18,9}$ leaves 17 rays, 7 bases, 14 uncovered pairs, and 12 core states whose abstract completion $C_{45,21}$ has $\chi = 5$ and $|\mathcal{S}_{\mathrm{ctx}}(C_{45,21})| = 3{,}584$ unital separating context states (24 canonical). However, $v_{16} \in \mathrm{span}(v_1, v_2)$ and $v_{17} \in \mathrm{span}(v_7, v_8)$ violate Eq. (22), forcing four extra orthogonalities that reduce the geometric state space to $|\mathcal{S}_{\mathrm{geom}}(C_{45,21})| = 2{,}048$ with $s(9) = s(16) = s(17) = 0$, destroying unitality and separation (Table V).*

Proof. For $C_{18,9}$, the 9 uncovered pairs are the 6 core bridges $\{1, 13\}$, $\{2, 14\}$, $\{3, 16\}$, $\{6, 17\}$, $\{7, 15\}$, $\{8, 12\}$ from H_{65} plus $\{11, a\}$, $\{10, b\}$, $\{9, c\}$. They satisfy Eq. (22) and are completed in Supplemental Tables S28–S29. Deleting c, D_3, D_6 leaves 12 core supports: 10 miss 8 of the 14 uncovered pairs and 2 miss 9 (Table S84), giving $10 \cdot 2^8 + 2 \cdot 2^9 = 3{,}584$ context states. Since $v_{16} = v_1 + v_2$ and $v_{17} = \frac{1}{2}(v_7 - v_8)$, completing $\{1, 2\}$ by $\{21, 22\}$ and $\{7, 8\}$ by $\{35, 36\}$ forces $v_{16} \perp v_{21}, v_{22}$ and $v_{17} \perp v_{35}, v_{36}$ in the local labels of Tables S31–S32. If $s(16) = 1$, neither 1 nor 2 is selected in any admissible core support, so $s(21) + s(22) = 1$, contradicting geometric exclusion; the same argument gives $s(17) = 0$. The retained hub bases then force $s(9) = 0$. Exactly 8 core supports satisfy $s(9) = s(16) = s(17) = 0$, each missing 8 uncovered pairs. All their fresh-pair choices obey the four extra exclusions, yielding $8 \cdot 2^8 = 2{,}048$ geometric states. □

VIII. DEPENDENCIES, COUNTEREXAMPLES, AND REMAINING QUESTIONS

The common dependency structure is incidence → states and colorings, followed by independent tests of geometry, event order, and quantum probabilities. A chromatic argument can be made on an abstract hypergraph; its operational interpretation requires a ray realization. Conversely, a realization does not select an ontic state family or a classical probability mixture. The exact-palette assumption, that the outcomes admit a noncontextual assignment of d eigenvalues, adds a precise completion test at the level of whole truth assignments: a truth assignment is compatible with noncontextual eigenvalues exactly when the outcomes it leaves false can carry the remaining $d - 1$ eigenvalues.

Table VI also explains why the probability and gadget examples carry different force. A one-way ordinary TITS already gives a symmetric color identity; a two-way purely chromatic identity requires counterstates in both truth directions. A ternary color exclusion may survive even when every pair can be equal. A strict inclusion $\mathcal{P}_{\mathrm{ext}} \subsetneq \mathcal{P}$ therefore need not be detectable through pair equalities alone.

Open problems concern global minimum faithful examples beyond the specified inventory, economical rational completions with fewer degree-one rays, residual-coloring certificates in dimensions above three, and geometric ancestry in which the retained rays are essential to the contradiction. The current statements establish existence and the recorded finite classifications, without asserting these stronger minima or essential-ancestry properties. All complete enumerations and the detailed verification scope remain in the supplement and electronic records.

IX. CONCLUSION

We started from a simple physical requirement on intertwined complete experiments: that each is the measurement of a maximal, nondegenerate observable with one common spectrum of d eigenvalues, and that the eigenvalue carried by an outcome is an element of physical reality in the sense of Einstein, Podolsky, and Rosen [3], the same in every experiment in which the outcome occurs. This requirement of noncontextual eigenvalues is exactly strong d-colorability of the context hypergraph. It asks for more than Kochen–Specker noncontextuality: not one classical truth assignment, but d of them with disjoint supports covering every outcome. Its failure, $\chi(H) > d$, is therefore a weaker but more widespread form of contextuality. Completed Yu–Oh has 24 truth assignments separating all of its rays, yet its outcomes cannot carry three eigenvalues noncontextually.

By unifying these common-spectrum chromatic obstructions ($\chi(H) > d$) with exact-palette chromatic separability constraints ($\chi(H) = d$, $\mathcal{S}_{\mathrm{ext}}(H) \subsetneq \mathcal{S}(H)$), we have distinguished the requirements across noncontextual eigenvalue assignments, geometric ray realizations, orthomodular event representations, and quantum probability models. At the foundation, $\chi(H) > d$ excludes noncontextual eigenvalues for uniform maximal context observables, whereas the exclusion of an injective realization for H_8 and of every ray assignment for

TABLE V. Context and geometric state spaces in the four-dimensional examples. Linearity of the listed incidence structure does not imply edge-faithfulness.

Configuration	$\lvert V\rvert/\lvert E\rvert$	$\lvert\mathcal{S}_{\mathrm{ctx}}\rvert$	$\lvert\mathcal{S}_{\mathrm{geom}}\rvert$	Scope
$C_{18,9}$	18/9	0	0	Nine uncovered geometric pairs; parity KS proof.
$C_{36,18}$	36/18	0	0	Faithful linear completion.
$C_{45,21}$	45/21	3,584	2,048	Four additional pairs destroy geometric unitality and separation.
H_{65}	65/29	393,216	393,216	Edge-faithful; all context states respect the geometry.

TABLE VI. Logical boundaries of the integrated results. Each row prevents transfer of a conclusion between different assignment or representation problems.

Tempting unrestricted inference	Counterexample or necessary qualification
$\chi > d$ excludes states or ray realizations	Completed Yu–Oh has faithful real rays and 24 unital separating states.
Ordinary states separate ⇒ colors separate	Faithful $H_{17,10}$ and $H_{21,13}$ have forced color identities.
Colors separate ⇒ every state extends	The pentagon separates all vertex pairs but has one nonextendable state.
Vertex separation ⇒ event-order determination	$H_{17,10}$ has 10 ⇒ 24; H_{65} has 16 ⇒ 9 and ray–plane collapse.
Order determination ⇒ all Born mixtures	The specified event order of H_{37} is represented, while $\rho_{33} > 1/3$ violates the state inequality.
$\chi_f > d$ establishes fixed-measurement state independence	Ref. [26, Thm. 1] (Yu–Oh plus apex); linear and apex-free: H_{65} admits the Born model of $(I_4 - Q)/3$ on all 65 rays.
Abstract stateless extension ⇒ geometric KS parent	Closing a color-identical pair changes orthogonality; transverse four-loops exclude injective triadic geometry.
Generic KS ancestry explains the original chromatic gap	Unrestricted extension holds even for colorable sets and need not use the old rays essentially.

H_9, G_{32} requires direct Hermitian projection forcing, correcting the earlier spectral argument for G_{32} [6] and establishing four-quadrant independence even at Berge girth five. Within chromatically complete hypergraphs ($\chi(H) = d$), where noncontextual eigenvalue assignments exist, our completion criterion ($\chi(\Gamma(H)[V \setminus \operatorname{supp} s]) \leq d - 1$) characterizes which truth assignments arise from noncontextual eigenvalue assignments and explains the specified examples of chromatic identities ($u \sim_c v$) and purely chromatic TIFS/TITS/ternary constraints ($H_{17,10}, H_{21,13}, D_{16,9}, T_{19,11}, W^{(4)}_{16,6}$) and yields measurable nonextendable weights 1/9 and 1/3 for quantum ray preparations on $H_{17,10}$, the latter being the largest value a single chromatic identity can certify there. In higher complexity, the Hesse hypergraph H_{37} (6/7/18 OML minima) and our rational four-dimensional completion H_{65} (393,216 states) demonstrate that neither exact order determination nor a fractional chromatic gap decides classicality for all Born preparations, while the elementary KS ancestry result, the 68-ray KS parents of H_{65}, and the completions $C_{36,18}$ and $C_{45,21}$ of $C_{18,9}$ delineate unrestricted embedding from specified critical parents and geometric pair exclusions.

The common thread is that a classical account of quantum outcomes can succeed at the level of truth values and still fail at the level of eigenvalues. For the preparation along z_3, the quantum predictions on $H_{17,10}$ admit noncontextual truth-assignment models, but every such model must place at least one third of its weight on truth assignments that no noncontextual eigenvalue assignment produces.

ACKNOWLEDGMENTS

I thank Mohammad Hadi Shekarriz for valuable comments and suggestions. This research was funded in part by the Austrian Science Fund (FWF), Grant DOI 10.55776/PIN5424624. The author acknowledges FWF and TU Wien Bibliothek for support through the Open Access Funding Programme. AI tools assisted with editorial work and computational checks. The author retains responsibility.

[1] J. von Neumann, Annals of Mathematics **32**, 191 (1931).
[2] P. R. Halmos, *Finite-Dimensional Vector Spaces*, Undergraduate Texts in Mathematics (Springer, New York, 1974).

[3] A. Einstein, B. Podolsky, and N. Rosen, Physical Review **47**, 777 (1935).
[4] S. Kochen and E. P. Specker, Journal of Mathematics and Mechanics **17**, 59 (1967).
[5] K. Svozil, Quantum Reports **2**, 278 (2020), arXiv:1808.00813.
[6] M. H. Shekarriz and K. Svozil, Journal of Mathematical Physics **63**, 032104 (2022), arXiv:2105.08520.
[7] K. Svozil, Entropy **27**, 387 (2025), arXiv:2501.15261.
[8] R. J. Greechie, Journal of Combinatorial Theory. Series A **10**, 119 (1971).
[9] S. Yu and C. H. Oh, Physical Review Letters **108**, 030402 (2012), arXiv:1109.4396.
[10] J. A. Martin H., Graph-chromatic implicit relations (2009), arXiv:0901.1255 [math.CO].
[11] M. Navara and K. Svozil, Construction of Kochen-Specker sets from mutually unbiased bases (2025), arXiv:2509.08636 [quant-ph].
[12] G. Haynes, C. Park, A. Schaeffer, J. Webster, and L. H. Mitchell, The Electronic Journal of Combinatorics **17**, R55 (2010).
[13] J. Bub, Foundations of Physics **26**, 787 (1996).
[14] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
[15] M. Kernaghan, Journal of Physics A: Mathematical and General **27**, L829 (1994).
[16] K. Svozil, Supplemental material for "Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments" (2026), contains all explicit unnormalized vector and context inventories, complete two-valued state and coloring enumerations, odd-cycle and separation profile certificates, and supplementary proofs and verification reductions.
[17] A. Silva and A. Ibiapina, On orthogonal vector edge coloring (2019), arXiv:1909.01918 [cs.DM].
[18] G. Brinkmann, M. Preissmann, and D. Sasaki, Discrete Mathematics & Theoretical Computer Science **17**, 369 (2015).
[19] G. Schrag, Discrete Mathematics **70**, 185 (1988).
[20] K. Svozil and J. Tkadlec, Journal of Mathematical Physics **37**, 5380 (1996).
[21] L. Hogben, K. F. Palmowski, D. E. Roberson, and S. Severini, Electronic Journal of Linear Algebra **32**, 98 (2017).
[22] M. Waegell and P. K. Aravind, Foundations of Physics **41**, 1786 (2011), arXiv:1103.6058 [quant-ph].
[23] R. Isaacs, The American Mathematical Monthly **82**, 221 (1975).
[24] D. Blanuša, Glasnik Matematičko-Fizički i Astronomski, Serija II **1**, 31 (1946).
[25] M. Pavičić, Entropy **21**, 1107 (2019).
[26] A. Cabello, M. Kleinmann, and C. Budroni, Physical Review Letters **114**, 250402 (2015), arXiv:1501.03432.
[27] S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
[28] A. Cabello, J. R. Portillo, A. Solís, and K. Svozil, Physical Review A **98**, 012106 (2018), arXiv:1805.00796.
[29] J. Harding and R. Salinas Schmeis, International Journal of Theoretical Physics **64**, 211 (2025), arXiv:2505.13871 [math-ph].
[30] A. Peres, Journal of Physics A: Mathematical and General **24**, L175 (1991).
[31] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **339**, 425 (2005).
[32] M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
[33] A. Cabello, Physical Review Letters **135**, 190203 (2025), arXiv:2508.07335 [quant-ph].
[34] M. Pavičić, On simplest Kochen-Specker sets (2025), arXiv:2512.10483 [quant-ph].
[35] M. Pavičić, Quantum **7**, 953 (2023).

Supplemental Material for "Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments"

Karl Svozil
Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, 1040 Vienna, Austria

(Dated: November 5, 2026)

This Supplemental Material provides the complete vector and context enumerations, two-valued state tables, normalized strong coloring tables, odd-cycle nonextendability certificates, chromatic signature and separation profile tables, parent-embedding dictionaries, and supplementary proofs and finite verification certificates supporting the main text. Section S1 lists the explicit unnormalized coordinates and context tables for the completed Yu–Oh hypergraph H^{+}_{YO} and its faithful restrictions $H_{17,10}$ and $H_{21,13}$, the three- and four-dimensional logical gadgets, the 37-ray complex Hesse hypergraph H_{37}, the 26-ray suspension $H^{(4)}_{\mathrm{YO}}$ and 65-ray canonical linear completion H_{65}, the Cabello completions $C_{36,18}$ and $C_{45,21}$, and the specified finite KS parents (49-ray Schütte–Bub, 87-ray MUB-inventory union, 40- and 72-ray Peres unions, and 68-ray three-ray extensions). Section S2 gives the complete two-valued state tables, extendable-state classifications, normalized coloring tables, odd-cycle certificates, and separation profiles across both regimes. Section S3 presents the supplementary proofs and exact reductions, including Isaacs dot-product snarks and quantitative Δ_3 robustness bounds for G_{32}, the 540,672-candidate reduction for the minimum 18-state order-determining family on $L(H_{37})$, the canonical completion selection rule for H_{65}, the forced-zero and ray-criticality certificates for the four-dimensional Cabello completions, transverse criticality deletion colorings, and historical Kochen–Specker gadget certificates.

READING GUIDE AND VERIFICATION SCOPE

This supplement collects complete vector and context inventories (S1), state and coloring enumerations (S2), and extended proofs and certificates (S3). Labels are local to each configuration. "Exact palette" means $\chi(H) = d$, with no additional density or geometric assumption: the outcomes admit a noncontextual assignment of d eigenvalues, that is, every complete experiment can be measured by a maximal observable with one common spectrum of d eigenvalues, with each outcome carrying the same eigenvalue in every context. Colorings are such eigenvalue assignments, and extendable states are the truth assignments "the outcome with eigenvalue λ_a occurs" that they produce.

The five-state and four-state $H_{17,10}$ mixtures and both-direction gadget witnesses use the complete state families. Every selected support satisfies every context equation, and rational arithmetic verifies every Born marginal and the attained nonextendable weights $1/9$ and $1/3$. The ordinary implication $10 \Rightarrow 24$ on $H_{17,10}$ gives a false inclusion between distinct rays and hence a failure of order determination.

The companion electronic data include all coordinates, state enumerations (including the compressed 393,216-state H_{65} inventory), verification programs, and the exhaustive certificates for the eighteen-state Hesse minimum and the 65,535-subset Yu–Oh classification. Checking an eighteen-state family establishes an upper bound; the matching lower bound requires the exhaustive certificate in Sec. S3.3. The directories `part-a/data`, `part-a/verification`, `part-b/data`, and `part-b/verification` contain the underlying data and verifiers. Reports and source-integrity hashes are in `audit/`; Sec. S3.7 describes the joint verification entry point and its scope.

S1. EXPLICIT UNNORMALIZED VECTORS AND CONTEXTS

All numerical vertex labels in this section are local to the named configuration and match the figures and theorems of the main text.

S1.1. Completed Yu–Oh and the Petersen-derived contexts

The three-dimensional vectors below give all projective rays used in Sec. III C; signs are chosen to make the first nonzero integer coordinate positive. The 26-ray suspension appends zero and adds e_4. The abstract G_{32}, H_9, H_8 context lists in Eq. (10) and Theorem 11 specify their incidences completely. The geometric obstructions exclude a corresponding injective three-dimensional vector enumeration for these abstract examples.

TABLE S1: All 25 unnormalized completed Yu–Oh vectors.

i	r_i	i	r_i
1	$(0,1,1)$	14	$(2,1,1)$
2	$(1,0,0)$	15	$(1,2,-1)$
3	$(0,1,-1)$	16	$(1,-2,1)$
4	$(1,1,0)$	17	$(1,1,-2)$
5	$(0,0,1)$	18	$(1,-1,-2)$
6	$(1,-1,0)$	19	$(2,1,-1)$
7	$(1,0,1)$	20	$(2,-1,1)$
8	$(0,1,0)$	21	$(1,-2,-1)$
9	$(1,0,-1)$	22	$(1,1,2)$
10	$(1,-1,1)$	23	$(1,2,1)$
11	$(1,-1,-1)$	24	$(1,-1,2)$
12	$(1,1,1)$	25	$(2,-1,-1)$
13	$(1,1,-1)$		

S1.2. Faithful Yu–Oh restrictions $H_{17,10}$ and $H_{21,13}$ and logical gadgets in dimensions three and four

Tables S2 and S3 record the restriction membership of the 25 parent rays and 16 parent contexts of H_{YO}^{+} in $H_{17,10}$ and $H_{21,13}$. Tables S4–S14 provide the explicit unnormalized integer coordinates and context lists for the faithful real 3-dimensional gadgets P_5, $B_{13,7}$, $D_{16,9}$, $I_{16,9}$, and $T_{19,11}$ and the faithful real 4-dimensional gadgets $W_{11,5}^{(4)}$, $W_{16,6}^{(4)}$, $B_{14,7}^{(4)}$, $L_{18,7}^{(4)}$, $D_{17,9}^{(4)}$, and $L_{22,9}^{(4)}$. Note that for the standalone gadgets P_5, $D_{16,9}$, $I_{16,9}$, and $T_{19,11}$, the vertex numbering is local to each gadget panel; in $L_{18,7}^{(4)}$ [Fig. 3(d)], $r_{13} = (1,0,1,1)$ and $r_{14} = (2,0,-1,-1)$ are chosen so that $r_{13} \not\perp r_{15}$, whereas in $L_{22,9}^{(4)}$ [Fig. 3(f)], $r_{13} = (1,0,4,4)$ and $r_{14} = (8,0,-1,-1)$ are used so that $r_{13} \perp r_{15}$ in the added context $\{13,20,15,19\}$.

TABLE S2: All 25 unnormalized parent vectors; membership of the two geometric examples.

i	r_i	$H_{17,10}$	$H_{21,13}$	i	r_i	$H_{17,10}$	$H_{21,13}$
1	$(0,1,1)$	yes	yes	14	$(2,1,1)$	yes	yes
2	$(1,0,0)$	yes	yes	15	$(1,2,-1)$	yes	yes
3	$(0,1,-1)$	yes	yes	16	$(1,-2,1)$	–	–
4	$(1,1,0)$	yes	yes	17	$(1,1,-2)$	–	–
5	$(0,0,1)$	yes	yes	18	$(1,-1,-2)$	yes	yes
6	$(1,-1,0)$	yes	yes	19	$(2,1,-1)$	yes	yes
7	$(1,0,1)$	yes	yes	20	$(2,-1,1)$	–	yes
8	$(0,1,0)$	yes	yes	21	$(1,-2,-1)$	–	yes
9	$(1,0,-1)$	yes	yes	22	$(1,1,2)$	–	yes
10	$(1,-1,1)$	yes	yes	23	$(1,2,1)$	yes	yes
11	$(1,-1,-1)$	yes	yes	24	$(1,-1,2)$	yes	yes
12	$(1,1,1)$	–	–	25	$(2,-1,-1)$	–	–
13	$(1,1,-1)$	–	yes				

TABLE S3: The 16 parent contexts and the two selected restrictions.

j	context	$H_{17,10}$	$H_{21,13}$
1	$\{1,2,3\}$	yes	yes
2	$\{7,8,9\}$	yes	yes
3	$\{4,5,6\}$	yes	yes
4	$\{2,8,5\}$	yes	yes
5	$\{3,12,25\}$	–	–
6	$\{9,12,16\}$	–	–
7	$\{6,12,17\}$	–	–
8	$\{3,11,14\}$	yes	yes
9	$\{7,11,15\}$	yes	yes
10	$\{4,11,24\}$	yes	yes
11	$\{9,10,23\}$	yes	yes
12	$\{1,10,19\}$	yes	yes
13	$\{4,10,18\}$	yes	yes
14	$\{1,13,20\}$	–	yes
15	$\{7,13,21\}$	–	yes
16	$\{6,13,22\}$	–	yes

S1.3. 3D panel (a): P_5

TABLE S4: Unnormalized vectors and all contexts of P_5.

i	r_i	j	Context
1	$(1,0,0)$	1	$\{1,6,2\}$
2	$(0,1,1)$	2	$\{2,7,3\}$
3	$(1,1,-1)$	3	$\{3,8,4\}$
4	$(1,-2,-1)$	4	$\{4,9,5\}$
5	$(0,1,-2)$	5	$\{5,10,1\}$
6	$(0,1,-1)$		
7	$(2,-1,1)$		
8	$(1,0,1)$		
9	$(5,2,1)$		
10	$(0,2,1)$		

S1.4. 3D panel (b): $B_{13,7}$

TABLE S5: Unnormalized vectors and all contexts of $B_{13,7}$.

i	r_i	j	Context
1	$(0,1,1)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,1,-1)$	3	$\{2,8,5\}$
5	$(0,0,1)$	4	$\{3,11,14\}$
7	$(1,0,1)$	5	$\{7,11,15\}$
8	$(0,1,0)$	6	$\{9,10,23\}$
9	$(1,0,-1)$	7	$\{1,10,19\}$
10	$(1,-1,1)$		
11	$(1,-1,-1)$		
14	$(2,1,1)$		
15	$(1,2,-1)$		
19	$(2,1,-1)$		
23	$(1,2,1)$		

S1.5. 3D panel (c): $D_{16,9}$

TABLE S6: Unnormalized vectors and all contexts of $D_{16,9}$.

i	r_i	j	Context
1	$(0,1,2)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,2,-1)$	3	$\{4,5,6\}$
4	$(1,2,0)$	4	$\{2,8,5\}$
5	$(0,0,1)$	5	$\{3,11,14\}$
6	$(2,-1,0)$	6	$\{7,11,15\}$
7	$(1,0,1)$	7	$\{4,11,24\}$
8	$(0,1,0)$	8	$\{9,10,23\}$
9	$(1,0,-1)$	9	$\{4,10,18\}$
10	$(2,-1,2)$		
11	$(2,-1,-2)$		
14	$(5,2,4)$		
15	$(1,4,-1)$		
18	$(4,-2,-5)$		
23	$(1,4,1)$		
24	$(4,-2,5)$		

S1.6. 3D panel (d): $I_{16,9}$

TABLE S7: Unnormalized vectors and all contexts of $I_{16,9}$.

i	r_i	j	Context
1	$(0,1,1)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,1,-1)$	3	$\{2,8,5\}$
4	$(4,5,3)$	4	$\{3,11,14\}$
5	$(0,0,1)$	5	$\{7,11,15\}$
7	$(1,0,2)$	6	$\{4,11,24\}$
8	$(0,1,0)$	7	$\{9,10,23\}$
9	$(2,0,-1)$	8	$\{1,10,19\}$
10	$(1,-2,2)$	9	$\{4,10,18\}$
11	$(2,-1,-1)$		
14	$(1,1,1)$		
15	$(2,5,-1)$		
18	$(16,-5,-13)$		
19	$(4,1,-1)$		
23	$(2,5,4)$		
24	$(1,-5,7)$		

S1.7. 3D panel (e): $T_{19,11}$

TABLE S8: Unnormalized vectors and all contexts of $T_{19,11}$.

i	r_i	j	Context
1	$(0,1,2)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,2,-1)$	3	$\{4,5,6\}$
4	$(1,2,0)$	4	$\{2,8,5\}$
5	$(0,0,1)$	5	$\{3,11,14\}$
6	$(2,-1,0)$	6	$\{7,11,15\}$
7	$(1,0,1)$	7	$\{4,11,24\}$
8	$(0,1,0)$	8	$\{9,10,23\}$
9	$(1,0,-1)$	9	$\{4,10,18\}$
10	$(2,-1,2)$	10	$\{14,26,27\}$
11	$(2,-1,-2)$	11	$\{18,26,28\}$
14	$(5,2,4)$		
15	$(1,4,-1)$		
18	$(4,-2,-5)$		
23	$(1,4,1)$		
24	$(4,-2,5)$		
26	$(2,-41,18)$		
27	$(200,-82,-209)$		
28	$(241,82,160)$		

S1.8. 4D panel (a): $W^{(4)}_{11,5}$

TABLE S9: Unnormalized vectors and all contexts of $W^{(4)}_{11,5}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{0,1,6,2\}$
1	$(1,0,0,0)$	2	$\{0,2,7,3\}$
2	$(0,1,1,0)$	3	$\{0,3,8,4\}$
3	$(1,1,-1,0)$	4	$\{0,4,9,5\}$
4	$(1,-2,-1,0)$	5	$\{0,5,10,1\}$
5	$(0,1,-2,0)$		
6	$(0,1,-1,0)$		
7	$(2,-1,1,0)$		
8	$(1,0,1,0)$		
9	$(5,2,1,0)$		
10	$(0,2,1,0)$		

S1.9. 4D panel (b): $W^{(4)}_{16,6}$

TABLE S10: Unnormalized vectors and all contexts of $W^{(4)}_{16,6}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{2,0,6,1\}$
1	$(1,0,0,0)$	2	$\{2,7,13,3\}$
2	$(0,1,1,0)$	3	$\{3,0,8,4\}$
3	$(1,1,-1,0)$	4	$\{4,9,14,5\}$
4	$(1,-2,-1,0)$	5	$\{5,10,15,1\}$
5	$(0,1,-2,0)$	6	$\{0,11,12,5\}$
6	$(0,1,-1,0)$		
7	$(2,-1,1,1)$		
8	$(1,0,1,0)$		
9	$(5,2,1,1)$		
10	$(0,4,2,1)$		
11	$(1,2,1,0)$		
12	$(5,-2,-1,0)$		
13	$(2,-1,1,-6)$		
14	$(5,2,1,-30)$		
15	$(0,2,1,-10)$		

S1.10. 4D panel (c): $B^{(4)}_{14,7}$

TABLE S11: Unnormalized vectors and all contexts of $B^{(4)}_{14,7}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{0,1,2,3\}$
1	$(0,1,1,0)$	2	$\{0,7,8,9\}$
2	$(1,0,0,0)$	3	$\{0,2,8,5\}$
3	$(0,1,-1,0)$	4	$\{0,3,11,14\}$
5	$(0,0,1,0)$	5	$\{0,7,11,15\}$
7	$(1,0,1,0)$	6	$\{0,1,10,19\}$
8	$(0,1,0,0)$	7	$\{0,9,10,23\}$
9	$(1,0,-1,0)$		
10	$(1,-1,1,0)$		
11	$(1,-1,-1,0)$		
14	$(2,1,1,0)$		
15	$(1,2,-1,0)$		
19	$(2,1,-1,0)$		
23	$(1,2,1,0)$		

S1.11. 4D panel (d): $L^{(4)}_{18,7}$

TABLE S12: Unnormalized vectors and all contexts of $L^{(4)}_{18,7}$.

i	r_i	j	Context
1	$(1,0,0,0)$	1	$\{1,2,4,3\}$
2	$(0,1,0,0)$	2	$\{5,8,7,6\}$
3	$(0,0,1,0)$	3	$\{4,9,10,8\}$
4	$(0,0,0,1)$	4	$\{11,1,12,5\}$
5	$(0,1,-1,-1)$	5	$\{11,14,13,2\}$
6	$(4,-5,-11,6)$	6	$\{15,3,16,7\}$
7	$(1,-4,0,-4)$	7	$\{15,17,18,6\}$
8	$(4,1,1,0)$		
9	$(1,-5,1,0)$		
10	$(2,-1,-7,0)$		
11	$(0,0,1,-1)$		
12	$(0,2,1,1)$		
13	$(1,0,1,1)$		
14	$(2,0,-1,-1)$		
15	$(4,2,0,-1)$		
16	$(4,-5,0,6)$		
17	$(5,-9,7,2)$		
18	$(2,113,77,234)$		

S1.12. 4D panel (e): $D^{(4)}_{17,9}$

TABLE S13: Unnormalized vectors and all contexts of $D^{(4)}_{17,9}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{0,1,2,3\}$
1	$(0,1,2,0)$	2	$\{0,7,8,9\}$
2	$(1,0,0,0)$	3	$\{0,4,5,6\}$
3	$(0,2,-1,0)$	4	$\{0,2,8,5\}$
4	$(1,2,0,0)$	5	$\{0,3,11,14\}$
5	$(0,0,1,0)$	6	$\{0,7,11,15\}$
6	$(2,-1,0,0)$	7	$\{0,9,10,23\}$
7	$(1,0,1,0)$	8	$\{0,4,11,24\}$
8	$(0,1,0,0)$	9	$\{0,4,10,18\}$
9	$(1,0,-1,0)$		
10	$(2,-1,2,0)$		
11	$(2,-1,-2,0)$		
14	$(5,2,4,0)$		
15	$(1,4,-1,0)$		
18	$(4,-2,-5,0)$		
23	$(1,4,1,0)$		
24	$(4,-2,5,0)$		

S1.13. 4D panel (f): $L^{(4)}_{22,9}$

TABLE S14: Unnormalized vectors and all contexts of $L^{(4)}_{22,9}$.

i	r_i	j	Context
1	$(1,0,0,0)$	1	$\{1,2,4,3\}$
2	$(0,1,0,0)$	2	$\{5,8,7,6\}$
3	$(0,0,1,0)$	3	$\{4,9,10,8\}$
4	$(0,0,0,1)$	4	$\{11,1,12,5\}$
5	$(0,1,-1,-1)$	5	$\{11,14,13,2\}$
6	$(4,-5,-11,6)$	6	$\{15,3,16,7\}$
7	$(1,-4,0,-4)$	7	$\{15,17,18,6\}$
8	$(4,1,1,0)$	8	$\{11,21,22,19\}$
9	$(1,-5,1,0)$	9	$\{13,20,15,19\}$
10	$(2,-1,-7,0)$		
11	$(0,0,1,-1)$		
12	$(0,2,1,1)$		
13	$(1,0,4,4)$		
14	$(8,0,-1,-1)$		
15	$(4,2,0,-1)$		
16	$(4,-5,0,6)$		
17	$(5,-9,7,2)$		
18	$(2,113,77,234)$		
19	$(16,-33,-2,-2)$		
20	$(4,2,-21,20)$		
21	$(22,12,-11,-11)$		
22	$(129,44,153,153)$		

S1.14. Hesse H_{37}: vectors, contexts, and non-collinearity

This appendix uses exactly the vertex numbers and context numbers of Fig. 4. All indices a, k in the family notation are in $\mathbb{Z}_3$. The vectors are unnormalized and the inner product is Hermitian. Table S15 lists every vector, squared norm, color, and incident context. Table S16 lists all 24 contexts with their actual vectors. Together they specify the faithful orthogonal representation completely.

S1.14.1. All 37 vectors, their colors, and incident contexts

TABLE S15: Complete enumeration of the 37 vectors of H_{37}. The color $c(i) \in \{0,1,2,3\}$ is the explicit four-coloring in the main text. Context indices refer to Table S16.

i	Ray	v_i	$\|v_i\|^2$	$c(i)$	Incident contexts
1	z	$(1,0,0)$	1	0	$\{4,5,6\}$
2	M_{00}	$(1,1,1)$	3	0	$\{1,7,16\}$
3	M_{01}	$(1,\zeta,\zeta^2)$	3	1	$\{1,8,17\}$
4	M_{02}	$(1,\zeta^2,\zeta)$	3	2	$\{1,9,18\}$
5	M_{10}	$(1,\zeta,\zeta)$	3	0	$\{2,10,21\}$
6	M_{11}	$(1,\zeta^2,1)$	3	1	$\{2,11,19\}$
7	M_{12}	$(1,1,\zeta^2)$	3	2	$\{2,12,20\}$
8	M_{20}	$(1,\zeta^2,\zeta^2)$	3	0	$\{3,13,23\}$
9	M_{21}	$(1,1,\zeta)$	3	1	$\{3,14,24\}$
10	M_{22}	$(1,\zeta,1)$	3	2	$\{3,15,22\}$
11	P_0	$(0,1,-1)$	2	3	$\{4,7,10,13\}$
12	P_1	$(0,1,-\zeta)$	2	3	$\{5,8,11,14\}$
13	P_2	$(0,1,-\zeta^2)$	2	3	$\{6,9,12,15\}$
14	Q_0	$(-1,0,1)$	2	3	$\{16,19,22\}$
15	Q_1	$(-\zeta,0,1)$	2	3	$\{17,20,23\}$
16	Q_2	$(-\zeta^2,0,1)$	2	3	$\{18,21,24\}$
17	A_0	$(0,1,1)$	2	1	$\{4\}$
18	A_1	$(0,1,\zeta)$	2	1	$\{5\}$
19	A_2	$(0,1,\zeta^2)$	2	1	$\{6\}$
20	B_{00}	$(-2,1,1)$	6	1	$\{7\}$
21	B_{01}	$(-2,\zeta,\zeta^2)$	6	2	$\{8\}$
22	B_{02}	$(-2,\zeta^2,\zeta)$	6	0	$\{9\}$

Continued on the next page.

TABLE S15: Complete enumeration of the 37 vectors of H_{37}. The color $c(i) \in \{0,1,2,3\}$ is the explicit four-coloring in the main text. Context indices refer to Table S16. (continued).

i	Ray	v_i	$\|v_i\|^2$	$c(i)$	Incident contexts
23	B_{10}	$(-2,\zeta,\zeta)$	6	1	{10}
24	B_{11}	$(-2,\zeta^2,1)$	6	2	{11}
25	B_{12}	$(-2,1,\zeta^2)$	6	0	{12}
26	B_{20}	$(-2,\zeta^2,\zeta^2)$	6	1	{13}
27	B_{21}	$(-2,1,\zeta)$	6	2	{14}
28	B_{22}	$(-2,\zeta,1)$	6	0	{15}
29	C_{00}	$(1,-2,1)$	6	1	{16}
30	C_{01}	$(1,-2\zeta,\zeta^2)$	6	2	{17}
31	C_{02}	$(1,-2\zeta^2,\zeta)$	6	0	{18}
32	C_{10}	$(1,-2\zeta^2,1)$	6	2	{19}
33	C_{11}	$(1,-2,\zeta^2)$	6	0	{20}
34	C_{12}	$(1,-2\zeta,\zeta)$	6	1	{21}
35	C_{20}	$(1,-2\zeta,1)$	6	0	{22}
36	C_{21}	$(1,-2\zeta^2,\zeta^2)$	6	1	{23}
37	C_{22}	$(1,-2,\zeta)$	6	2	{24}

S1.14.2. All 24 contexts and their orthogonal vectors

TABLE S16: All 24 orthogonal triads, numbered in the same order as the context colors in Fig. 4. Vectors are given in increasing order of their vertex numbers. Each triad contributes three orthogonality edges; the 72 resulting pairs are all the orthogonal pairs.

Context	Family	Vertices	Unnormalized vectors
e_1	$\mathcal{M}_0$	$\{2,3,4\}$	$\{(1,1,1),\ (1,\zeta,\zeta^2),\ (1,\zeta^2,\zeta)\}$
e_2	$\mathcal{M}_1$	$\{5,6,7\}$	$\{(1,\zeta,\zeta),\ (1,\zeta^2,1),\ (1,1,\zeta^2)\}$
e_3	$\mathcal{M}_2$	$\{8,9,10\}$	$\{(1,\zeta^2,\zeta^2),\ (1,1,\zeta),\ (1,\zeta,1)\}$
e_4	$\mathcal{A}_0$	$\{1,11,17\}$	$\{(1,0,0),\ (0,1,-1),\ (0,1,1)\}$
e_5	$\mathcal{A}_1$	$\{1,12,18\}$	$\{(1,0,0),\ (0,1,-\zeta),\ (0,1,\zeta)\}$
e_6	$\mathcal{A}_2$	$\{1,13,19\}$	$\{(1,0,0),\ (0,1,-\zeta^2),\ (0,1,\zeta^2)\}$
e_7	$\mathcal{B}_{00}$	$\{2,11,20\}$	$\{(1,1,1),\ (0,1,-1),\ (-2,1,1)\}$
e_8	$\mathcal{B}_{01}$	$\{3,12,21\}$	$\{(1,\zeta,\zeta^2),\ (0,1,-\zeta),\ (-2,\zeta,\zeta^2)\}$
e_9	$\mathcal{B}_{02}$	$\{4,13,22\}$	$\{(1,\zeta^2,\zeta),\ (0,1,-\zeta^2),\ (-2,\zeta^2,\zeta)\}$
e_{10}	$\mathcal{B}_{10}$	$\{5,11,23\}$	$\{(1,\zeta,\zeta),\ (0,1,-1),\ (-2,\zeta,\zeta)\}$
e_{11}	$\mathcal{B}_{11}$	$\{6,12,24\}$	$\{(1,\zeta^2,1),\ (0,1,-\zeta),\ (-2,\zeta^2,1)\}$
e_{12}	$\mathcal{B}_{12}$	$\{7,13,25\}$	$\{(1,1,\zeta^2),\ (0,1,-\zeta^2),\ (-2,1,\zeta^2)\}$
e_{13}	$\mathcal{B}_{20}$	$\{8,11,26\}$	$\{(1,\zeta^2,\zeta^2),\ (0,1,-1),\ (-2,\zeta^2,\zeta^2)\}$
e_{14}	$\mathcal{B}_{21}$	$\{9,12,27\}$	$\{(1,1,\zeta),\ (0,1,-\zeta),\ (-2,1,\zeta)\}$
e_{15}	$\mathcal{B}_{22}$	$\{10,13,28\}$	$\{(1,\zeta,1),\ (0,1,-\zeta^2),\ (-2,\zeta,1)\}$
e_{16}	$\mathcal{C}_{00}$	$\{2,14,29\}$	$\{(1,1,1),\ (-1,0,1),\ (1,-2,1)\}$

Continued on the next page.

TABLE S16: All 24 orthogonal triads, numbered in the same order as the context colors in Fig. 4. Vectors are given in increasing order of their vertex numbers. Each triad contributes three orthogonality edges; the 72 resulting pairs are all the orthogonal pairs. (continued).

Context	Family	Vertices	Unnormalized vectors
e_{17}	$\mathcal{C}_{01}$	$\{3,15,30\}$	$\{(1,\zeta,\zeta^2),\ (-\zeta,0,1),\ (1,-2\zeta,\zeta^2)\}$
e_{18}	$\mathcal{C}_{02}$	$\{4,16,31\}$	$\{(1,\zeta^2,\zeta),\ (-\zeta^2,0,1),\ (1,-2\zeta^2,\zeta)\}$
e_{19}	$\mathcal{C}_{10}$	$\{6,14,32\}$	$\{(1,\zeta^2,1),\ (-1,0,1),\ (1,-2\zeta^2,1)\}$
e_{20}	$\mathcal{C}_{11}$	$\{7,15,33\}$	$\{(1,1,\zeta^2),\ (-\zeta,0,1),\ (1,-2,\zeta^2)\}$
e_{21}	$\mathcal{C}_{12}$	$\{5,16,34\}$	$\{(1,\zeta,\zeta),\ (-\zeta^2,0,1),\ (1,-2\zeta,\zeta)\}$
e_{22}	$\mathcal{C}_{20}$	$\{10,14,35\}$	$\{(1,\zeta,1),\ (-1,0,1),\ (1,-2\zeta,1)\}$
e_{23}	$\mathcal{C}_{21}$	$\{8,15,36\}$	$\{(1,\zeta^2,\zeta^2),\ (-\zeta,0,1),\ (1,-2\zeta^2,\zeta^2)\}$
e_{24}	$\mathcal{C}_{22}$	$\{9,16,37\}$	$\{(1,1,\zeta),\ (-\zeta^2,0,1),\ (1,-2,\zeta)\}$

S1.14.3. Norm shells and a non-collinearity certificate

The four squared-norm shells have sizes

$$\#\{i:\|v_i\|^2=1\}=1,\quad \#\{i:\|v_i\|^2=2\}=9,\quad \#\{i:\|v_i\|^2=3\}=9,\quad \#\{i:\|v_i\|^2=6\}=18. \tag{S1}$$

Exact evaluation over the $\binom{37}{2}=666$ unordered pairs gives

$$\max_{i<j}\frac{|\langle v_i,v_j\rangle|^2}{\|v_i\|^2\|v_j\|^2}=\frac{3}{4}<1. \tag{S2}$$

For example, the maximum is attained by $v_{11}=P_0=(0,1,-1)$ and $v_{29}=C_{00}=(1,-2,1)$:

$$\frac{|\langle v_{11},v_{29}\rangle|^2}{\|v_{11}\|^2\|v_{29}\|^2}=\frac{(-3)^2}{2\cdot 6}=\frac{3}{4}. \tag{S3}$$

Thus no two printed vectors represent the same ray. The maximum and all 72 zero inner products are checked exactly by the accompanying verifier; the family-wise argument in the main text explains why there are no further orthogonalities.

S1.15. Four-dimensional $H^{(4)}_{\mathrm{YO}}$ and H_{65}: vectors and contexts

For direct human and computational verification, this appendix enumerates all unnormalized integer vectors in $\mathbb{Z}^4$ and all four-element contexts for the two four-dimensional examples H_{65} and $H^{(4)}_{\mathrm{YO}}$:

1. The *canonical linear hypergraph* $H_{65}=(V_{65},E_{29})$ (Theorem 22, Appendix S1.15.1), which has exactly 65 **pairwise non-collinear integer vectors** (17 core vectors $v_1,\dots,v_{17}$ and 48 degree-1 completion vectors $v_{18},\dots,v_{65}$), 29 **four-element contexts** (5 core contexts $C_1,\dots,C_5$ and 24 bridge-completion contexts $E_1,\dots,E_{24}$), and 174 **orthogonality edges** ($29\times\binom{4}{2}=174$).

2. The *non-linear warm-up apex suspension* $H^{(4)}_{\mathrm{YO}}=(V_{26},E^{(4)}_{16})$ (Proposition 20, Appendix S1.15.2), which has exactly 26 **pairwise non-collinear integer vectors** (13 Yu–Oh rays $v_1,\dots,v_{13}$, 12 completion rays $v_{14},\dots,v_{25}$ in $\mathbb{R}^3\times\{0\}$, and 1 apex ray v_{26}), 16 **four-element contexts**, and 73 **orthogonality edges** ($16\times\binom{3}{2}=48$ edges in $\mathbb{R}^3\times\{0\}$ plus 25 apex edges).

Canonical primitive normal form and human-verifiable non-collinearity. Every vector $v_i=(x_{i,1},x_{i,2},x_{i,3},x_{i,4})^T\in\mathbb{Z}^4$ listed below in both H_{65} and $H^{(4)}_{\mathrm{YO}}$ is written in *canonical primitive projective form*:

$$\begin{gathered}\gcd\big(|x_{i,1}|,|x_{i,2}|,|x_{i,3}|,|x_{i,4}|\big)=1,\\ \text{first nonzero entry of } v_i>0.\end{gathered} \tag{S4}$$

Because every projective ray $[v]\in\mathbb{P}(\mathbb{Q}^4)$ possesses a *unique* representative in $\mathbb{Z}^4$ satisfying Eq. (S4), two vectors v_i,v_j in canonical primitive form are collinear (proportional over $\mathbb{R}$ or $\mathbb{C}$) *if and only if they are identical* 4*-tuples* ($v_i=v_j$) and, in particular, have the same squared Euclidean norm $\|v_i\|^2=\|v_j\|^2$.

S1.15.1. The canonical linear 65-vertex, 29-context hypergraph H_{65}

Table S17 enumerates all 65 unnormalized integer vectors $v_1, \dots, v_{65} \in \mathbb{Z}^4$ of H_{65}, together with their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0,1,2,3,4\}$ in the strong 5-coloring extending Eq. (S28), and the contexts in E_{29} to which each vertex belongs. Table S18 enumerates all 29 contexts $C_1, \dots, C_5, E_1, \dots, E_{24} \in E_{29}$, displaying side by side both the vertex indices $\{i,j,k,\ell\}$ and the four mutually orthogonal integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

To verify by hand that all 65 vectors in Table S17 are pairwise non-collinear across all $\binom{65}{2} = 2{,}080$ pairs, observe that all 65 tuples satisfy Eq. (S4) and partition by squared norm $\|v_i\|^2$ into seven small, disjoint norm shells ($4+6+7+12+12+12+12=65$):

- $\|v_i\|^2 = 1$ (4 vectors): indices $\{1,2,3,4\}$, the 4 standard coordinate basis vectors;
- $\|v_i\|^2 = 2$ (6 vectors): indices $\{12,\dots,17\}$, one vector for each of the $\binom{4}{2} = 6$ coordinate-pair supports;
- $\|v_i\|^2 = 4$ (7 vectors): indices $\{5,\dots,11\}$, seven distinct ± 1 vectors with $x_{i,1} = +1$;
- $\|v_i\|^2 = 11$ (12 vectors): indices $\{18,20,22,25,26,28,30,33,34,36,39,41\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct sign/entry patterns of $\{0, \pm 1, \pm 1, \pm 3\}$;
- $\|v_i\|^2 = 18$ (12 vectors): indices $\{43,45,46,49,51,53,54,56,59,61,62,65\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 2, \pm 2, \pm 3\}$;
- $\|v_i\|^2 = 22$ (12 vectors): indices $\{19,21,23,24,27,29,31,32,35,37,38,40\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct patterns of $\{0, \pm 2, \pm 3, \pm 3\}$;
- $\|v_i\|^2 = 36$ (12 vectors): indices $\{42,44,47,48,50,52,55,57,58,60,63,64\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 1, \pm 3, \pm 5\}$.

Because no two 4-tuples within any of the seven norm shells coincide, no two of the 65 vectors are collinear; quantitatively, the maximum squared cosine over all $\binom{65}{2} = 2{,}080$ pairs is $\max_{1 \le i < j \le 65} (v_i \cdot v_j)^2 / (\|v_i\|^2 \|v_j\|^2) = 19^2/(11 \times 36) = 361/396 \approx 0.9116 < 1$, attained by (v_{41}, v_{47}), whose squared norms are 11 and 36 and whose inner product is 19.

TABLE S17: Complete enumeration of the 65 unnormalized integer vectors $v_i \in \mathbb{Z}^4$ of H_{65}, their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0,1,2,3,4\}$, and the contexts in E_{29} containing vertex i.

i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i	i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i
1	$(1,0,0,0)$	1	0	C_1, E_3, E_7, E_{11}	34	$(1,-1,0,3)$	11	1	$E_9 = \{3,16,34,35\}$
2	$(0,1,0,0)$	1	1	C_1, E_1, E_5, E_{12}	35	$(3,-3,0,-2)$	22	3	$E_9 = \{3,16,34,35\}$
3	$(0,0,1,0)$	1	2	C_1, E_2, E_8, E_9	36	$(1,-1,3,0)$	11	1	$E_{10} = \{4,16,36,37\}$
4	$(0,0,0,1)$	1	3	C_1, E_4, E_6, E_{10}	37	$(3,-3,-2,0)$	22	2	$E_{10} = \{4,16,36,37\}$
5	$(1,1,1,1)$	4	0	$C_2, E_{13}, E_{19}, E_{23}$	38	$(0,2,3,3)$	22	1	$E_{11} = \{1,17,38,39\}$
6	$(1,1,-1,-1)$	4	1	$C_2, E_{15}, E_{17}, E_{24}$	39	$(0,3,-1,-1)$	11	3	$E_{11} = \{1,17,38,39\}$
7	$(1,-1,1,-1)$	4	2	$C_2, E_{16}, E_{20}, E_{21}$	40	$(2,0,-3,-3)$	22	0	$E_{12} = \{2,17,40,41\}$
8	$(1,-1,-1,1)$	4	4	$C_2, E_{14}, E_{18}, E_{22}$	41	$(3,0,1,1)$	11	3	$E_{12} = \{2,17,40,41\}$
9	$(1,1,-1,1)$	4	0	C_3, C_4	42	$(1,-5,3,1)$	36	1	$E_{13} = \{5,12,42,43\}$
10	$(1,-1,1,1)$	4	1	C_3, C_5	43	$(2,-1,-3,2)$	18	2	$E_{13} = \{5,12,42,43\}$
11	$(1,-1,-1,-1)$	4	3	C_4, C_5	44	$(1,-3,5,1)$	36	0	$E_{14} = \{8,12,44,45\}$
12	$(1,0,0,-1)$	2	3	$C_3, E_1, E_2, E_{13}, E_{14}$	45	$(2,3,1,2)$	18	1	$E_{14} = \{8,12,44,45\}$
13	$(0,1,1,0)$	2	4	$C_3, E_3, E_4, E_{15}, E_{16}$	46	$(1,-2,2,-3)$	18	0	$E_{15} = \{6,13,46,47\}$
14	$(1,0,1,0)$	2	2	$C_4, E_5, E_6, E_{17}, E_{18}$	47	$(5,-1,1,3)$	36	2	$E_{15} = \{6,13,46,47\}$
15	$(0,1,0,-1)$	2	1	$C_4, E_7, E_8, E_{19}, E_{20}$	48	$(3,-1,1,5)$	36	0	$E_{16} = \{7,13,48,49\}$
16	$(1,1,0,0)$	2	0	$C_5, E_9, E_{10}, E_{21}, E_{22}$	49	$(3,2,-2,-1)$	18	1	$E_{16} = \{7,13,48,49\}$
17	$(0,0,1,-1)$	2	2	$C_5, E_{11}, E_{12}, E_{23}, E_{24}$	50	$(1,3,-1,5)$	36	0	$E_{17} = \{6,14,50,51\}$
18	$(1,0,-3,1)$	11	0	$E_1 = \{2,12,18,19\}$	51	$(2,-3,-2,1)$	18	3	$E_{17} = \{6,14,50,51\}$
19	$(3,0,2,3)$	22	2	$E_1 = \{2,12,18,19\}$	52	$(1,5,-1,3)$	36	0	$E_{18} = \{8,14,52,53\}$
20	$(1,3,0,1)$	11	0	$E_2 = \{3,12,20,21\}$	53	$(2,1,-2,-3)$	18	1	$E_{18} = \{8,14,52,53\}$
21	$(3,-2,0,3)$	22	1	$E_2 = \{3,12,20,21\}$	54	$(3,-2,1,-2)$	18	2	$E_{19} = \{5,15,54,55\}$
22	$(0,1,-1,3)$	11	1	$E_3 = \{1,13,22,23\}$	55	$(3,1,-5,1)$	36	3	$E_{19} = \{5,15,54,55\}$
23	$(0,3,-3,-2)$	22	2	$E_3 = \{1,13,22,23\}$	56	$(1,2,3,2)$	18	0	$E_{20} = \{7,15,56,57\}$
24	$(2,-3,3,0)$	22	0	$E_4 = \{4,13,24,25\}$	57	$(5,1,-3,1)$	36	3	$E_{20} = \{7,15,56,57\}$

Continued on the next page.

TABLE S17: All 65 vectors of H_{65} (continued).

i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i	i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i
25	$(3,1,-1,0)$	11	1	$E_4 = \{4,13,24,25\}$	58	$(1,-1,-5,-3)$	36	1	$E_{21} = \{7,16,58,59\}$
26	$(1,0,-1,-3)$	11	0	$E_5 = \{2,14,26,27\}$	59	$(2,-2,-1,3)$	18	3	$E_{21} = \{7,16,58,59\}$
27	$(3,0,-3,2)$	22	3	$E_5 = \{2,14,26,27\}$	60	$(1,-1,-3,-5)$	36	1	$E_{22} = \{8,16,60,61\}$
28	$(1,-3,-1,0)$	11	0	$E_6 = \{4,14,28,29\}$	61	$(2,-2,3,-1)$	18	2	$E_{22} = \{8,16,60,61\}$
29	$(3,2,-3,0)$	22	1	$E_6 = \{4,14,28,29\}$	62	$(1,3,-2,-2)$	18	1	$E_{23} = \{5,17,62,63\}$
30	$(0,1,3,1)$	11	2	$E_7 = \{1,15,30,31\}$	63	$(5,-3,-1,-1)$	36	3	$E_{23} = \{5,17,62,63\}$
31	$(0,3,-2,3)$	22	3	$E_7 = \{1,15,30,31\}$	64	$(3,-5,-1,-1)$	36	0	$E_{24} = \{6,17,64,65\}$
32	$(2,3,0,3)$	22	0	$E_8 = \{3,15,32,33\}$	65	$(3,1,2,2)$	18	3	$E_{24} = \{6,17,64,65\}$
33	$(3,-1,0,-1)$	11	3	$E_8 = \{3,15,32,33\}$					

TABLE S18: Complete enumeration of the 29 four-element contexts of $H_{65} = (V_{65}, E_{29})$, listing for each context both its vertex indices $\{i,j,k,\ell\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

Context	Vertex set	v_i	v_j	v_k	v_ℓ
Five core 4-contexts $C_1, \ldots, C_5$ on $V_{\text{core}} = \{1, \ldots, 17\}$:					
C_1	$\{1,2,3,4\}$	$v_1 = (1,0,0,0)$	$v_2 = (0,1,0,0)$	$v_3 = (0,0,1,0)$	$v_4 = (0,0,0,1)$
C_2	$\{5,6,7,8\}$	$v_5 = (1,1,1,1)$	$v_6 = (1,1,-1,-1)$	$v_7 = (1,-1,1,-1)$	$v_8 = (1,-1,-1,1)$
C_3	$\{9,10,12,13\}$	$v_9 = (1,1,-1,1)$	$v_{10} = (1,-1,1,1)$	$v_{12} = (1,0,0,-1)$	$v_{13} = (0,1,1,0)$
C_4	$\{9,11,14,15\}$	$v_9 = (1,1,-1,1)$	$v_{11} = (1,-1,-1,-1)$	$v_{14} = (1,0,1,0)$	$v_{15} = (0,1,0,-1)$
C_5	$\{10,11,16,17\}$	$v_{10} = (1,-1,1,1)$	$v_{11} = (1,-1,-1,-1)$	$v_{16} = (1,1,0,0)$	$v_{17} = (0,0,1,-1)$
Twelve coordinate-to-bridge completion contexts $E_1, \ldots, E_{12}$ (between C_1 and $B_1 \cup B_2 \cup B_3$):					
E_1	$\{2,12,18,19\}$	$v_2 = (0,1,0,0)$	$v_{12} = (1,0,0,-1)$	$v_{18} = (1,0,-3,1)$	$v_{19} = (3,0,2,3)$
E_2	$\{3,12,20,21\}$	$v_3 = (0,0,1,0)$	$v_{12} = (1,0,0,-1)$	$v_{20} = (1,3,0,1)$	$v_{21} = (3,-2,0,3)$
E_3	$\{1,13,22,23\}$	$v_1 = (1,0,0,0)$	$v_{13} = (0,1,1,0)$	$v_{22} = (0,1,-1,3)$	$v_{23} = (0,3,-3,-2)$
E_4	$\{4,13,24,25\}$	$v_4 = (0,0,0,1)$	$v_{13} = (0,1,1,0)$	$v_{24} = (2,-3,3,0)$	$v_{25} = (3,1,-1,0)$
E_5	$\{2,14,26,27\}$	$v_2 = (0,1,0,0)$	$v_{14} = (1,0,1,0)$	$v_{26} = (1,0,-1,-3)$	$v_{27} = (3,0,-3,2)$
E_6	$\{4,14,28,29\}$	$v_4 = (0,0,0,1)$	$v_{14} = (1,0,1,0)$	$v_{28} = (1,-3,-1,0)$	$v_{29} = (3,2,-3,0)$
E_7	$\{1,15,30,31\}$	$v_1 = (1,0,0,0)$	$v_{15} = (0,1,0,-1)$	$v_{30} = (0,1,3,1)$	$v_{31} = (0,3,-2,3)$
E_8	$\{3,15,32,33\}$	$v_3 = (0,0,1,0)$	$v_{15} = (0,1,0,-1)$	$v_{32} = (2,3,0,3)$	$v_{33} = (3,-1,0,-1)$
E_9	$\{3,16,34,35\}$	$v_3 = (0,0,1,0)$	$v_{16} = (1,1,0,0)$	$v_{34} = (1,-1,0,3)$	$v_{35} = (3,-3,0,-2)$
E_{10}	$\{4,16,36,37\}$	$v_4 = (0,0,0,1)$	$v_{16} = (1,1,0,0)$	$v_{36} = (1,-1,3,0)$	$v_{37} = (3,-3,-2,0)$
E_{11}	$\{1,17,38,39\}$	$v_1 = (1,0,0,0)$	$v_{17} = (0,0,1,-1)$	$v_{38} = (0,2,3,3)$	$v_{39} = (0,3,-1,-1)$
E_{12}	$\{2,17,40,41\}$	$v_2 = (0,1,0,0)$	$v_{17} = (0,0,1,-1)$	$v_{40} = (2,0,-3,-3)$	$v_{41} = (3,0,1,1)$
Twelve Hadamard-to-bridge completion contexts $E_{13}, \ldots, E_{24}$ (between C_2 and $B_1 \cup B_2 \cup B_3$):					
E_{13}	$\{5,12,42,43\}$	$v_5 = (1,1,1,1)$	$v_{12} = (1,0,0,-1)$	$v_{42} = (1,-5,3,1)$	$v_{43} = (2,-1,-3,2)$
E_{14}	$\{8,12,44,45\}$	$v_8 = (1,-1,-1,1)$	$v_{12} = (1,0,0,-1)$	$v_{44} = (1,-3,5,1)$	$v_{45} = (2,3,1,2)$
E_{15}	$\{6,13,46,47\}$	$v_6 = (1,1,-1,-1)$	$v_{13} = (0,1,1,0)$	$v_{46} = (1,-2,2,-3)$	$v_{47} = (5,-1,1,3)$
E_{16}	$\{7,13,48,49\}$	$v_7 = (1,-1,1,-1)$	$v_{13} = (0,1,1,0)$	$v_{48} = (3,-1,1,5)$	$v_{49} = (3,2,-2,-1)$
E_{17}	$\{6,14,50,51\}$	$v_6 = (1,1,-1,-1)$	$v_{14} = (1,0,1,0)$	$v_{50} = (1,3,-1,5)$	$v_{51} = (2,-3,-2,1)$
E_{18}	$\{8,14,52,53\}$	$v_8 = (1,-1,-1,1)$	$v_{14} = (1,0,1,0)$	$v_{52} = (1,5,-1,3)$	$v_{53} = (2,1,-2,-3)$
E_{19}	$\{5,15,54,55\}$	$v_5 = (1,1,1,1)$	$v_{15} = (0,1,0,-1)$	$v_{54} = (3,-2,1,-2)$	$v_{55} = (3,1,-5,1)$
E_{20}	$\{7,15,56,57\}$	$v_7 = (1,-1,1,-1)$	$v_{15} = (0,1,0,-1)$	$v_{56} = (1,2,3,2)$	$v_{57} = (5,1,-3,1)$
E_{21}	$\{7,16,58,59\}$	$v_7 = (1,-1,1,-1)$	$v_{16} = (1,1,0,0)$	$v_{58} = (1,-1,-5,-3)$	$v_{59} = (2,-2,-1,3)$
E_{22}	$\{8,16,60,61\}$	$v_8 = (1,-1,-1,1)$	$v_{16} = (1,1,0,0)$	$v_{60} = (1,-1,-3,-5)$	$v_{61} = (2,-2,3,-1)$
E_{23}	$\{5,17,62,63\}$	$v_5 = (1,1,1,1)$	$v_{17} = (0,0,1,-1)$	$v_{62} = (1,3,-2,-2)$	$v_{63} = (5,-3,-1,-1)$
E_{24}	$\{6,17,64,65\}$	$v_6 = (1,1,-1,-1)$	$v_{17} = (0,0,1,-1)$	$v_{64} = (3,-5,-1,-1)$	$v_{65} = (3,1,2,2)$

S1.15.2. The non-linear warm-up 26-vertex, 16-context apex suspension $H_{\mathrm{YO}}^{(4)}$

For completeness, the 26 unnormalized integer vectors $v_1, \dots, v_{26} \in \mathbb{Z}^4$ of the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$ in Proposition 20 (written in canonical primitive form satisfying Eq. (S4)) are:

$$\begin{aligned}
v_1 &= (0,1,1,0), & v_2 &= (1,0,0,0),\\
v_3 &= (0,1,-1,0), & v_4 &= (1,1,0,0),\\
v_5 &= (0,0,1,0), & v_6 &= (1,-1,0,0),\\
v_7 &= (1,0,1,0), & v_8 &= (0,1,0,0),\\
v_9 &= (1,0,-1,0), & v_{10} &= (1,-1,1,0),\\
v_{11} &= (1,-1,-1,0), & v_{12} &= (1,1,1,0),\\
v_{13} &= (1,1,-1,0), & v_{14} &= (2,1,1,0),\\
v_{15} &= (1,2,-1,0), & v_{16} &= (1,-2,1,0),\\
v_{17} &= (1,1,-2,0), & v_{18} &= (1,-1,-2,0),\\
v_{19} &= (2,1,-1,0), & v_{20} &= (2,-1,1,0),\\
v_{21} &= (1,-2,-1,0), & v_{22} &= (1,1,2,0),\\
v_{23} &= (1,2,1,0), & v_{24} &= (1,-1,2,0),\\
v_{25} &= (2,-1,-1,0), & v_{26} &= (0,0,0,1).
\end{aligned} \tag{S5}$$

These 26 canonical primitive vectors partition by squared norm $\|v_i\|^2$ into four disjoint norm shells ($4+6+4+12=26$):

- $\|v_i\|^2 = 1$ (4 vectors): $\{v_2, v_5, v_8, v_{26}\}$, the four coordinate basis vectors of $\mathbb{R}^4$;
- $\|v_i\|^2 = 2$ (6 vectors): $\{v_1, v_3, v_4, v_6, v_7, v_9\}$, the six binary rays in $\mathbb{R}^3 \times \{0\}$;
- $\|v_i\|^2 = 3$ (4 vectors): $\{v_{10}, v_{11}, v_{12}, v_{13}\}$, the four triad rays $(1, \pm 1, \pm 1, 0)^T$;
- $\|v_i\|^2 = 6$ (12 vectors): $\{v_{14}, \dots, v_{25}\}$, all $3 \times 4 = 12$ canonical primitive permutations and sign patterns of $(\pm 2, \pm 1, \pm 1, 0)^T$.

Since all 26 canonical 4-tuples are distinct, all $\binom{26}{2} = 325$ pairs of vectors are strictly non-collinear, with maximum squared cosine $\max_{1 \le i < j \le 26} (v_i \cdot v_j)^2 / (\|v_i\|^2 \|v_j\|^2) = 4^2/(3 \times 6) = 8/9 \approx 0.8889 < 1$. Table S19 lists all 16 four-element contexts of $E_{16}^{(4)}$ side by side with their four mutually orthogonal integer vectors.

TABLE S19: Complete enumeration of the 16 four-element contexts of the non-linear warm-up apex suspension $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, listing for each context both its vertex indices $\{i, j, k, 26\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_{26}\} \subset \mathbb{Z}^4$.

Context	Vertex set	v_i	v_j	v_k	v_{26}
$e_1^{(4)}$	$\{1,2,3,26\}$	$v_1 = (0,1,1,0)$	$v_2 = (1,0,0,0)$	$v_3 = (0,1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_2^{(4)}$	$\{7,8,9,26\}$	$v_7 = (1,0,1,0)$	$v_8 = (0,1,0,0)$	$v_9 = (1,0,-1,0)$	$v_{26} = (0,0,0,1)$
$e_3^{(4)}$	$\{4,5,6,26\}$	$v_4 = (1,1,0,0)$	$v_5 = (0,0,1,0)$	$v_6 = (1,-1,0,0)$	$v_{26} = (0,0,0,1)$
$e_4^{(4)}$	$\{2,5,8,26\}$	$v_2 = (1,0,0,0)$	$v_5 = (0,0,1,0)$	$v_8 = (0,1,0,0)$	$v_{26} = (0,0,0,1)$
$e_5^{(4)}$	$\{3,12,25,26\}$	$v_3 = (0,1,-1,0)$	$v_{12} = (1,1,1,0)$	$v_{25} = (2,-1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_6^{(4)}$	$\{9,12,16,26\}$	$v_9 = (1,0,-1,0)$	$v_{12} = (1,1,1,0)$	$v_{16} = (1,-2,1,0)$	$v_{26} = (0,0,0,1)$
$e_7^{(4)}$	$\{6,12,17,26\}$	$v_6 = (1,-1,0,0)$	$v_{12} = (1,1,1,0)$	$v_{17} = (1,1,-2,0)$	$v_{26} = (0,0,0,1)$
$e_8^{(4)}$	$\{3,11,14,26\}$	$v_3 = (0,1,-1,0)$	$v_{11} = (1,-1,-1,0)$	$v_{14} = (2,1,1,0)$	$v_{26} = (0,0,0,1)$
$e_9^{(4)}$	$\{7,11,15,26\}$	$v_7 = (1,0,1,0)$	$v_{11} = (1,-1,-1,0)$	$v_{15} = (1,2,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{10}^{(4)}$	$\{4,11,24,26\}$	$v_4 = (1,1,0,0)$	$v_{11} = (1,-1,-1,0)$	$v_{24} = (1,-1,2,0)$	$v_{26} = (0,0,0,1)$
$e_{11}^{(4)}$	$\{9,10,23,26\}$	$v_9 = (1,0,-1,0)$	$v_{10} = (1,-1,1,0)$	$v_{23} = (1,2,1,0)$	$v_{26} = (0,0,0,1)$
$e_{12}^{(4)}$	$\{1,10,19,26\}$	$v_1 = (0,1,1,0)$	$v_{10} = (1,-1,1,0)$	$v_{19} = (2,1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{13}^{(4)}$	$\{4,10,18,26\}$	$v_4 = (1,1,0,0)$	$v_{10} = (1,-1,1,0)$	$v_{18} = (1,-1,-2,0)$	$v_{26} = (0,0,0,1)$
$e_{14}^{(4)}$	$\{1,13,20,26\}$	$v_1 = (0,1,1,0)$	$v_{13} = (1,1,-1,0)$	$v_{20} = (2,-1,1,0)$	$v_{26} = (0,0,0,1)$
$e_{15}^{(4)}$	$\{7,13,21,26\}$	$v_7 = (1,0,1,0)$	$v_{13} = (1,1,-1,0)$	$v_{21} = (1,-2,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{16}^{(4)}$	$\{6,13,22,26\}$	$v_6 = (1,-1,0,0)$	$v_{13} = (1,1,-1,0)$	$v_{22} = (1,1,2,0)$	$v_{26} = (0,0,0,1)$

S1.16. Explicit parent configurations and inclusion maps

S1.16.1. The 49-ray real Yu–Oh parent and published Bub labels

Table S20 lists all rays, including the original 25. The final column gives the exact labels in the published full 49-ray encoding [1, p. 14, Sec. 5(xi)] of Bub's set [2]. Under this map the 36 bases in Table S21 equal all 36 published contexts; their 108 pairs equal all geometric orthogonalities. In this labeling 26–33 are Eq. (S31), and 34–49 complete the sixteen remaining orthogonal pairs. The first 25 vectors are those of Table S1.

TABLE S20: All unnormalized vectors of the real 49-ray parent, with the verified published Bub labels.

i	v_i	Published label
1	$(0,1,1)$	G
2	$(1,0,0)$	1
3	$(0,1,-1)$	P
4	$(1,1,0)$	H
5	$(0,0,1)$	3
6	$(1,-1,0)$	U
7	$(1,0,1)$	Q
8	$(0,1,0)$	2
9	$(1,0,-1)$	e
10	$(1,-1,1)$	D
11	$(1,-1,-1)$	h
12	$(1,1,1)$	S
13	$(1,1,-1)$	a
14	$(2,1,1)$	g
15	$(1,2,-1)$	n
16	$(1,-2,1)$	d
17	$(1,1,-2)$	T
18	$(1,-1,-2)$	F
19	$(2,1,-1)$	E
20	$(2,-1,1)$	I
21	$(1,-2,-1)$	k
22	$(1,1,2)$	b
23	$(1,2,1)$	A
24	$(1,-1,2)$	N
25	$(2,-1,-1)$	R
26	$(0,1,2)$	i
27	$(0,2,-1)$	O
28	$(0,1,-2)$	6
29	$(0,2,1)$	7
30	$(1,2,0)$	5
31	$(2,-1,0)$	4
32	$(1,-2,0)$	V
33	$(2,1,0)$	j
34	$(2,1,-5)$	Z
35	$(5,-2,1)$	m
36	$(1,2,5)$	W
37	$(5,2,-1)$	c
38	$(1,-2,-5)$	f
39	$(5,-1,2)$	L
40	$(5,1,2)$	9
41	$(2,1,5)$	8
42	$(2,-1,-5)$	J
43	$(5,2,1)$	X
44	$(1,-2,5)$	l
45	$(5,-1,-2)$	Y
46	$(5,-2,-1)$	B
47	$(1,2,-5)$	C
48	$(5,1,-2)$	M
49	$(2,-1,5)$	K

TABLE S21: All 36 complete bases of the real parent.

e	Context	e	Context	e	Context
1	$\{1,2,3\}$	2	$\{1,10,19\}$	3	$\{1,13,20\}$
4	$\{2,5,8\}$	5	$\{2,26,27\}$	6	$\{2,28,29\}$
7	$\{3,11,14\}$	8	$\{3,12,25\}$	9	$\{4,5,6\}$
10	$\{4,10,18\}$	11	$\{4,11,24\}$	12	$\{5,30,31\}$
13	$\{5,32,33\}$	14	$\{6,12,17\}$	15	$\{6,13,22\}$
16	$\{7,8,9\}$	17	$\{7,11,15\}$	18	$\{7,13,21\}$
19	$\{9,10,23\}$	20	$\{9,12,16\}$	21	$\{14,32,34\}$
22	$\{15,26,35\}$	23	$\{15,31,36\}$	24	$\{16,26,37\}$
25	$\{16,33,38\}$	26	$\{17,29,39\}$	27	$\{18,27,40\}$
28	$\{19,32,41\}$	29	$\{20,30,42\}$	30	$\{21,28,43\}$
31	$\{21,33,44\}$	32	$\{22,27,45\}$	33	$\{23,28,46\}$
34	$\{23,31,47\}$	35	$\{24,29,48\}$	36	$\{25,30,49\}$

S1.16.2. The 87-ray Hesse parent

Use $\zeta = \exp(2\pi i/3)$ and $\zeta^2+\zeta+1 = 0$. The first 37 rays retain exactly the Hesse numbering; projective representatives have their first nonzero coordinate one. The inventory labels use the generation convention of Sec. VII B. Tables S22 and S23 list all 87 rays and all 64 complete bases. The six additional orthogonal pairs outside the 64 complete bases are listed separately: this larger parent's full orthogonality graph need not equal its context two-section. The retained Hesse restriction has all its original orthogonalities covered by contexts.

TABLE S22: All 87 unnormalized rays of the first Hesse/lineage union.

i	v_i	Inventory label
1	$(1,0,0)$	a_{11}
2	$(1,1,1)$	u_1
3	$(1,\zeta,-1-\zeta)$	u_2
4	$(1,-1-\zeta,\zeta)$	u_3
5	$(1,\zeta,\zeta)$	u_4
6	$(1,-1-\zeta,1)$	u_5
7	$(1,1,-1-\zeta)$	u_6
8	$(1,-1-\zeta,-1-\zeta)$	u_7
9	$(1,1,\zeta)$	u_9
10	$(1,\zeta,1)$	u_8
11	$(0,1,-1)$	c_{11}
12	$(0,1,-\zeta)$	c_{12}
13	$(0,1,1+\zeta)$	c_{13}
14	$(1,0,-1)$	c_{21}
15	$(1,0,1+\zeta)$	c_{22}
16	$(1,0,-\zeta)$	c_{23}
17	$(0,1,1)$	b_{11}
18	$(0,1,\zeta)$	b_{12}
19	$(0,1,-1-\zeta)$	b_{13}
20	$(1,\frac{-1}{2},\frac{-1}{2})$	d_{11}
21	$(1,-\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$	d_{12}
22	$(1,\frac{1}{2}+\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$	d_{13}
23	$(1,-\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$	d_{14}
24	$(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{-1}{2})$	d_{15}
25	$(1,\frac{-1}{2},\frac{1}{2}+\frac{1}{2}\zeta)$	d_{16}
26	$(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$	d_{17}
27	$(1,\frac{-1}{2},-\frac{1}{2}\zeta)$	d_{19}
28	$(1,-\frac{1}{2}\zeta,\frac{-1}{2})$	d_{18}
29	$(1,-2,1)$	d_{21}
30	$(1,-2\zeta,-1-\zeta)$	d_{22}
31	$(1,2+2\zeta,\zeta)$	d_{23}
32	$(1,2+2\zeta,1)$	d_{25}
33	$(1,-2,-1-\zeta)$	d_{26}
34	$(1,-2\zeta,\zeta)$	d_{24}
35	$(1,-2\zeta,1)$	d_{28}
36	$(1,2+2\zeta,-1-\zeta)$	d_{27}
37	$(1,-2,\zeta)$	d_{29}
38	$(0,1,0)$	a_{21}
39	$(0,0,1)$	a_{31}
40	$(1,\frac{-1}{2},\frac{1}{2})$	b_{1121}

i	v_i	Inventory label
41	$(1, -\frac{1}{2}\zeta, \frac{-1}{2} - \frac{1}{2}\zeta)$	b_{1122}
42	$(1, \frac{1}{2} + \frac{1}{2}\zeta, \frac{1}{2}\zeta)$	b_{1123}
43	$(1, \frac{1}{2}, \frac{-1}{2})$	b_{1131}
44	$(1, \frac{1}{2}\zeta, \frac{1}{2} + \frac{1}{2}\zeta)$	b_{1132}
45	$(1, \frac{-1}{2} - \frac{1}{2}\zeta, -\frac{1}{2}\zeta)$	b_{1133}
46	$(1, 1, -1)$	b_{121}
47	$(1, \zeta, 1 + \zeta)$	b_{122}
48	$(1, -1 - \zeta, -\zeta)$	b_{123}
49	$(1, -1, 1)$	b_{131}
50	$(1, -\zeta, -1 - \zeta)$	b_{132}
51	$(1, 1 + \zeta, \zeta)$	b_{133}
52	$(1, 0, 1)$	b_{21}
53	$(1, -2, -1)$	b_{2121}
54	$(1, -2\zeta, 1 + \zeta)$	b_{2122}
55	$(1, 2 + 2\zeta, -\zeta)$	b_{2123}
56	$(1, 0, -1 - \zeta)$	b_{22}
57	$(1, 2, -1)$	b_{2231}
58	$(1, 2\zeta, 1 + \zeta)$	b_{2232}
59	$(1, -2 - 2\zeta, -\zeta)$	b_{2233}
60	$(1, 0, \zeta)$	b_{23}
61	$(1, -1, -1)$	b_{231}
62	$(1, -\zeta, 1 + \zeta)$	b_{232}
63	$(1, 1 + \zeta, -\zeta)$	b_{233}
64	$(1, 1, 0)$	b_{31}
65	$(1, -1, -2)$	b_{3131}
66	$(1, -\zeta, 2 + 2\zeta)$	b_{3132}
67	$(1, 1 + \zeta, -2\zeta)$	b_{3133}
68	$(1, \zeta, 0)$	b_{32}
69	$(1, -1, 2)$	b_{3231}
70	$(1, -\zeta, -2 - 2\zeta)$	b_{3232}
71	$(1, 1 + \zeta, 2\zeta)$	b_{3233}
72	$(1, -1 - \zeta, 0)$	b_{33}
73	$(1, -1, 0)$	c_{31}
74	$(1, -\zeta, 0)$	c_{32}
75	$(1, 1 + \zeta, 0)$	c_{33}
76	$(1, 1, -2)$	d_{31}
77	$(1, \zeta, 2 + 2\zeta)$	d_{32}
78	$(1, -1 - \zeta, -2\zeta)$	d_{33}
79	$(1, 2, 1)$	e_{11}
80	$(1, 2\zeta, -1 - \zeta)$	e_{12}
81	$(1, -2 - 2\zeta, \zeta)$	e_{13}
82	$(1, 1, 2)$	e_{21}
83	$(1, \zeta, -2 - 2\zeta)$	e_{22}
84	$(1, -1 - \zeta, 2\zeta)$	e_{23}
85	$(1, \frac{1}{2}, \frac{1}{2})$	e_{31}
86	$(1, \frac{1}{2}\zeta, \frac{-1}{2} - \frac{1}{2}\zeta)$	e_{32}
87	$(1, \frac{-1}{2} - \frac{1}{2}\zeta, \frac{1}{2}\zeta)$	e_{33}

TABLE S23: All 64 complete bases of the first 87-ray Hesse parent.

e	Context	e	Context	e	Context
1	$\{1,11,17\}$	2	$\{1,12,18\}$	3	$\{1,13,19\}$
4	$\{1,38,39\}$	5	$\{2,3,4\}$	6	$\{2,11,20\}$
7	$\{2,14,29\}$	8	$\{2,73,76\}$	9	$\{3,12,21\}$
10	$\{3,15,30\}$	11	$\{3,74,77\}$	12	$\{4,13,22\}$
13	$\{4,16,31\}$	14	$\{4,75,78\}$	15	$\{5,6,7\}$
16	$\{5,11,23\}$	17	$\{5,16,34\}$	18	$\{6,12,24\}$
19	$\{6,14,32\}$	20	$\{7,13,25\}$	21	$\{7,15,33\}$
22	$\{8,9,10\}$	23	$\{8,11,26\}$	24	$\{8,15,36\}$
25	$\{9,12,27\}$	26	$\{9,16,37\}$	27	$\{10,13,28\}$
28	$\{10,14,35\}$	29	$\{11,61,85\}$	30	$\{12,62,86\}$
31	$\{13,63,87\}$	32	$\{14,38,52\}$	33	$\{14,49,79\}$
34	$\{15,38,56\}$	35	$\{15,50,80\}$	36	$\{16,38,60\}$
37	$\{16,51,81\}$	38	$\{17,40,46\}$	39	$\{17,43,49\}$
40	$\{18,41,47\}$	41	$\{18,44,50\}$	42	$\{19,42,48\}$
43	$\{19,45,51\}$	44	$\{39,64,73\}$	45	$\{39,68,74\}$
46	$\{39,72,75\}$	47	$\{46,47,48\}$	48	$\{46,52,53\}$
49	$\{46,73,82\}$	50	$\{47,54,56\}$	51	$\{47,74,83\}$
52	$\{48,55,60\}$	53	$\{48,75,84\}$	54	$\{49,50,51\}$
55	$\{49,64,65\}$	56	$\{50,66,68\}$	57	$\{51,67,72\}$
58	$\{52,57,61\}$	59	$\{56,58,62\}$	60	$\{59,60,63\}$
61	$\{61,62,63\}$	62	$\{61,64,69\}$	63	$\{62,68,70\}$
64	$\{63,71,72\}$				

Its six further orthogonal pairs are $\{5,74\},\ \{6,75\},\ \{7,73\},\ \{8,75\},\ \{9,73\},\ \{10,74\}$.

S1.16.3. Peres24 and the complete four-dimensional extensions

The coordinates of H_{65} and $H_{\mathrm{YO}}^{(4)}$ are those already listed above. Table S24 explicitly enumerates the known Peres rays. In each union, retain the original local labels and append the missing rays in the order displayed in Table S25. Tables S26 and S27 list every available complete basis. Full orthogonality-pair lists accompany the coordinates in the data; the 40-ray union has additional pairs outside its complete contexts.

TABLE S24: All 24 primitive unnormalized Peres rays, in the certificate order.

i	p_i
1	$(1,0,0,0)$
2	$(0,1,0,0)$
3	$(0,0,1,0)$
4	$(0,0,0,1)$
5	$(1,-1,0,0)$
6	$(1,1,0,0)$
7	$(1,0,-1,0)$
8	$(1,0,1,0)$
9	$(1,0,0,-1)$
10	$(1,0,0,1)$
11	$(0,1,-1,0)$
12	$(0,1,1,0)$
13	$(0,1,0,-1)$
14	$(0,1,0,1)$
15	$(0,0,1,-1)$
16	$(0,0,1,1)$
17	$(1,-1,-1,-1)$
18	$(1,-1,-1,1)$
19	$(1,-1,1,-1)$
20	$(1,-1,1,1)$
21	$(1,1,-1,-1)$
22	$(1,1,-1,1)$
23	$(1,1,1,-1)$
24	$(1,1,1,1)$

TABLE S25: Every ray added to the original four-dimensional examples to obtain the stated parents.

Original	New label	Added vector
$H^{(4)}_{\mathrm{YO}}$	27	$(1,0,0,-1)$
$H^{(4)}_{\mathrm{YO}}$	28	$(1,0,0,1)$
$H^{(4)}_{\mathrm{YO}}$	29	$(0,1,0,-1)$
$H^{(4)}_{\mathrm{YO}}$	30	$(0,1,0,1)$
$H^{(4)}_{\mathrm{YO}}$	31	$(0,0,1,-1)$
$H^{(4)}_{\mathrm{YO}}$	32	$(0,0,1,1)$
$H^{(4)}_{\mathrm{YO}}$	33	$(1,-1,-1,-1)$
$H^{(4)}_{\mathrm{YO}}$	34	$(1,-1,-1,1)$
$H^{(4)}_{\mathrm{YO}}$	35	$(1,-1,1,-1)$
$H^{(4)}_{\mathrm{YO}}$	36	$(1,-1,1,1)$
$H^{(4)}_{\mathrm{YO}}$	37	$(1,1,-1,-1)$
$H^{(4)}_{\mathrm{YO}}$	38	$(1,1,-1,1)$
$H^{(4)}_{\mathrm{YO}}$	39	$(1,1,1,-1)$
$H^{(4)}_{\mathrm{YO}}$	40	$(1,1,1,1)$
H_{65}	66	$(1,-1,0,0)$
H_{65}	67	$(1,0,-1,0)$
H_{65}	68	$(1,0,0,1)$
H_{65}	69	$(0,1,-1,0)$
H_{65}	70	$(0,1,0,1)$
H_{65}	71	$(0,0,1,1)$
H_{65}	72	$(1,1,1,-1)$

TABLE S26: All 36 complete bases of the 40-ray parent.

e	Context	e	Context
1	$\{1,2,3,26\}$	2	$\{1,3,27,28\}$
3	$\{1,10,19,26\}$	4	$\{1,13,20,26\}$
5	$\{1,27,36,38\}$	6	$\{1,28,35,37\}$
7	$\{2,5,8,26\}$	8	$\{2,5,29,30\}$
9	$\{2,8,31,32\}$	10	$\{3,11,14,26\}$
11	$\{3,12,25,26\}$	12	$\{3,27,34,40\}$
13	$\{3,28,33,39\}$	14	$\{4,5,6,26\}$
15	$\{4,6,31,32\}$	16	$\{4,10,18,26\}$
17	$\{4,11,24,26\}$	18	$\{4,31,33,36\}$
19	$\{4,32,34,35\}$	20	$\{5,8,27,28\}$
21	$\{6,12,17,26\}$	22	$\{6,13,22,26\}$
23	$\{6,31,37,40\}$	24	$\{6,32,38,39\}$
25	$\{7,8,9,26\}$	26	$\{7,9,29,30\}$
27	$\{7,11,15,26\}$	28	$\{7,13,21,26\}$
29	$\{7,29,33,38\}$	30	$\{7,30,34,37\}$
31	$\{9,10,23,26\}$	32	$\{9,12,16,26\}$
33	$\{9,29,35,40\}$	34	$\{9,30,36,39\}$
35	$\{33,36,38,39\}$	36	$\{34,35,37,40\}$

TABLE S27: All 48 complete bases of the 72-ray parent.

e	Context	e	Context
1	$\{1,2,3,4\}$	2	$\{1,2,17,71\}$
3	$\{1,3,15,70\}$	4	$\{1,4,13,69\}$
5	$\{1,13,22,23\}$	6	$\{1,15,30,31\}$
7	$\{1,17,38,39\}$	8	$\{2,3,12,68\}$
9	$\{2,4,14,67\}$	10	$\{2,12,18,19\}$
11	$\{2,14,26,27\}$	12	$\{2,17,40,41\}$
13	$\{3,4,16,66\}$	14	$\{3,12,20,21\}$
15	$\{3,15,32,33\}$	16	$\{3,16,34,35\}$
17	$\{4,13,24,25\}$	18	$\{4,14,28,29\}$
19	$\{4,16,36,37\}$	20	$\{5,6,7,8\}$
21	$\{5,6,17,66\}$	22	$\{5,7,15,67\}$
23	$\{5,8,12,69\}$	24	$\{5,12,42,43\}$
25	$\{5,15,54,55\}$	26	$\{5,17,62,63\}$
27	$\{6,7,13,68\}$	28	$\{6,8,14,70\}$
29	$\{6,13,46,47\}$	30	$\{6,14,50,51\}$
31	$\{6,17,64,65\}$	32	$\{7,8,16,71\}$
33	$\{7,13,48,49\}$	34	$\{7,15,56,57\}$
35	$\{7,16,58,59\}$	36	$\{8,12,44,45\}$
37	$\{8,14,52,53\}$	38	$\{8,16,60,61\}$
39	$\{9,10,11,72\}$	40	$\{9,10,12,13\}$
41	$\{9,11,14,15\}$	42	$\{9,66,71,72\}$
43	$\{10,11,16,17\}$	44	$\{10,67,70,72\}$
45	$\{11,68,69,72\}$	46	$\{12,13,68,69\}$
47	$\{14,15,67,70\}$	48	$\{16,17,66,71\}$

S1.17. Cabello completions and the two 68-ray parents

S1.17.1. The eighteen-ray input and faithful 36-ray completion

All labels in this subsection are local. In $C_{18,9}$ and $C_{36,18}$, $18 = a$, $19 = b$, and $20 = c$; labels 4 and 5 are absent. They do not denote the completion rays of H_{65}. The core is the first eighteen rows of Table S28; labels 21–38 are fresh. The contexts of $C_{18,9}$ are the first nine rows of Table S29.

TABLE S28: All 36 unnormalized vectors of the faithful Cabello completion.

i	v_i
1	$(1,0,0,0)$
2	$(0,1,0,0)$
3	$(0,0,1,0)$
6	$(1,1,-1,-1)$
7	$(1,-1,1,-1)$
8	$(1,-1,-1,1)$
9	$(1,1,-1,1)$
10	$(1,-1,1,1)$
11	$(1,-1,-1,-1)$
12	$(1,0,0,-1)$
13	$(0,1,1,0)$
14	$(1,0,1,0)$
15	$(0,1,0,-1)$
16	$(1,1,0,0)$
17	$(0,0,1,-1)$
18	$(1,0,0,1)$
19	$(0,1,0,1)$
20	$(0,0,1,1)$
21	$(0,1,-1,3)$
22	$(0,3,-3,-2)$
23	$(1,0,-1,3)$
24	$(3,0,-3,-2)$
25	$(1,-1,0,3)$
26	$(3,-3,0,-2)$
27	$(3,1,2,2)$
28	$(3,-5,-1,-1)$
29	$(3,2,1,2)$
30	$(3,-1,-5,-1)$

i	v_i
31	$(2,3,1,2)$
32	$(1,-3,5,1)$
33	$(4,2,3,-3)$
34	$(5,-7,-1,1)$
35	$(4,3,2,-3)$
36	$(5,-1,-7,1)$
37	$(3,4,2,-3)$
38	$(1,-5,7,-1)$

TABLE S29: All eighteen full bases of the faithful Cabello completion.

e	Context
1	$\{9,10,12,13\}$
2	$\{9,11,14,15\}$
3	$\{10,11,16,17\}$
4	$\{2,3,12,18\}$
5	$\{1,3,15,19\}$
6	$\{1,2,17,20\}$
7	$\{6,7,13,18\}$
8	$\{6,8,14,19\}$
9	$\{7,8,16,20\}$
10	$\{1,13,21,22\}$
11	$\{2,14,23,24\}$
12	$\{3,16,25,26\}$
13	$\{6,17,27,28\}$
14	$\{7,15,29,30\}$
15	$\{8,12,31,32\}$
16	$\{9,20,33,34\}$
17	$\{10,19,35,36\}$
18	$\{11,18,37,38\}$

The orthogonal matrix $Q = M/\sqrt{2}$, with

$$M = \begin{pmatrix} 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \qquad M^T M = 2I_4, \tag{S6}$$

maps the published eighteen-ray set and all nine bases of Ref. [3, Eqs. (1)–(9)] onto this realization. Table S30 gives the complete ray map; the input representatives are unnormalized and signs are immaterial.

TABLE S30: Exact ray map from the originally published Cabello coordinates.

Published vector u	Local label of $[Qu]$
$(0,0,0,1)$	12
$(0,0,1,0)$	13
$(0,0,1,1)$	11
$(0,1,-1,0)$	8
$(0,1,0,-1)$	1
$(0,1,0,0)$	18
$(1,-1,-1,-1)$	19
$(1,-1,-1,1)$	16
$(1,-1,0,0)$	9
$(1,-1,1,-1)$	17
$(1,0,-1,0)$	2
$(1,0,0,-1)$	7
$(1,0,0,1)$	6
$(1,0,1,0)$	3
$(1,1,-1,1)$	15
$(1,1,0,0)$	10
$(1,1,1,-1)$	14
$(1,1,1,1)$	20

S1.17.2. An injective 45-ray completion after deleting c

Delete local ray 20 and contexts 6 and 9 from the first nine contexts above. The seven remaining bases are followed by fourteen fresh-pair contexts in Table S32. The next table prints all 45 vectors; 4, 5, and 20 are absent and 21–48 are fresh. Its only uncovered orthogonal pairs are $\{16,21\},\{16,22\},\{17,35\},\{17,36\}$.

TABLE S31: All 45 unnormalized vectors of the vertex-deleted completion.

i	v_i
1	$(1,0,0,0)$
2	$(0,1,0,0)$
3	$(0,0,1,0)$
6	$(1,1,-1,-1)$
7	$(1,-1,1,-1)$
8	$(1,-1,-1,1)$
9	$(1,1,-1,1)$
10	$(1,-1,1,1)$
11	$(1,-1,-1,-1)$
12	$(1,0,0,-1)$
13	$(0,1,1,0)$
14	$(1,0,1,0)$
15	$(0,1,0,-1)$
16	$(1,1,0,0)$
17	$(0,0,1,-1)$
18	$(1,0,0,1)$
19	$(0,1,0,1)$
21	$(0,0,1,2)$
22	$(0,0,2,-1)$
23	$(0,1,-1,3)$
24	$(0,3,-3,-2)$
25	$(0,1,2,2)$
26	$(0,4,-1,-1)$
27	$(1,0,-1,3)$
28	$(3,0,-3,-2)$
29	$(1,0,2,2)$
30	$(4,0,-1,-1)$
31	$(1,-1,0,3)$
32	$(3,-3,0,-2)$
33	$(5,3,4,4)$
34	$(7,-9,-1,-1)$
35	$(1,1,3,3)$
36	$(3,3,-1,-1)$
37	$(5,4,3,4)$
38	$(7,-1,-9,-1)$
39	$(2,-2,-1,3)$

i	v_i
40	$(1,-1,-5,-3)$
41	$(4,5,3,4)$
42	$(1,-7,9,1)$
43	$(4,-4,-1,-9)$
44	$(5,-5,13,3)$
45	$(6,5,4,-5)$
46	$(9,-1,-11,1)$
47	$(3,4,2,-3)$
48	$(1,-5,7,-1)$

TABLE S32: All 21 full bases of the vertex-deleted completion.

e	Context
1	$\{9,10,12,13\}$
2	$\{9,11,14,15\}$
3	$\{10,11,16,17\}$
4	$\{2,3,12,18\}$
5	$\{1,3,15,19\}$
6	$\{6,7,13,18\}$
7	$\{6,8,14,19\}$
8	$\{1,2,21,22\}$
9	$\{1,13,23,24\}$
10	$\{1,17,25,26\}$
11	$\{2,14,27,28\}$
12	$\{2,17,29,30\}$
13	$\{3,16,31,32\}$
14	$\{6,17,33,34\}$
15	$\{7,8,35,36\}$
16	$\{7,15,37,38\}$
17	$\{7,16,39,40\}$
18	$\{8,12,41,42\}$
19	$\{8,16,43,44\}$
20	$\{10,19,45,46\}$
21	$\{11,18,47,48\}$

S1.17.3. All vectors and bases of the two 68-ray parents

In each parent retain all 65 vectors of Appendix S1.15.1, with their original labels, and add the three vectors in the corresponding rows of Table S33; Tables S34 and S35 list all 35 bases of each parent. Thus the six rows specify every added ray; the two parent inventories use distinct local meanings for labels 66–68. These labels also differ from the 72-ray parent numbering in Appendix S1.16.

TABLE S33: The three added rays in each local 68-ray parent inventory.

Parent variant	i	v_i
18–9 subsystem	66	$(1,0,0,1)$
18–9 subsystem	67	$(0,1,0,1)$
18–9 subsystem	68	$(0,0,1,1)$
20–11 subsystem	66	$(0,1,-1,0)$
20–11 subsystem	67	$(1,0,-1,0)$
20–11 subsystem	68	$(1,-1,0,0)$

TABLE S34: All 35 full bases of 68-ray parent variant 1.

e	Context	e	Context
1	$\{1,2,3,4\}$	2	$\{1,2,17,68\}$
3	$\{1,3,15,67\}$	4	$\{1,13,22,23\}$
5	$\{1,15,30,31\}$	6	$\{1,17,38,39\}$
7	$\{2,3,12,66\}$	8	$\{2,12,18,19\}$
9	$\{2,14,26,27\}$	10	$\{2,17,40,41\}$
11	$\{3,12,20,21\}$	12	$\{3,15,32,33\}$
13	$\{3,16,34,35\}$	14	$\{4,13,24,25\}$
15	$\{4,14,28,29\}$	16	$\{4,16,36,37\}$
17	$\{5,6,7,8\}$	18	$\{5,12,42,43\}$
19	$\{5,15,54,55\}$	20	$\{5,17,62,63\}$
21	$\{6,7,13,66\}$	22	$\{6,8,14,67\}$
23	$\{6,13,46,47\}$	24	$\{6,14,50,51\}$
25	$\{6,17,64,65\}$	26	$\{7,8,16,68\}$
27	$\{7,13,48,49\}$	28	$\{7,15,56,57\}$
29	$\{7,16,58,59\}$	30	$\{8,12,44,45\}$
31	$\{8,14,52,53\}$	32	$\{8,16,60,61\}$
33	$\{9,10,12,13\}$	34	$\{9,11,14,15\}$
35	$\{10,11,16,17\}$		

TABLE S35: All 35 full bases of 68-ray parent variant 2.

e	Context	e	Context
1	$\{1,2,3,4\}$	2	$\{1,4,13,66\}$
3	$\{1,13,22,23\}$	4	$\{1,15,30,31\}$
5	$\{1,17,38,39\}$	6	$\{2,4,14,67\}$
7	$\{2,12,18,19\}$	8	$\{2,14,26,27\}$
9	$\{2,17,40,41\}$	10	$\{3,4,16,68\}$
11	$\{3,12,20,21\}$	12	$\{3,15,32,33\}$
13	$\{3,16,34,35\}$	14	$\{4,13,24,25\}$
15	$\{4,14,28,29\}$	16	$\{4,16,36,37\}$
17	$\{5,6,7,8\}$	18	$\{5,6,17,68\}$
19	$\{5,7,15,67\}$	20	$\{5,8,12,66\}$
21	$\{5,12,42,43\}$	22	$\{5,15,54,55\}$
23	$\{5,17,62,63\}$	24	$\{6,13,46,47\}$
25	$\{6,14,50,51\}$	26	$\{6,17,64,65\}$
27	$\{7,13,48,49\}$	28	$\{7,15,56,57\}$
29	$\{7,16,58,59\}$	30	$\{8,12,44,45\}$
31	$\{8,14,52,53\}$	32	$\{8,16,60,61\}$
33	$\{9,10,12,13\}$	34	$\{9,11,14,15\}$
35	$\{10,11,16,17\}$		

For variant 1 the nine-context parity subsystem consists of exactly $C_3, C_4, C_5, D_1, \ldots, D_6$ with $a = 66, b = 67, c = 68$. For variant 2 the twenty-ray subsystem consists of all rows whose labels lie in $\{1, \ldots, 17, 66, 67, 68\}$. The machine-readable inventories repeat every coordinate, basis, and inclusion map.

S2. TWO-VALUED STATES, EXTENDABILITY CERTIFICATES, COLORINGS, AND SEPARATION PROFILES

S2.1. State, coloring, odd-cycle certificate, and profile tables for chromatically complete configurations ($\chi(H) = d$)

This subsection provides the complete state and coloring enumerations and odd-cycle nonextendability certificates for $H_{6,3}$ (Table S36), P_5 (Table S37), H_8 (Tables S38 and S39), $H_{17,10}$ (Tables S40 and S41), $H_{21,13}$ (Tables S42, S43, and S44), the vertex separation profiles and Born probability decomposition on $H_{17,10}$ and $H_{21,13}$ (Tables S45–S47), the complete state and normalized coloring tables for all faithful 3- and 4-dimensional logical gadgets (Tables S48–S69 and S70), and the ten chromatically inseparable faithful restrictions of H_{YO}^{+} (Table S71).

TABLE S36: All two-valued states of $H_{6,3}$. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4\}$	yes	–
2	$\{2,4,6\}$	no	$(3,1,5)$
3	$\{2,5\}$	yes	–
4	$\{3,6\}$	yes	–

TABLE S37: All two-valued states of P_5. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,3,9\}$	yes	–
2	$\{1,4,7\}$	yes	–
3	$\{1,7,8,9\}$	yes	–
4	$\{2,4,10\}$	yes	–
5	$\{2,5,8\}$	yes	–
6	$\{2,8,9,10\}$	yes	–
7	$\{3,5,6\}$	yes	–
8	$\{3,6,9,10\}$	yes	–
9	$\{4,6,7,10\}$	yes	–
10	$\{5,6,7,8\}$	yes	–
11	$\{6,7,8,9,10\}$	no	$(3,2,1,5,4)$

TABLE S38: All two-valued states of H_8. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4,6,9\}$	yes	–
2	$\{1,5,8,10,14\}$	no	$(9,7,6,12,11)$
3	$\{1,5,9,13,14\}$	yes	–
4	$\{2,4,6,8,11\}$	yes	–
5	$\{2,4,7,11,14\}$	no	$(5,3,1,12,6)$
6	$\{2,5,8,10,12\}$	yes	–
7	$\{2,5,8,11,13,14\}$	yes	–
8	$\{2,5,9,12,13\}$	no	$(4,3,1,11,10)$
9	$\{3,6,8,11,13\}$	no	$(7,5,4,10,9)$
10	$\{3,7,10,12\}$	yes	–
11	$\{3,7,11,13,14\}$	yes	–

TABLE S39. All strong three-colorings of H_8 up to color permutation. The anchor is $(1,2,3) = (0,1,2)$.

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14
c_1	0	1	2	0	1	0	2	1	0	1	2	1	2	2
c_2	0	1	2	0	1	0	2	1	0	2	1	2	1	1
c_3	0	1	2	1	0	1	2	1	0	2	1	2	0	0

TABLE S40: All two-valued states of $H_{17,10}$. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4,8,14,15,23\}$	yes	–
2	$\{1,5,7,14,18,23,24\}$	no	$(4,11,3,2,8,9,10)$
3	$\{1,5,9,11,18\}$	yes	–
4	$\{1,5,9,14,15,18,24\}$	no	$(11,3,2,8,7)$
5	$\{1,6,8,11,18,23\}$	yes	–
6	$\{1,6,8,14,15,18,23,24\}$	no	$(11,3,2,5,4)$
7	$\{2,4,7,14,19,23\}$	yes	–
8	$\{2,4,9,14,15,19\}$	yes	–
9	$\{2,6,7,10,14,24\}$	yes	–
10	$\{2,6,7,14,18,19,23,24\}$	no	$(11,3,1,10,4)$
11	$\{2,6,9,11,18,19\}$	yes	–
12	$\{2,6,9,14,15,18,19,24\}$	no	$(11,3,1,10,4)$
13	$\{3,4,8,15,19,23\}$	yes	–
14	$\{3,5,7,10,24\}$	yes	–
15	$\{3,5,7,18,19,23,24\}$	no	$(8,2,1,10,9)$
16	$\{3,5,9,15,18,19,24\}$	no	$(7,8,2,1,10,4,11)$
17	$\{3,6,8,10,15,24\}$	yes	–
18	$\{3,6,8,15,18,19,23,24\}$	no	$(5,2,1,10,4)$

TABLE S41. All strong three-colorings of $H_{17,10}$ up to color permutation. The anchor is $(2,8,5)=(0,1,2)$.

i	1	2	3	4	5	6	7	8	9	10	11	14	15	18	19	23	24
c_1	1	0	2	0	2	1	2	1	0	2	1	0	0	1	0	1	2
c_2	1	0	2	1	2	0	2	1	0	2	0	1	1	0	0	1	2
c_3	2	0	1	0	2	1	0	1	2	1	2	0	1	2	0	0	1
c_4	2	0	1	1	2	0	0	1	2	0	2	0	1	2	1	1	0

TABLE S42: All two-valued states of $H_{21,13}$. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4,8,14,15,21,22,23\}$	no	$(7,11,3,2,5,6,13)$
2	$\{1,5,7,14,18,22,23,24\}$	no	$(4,11,3,2,8,9,10)$
3	$\{1,5,9,11,18,21,22\}$	yes	–
4	$\{1,5,9,14,15,18,21,22,24\}$	no	$(11,3,2,8,7)$
5	$\{1,6,8,11,18,21,23\}$	yes	–
6	$\{1,6,8,14,15,18,21,23,24\}$	no	$(11,3,2,5,4)$
7	$\{2,4,7,14,19,20,22,23\}$	no	$(8,9,10,1,13,6,5)$
8	$\{2,4,9,13,14,15,19\}$	yes	–
9	$\{2,4,9,14,15,19,20,21,22\}$	no	$(11,3,1,13,7)$
10	$\{2,6,7,10,14,20,24\}$	yes	–
11	$\{2,6,7,14,18,19,20,23,24\}$	no	$(11,3,1,10,4)$
12	$\{2,6,9,11,18,19,20,21\}$	no	$(5,4,10,1,13,7,8)$
13	$\{2,6,9,14,15,18,19,20,21,24\}$	no	$(11,3,1,10,4)$
14	$\{3,4,8,13,15,19,23\}$	yes	–
15	$\{3,4,8,15,19,20,21,22,23\}$	no	$(5,2,1,13,6)$
16	$\{3,5,7,10,20,22,24\}$	yes	–
17	$\{3,5,7,18,19,20,22,23,24\}$	no	$(8,2,1,10,9)$
18	$\{3,5,9,13,15,18,19,24\}$	no	$(7,8,2,1,10,4,11)$
19	$\{3,5,9,15,18,19,20,21,22,24\}$	no	$(8,2,1,13,7)$
20	$\{3,6,8,10,15,20,21,24\}$	no	$(4,5,2,1,13,7,11)$
21	$\{3,6,8,15,18,19,20,21,23,24\}$	no	$(5,2,1,10,4)$

TABLE S43. All strong three-colorings of $H_{21,13}$ up to color permutation. The anchor is $(2,8,5)=(0,1,2)$.

i	1	2	3	4	5	6	7	8	9	10	11	13	14	15	18	19	20	21	22	23	24
c_1	1	0	2	0	2	1	2	1	0	2	1	0	0	0	1	0	2	1	2	1	2
c_2	2	0	1	1	2	0	0	1	2	0	2	1	0	1	2	1	0	2	2	1	0

TABLE S44. The nine chromatic signatures of $H_{21,13}$. Columns give c_1 and rows give c_2 for the two normalized colorings in Table S43. Each cell is one inseparability class: its vertices have the same color in every admissible coloring. Ordinary two-valued states distinguish every pair, including vertices in the same cell.

$c_2 \backslash c_1$	0	1	2
2	$\{9\}$	$\{1, 11, 18, 21\}$	$\{5, 22\}$
1	$\{4, 13, 15, 19\}$	$\{8, 23\}$	$\{3\}$
0	$\{2, 14\}$	$\{6\}$	$\{7, 10, 20, 24\}$

TABLE S46. Valid states and exact weights reproducing two $H_{17,10}$ preparations: the five-state mixture for P_{24} (upper block, Eq. (27)) and the four-state mixture for P_{z_3}, the preparation along $z_3 = r_5$ (lower block). State numbering is that of Table S40. Every context equation and every Born marginal are checked with rational arithmetic.

State	Weight	Support	Extendable?
Preparation P_{24}, nonextendable weight $\frac{1}{9}$			
s_4	$\frac{1}{12}$	$\{1, 5, 9, 14, 15, 18, 24\}$	no
s_9	$\frac{5}{36}$	$\{2, 6, 7, 10, 14, 24\}$	yes
s_{10}	$\frac{1}{36}$	$\{2, 6, 7, 14, 18, 19, 23, 24\}$	no
s_{14}	$\frac{7}{12}$	$\{3, 5, 7, 10, 24\}$	yes
s_{17}	$\frac{1}{6}$	$\{3, 6, 8, 10, 15, 24\}$	yes
Preparation P_{z_3}, nonextendable weight $\frac{1}{3}$			
s_3	$\frac{1}{3}$	$\{1, 5, 9, 11, 18\}$	yes
s_4	$\frac{1}{6}$	$\{1, 5, 9, 14, 15, 18, 24\}$	no
s_{14}	$\frac{1}{3}$	$\{3, 5, 7, 10, 24\}$	yes
s_{15}	$\frac{1}{6}$	$\{3, 5, 7, 18, 19, 23, 24\}$	no

TABLE S45: Complete state profiles in hexadecimal, and all Born probabilities for the 17-ray preparation in Eq. (26). The probabilities are also reproduced by Eq. (27).

i	$H_{17,10}$ profile	$H_{21,13}$ profile	p_i on $H_{17,10}$
1	3F	3F	1/12
2	FC0	1FC0	1/6
3	3F000	1FE000	3/4
4	10C1	61C1	0
5	E00E	7800E	2/3
6	30F30	181E30	1/3
7	6342	18642	3/4
8	31031	186031	1/6
9	8C8C	6198C	1/12
10	12100	88200	8/9
11	414	814	0
13	-	22080	–
14	BEB	17EB	1/4
15	398A9	1E71A9	1/4
18	2CE3E	171C3E	1/9
19	2DEC0	177DC0	1/36
20	-	1DDF40	–
21	-	1C593D	–
22	-	5C14F	–
23	25273	116473	1/36
24	3EB2A	1F962A	1

TABLE S47. Exact Born probabilities $p_i = (r_{24} \cdot r_i)^2/(6\|r_i\|^2)$ for P_{24} and $p'_i = (r_5 \cdot r_i)^2/\|r_i\|^2$ for P_{z_3}, also reproduced by the two mixtures of Table S46.

i	p_i	p'_i	i	p_i	p'_i
1	$\frac{1}{12}$	$\frac{1}{2}$	10	$\frac{8}{9}$	$\frac{1}{3}$
2	$\frac{1}{6}$	0	11	0	$\frac{1}{3}$
3	$\frac{3}{4}$	$\frac{1}{2}$	14	$\frac{1}{4}$	$\frac{1}{6}$
4	0	0	15	$\frac{1}{4}$	$\frac{1}{6}$
5	$\frac{2}{3}$	1	18	$\frac{1}{9}$	$\frac{2}{3}$
6	$\frac{1}{3}$	0	19	$\frac{1}{36}$	$\frac{1}{6}$
7	$\frac{3}{4}$	$\frac{1}{2}$	23	$\frac{1}{36}$	$\frac{1}{6}$
8	$\frac{1}{6}$	0	24	1	$\frac{2}{3}$
9	$\frac{1}{12}$	$\frac{1}{2}$			

S2.2. State conventions for the additional gadgets

The state tables below are local to each labeled configuration. Supports are in lexicographic order; E is yes precisely for color-class supports. The three-dimensional pentagon and $H_{21,13}$ use the complete earlier tables and are not repeated. The four-dimensional apex-only states are included explicitly. All ordinary state families listed here are unital and separating.

S2.3. All states of $B_{13,7}$

TABLE S48: All two-valued states of $B_{13,7}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,5,7,14,23\}$	yes	8	$\{2,9,11,19\}$	yes
2	$\{1,5,9,11\}$	yes	9	$\{2,9,14,15,19\}$	yes
3	$\{1,5,9,14,15\}$	no	10	$\{3,5,7,10\}$	yes
4	$\{1,8,11,23\}$	yes	11	$\{3,5,7,19,23\}$	no
5	$\{1,8,14,15,23\}$	yes	12	$\{3,5,9,15,19\}$	yes
6	$\{2,7,10,14\}$	yes	13	$\{3,8,10,15\}$	yes
7	$\{2,7,14,19,23\}$	yes	14	$\{3,8,15,19,23\}$	yes

S2.4. All states of $D_{16,9}$

TABLE S49: All two-valued states of $D_{16,9}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,4,8,14,15,23\}$	yes	12	$\{2,6,7,10,14,24\}$	yes
2	$\{1,5,7,10,14,24\}$	yes	13	$\{2,6,7,14,18,23,24\}$	no
3	$\{1,5,7,14,18,23,24\}$	no	14	$\{2,6,9,11,18\}$	yes
4	$\{1,5,9,11,18\}$	yes	15	$\{2,6,9,14,15,18,24\}$	no
5	$\{1,5,9,14,15,18,24\}$	no	16	$\{3,4,8,15,23\}$	yes
6	$\{1,6,8,10,11\}$	yes	17	$\{3,5,7,10,24\}$	yes
7	$\{1,6,8,10,14,15,24\}$	no	18	$\{3,5,7,18,23,24\}$	yes
8	$\{1,6,8,11,18,23\}$	yes	19	$\{3,5,9,15,18,24\}$	yes
9	$\{1,6,8,14,15,18,23,24\}$	no	20	$\{3,6,8,10,15,24\}$	yes
10	$\{2,4,7,14,23\}$	yes	21	$\{3,6,8,15,18,23,24\}$	no
11	$\{2,4,9,14,15\}$	yes			

S2.5. All states of $I_{16,9}$

TABLE S50: All two-valued states of $I_{16,9}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,4,5,7,14,23\}$	yes	12	$\{2,7,14,18,19,23,24\}$	no
2	$\{1,4,5,9,14,15\}$	no	13	$\{2,9,11,18,19\}$	yes
3	$\{1,4,8,14,15,23\}$	yes	14	$\{2,9,14,15,18,19,24\}$	no
4	$\{1,5,7,14,18,23,24\}$	no	15	$\{3,4,5,7,19,23\}$	no
5	$\{1,5,9,11,18\}$	yes	16	$\{3,4,5,9,15,19\}$	yes
6	$\{1,5,9,14,15,18,24\}$	no	17	$\{3,4,8,15,19,23\}$	yes
7	$\{1,8,11,18,23\}$	yes	18	$\{3,5,7,10,24\}$	yes
8	$\{1,8,14,15,18,23,24\}$	no	19	$\{3,5,7,18,19,23,24\}$	no
9	$\{2,4,7,14,19,23\}$	yes	20	$\{3,5,9,15,18,19,24\}$	no
10	$\{2,4,9,14,15,19\}$	yes	21	$\{3,8,10,15,24\}$	yes
11	$\{2,7,10,14,24\}$	yes	22	$\{3,8,15,18,19,23,24\}$	no

S2.6. All states of $T_{19,11}$

TABLE S51: All two-valued states of $T_{19,11}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,4,8,14,15,23,28\}$	yes	14	$\{2,6,7,14,18,23,24\}$	no
2	$\{1,5,7,10,14,24,28\}$	yes	15	$\{2,6,9,11,18,27\}$	yes
3	$\{1,5,7,14,18,23,24\}$	no	16	$\{2,6,9,14,15,18,24\}$	no
4	$\{1,5,9,11,18,27\}$	yes	17	$\{3,4,8,15,23,26\}$	yes
5	$\{1,5,9,14,15,18,24\}$	no	18	$\{3,4,8,15,23,27,28\}$	no
6	$\{1,6,8,10,11,26\}$	yes	19	$\{3,5,7,10,24,26\}$	yes
7	$\{1,6,8,10,11,27,28\}$	no	20	$\{3,5,7,10,24,27,28\}$	no
8	$\{1,6,8,10,14,15,24,28\}$	no	21	$\{3,5,7,18,23,24,27\}$	yes
9	$\{1,6,8,11,18,23,27\}$	yes	22	$\{3,5,9,15,18,24,27\}$	yes
10	$\{1,6,8,14,15,18,23,24\}$	no	23	$\{3,6,8,10,15,24,26\}$	yes
11	$\{2,4,7,14,23,28\}$	yes	24	$\{3,6,8,10,15,24,27,28\}$	no
12	$\{2,4,9,14,15,28\}$	yes	25	$\{3,6,8,15,18,23,24,27\}$	no
13	$\{2,6,7,10,14,24,28\}$	yes			

S2.7. All states of $W^{(4)}_{11,5}$

TABLE S52: All two-valued states of $W^{(4)}_{11,5}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0\}$	yes	7	$\{2,8,9,10\}$	yes
2	$\{1,3,9\}$	yes	8	$\{3,5,6\}$	yes
3	$\{1,4,7\}$	yes	9	$\{3,6,9,10\}$	yes
4	$\{1,7,8,9\}$	yes	10	$\{4,6,7,10\}$	yes
5	$\{2,4,10\}$	yes	11	$\{5,6,7,8\}$	yes
6	$\{2,5,8\}$	yes	12	$\{6,7,8,9,10\}$	no

S2.8. All states of $W_{16,6}^{(4)}$

TABLE S53: All two-valued states of $W_{16,6}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0,7,9,10\}$	yes	37	$\{2,8,12,14,15\}$	yes
2	$\{0,7,9,15\}$	yes	38	$\{3,5,6\}$	yes
3	$\{0,7,10,14\}$	yes	39	$\{3,6,9,10,11\}$	yes
4	$\{0,7,14,15\}$	yes	40	$\{3,6,9,10,12\}$	yes
5	$\{0,9,10,13\}$	yes	41	$\{3,6,9,11,15\}$	yes
6	$\{0,9,13,15\}$	yes	42	$\{3,6,9,12,15\}$	yes
7	$\{0,10,13,14\}$	yes	43	$\{3,6,10,11,14\}$	yes
8	$\{0,13,14,15\}$	yes	44	$\{3,6,10,12,14\}$	yes
9	$\{1,3,9,11\}$	yes	45	$\{3,6,11,14,15\}$	yes
10	$\{1,3,9,12\}$	yes	46	$\{3,6,12,14,15\}$	yes
11	$\{1,3,11,14\}$	yes	47	$\{4,6,7,10,11\}$	yes
12	$\{1,3,12,14\}$	yes	48	$\{4,6,7,10,12\}$	yes
13	$\{1,4,7,11\}$	yes	49	$\{4,6,7,11,15\}$	yes
14	$\{1,4,7,12\}$	yes	50	$\{4,6,7,12,15\}$	yes
15	$\{1,4,11,13\}$	yes	51	$\{4,6,10,11,13\}$	yes
16	$\{1,4,12,13\}$	yes	52	$\{4,6,10,12,13\}$	yes
17	$\{1,7,8,9,11\}$	yes	53	$\{4,6,11,13,15\}$	yes
18	$\{1,7,8,9,12\}$	yes	54	$\{4,6,12,13,15\}$	yes
19	$\{1,7,8,11,14\}$	yes	55	$\{5,6,7,8\}$	yes
20	$\{1,7,8,12,14\}$	yes	56	$\{5,6,8,13\}$	yes
21	$\{1,8,9,11,13\}$	yes	57	$\{6,7,8,9,10,11\}$	no
22	$\{1,8,9,12,13\}$	yes	58	$\{6,7,8,9,10,12\}$	no
23	$\{1,8,11,13,14\}$	yes	59	$\{6,7,8,9,11,15\}$	no
24	$\{1,8,12,13,14\}$	yes	60	$\{6,7,8,9,12,15\}$	no
25	$\{2,4,10,11\}$	yes	61	$\{6,7,8,10,11,14\}$	no
26	$\{2,4,10,12\}$	yes	62	$\{6,7,8,10,12,14\}$	no
27	$\{2,4,11,15\}$	yes	63	$\{6,7,8,11,14,15\}$	no
28	$\{2,4,12,15\}$	yes	64	$\{6,7,8,12,14,15\}$	no
29	$\{2,5,8\}$	yes	65	$\{6,8,9,10,11,13\}$	no
30	$\{2,8,9,10,11\}$	yes	66	$\{6,8,9,10,12,13\}$	no
31	$\{2,8,9,10,12\}$	yes	67	$\{6,8,9,11,13,15\}$	no
32	$\{2,8,9,11,15\}$	yes	68	$\{6,8,9,12,13,15\}$	no
33	$\{2,8,9,12,15\}$	yes	69	$\{6,8,10,11,13,14\}$	no
34	$\{2,8,10,11,14\}$	yes	70	$\{6,8,10,12,13,14\}$	no
35	$\{2,8,10,12,14\}$	yes	71	$\{6,8,11,13,14,15\}$	no
36	$\{2,8,11,14,15\}$	yes	72	$\{6,8,12,13,14,15\}$	no

S2.9. All states of $B_{14,7}^{(4)}$

TABLE S54: All two-valued states of $B_{14,7}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0\}$	yes	9	$\{2,9,11,19\}$	yes
2	$\{1,5,7,14,23\}$	yes	10	$\{2,9,14,15,19\}$	yes
3	$\{1,5,9,11\}$	yes	11	$\{3,5,7,10\}$	yes
4	$\{1,5,9,14,15\}$	no	12	$\{3,5,7,19,23\}$	no
5	$\{1,8,11,23\}$	yes	13	$\{3,5,9,15,19\}$	yes
6	$\{1,8,14,15,23\}$	yes	14	$\{3,8,10,15\}$	yes
7	$\{2,7,10,14\}$	yes	15	$\{3,8,15,19,23\}$	yes
8	$\{2,7,14,19,23\}$	yes			

S2.10. All states of $L_{18,7}^{(4)}$

TABLE S55: All two-valued states of $L_{18,7}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,6,9,13,16\}$	yes	35	$\{3,5,9,13,18\}$	yes
2	$\{1,6,9,14,16\}$	yes	36	$\{3,5,9,14,17\}$	yes
3	$\{1,6,10,13,16\}$	yes	37	$\{3,5,9,14,18\}$	yes
4	$\{1,6,10,14,16\}$	yes	38	$\{3,5,10,13,17\}$	yes
5	$\{1,7,9,13,17\}$	yes	39	$\{3,5,10,13,18\}$	yes
6	$\{1,7,9,13,18\}$	yes	40	$\{3,5,10,14,17\}$	yes
7	$\{1,7,9,14,17\}$	yes	41	$\{3,5,10,14,18\}$	yes
8	$\{1,7,9,14,18\}$	yes	42	$\{3,6,9,11\}$	yes
9	$\{1,7,10,13,17\}$	yes	43	$\{3,6,9,12,13\}$	yes
10	$\{1,7,10,13,18\}$	yes	44	$\{3,6,9,12,14\}$	yes
11	$\{1,7,10,14,17\}$	yes	45	$\{3,6,10,11\}$	yes
12	$\{1,7,10,14,18\}$	yes	46	$\{3,6,10,12,13\}$	yes
13	$\{1,8,13,15\}$	yes	47	$\{3,6,10,12,14\}$	yes
14	$\{1,8,13,16,17\}$	yes	48	$\{3,8,11,17\}$	yes
15	$\{1,8,13,16,18\}$	yes	49	$\{3,8,11,18\}$	yes
16	$\{1,8,14,15\}$	yes	50	$\{3,8,12,13,17\}$	yes
17	$\{1,8,14,16,17\}$	yes	51	$\{3,8,12,13,18\}$	yes
18	$\{1,8,14,16,18\}$	yes	52	$\{3,8,12,14,17\}$	yes
19	$\{2,5,9,15\}$	yes	53	$\{3,8,12,14,18\}$	yes
20	$\{2,5,9,16,17\}$	yes	54	$\{4,5,13,15\}$	yes
21	$\{2,5,9,16,18\}$	yes	55	$\{4,5,13,16,17\}$	yes
22	$\{2,5,10,15\}$	yes	56	$\{4,5,13,16,18\}$	yes
23	$\{2,5,10,16,17\}$	yes	57	$\{4,5,14,15\}$	yes
24	$\{2,5,10,16,18\}$	yes	58	$\{4,5,14,16,17\}$	yes
25	$\{2,6,9,12,16\}$	yes	59	$\{4,5,14,16,18\}$	yes
26	$\{2,6,10,12,16\}$	yes	60	$\{4,6,11,16\}$	yes
27	$\{2,7,9,12,17\}$	yes	61	$\{4,6,12,13,16\}$	yes
28	$\{2,7,9,12,18\}$	yes	62	$\{4,6,12,14,16\}$	yes
29	$\{2,7,10,12,17\}$	yes	63	$\{4,7,11,17\}$	yes
30	$\{2,7,10,12,18\}$	yes	64	$\{4,7,11,18\}$	yes
31	$\{2,8,12,15\}$	yes	65	$\{4,7,12,13,17\}$	yes
32	$\{2,8,12,16,17\}$	yes	66	$\{4,7,12,13,18\}$	yes
33	$\{2,8,12,16,18\}$	yes	67	$\{4,7,12,14,17\}$	yes
34	$\{3,5,9,13,17\}$	yes	68	$\{4,7,12,14,18\}$	yes

S2.11. All states of $D_{17,9}^{(4)}$

TABLE S56: All two-valued states of $D_{17,9}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0\}$	yes	12	$\{2,4,9,14,15\}$	yes
2	$\{1,4,8,14,15,23\}$	yes	13	$\{2,6,7,10,14,24\}$	yes
3	$\{1,5,7,10,14,24\}$	yes	14	$\{2,6,7,14,18,23,24\}$	no
4	$\{1,5,7,14,18,23,24\}$	no	15	$\{2,6,9,11,18\}$	yes
5	$\{1,5,9,11,18\}$	yes	16	$\{2,6,9,14,15,18,24\}$	no
6	$\{1,5,9,14,15,18,24\}$	no	17	$\{3,4,8,15,23\}$	yes
7	$\{1,6,8,10,11\}$	yes	18	$\{3,5,7,10,24\}$	yes
8	$\{1,6,8,10,14,15,24\}$	no	19	$\{3,5,7,18,23,24\}$	yes
9	$\{1,6,8,11,18,23\}$	yes	20	$\{3,5,9,15,18,24\}$	yes
10	$\{1,6,8,14,15,18,23,24\}$	no	21	$\{3,6,8,10,15,24\}$	yes
11	$\{2,4,7,14,23\}$	yes	22	$\{3,6,8,15,18,23,24\}$	no

S2.12. All states of $L_{22,9}^{(4)}$

TABLE S57: All two-valued states of $L_{22,9}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,6,9,13,16,21\}$	yes	80	$\{2,8,12,16,17,20,21\}$	no
2	$\{1,6,9,13,16,22\}$	yes	81	$\{2,8,12,16,17,20,22\}$	no
3	$\{1,6,9,14,16,19\}$	yes	82	$\{2,8,12,16,18,19\}$	yes
4	$\{1,6,9,14,16,20,21\}$	no	83	$\{2,8,12,16,18,20,21\}$	no
5	$\{1,6,9,14,16,20,22\}$	no	84	$\{2,8,12,16,18,20,22\}$	no
6	$\{1,6,10,13,16,21\}$	yes	85	$\{3,5,9,13,17,21\}$	yes
7	$\{1,6,10,13,16,22\}$	yes	86	$\{3,5,9,13,17,22\}$	yes
8	$\{1,6,10,14,16,19\}$	yes	87	$\{3,5,9,13,18,21\}$	yes
9	$\{1,6,10,14,16,20,21\}$	no	88	$\{3,5,9,13,18,22\}$	yes
10	$\{1,6,10,14,16,20,22\}$	no	89	$\{3,5,9,14,17,19\}$	yes
11	$\{1,7,9,13,17,21\}$	yes	90	$\{3,5,9,14,17,20,21\}$	no
12	$\{1,7,9,13,17,22\}$	yes	91	$\{3,5,9,14,17,20,22\}$	no
13	$\{1,7,9,13,18,21\}$	yes	92	$\{3,5,9,14,18,19\}$	yes
14	$\{1,7,9,13,18,22\}$	yes	93	$\{3,5,9,14,18,20,21\}$	no
15	$\{1,7,9,14,17,19\}$	yes	94	$\{3,5,9,14,18,20,22\}$	no
16	$\{1,7,9,14,17,20,21\}$	no	95	$\{3,5,10,13,17,21\}$	yes
17	$\{1,7,9,14,17,20,22\}$	no	96	$\{3,5,10,13,17,22\}$	yes
18	$\{1,7,9,14,18,19\}$	yes	97	$\{3,5,10,13,18,21\}$	yes
19	$\{1,7,9,14,18,20,21\}$	no	98	$\{3,5,10,13,18,22\}$	yes
20	$\{1,7,9,14,18,20,22\}$	no	99	$\{3,5,10,14,17,19\}$	yes
21	$\{1,7,10,13,17,21\}$	yes	100	$\{3,5,10,14,17,20,21\}$	no
22	$\{1,7,10,13,17,22\}$	yes	101	$\{3,5,10,14,17,20,22\}$	no
23	$\{1,7,10,13,18,21\}$	yes	102	$\{3,5,10,14,18,19\}$	yes
24	$\{1,7,10,13,18,22\}$	yes	103	$\{3,5,10,14,18,20,21\}$	no
25	$\{1,7,10,14,17,19\}$	yes	104	$\{3,5,10,14,18,20,22\}$	no
26	$\{1,7,10,14,17,20,21\}$	no	105	$\{3,6,9,11,20\}$	yes
27	$\{1,7,10,14,17,20,22\}$	no	106	$\{3,6,9,12,13,21\}$	yes
28	$\{1,7,10,14,18,19\}$	yes	107	$\{3,6,9,12,13,22\}$	yes
29	$\{1,7,10,14,18,20,21\}$	no	108	$\{3,6,9,12,14,19\}$	yes
30	$\{1,7,10,14,18,20,22\}$	no	109	$\{3,6,9,12,14,20,21\}$	no
31	$\{1,8,13,16,17,21\}$	yes	110	$\{3,6,9,12,14,20,22\}$	no
32	$\{1,8,13,16,17,22\}$	yes	111	$\{3,6,10,11,20\}$	yes
33	$\{1,8,13,16,18,21\}$	yes	112	$\{3,6,10,12,13,21\}$	yes
34	$\{1,8,13,16,18,22\}$	yes	113	$\{3,6,10,12,13,22\}$	yes
35	$\{1,8,14,15,21\}$	yes	114	$\{3,6,10,12,14,19\}$	yes
36	$\{1,8,14,15,22\}$	yes	115	$\{3,6,10,12,14,20,21\}$	no
37	$\{1,8,14,16,17,19\}$	yes	116	$\{3,6,10,12,14,20,22\}$	no
38	$\{1,8,14,16,17,20,21\}$	no	117	$\{3,8,11,17,20\}$	yes
39	$\{1,8,14,16,17,20,22\}$	no	118	$\{3,8,11,18,20\}$	yes
40	$\{1,8,14,16,18,19\}$	yes	119	$\{3,8,12,13,17,21\}$	yes
41	$\{1,8,14,16,18,20,21\}$	no	120	$\{3,8,12,13,17,22\}$	yes
42	$\{1,8,14,16,18,20,22\}$	no	121	$\{3,8,12,13,18,21\}$	yes
43	$\{2,5,9,15,21\}$	yes	122	$\{3,8,12,13,18,22\}$	yes
44	$\{2,5,9,15,22\}$	yes	123	$\{3,8,12,14,17,19\}$	yes
45	$\{2,5,9,16,17,19\}$	yes	124	$\{3,8,12,14,17,20,21\}$	no
46	$\{2,5,9,16,17,20,21\}$	no	125	$\{3,8,12,14,17,20,22\}$	no
47	$\{2,5,9,16,17,20,22\}$	no	126	$\{3,8,12,14,18,19\}$	yes
48	$\{2,5,9,16,18,19\}$	yes	127	$\{3,8,12,14,18,20,21\}$	no
49	$\{2,5,9,16,18,20,21\}$	no	128	$\{3,8,12,14,18,20,22\}$	no
50	$\{2,5,9,16,18,20,22\}$	no	129	$\{4,5,13,16,17,21\}$	yes
51	$\{2,5,10,15,21\}$	yes	130	$\{4,5,13,16,17,22\}$	yes
52	$\{2,5,10,15,22\}$	yes	131	$\{4,5,13,16,18,21\}$	yes
53	$\{2,5,10,16,17,19\}$	yes	132	$\{4,5,13,16,18,22\}$	yes
54	$\{2,5,10,16,17,20,21\}$	no	133	$\{4,5,14,15,21\}$	yes
55	$\{2,5,10,16,17,20,22\}$	no	134	$\{4,5,14,15,22\}$	yes
56	$\{2,5,10,16,18,19\}$	yes	135	$\{4,5,14,16,17,19\}$	yes
57	$\{2,5,10,16,18,20,21\}$	no	136	$\{4,5,14,16,17,20,21\}$	no
58	$\{2,5,10,16,18,20,22\}$	no	137	$\{4,5,14,16,17,20,22\}$	no
59	$\{2,6,9,12,16,19\}$	yes	138	$\{4,5,14,16,18,19\}$	yes
60	$\{2,6,9,12,16,20,21\}$	no	139	$\{4,5,14,16,18,20,21\}$	no
61	$\{2,6,9,12,16,20,22\}$	no	140	$\{4,5,14,16,18,20,22\}$	no
62	$\{2,6,10,12,16,19\}$	yes	141	$\{4,6,11,16,20\}$	yes
63	$\{2,6,10,12,16,20,21\}$	no	142	$\{4,6,12,13,16,21\}$	yes
64	$\{2,6,10,12,16,20,22\}$	no	143	$\{4,6,12,13,16,22\}$	yes
65	$\{2,7,9,12,17,19\}$	yes	144	$\{4,6,12,14,16,19\}$	yes
66	$\{2,7,9,12,17,20,21\}$	no	145	$\{4,6,12,14,16,20,21\}$	no
67	$\{2,7,9,12,17,20,22\}$	no	146	$\{4,6,12,14,16,20,22\}$	no

j	Support	E	j	Support	E
68	$\{2,7,9,12,18,19\}$	yes	147	$\{4,7,11,17,20\}$	yes
69	$\{2,7,9,12,18,20,21\}$	no	148	$\{4,7,11,18,20\}$	yes
70	$\{2,7,9,12,18,20,22\}$	no	149	$\{4,7,12,13,17,21\}$	yes
71	$\{2,7,10,12,17,19\}$	yes	150	$\{4,7,12,13,17,22\}$	yes
72	$\{2,7,10,12,17,20,21\}$	no	151	$\{4,7,12,13,18,21\}$	yes
73	$\{2,7,10,12,17,20,22\}$	no	152	$\{4,7,12,13,18,22\}$	yes
74	$\{2,7,10,12,18,19\}$	yes	153	$\{4,7,12,14,17,19\}$	yes
75	$\{2,7,10,12,18,20,21\}$	no	154	$\{4,7,12,14,17,20,21\}$	no
76	$\{2,7,10,12,18,20,22\}$	no	155	$\{4,7,12,14,17,20,22\}$	no
77	$\{2,8,12,15,21\}$	yes	156	$\{4,7,12,14,18,19\}$	yes
78	$\{2,8,12,15,22\}$	yes	157	$\{4,7,12,14,18,20,21\}$	no
79	$\{2,8,12,16,17,19\}$	yes	158	$\{4,7,12,14,18,20,22\}$	no

S2.13. Complete normalized colorings of the gadget panels

Each color word gives one representative under global color permutation; the vertices and the anchor convention are specified separately for each table. The full labeled family is recovered by all palette permutations. Equal color labels across different colorings carry no extra constraint.

S2.14. Colorings of P_5

The columns of each word correspond, in order, to vertices $1,2,3,4,5,6,7,8,9,10$. The anchor context $(1,6,2)$ has colors $(0,\dots,2)$.

TABLE S58: Every normalized coloring of P_5.

j	Color word	j	Color word
1	0201211201	4	0210210211
2	0202111102	5	0212110002
3	0210110222		

S2.15. Colorings of $B_{13,7}$

The columns of each word correspond, in order, to vertices $1,2,3,5,7,8,9,10,11,14,15,19,23$. The anchor context $(2,8,5)$ has colors $(0,\dots,2)$.

TABLE S59: Every normalized coloring of $B_{13,7}$.

j	Color word	j	Color word
1	1022012010221	4	2012012020111
2	1022210201101	5	2012012120100
3	1022210210001	6	2012210102102

S2.16. Colorings of $D_{16,9}$

The columns of each word correspond, in order, to vertices $1,2,3,4,5,6,7,8,9,10,11,14,15,18,23,24$. The anchor context $(2,8,5)$ has colors $(0,\dots,2)$.

TABLE S60: Every normalized coloring of $D_{16,9}$.

j	Color word	j	Color word
1	1020210121102202	5	2010210121201201
2	1020212101100222	6	2011200120201210
3	1020212102100112	7	2011202102021012
4	1021202102011012		

S2.17. Colorings of $I_{16,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 14, 15, 18, 19, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S61: Every normalized coloring of $I_{16,9}$.

j	Color word	j	Color word
1	1020221021001012	4	2010201212012001
2	1021221020110012	5	2011201202012110
3	1022201201021210	6	2012221010210021

S2.18. Colorings of $T_{19,11}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24, 26, 27, 28$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S62: Every normalized coloring of $T_{19,11}$.

j	Color word	j	Color word
1	10202101211022021 20	5	20102101212012011 20
2	10202121011002221 20	6	20112001202012101 20
3	10202121021001122 10	7	20112021020210121 02
4	10212021020110122 01		

S2.19. Colorings of $H_{21,13}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 18, 19, 20, 21, 22, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S63: Every normalized coloring of $H_{21,13}$.

j	Color word	j	Color word
1	102021210210001021212	2	201120012021012102210

S2.20. Colorings of $W^{(4)}_{11,5}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10$. The anchor context $(0, 1, 6, 2)$ has colors $(0, \ldots, 3)$.

TABLE S64: Every normalized coloring of $W^{(4)}_{11,5}$.

j	Color word	j	Color word
1	01312322312	4	01321321322
2	01313222213	5	01323221113
3	01321221333		

S2.21. Colorings of $W^{(4)}_{16,6}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15$. The anchor context $(2, 0, 6, 1)$ has colors $(0, \ldots, 3)$.

TABLE S65: Every normalized coloring of $W^{(4)}_{16,6}$.

j	Color word	j	Color word
1	1302302101123322	41	1302022331003131
2	1302302101132322	42	1302022331030131
3	1302302101223321	43	1302022331103130
4	1302302101232321	44	1302022331130130
5	1302302102123312	45	1302022333003111
6	1302302102132312	46	1302022333030111
7	1302302102223311	47	1302022333103110
8	1302302102232311	48	1302022333130110
9	1302302301123122	49	1302322100003311
10	1302302301132122	50	1302322100030311
11	1302302301223121	51	1302322100103310
12	1302302301232121	52	1302322100130310
13	1302302302123112	53	1302322101003301
14	1302302302132112	54	1302322101030301
15	1302302302223111	55	1302322101103300
16	1302302302232111	56	1302322101130300
17	1303202101123232	57	1302322300003111
18	1303202101132232	58	1302322300030111
19	1303202101223231	59	1302322300103110
20	1303202101232231	60	1302322300130110
21	1303202103123212	61	1302322301003101
22	1303202103132212	62	1302322301030101
23	1303202103223211	63	1302322301103100
24	1303202103232211	64	1302322301130100
25	1303202201123132	65	1303022121003231
26	1303202201132132	66	1303022121030231
27	1303202201223131	67	1303022121103230
28	1303202201232131	68	1303022121130230
29	1303202203123112	69	1303022123003211
30	1303202203132112	70	1303022123030211
31	1303202203223111	71	1303022123103210
32	1303202203232111	72	1303022123130210
33	1302022131003331	73	1303022221003131
34	1302022131030331	74	1303022221030131
35	1302022131103330	75	1303022221103130
36	1302022131130330	76	1303022221130130
37	1302022133003311	77	1303022223003111
38	1302022133030311	78	1303022223030111
39	1302022133103310	79	1303022223103110
40	1302022133130310	80	1303022223130110

S2.22. Colorings of $B^{(4)}_{14,7}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 5, 7, 8, 9, 10, 11, 14, 15, 19, 23$. The anchor context $(0, 1, 2, 3)$ has colors $(0, \ldots, 3)$.

TABLE S66: Every normalized coloring of $B^{(4)}_{14,7}$.

j	Color word	j	Color word
1	01233213212331	4	01231132321321
2	01233312312221	5	01231231212333
3	01233312321121	6	01231231312322

S2.23. Colorings of $L^{(4)}_{18,7}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18$. The anchor context $(1, 2, 4, 3)$ has colors $(0, \ldots, 3)$.

TABLE S67: Every normalized coloring of $L^{(4)}_{18,7}$.

j	Color word	j	Color word
1	013212030123031203	73	013210230132021023
2	013212030123031230	74	013210230132021032
3	013212030123301203	75	013210230132201023
4	013212030123301230	76	013210230132201032
5	013212031023031203	77	013210231032021023
6	013212031023031230	78	013210231032021032
7	013212031023301203	79	013210231032201023
8	013212031023301230	80	013210231032201032
9	013210230123031023	81	013213201332020112
10	013210230123031032	82	013213201332020121
11	013210230123301023	83	013213201332200112
12	013210230123301032	84	013213201332200121
13	013210231023031023	85	013213203132020112
14	013210231023031032	86	013213203132020121
15	013210231023301023	87	013213203132200112
16	013210231023301032	88	013213203132200121
17	013213201323030112	89	013213201332021002
18	013213201323030121	90	013213201332021020
19	013213201323300112	91	013213201332201002
20	013213201323300121	92	013213201332201020
21	013213203123030112	93	013213203132021002
22	013213203123030121	94	013213203132021020
23	013213203123300112	95	013213203132201002
24	013213203123300121	96	013213203132201020
25	013213201323031002	97	013221030131022103
26	013213201323031020	98	013221030131022130
27	013213201323301002	99	013221030131202103
28	013213201323301020	100	013221030131202130
29	013213203123031002	101	013221031031022103
30	013213203123031020	102	013221031031022130
31	013213203123301002	103	013221031031202103
32	013213203123301020	104	013221031031202130
33	013231201321030123	105	013223010331021202

j	Color word	j	Color word
34	013231201321030132	106	013223010331021220
35	013231201321300123	107	013223010331201202
36	013231201321300132	108	013223010331201220
37	013231203121030123	109	013223013031021202
38	013231203121030132	110	013223013031021220
39	013231203121300123	111	013223013031201202
40	013231203121300132	112	013223013031201220
41	013232101321030213	113	013223010331022101
42	013232101321030231	114	013223010331022110
43	013232101321300213	115	013223010331202101
44	013232101321300231	116	013223010331202110
45	013232103121030213	117	013223013031022101
46	013232103121030231	118	013223013031022110
47	013232103121300213	119	013223013031202101
48	013232103121300231	120	013223013031202110
49	013230210321031023	121	013220130131022013
50	013230210321031032	122	013220130131022031
51	013230210321301023	123	013220130131202013
52	013230210321301032	124	013220130131202031
53	013230213021031023	125	013220131031022013
54	013230213021031032	126	013220131031022031
55	013230213021301023	127	013220131031202013
56	013230213021301032	128	013220131031202031
57	013232010321031203	129	013223101331020212
58	013232010321031230	130	013223101331020221
59	013232010321301203	131	013223101331200212
60	013232010321301230	132	013223101331200221
61	013232013021031203	133	013223103131020212
62	013232013021031230	134	013223103131020221
63	013232013021301203	135	013223103131200212
64	013232013021301230	136	013223103131200221
65	013212030132021203	137	013223101331022001
66	013212030132021230	138	013223101331022010
67	013212030132201203	139	013223101331202001
68	013212030132201230	140	013223101331202010
69	013212031032021203	141	013223103131022001
70	013212031032021230	142	013223103131022010
71	013212031032201203	143	013223103131202001
72	013212031032201230	144	013223103131202010

S2.24. Colorings of $D^{(4)}_{17,9}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24$. The anchor context $(0, 1, 2, 3)$ has colors $(0, \ldots, 3)$.

TABLE S68: Every normalized coloring of $D^{(4)}_{17,9}$.

j	Color word	j	Color word
1	01232132313123123	5	01233122312123132
2	01232312131123323	6	01231323123211213
3	01232313121122333	7	01233121321213231
4	01232313123122113		

S2.25. Colorings of $L^{(4)}_{22,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22$. The anchor context $(1, 2, 4, 3)$ has colors $(0, \ldots, 3)$.

TABLE S69: Every normalized coloring of $L^{(4)}_{22,9}$.

j	Color word	j	Color word
1	0132120301230312033201	109	0132120310320212032301
2	0132120301230312033210	110	0132120310320212032310
3	0132120301230312303201	111	0132120310320212302301
4	0132120301230312303210	112	0132120310320212302310
5	0132120310230312033201	113	0132120301322012030312
6	0132120310230312033210	114	0132120301322012030321
7	0132120310230312303201	115	0132120301322012300312
8	0132120310230312303210	116	0132120301322012300321
9	0132120301233012030213	117	0132120310322012030312
10	0132120301233012030231	118	0132120310322012030321
11	0132120301233012300213	119	0132120310322012300312
12	0132120301233012300231	120	0132120310322012300321
13	0132120310233012030213	121	0132132013322001121302
14	0132120310233012030231	122	0132132013322001121320
15	0132120310233012300213	123	0132132013322001211302
16	0132120310233012300231	124	0132132013322001211320
17	0132132013233001121203	125	0132132031322001121302
18	0132132013233001121230	126	0132132031322001121320
19	0132132013233001211203	127	0132132031322001211302
20	0132132013233001211230	128	0132132031322001211320
21	0132132031233001121203	129	0132102301320210232301
22	0132132031233001121230	130	0132102301320210232310
23	0132132031233001211203	131	0132102301320210322301
24	0132132031233001211230	132	0132102301320210322310
25	0132102301230310233201	133	0132102310320210232301
26	0132102301230310233210	134	0132102310320210232310
27	0132102301230310323201	135	0132102310320210322301
28	0132102301230310323210	136	0132102310320210322310
29	0132102310230310233201	137	0132102301322010230312
30	0132102310230310233210	138	0132102301322010230321
31	0132102310230310323201	139	0132102301322010320312
32	0132102310230310323210	140	0132102301322010320321

j	Color word	j	Color word
33	013210230l233010230213	141	013210231O322010230312
34	013210230l233010230231	142	013210231O322010230321
35	013210230l233010320213	143	013210231O322010320312
36	013210230l233010320231	144	013210231O322010320321
37	013210231O233010230213	145	013213201332021002230l
38	013210231O233010230231	146	013213201332021002231O
39	013210231O233010320213	147	013213201332021020230l
40	013210231O233010320231	148	013213201332021020231O
41	013213201323031002320l	149	013213203132021002230l
42	013213201323031002321O	150	013213203132021002231O
43	013213201323031020320l	151	013213203132021020230l
44	013213201323031020321O	152	013213203132021020231O
45	013213203123031002320l	153	013213201332201002031l
46	013213203123031002321O	154	013213201332201002032l
47	013213203123031020320l	155	013213201332201020031l
48	013213203123031020321O	156	013213201332201020032l
49	013213201323301002021l	157	013213203132201002031l
50	013213201323301002023l	158	013213203132201002032l
51	013213201323301020021l	159	013213203132201020031l
52	013213201323301020023l	160	013213203132201020032l
53	013213203123301002021l	161	013223010331021202230l
54	013213203123301002023l	162	013223010331021202231O
55	013213203123301020021l	163	013223010331021220230l
56	013213203123301020023l	164	013223010331021220231O
57	013231201321300123120l	165	013223013031021202230l
58	013231201321300123123O	166	013223013031021202231O
59	013231201321300132120l	167	013223013031021220230l
60	013231201321300132123O	168	013223013031021220231O
61	013231203121300123120l	169	013223010331201202031l
62	013231203121300123123O	170	013223010331201202032l
63	013231203121300132120l	171	013223010331201220031l
64	013231203121300132123O	172	013223010331201220032l
65	013232101321300213120l	173	013223013031201202031l
66	013232101321300213123O	174	013223013031201202032l
67	013232101321300231120l	175	013223013031201220031l
68	013232101321300231123O	176	013223013031201220032l
69	013232103121300213120l	177	013221030131022103130l
70	013232103121300213123O	178	013221030131022103132O
71	013232103121300231120l	179	013221030131022130130l
72	013232103121300231123O	180	013221030131022130132O
73	013230210321031023320l	181	013221031031022103130l
74	013230210321031023321O	182	013221031031022103132O
75	013230210321031032320l	183	013221031031022130130l
76	013230210321031032321O	184	013221031031022130132O
77	013230213021031023320l	185	013223010331022101130l
78	013230213021031023321O	186	013223010331022101132O
79	013230213021031032320l	187	013223010331022110130l
80	013230213021031032321O	188	013223010331022110132O
81	013230210321301023021l	189	013223013031022101130l
82	013230210321301023023l	190	013223013031022101132O
83	013230210321301032021l	191	013223013031022110130l

j	Color word	j	Color word
84	013230210321301032023 1	192	013223013031022110132 0
85	013230213021301023021 3	193	013223101331200212130 2
86	013230213021301023023 1	194	013223101331200212132 0
87	013230213021301032021 3	195	013223101331200221130 2
88	013230213021301032023 1	196	013223101331200221132 0
89	013232010321031203320 1	197	013223103131200212130 2
90	013232010321031203321 0	198	013223103131200212132 0
91	013232010321031230320 1	199	013223103131200221130 2
92	013232010321031230321 0	200	013223103131200221132 0
93	013232013021031203320 1	201	013220130131022013130 2
94	013232013021031203321 0	202	013220130131022013132 0
95	013232013021031230320 1	203	013220130131022031130 2
96	013232013021031230321 0	204	013220130131022031132 0
97	013232010321301203021 3	205	013220131031022013130 2
98	013232010321301203023 1	206	013220131031022013132 0
99	013232010321301230021 3	207	013220131031022031130 2
100	013232010321301230023 1	208	013220131031022031132 0
101	013232013021301203021 3	209	013223101331022001130 2
102	013232013021301203023 1	210	013223101331022001132 0
103	013232013021301230021 3	211	013223101331022010130 2
104	013232013021301230023 1	212	013223101331022010132 0
105	013212030132021203230 1	213	013223103131022001130 2
106	013212030132021203231 0	214	013223103131022001132 0
107	013212030132021230230 1	215	013223103131022010130 2
108	013212030132021230231 0	216	013223103131022010132 0

Exhaustive classification of chromatic identities. For a chromatically inseparable pair (u, v), form the two directed witness sets

$$W_{10}(u,v) = \{j : s_j(u) = 1,\ s_j(v) = 0\},$$
$$W_{01}(u,v) = \{j : s_j(u) = 0,\ s_j(v) = 1\}.$$

Both sets are nonempty exactly for a two-way purely chromatic identity. If only W_{10} is empty, $u \Rightarrow v$ holds and its converse fails; if only W_{01} is empty, the implication is reversed. If both are empty, the pair is inseparable under all ordinary states. Any member of either witness set is automatically nonextendable, since every color-class state has equal values on a chromatically inseparable pair.

The following Python standard-library procedure reconstructs $H_{21,13}$ directly from its 13 contexts in Table S3, enumerates all binary assignments, and reconstructs normalized strong colorings as partitions into three disjoint state supports. Lexicographic support order agrees with Table S42. It prints both complete directed witness sets for every chromatic identity, rather than only the representatives in Table S70.

TABLE S70. Both directed truth-pattern witnesses for all 15 two-way purely chromatic pairs of $H_{21,13}$ and both displayed pairs of $T_{19,11}$. For $H_{21,13}$, s_j is the state in Table S42; for $T_{19,11}$, entries give supports. Every witness is nonextendable and satisfies every context equation.

Configuration	Pair (u, v)	$(s(u), s(v)) = (1, 0)$	$(s(u), s(v)) = (0, 1)$
$T_{19,11}$	$(14, 28)$	$\{1, 5, 7, 14, 18, 23, 24\}$	$\{1, 6, 8, 10, 11, 27, 28\}$
$T_{19,11}$	$(18, 27)$	$\{1, 5, 7, 14, 18, 23, 24\}$	$\{1, 6, 8, 10, 11, 27, 28\}$
$H_{21,13}$	$(1, 11)$	s_1	s_{12}
$H_{21,13}$	$(1, 18)$	s_1	s_{11}
$H_{21,13}$	$(1, 21)$	s_2	s_9
$H_{21,13}$	$(2, 14)$	s_{12}	s_1
$H_{21,13}$	$(4, 13)$	s_1	s_{18}
$H_{21,13}$	$(4, 15)$	s_7	s_4
$H_{21,13}$	$(4, 19)$	s_1	s_{11}
$H_{21,13}$	$(5, 22)$	s_{18}	s_1
$H_{21,13}$	$(7, 10)$	s_2	s_{20}
$H_{21,13}$	$(7, 20)$	s_2	s_9
$H_{21,13}$	$(7, 24)$	s_7	s_4
$H_{21,13}$	$(8, 23)$	s_{20}	s_2
$H_{21,13}$	$(15, 19)$	s_1	s_7
$H_{21,13}$	$(18, 21)$	s_2	s_1
$H_{21,13}$	$(20, 24)$	s_7	s_2

```
from itertools import combinations
E = [(1,2,3),(7,8,9),(4,5,6),(2,8,5),
     (3,11,14),(7,11,15),(4,11,24),
     (9,10,23),(1,10,19),(4,10,18),
     (1,13,20),(7,13,21),(6,13,22)]
V = sorted(set().union(*map(set, E)))
M = [sum(1 << V.index(v) for v in e) for e in E]
S = sorted(tuple(v for i,v in enumerate(V) if b >> i & 1)
           for b in range(1 << len(V))
           if all((b & m).bit_count() == 1 for m in M))
S = list(map(frozenset, S))
C = []
for blocks in combinations(S, 3):
    if any(a & b for a,b in combinations(blocks, 2)):
        continue
    blocks = [next(s for s in blocks if a in s) for a in (2,8,5)]
    C.append({v:k for k,s in enumerate(blocks) for v in s})
assert (len(S), len(C)) == (21, 2)
counts = [0, 0, 0]
for u,v in combinations(V, 2):
    if not all(c[u] == c[v] for c in C):
        continue
    W10 = [j for j,s in enumerate(S, 1) if u in s and v not in s]
    W01 = [j for j,s in enumerate(S, 1) if v in s and u not in s]
    k = 0 if W10 and W01 else 1 if W10 or W01 else 2
    counts[k] += 1
    print((u,v), W10, W01)
assert counts == [15, 6, 0]
```

TABLE S71: All ten connected, edge-faithful, colorable restrictions with a forced color identity in the specified inventory.

retained diagonals	rays	contexts	colorings	inseparable pairs
$\{10,11\}$	17	10	4	$(10,24)$, $(11,18)$
$\{10,12\}$	17	10	4	$(10,16)$, $(12,23)$
$\{10,13\}$	17	10	4	$(10,20)$, $(13,19)$
$\{11,12\}$	17	10	4	$(11,25)$, $(12,14)$
$\{11,13\}$	17	10	4	$(11,21)$, $(13,15)$
$\{12,13\}$	17	10	4	$(12,22)$, $(13,17)$
$\{10,11,12\}$	21	13	2	21 pairs
$\{10,11,13\}$	21	13	2	21 pairs
$\{10,12,13\}$	21	13	2	21 pairs
$\{11,12,13\}$	21	13	2	21 pairs

S2.26. An explicit state certificate for the completed Yu–Oh hypergraph

The first four contexts of Eq. (15) force exactly one of vertices 2, 5, and 8 to have value one. In the triad containing that coordinate vertex, both other vertices have value zero. In each of the other two triads, one of the two remaining vertices has value one. There are therefore $3 \cdot 2^2 = 12$ possible supports T_k on vertices $1, \ldots, 9$, listed in Table S72.

For each such support, exactly one of vertices $10, 11, 12, 13$ has no orthogonal neighbor in T_k; call it η_k. The other three are forced to zero, while $s(\eta_k)$ may be either zero or one. This gives precisely 24 assignments on the original 13 vertices. Every completion vertex w belongs to one context $\{i, j, w\}$ and has the uniquely determined value

$$s(w) = 1 - s(i) - s(j). \tag{S7}$$

The listed assignments have $s(i) + s(j) \leq 1$ on every original orthogonal pair, so these extensions are valid two-valued states. Conversely, every state of the completed hypergraph restricts to one of these assignments.

For a compact separation certificate, encode the profile of vertex i as the nonnegative integer

$$m_i = \sum_{k=1}^{12} \sum_{b=0}^{1} s_{k,b}(i)\, 2^{2(k-1)+b}.$$

TABLE S72. The 24 states are indexed by (k, b), where $1 \leq k \leq 12$ and $b \in \{0, 1\}$. Their support on vertices $1, \ldots, 13$ is T_k when $b = 0$ and $T_k \cup \{\eta_k\}$ when $b = 1$. Equation (S7) determines all remaining values.

k	T_k	η_k
1	$\{1,4,8\}$	12
2	$\{1,5,7\}$	12
3	$\{1,5,9\}$	11
4	$\{1,6,8\}$	11
5	$\{2,4,7\}$	12
6	$\{2,4,9\}$	13
7	$\{2,6,7\}$	10
8	$\{2,6,9\}$	11
9	$\{3,4,8\}$	13
10	$\{3,5,7\}$	10
11	$\{3,5,9\}$	13
12	$\{3,6,8\}$	10

In hexadecimal notation these 25 integers are

i	1	2	3	4	5
m_i	0000FF	00FF00	FF0000	030F03	3C003C

i	6	7	8	9	10
m_i	C0F0C0	0C330C	C300C3	30CC30	882000

i	11	12	13	14	15
m_i	0080A0	00020A	220800	007F5F	F34C53

i	16	17	18	19	20
m_i	CF31C5	3F0D35	74D0FC	77DF00	DDF700

i	21	22	23	24	25
m_i	D1C4F3	1D073F	4713CF	FC705C	00FDF5

(S8)

All 25 integers are nonzero and pairwise distinct, certifying unitality and separation. Notably, the Hamming weights (number of states in which $s(i) = 1$) partition the vertices into three classes: 8 for $i \in \{1, \ldots, 9\}$, 3 for $i \in \{10, \ldots, 13\}$, and 13 for the completion vertices $i \in \{14, \ldots, 25\}$ (with $8 + 3 + 13 = 24$ in each completion context), in contrast to the uniform state weight 2 across all 15 vertices of G_{32} and consistent with the fractional chromatic–clique gap $\chi_f(\Gamma(H_{\mathrm{YO}}^+)) = 35/11 > 3 = \omega(\Gamma(H_{\mathrm{YO}}^+))$.

S2.27. Hesse H_{37}: all states and separating profiles

The canonical states $s_1, \ldots, s_{153}$ are listed in Table S74. A two-valued coloring is a state with exactly one value one per context. Vertex numbers are those of Appendix S1.14 and Fig. 4: vertices $1, \ldots, 16$ form the core, and $17, \ldots, 37$ are completions. We list every state, a minimum separating family, and every vertex profile.

S2.27.1. State parametrization and all 27 branches

Choose $t = (t_0, t_1, t_2) \in \mathbb{Z}_3^3$ and $\varepsilon \in \{0, 1\}$. Put

$$s(M_{ak}) = \mathbf{1}_{k=t_a}, \qquad s(z) = \varepsilon, \tag{S9}$$

$$U(t) = \mathbb{Z}_3 \setminus \{t_0, t_1, t_2\}, \qquad W(t) = \mathbb{Z}_3 \setminus \{t_a - a : a \in \mathbb{Z}_3\}. \tag{S10}$$

For $\varepsilon = 0$, choose the values of P_k freely for $k \in U(t)$, with other P values zero. For $\varepsilon = 1$, all P values are zero. Independently, choose the values of Q_k freely for $k \in W(t)$, with other Q values zero. All completion values are forced:

$$s(A_k) = 1 - \varepsilon - s(P_k), \tag{S11}$$

$$s(B_{ak}) = 1 - s(M_{ak}) - s(P_k), \tag{S12}$$

$$s(C_{ak}) = 1 - s(M_{a,a+k}) - s(Q_k). \tag{S13}$$

Enumerate t lexicographically with $0 < 1 < 2$, then $\varepsilon = 1, 0$. Within each branch, list the available P bits in increasing k, followed by the available Q bits in increasing k, and order the resulting bit strings with 1 before 0. This rule fixes all state indices used in the text, tables, and verification data. Table S73 gives the index ranges of all 27 branches.

TABLE S73: All 27 state branches. The sets $U(t), W(t)$ identify the free core bits. The two choices of ε give $N_0 = 2^{|U|+|W|}$ and $N_1 = 2^{|W|}$ states; their index ranges refer to Table S74.

t	$U(t)$	$W(t)$	N_0	r for $\varepsilon = 0$	N_1	r for $\varepsilon = 1$
$(0,0,0)$	$\{1,2\}$	$\varnothing$	4	2–5	1	1
$(0,0,1)$	$\{2\}$	$\{1\}$	4	8–11	2	6–7
$(0,0,2)$	$\{1\}$	$\{1\}$	4	14–17	2	12–13
$(0,1,0)$	$\{2\}$	$\{2\}$	4	20–23	2	18–19
$(0,1,1)$	$\{2\}$	$\{1\}$	4	26–29	2	24–25
$(0,1,2)$	$\varnothing$	$\{1,2\}$	4	34–37	4	30–33
$(0,2,0)$	$\{1\}$	$\{2\}$	4	40–43	2	38–39
$(0,2,1)$	$\varnothing$	$\varnothing$	1	45	1	44
$(0,2,2)$	$\{1\}$	$\{2\}$	4	48–51	2	46–47
$(1,0,0)$	$\{2\}$	$\{0\}$	4	54–57	2	52–53
$(1,0,1)$	$\{2\}$	$\{0\}$	4	60–63	2	58–59
$(1,0,2)$	$\varnothing$	$\varnothing$	1	65	1	64
$(1,1,0)$	$\{2\}$	$\{2\}$	4	68–71	2	66–67
$(1,1,1)$	$\{0,2\}$	$\varnothing$	4	73–76	1	72
$(1,1,2)$	$\{0\}$	$\{2\}$	4	79–82	2	77–78
$(1,2,0)$	$\varnothing$	$\{0,2\}$	4	87–90	4	83–86
$(1,2,1)$	$\{0\}$	$\{0\}$	4	93–96	2	91–92
$(1,2,2)$	$\{0\}$	$\{2\}$	4	99–102	2	97–98
$(2,0,0)$	$\{1\}$	$\{0\}$	4	105–108	2	103–104
$(2,0,1)$	$\varnothing$	$\{0,1\}$	4	113–116	4	109–112
$(2,0,2)$	$\{1\}$	$\{1\}$	4	119–122	2	117–118
$(2,1,0)$	$\varnothing$	$\varnothing$	1	124	1	123
$(2,1,1)$	$\{0\}$	$\{1\}$	4	127–130	2	125–126
$(2,1,2)$	$\{0\}$	$\{1\}$	4	133–136	2	131–132
$(2,2,0)$	$\{1\}$	$\{0\}$	4	139–142	2	137–138
$(2,2,1)$	$\{0\}$	$\{0\}$	4	145–148	2	143–144
$(2,2,2)$	$\{0,1\}$	$\varnothing$	4	150–153	1	149
Total			99		54	

S2.27.2. Canonical two-valued colorings: all 153 states

Table S74 explicitly lists all 153 states. The last column is the complete binary vector in vertex order $1,\dots,37$, separated after the 16 core entries. Every row has context sum one in all 24 contexts. The support-size distribution is

$$\begin{array}{c|cccccc} m & 12 & 13 & 14 & 15 & 16 & 18 \\ \#\{s : |\operatorname{supp} s| = m\} & 6 & 18 & 27 & 24 & 51 & 27 \end{array} \tag{S14}$$

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1),\dots,s_r(16)\,;\, s_r(17),\dots,s_r(37))$.

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
1	$\{1,2,5,8\}$	$\{21,22,24,25,27,28,30,31,32,33,35,37\}$	16	1100100100000000 00001101101101110101
2	$\{2,5,8,12,13\}$	$\{17,30,31,32,33,35,37\}$	12	0100100100011000 10000000000011110101
3	$\{2,5,8,12\}$	$\{17,19,22,25,28,30,31,32,33,35,37\}$	15	0100100100010000 10100100100101110101
4	$\{2,5,8,13\}$	$\{17,18,21,24,27,30,31,32,33,35,37\}$	15	0100100100001000 11001001001001110101
5	$\{2,5,8\}$	$\{17,18,19,21,22,24,25,27,28,30,31,32,33,35,37\}$	18	0100100100000000 11101101101101110101
6	$\{1,2,5,9,15\}$	$\{21,22,24,25,26,28,31,32,35\}$	14	1100100010000010 00001101110100110010 0
7	$\{1,2,5,9\}$	$\{21,22,24,25,26,28,30,31,32,33,35,36\}$	16	1100100010000000 00001101110101110110
8	$\{2,5,9,13,15\}$	$\{17,18,21,24,26,31,32,35\}$	13	0100100010001010 11001001010000110010 0
9	$\{2,5,9,13\}$	$\{17,18,21,24,26,30,31,32,33,35,36\}$	15	0100100010001000 11001001010001110110
10	$\{2,5,9,15\}$	$\{17,18,19,21,22,24,25,26,28,31,32,35\}$	16	0100100010000010 11101101110100110010 0
11	$\{2,5,9\}$	$\{17,18,19,21,22,24,25,26,28,30,31,32,33,35,36\}$	18	0100100010000000 11101101110101110110
12	$\{1,2,5,10,15\}$	$\{21,22,24,25,26,27,31,32,37\}$	14	1100100001000010 00001101111000110000 1
13	$\{1,2,5,10\}$	$\{21,22,24,25,26,27,30,31,32,33,36,37\}$	16	1100100001000000 00001101111001110011
14	$\{2,5,10,12,15\}$	$\{17,19,22,25,26,31,32,37\}$	13	0100100001010010 10100100110000110000 1
15	$\{2,5,10,12\}$	$\{17,19,22,25,26,30,31,32,33,36,37\}$	15	0100100001010000 10100100110000111001 1
16	$\{2,5,10,15\}$	$\{17,18,19,21,22,24,25,26,27,31,32,37\}$	16	0100100001000010 11101101111000110000 1
17	$\{2,5,10\}$	$\{17,18,19,21,22,24,25,26,27,30,31,32,33,36,37\}$	18	0100100001000000 11101101111001110011
18	$\{1,2,6,8,16\}$	$\{21,22,23,25,27,28,30,33,35\}$	14	1100010100000001 00001110101101001010 0

Continued on the next page.

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
19	$\{1,2,6,8\}$	$\{21,22,23,25,27,28,30,31,33,34,35,37\}$	16	1100010100000000 00001110101101101101
20	$\{2,6,8,13,16\}$	$\{17,18,21,23,27,30,33,35\}$	13	0100010100001001 11001010001001010100
21	$\{2,6,8,13\}$	$\{17,18,21,23,27,30,31,33,34,35,37\}$	15	0100010100001000 11001010001011011101
22	$\{2,6,8,16\}$	$\{17,18,19,21,22,23,25,27,28,30,33,35\}$	16	0100010100000001 11101110101101010100
23	$\{2,6,8\}$	$\{17,18,19,21,22,23,25,27,28,30,31,33,34,35,37\}$	18	0100010100000000 11101110101101101101
24	$\{1,2,6,9,15\}$	$\{21,22,23,25,26,28,31,34,35\}$	14	1100010010000010 00001110110100100100
25	$\{1,2,6,9\}$	$\{21,22,23,25,26,28,30,31,33,34,35,36\}$	16	1100010010000000 00001110110101101110
26	$\{2,6,9,13,15\}$	$\{17,18,21,23,26,31,34,35\}$	13	0100010010001010 11001010010000100100
27	$\{2,6,9,13\}$	$\{17,18,21,23,26,30,31,33,34,35,36\}$	15	0100010010001000 11001010010001101110
28	$\{2,6,9,15\}$	$\{17,18,19,21,22,23,25,26,28,31,34,35\}$	16	0100010010000010 11101110110100100100
29	$\{2,6,9\}$	$\{17,18,19,21,22,23,25,26,28,30,31,33,34,35,36\}$	18	0100010010000000 11101110110101101110
30	$\{1,2,6,10,15,16\}$	$\{21,22,23,25,26,27\}$	12	1100010001000011 00001110111000000000
31	$\{1,2,6,10,15\}$	$\{21,22,23,25,26,27,31,34,37\}$	14	1100010001000010 00001110111000100100
32	$\{1,2,6,10,16\}$	$\{21,22,23,25,26,27,30,33,36\}$	14	1100010001000001 00001110111001001010
33	$\{1,2,6,10\}$	$\{21,22,23,25,26,27,30,31,33,34,36,37\}$	16	1100010001000000 00001110111001101011
34	$\{2,6,10,15,16\}$	$\{17,18,19,21,22,23,25,26,27\}$	14	0100010001000011 11101110111000000000
35	$\{2,6,10,15\}$	$\{17,18,19,21,22,23,25,26,27,31,34,37\}$	16	0100010001000010 11101110111000100100
36	$\{2,6,10,16\}$	$\{17,18,19,21,22,23,25,26,27,30,33,36\}$	16	0100010001000001 11101110111001001010
37	$\{2,6,10\}$	$\{17,18,19,21,22,23,25,26,27,30,31,33,34,36,37\}$	18	0100010001000000 11101110111001101011
38	$\{1,2,7,8,16\}$	$\{21,22,23,24,27,28,30,32,35\}$	14	1100001100000001 00001111001101010100
39	$\{1,2,7,8\}$	$\{21,22,23,24,27,28,30,31,32,34,35,37\}$	16	1100001100000000 00001111001101101101
40	$\{2,7,8,12,16\}$	$\{17,19,22,23,28,30,32,35\}$	13	0100001100010001 10100110000101010100
41	$\{2,7,8,12\}$	$\{17,19,22,23,28,30,31,32,34,35,37\}$	15	0100001100010000 10100110000101101101
42	$\{2,7,8,16\}$	$\{17,18,19,21,22,23,24,27,28,30,32,35\}$	16	0100001100000001 11101110011010100100
43	$\{2,7,8\}$	$\{17,18,19,21,22,23,24,27,28,30,31,32,34,35,37\}$	18	0100001100000000 11101110011011101101
44	$\{1,2,7,9\}$	$\{21,22,23,24,26,28,30,31,32,34,35,36\}$	16	1100001010000000 00001111010101101110
45	$\{2,7,9\}$	$\{17,18,19,21,22,23,24,26,28,30,31,32,34,35,36\}$	18	0100001010000000 11101111010101101110
46	$\{1,2,7,10,16\}$	$\{21,22,23,24,26,27,30,32,36\}$	14	1100001001000001 00001111011001010010
47	$\{1,2,7,10\}$	$\{21,22,23,24,26,27,30,31,32,34,36,37\}$	16	1100001001000000 00001111011001101011
48	$\{2,7,10,12,16\}$	$\{17,19,22,23,26,30,32,36\}$	13	0100001001010001 10100110010010100010
49	$\{2,7,10,12\}$	$\{17,19,22,23,26,30,31,32,34,36,37\}$	15	0100001001010000 10100110010011101011
50	$\{2,7,10,16\}$	$\{17,18,19,21,22,23,24,26,27,30,32,36\}$	16	0100001001000001 11101111010010100010
51	$\{2,7,10\}$	$\{17,18,19,21,22,23,24,26,27,30,31,32,34,36,37\}$	18	0100001001000000 11101111010011101011
52	$\{1,3,5,8,14\}$	$\{20,22,24,25,27,28,31,33,37\}$	14	1010100100000100 00010101101001010001
53	$\{1,3,5,8\}$	$\{20,22,24,25,27,28,29,31,32,33,35,37\}$	16	1010100100000000 00010101101110110101
54	$\{3,5,8,13,14\}$	$\{17,18,20,24,27,31,33,37\}$	13	0010100100001100 11010001001001010001
55	$\{3,5,8,13\}$	$\{17,18,20,24,27,29,31,32,33,35,37\}$	15	0010100100001000 11010001001010110101
56	$\{3,5,8,14\}$	$\{17,18,19,20,22,24,25,27,28,31,33,37\}$	16	0010100100000100 11110101101001010001
57	$\{3,5,8\}$	$\{17,18,19,20,22,24,25,27,28,29,31,32,33,35,37\}$	18	0010100100000000 11110101101101110101
58	$\{1,3,5,9,14\}$	$\{20,22,24,25,26,28,31,33,36\}$	14	1010100010000100 00010101101001010010
59	$\{1,3,5,9\}$	$\{20,22,24,25,26,28,29,31,32,33,35,36\}$	16	1010100010000000 00010101101101110110
60	$\{3,5,9,13,14\}$	$\{17,18,20,24,26,31,33,36\}$	13	0010100010001100 11010001010000101010
61	$\{3,5,9,13\}$	$\{17,18,20,24,26,29,31,32,33,35,36\}$	15	0010100010001000 11010001010010110110
62	$\{3,5,9,14\}$	$\{17,18,19,20,22,24,25,26,28,31,33,36\}$	16	0010100010000100 11110101101001010010
63	$\{3,5,9\}$	$\{17,18,19,20,22,24,25,26,28,29,31,32,33,35,36\}$	18	0010100010000000 11110101101101110110
64	$\{1,3,5,10\}$	$\{20,22,24,25,26,27,29,31,32,33,36,37\}$	16	1010100001000000 00010101110101110011
65	$\{3,5,10\}$	$\{17,18,19,20,22,24,25,26,27,29,31,32,33,36,37\}$	18	0010100001000000 11110101110101110011
66	$\{1,3,6,8,16\}$	$\{20,22,23,25,27,28,29,33,35\}$	14	1010010100000001 00010110101110010100
67	$\{1,3,6,8\}$	$\{20,22,23,25,27,28,29,31,33,34,35,37\}$	16	1010010100000000 00010110101101011101
68	$\{3,6,8,13,16\}$	$\{17,18,20,23,27,29,33,35\}$	13	0010010100001001 11010010001010010100
69	$\{3,6,8,13\}$	$\{17,18,20,23,27,29,31,33,34,35,37\}$	15	0010010100001000 11010010001010110101
70	$\{3,6,8,16\}$	$\{17,18,19,20,22,23,25,27,28,29,33,35\}$	16	0010010100000001 11110110101110010100
71	$\{3,6,8\}$	$\{17,18,19,20,22,23,25,27,28,29,31,33,34,35,37\}$	18	0010010100000000 11110110101101011101
72	$\{1,3,6,9\}$	$\{20,22,23,25,26,28,29,31,33,34,35,36\}$	16	1010010010000000 00010110110110101110
73	$\{3,6,9,11,13\}$	$\{18,29,31,33,34,35,36\}$	12	0010010010101000 01000000000010101110
74	$\{3,6,9,11\}$	$\{18,19,22,25,28,29,31,33,34,35,36\}$	15	0010010010100000 01100100100110101110
75	$\{3,6,9,13\}$	$\{17,18,20,23,26,29,31,33,34,35,36\}$	15	0010010010001000 11010010010010110110
76	$\{3,6,9\}$	$\{17,18,19,20,22,23,25,26,28,29,31,33,34,35,36\}$	18	0010010010000000 11110110110110101110
77	$\{1,3,6,10,16\}$	$\{20,22,23,25,26,27,29,33,36\}$	14	1010010001000001 00010110110100010010
78	$\{1,3,6,10\}$	$\{20,22,23,25,26,27,29,31,33,34,36,37\}$	16	1010010001000000 00010110110101011011
79	$\{3,6,10,11,16\}$	$\{18,19,22,25,27,29,33,36\}$	13	0010010001100001 01100100101010010010
80	$\{3,6,10,11\}$	$\{18,19,22,25,27,29,31,33,34,36,37\}$	15	0010010001100000 01100100101010101011
81	$\{3,6,10,16\}$	$\{17,18,19,20,22,23,25,26,27,29,33,36\}$	16	0010010001000001 11110110111010010010
82	$\{3,6,10\}$	$\{17,18,19,20,22,23,25,26,27,29,31,33,34,36,37\}$	18	0010010001000000 11110110111010101011
83	$\{1,3,7,8,14,16\}$	$\{20,22,23,24,27,28\}$	12	1010001100000101 00010111001100000000
84	$\{1,3,7,8,14\}$	$\{20,22,23,24,27,28,31,34,37\}$	14	1010001100000100 00010111001101001001

Continued on the next page.

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
85	$\{1,3,7,8,16\}$	$\{20,22,23,24,27,28,29,32,35\}$	14	10100011000000001 00010111001110010 0100
86	$\{1,3,7,8\}$	$\{20,22,23,24,27,28,29,31,32,34,35,37\}$	16	10100011000000000 00010111001110110 1101
87	$\{3,7,8,14,16\}$	$\{17,18,19,20,22,23,24,27,28\}$	14	00100011000001010 11110111001100000 0000
88	$\{3,7,8,14\}$	$\{17,18,19,20,22,23,24,27,28,31,34,37\}$	16	00100011000001000 11110111001100100 1001
89	$\{3,7,8,16\}$	$\{17,18,19,20,22,23,24,27,28,29,32,35\}$	16	00100011000000010 11110111001110010 0100
90	$\{3,7,8\}$	$\{17,18,19,20,22,23,24,27,28,29,31,32,34,35,37\}$	18	00100011000000000 11110111001110110 1101
91	$\{1,3,7,9,14\}$	$\{20,22,23,24,26,28,31,34,36\}$	14	10100010100000100 00010111010100100 1010
92	$\{1,3,7,9\}$	$\{20,22,23,24,26,28,29,31,32,34,35,36\}$	16	10100010100000000 00010111010110110 1110
93	$\{3,7,9,11,14\}$	$\{18,19,22,24,28,31,34,36\}$	13	00100010101001000 11001010001001001 0010
94	$\{3,7,9,11\}$	$\{18,19,22,24,28,29,31,32,34,35,36\}$	15	00100010101000000 11001010001101101 1110
95	$\{3,7,9,14\}$	$\{17,18,19,20,22,23,24,26,28,31,34,36\}$	16	00100010100000100 11110111010100100 1010
96	$\{3,7,9\}$	$\{17,18,19,20,22,23,24,26,28,29,31,32,34,35,36\}$	18	00100010100000000 11110111010110110 1110
97	$\{1,3,7,10,16\}$	$\{20,22,23,24,26,27,29,32,36\}$	14	10100010010000001 00010111011010010 0010
98	$\{1,3,7,10\}$	$\{20,22,23,24,26,27,29,31,32,34,36,37\}$	16	10100010010000000 00010111011010110 1011
99	$\{3,7,10,11,16\}$	$\{18,19,22,24,27,29,32,36\}$	13	00100010011000010 11001010010100100 0010
100	$\{3,7,10,11\}$	$\{18,19,22,24,27,29,31,32,34,36,37\}$	15	00100010011000000 11001010010101101 1011
101	$\{3,7,10,16\}$	$\{17,18,19,20,22,23,24,26,27,29,32,36\}$	16	00100010010000001 11110111011010010 0010
102	$\{3,7,10\}$	$\{17,18,19,20,22,23,24,26,27,29,31,32,34,36,37\}$	18	00100010010000000 11110111011010110 1011
103	$\{1,4,5,8,14\}$	$\{20,21,24,25,27,28,30,33,37\}$	14	10011001000000100 00011001101101001 0001
104	$\{1,4,5,8\}$	$\{20,21,24,25,27,28,29,30,32,33,35,37\}$	16	10011001000000000 00011001101111011 0101
105	$\{4,5,8,12,14\}$	$\{17,19,20,25,28,30,33,37\}$	13	00011001000101001 01100001001010010 0001
106	$\{4,5,8,12\}$	$\{17,19,20,25,28,29,30,32,33,35,37\}$	15	00011001000100000 10110000100111011 0101
107	$\{4,5,8,14\}$	$\{17,18,19,20,21,24,25,27,28,30,33,37\}$	16	00011001000001000 11111001101101001 0001
108	$\{4,5,8\}$	$\{17,18,19,20,21,24,25,27,28,29,30,32,33,35,37\}$	18	00011001000000000 11111001101111011 0101
109	$\{1,4,5,9,14,15\}$	$\{20,21,24,25,26,28\}$	12	10011000100001100 00011001101000000 0000
110	$\{1,4,5,9,14\}$	$\{20,21,24,25,26,28,30,33,36\}$	14	10011000100001000 00011001101010010 0010
111	$\{1,4,5,9,15\}$	$\{20,21,24,25,26,28,29,32,35\}$	14	10011000100000010 00011001101100100 0100
112	$\{1,4,5,9\}$	$\{20,21,24,25,26,28,29,30,32,33,35,36\}$	16	10011000100000000 00011001101110110 0110
113	$\{4,5,9,14,15\}$	$\{17,18,19,20,21,24,25,26,28\}$	14	00011000100001100 11111001101000000 0000
114	$\{4,5,9,14\}$	$\{17,18,19,20,21,24,25,26,28,30,33,36\}$	16	00011000100001000 11111001101010010 0010
115	$\{4,5,9,15\}$	$\{17,18,19,20,21,24,25,26,28,29,32,35\}$	16	00011000100000010 11111001101100100 0100
116	$\{4,5,9\}$	$\{17,18,19,20,21,24,25,26,28,29,30,32,33,35,36\}$	18	00011000100000000 11111001101110110 0110
117	$\{1,4,5,10,15\}$	$\{20,21,24,25,26,27,29,32,37\}$	14	10011000010000100 00011001111010010 0001
118	$\{1,4,5,10\}$	$\{20,21,24,25,26,27,29,30,32,33,36,37\}$	16	10011000010000000 00011001111011011 0011
119	$\{4,5,10,12,15\}$	$\{17,19,20,25,26,29,32,37\}$	13	00011000010100100 10110000110010010 0001
120	$\{4,5,10,12\}$	$\{17,19,20,25,26,29,30,32,33,36,37\}$	15	00011000010100000 10110000110011011 0011
121	$\{4,5,10,15\}$	$\{17,18,19,20,21,24,25,26,27,29,32,37\}$	16	00011000010000100 11111001111010010 0001
122	$\{4,5,10\}$	$\{17,18,19,20,21,24,25,26,27,29,30,32,33,36,37\}$	18	00011000010000000 11111001111011011 0011
123	$\{1,4,6,8\}$	$\{20,21,23,25,27,28,29,30,33,34,35,37\}$	16	10010101000000000 00011010101111001 1101
124	$\{4,6,8\}$	$\{17,18,19,20,21,23,25,27,28,29,30,33,34,35,37\}$	18	00010101000000000 11111010101111001 1101
125	$\{1,4,6,9,15\}$	$\{20,21,23,25,26,28,29,34,35\}$	14	10010100100000100 00011010111001100 0100
126	$\{1,4,6,9\}$	$\{20,21,23,25,26,28,29,30,33,34,35,36\}$	16	10010100100000000 00011010111001110 0110
127	$\{4,6,9,11,15\}$	$\{18,19,21,25,28,29,34,35\}$	13	00010100101000100 01101000100110000 1100
128	$\{4,6,9,11\}$	$\{18,19,21,25,28,29,30,33,34,35,36\}$	15	00010100101000000 01101000100111001 1110
129	$\{4,6,9,15\}$	$\{17,18,19,20,21,23,25,26,28,29,34,35\}$	16	00010100100000010 11111010110110000 1100
130	$\{4,6,9\}$	$\{17,18,19,20,21,23,25,26,28,29,30,33,34,35,36\}$	18	00010100100000000 11111010110111001 1110
131	$\{1,4,6,10,15\}$	$\{20,21,23,25,26,27,29,34,37\}$	14	10010100010000100 00011010111010000 1001
132	$\{1,4,6,10\}$	$\{20,21,23,25,26,27,29,30,33,34,36,37\}$	16	10010100010000000 00011010111011001 1011
133	$\{4,6,10,11,15\}$	$\{18,19,21,25,27,29,34,37\}$	13	00010100011000100 01101000101010000 1001
134	$\{4,6,10,11\}$	$\{18,19,21,25,27,29,30,33,34,36,37\}$	15	00010100011000000 01101000101011001 1011
135	$\{4,6,10,15\}$	$\{17,18,19,20,21,23,25,26,27,29,34,37\}$	16	00010100010000100 11111010111010000 1001
136	$\{4,6,10\}$	$\{17,18,19,20,21,23,25,26,27,29,30,33,34,36,37\}$	18	00010100010000000 11111010111011001 1011
137	$\{1,4,7,8,14\}$	$\{20,21,23,24,27,28,30,34,37\}$	14	10010011000000100 00011011001101000 1001
138	$\{1,4,7,8\}$	$\{20,21,23,24,27,28,29,30,32,34,35,37\}$	16	10010011000000000 00011011001111010 1101
139	$\{4,7,8,12,14\}$	$\{17,19,20,23,28,30,34,37\}$	13	00010011000101001 01100100001010001 1001
140	$\{4,7,8,12\}$	$\{17,19,20,23,28,29,30,32,34,35,37\}$	15	00010011000100000 10110010000111010 1101
141	$\{4,7,8,14\}$	$\{17,18,19,20,21,23,24,27,28,30,34,37\}$	16	00010011000001000 11111011001101000 1001
142	$\{4,7,8\}$	$\{17,18,19,20,21,23,24,27,28,29,30,32,34,35,37\}$	18	00010011000000000 11111011001111010 1101
143	$\{1,4,7,9,14\}$	$\{20,21,23,24,26,28,30,34,36\}$	14	10010010100001000 00011011010101000 1010
144	$\{1,4,7,9\}$	$\{20,21,23,24,26,28,29,30,32,34,35,36\}$	16	10010010100000000 00011011010111010 1110
145	$\{4,7,9,11,14\}$	$\{18,19,21,24,28,30,34,36\}$	13	00010010101001000 11010010000101000 1010

Continued on the next page.

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
146	$\{4,7,9,11\}$	$\{18,19,21,24,28,29,30,32,34,35,36\}$	15	0001001010100000 01101001000111010 1110
147	$\{4,7,9,14\}$	$\{17,18,19,20,21,23,24,26,28,30,34,36\}$	16	0001001010000100 11111011010101000 1010
148	$\{4,7,9\}$	$\{17,18,19,20,21,23,24,26,28,29,30,32,34,35,36\}$	18	0001001010000000 11111011010111010 1110
149	$\{1,4,7,10\}$	$\{20,21,23,24,26,27,29,30,32,34,36,37\}$	16	1001001001000000 00011011011011010 1011
150	$\{4,7,10,11,12\}$	$\{19,29,30,32,34,36,37\}$	12	0001001001110000 00100000000011010 1011
151	$\{4,7,10,11\}$	$\{18,19,21,24,27,29,30,32,34,36,37\}$	15	0001001001100000 01101001001011010 1011
152	$\{4,7,10,12\}$	$\{17,19,20,23,26,29,30,32,34,36,37\}$	15	0001001001010000 10110010010011010 1011
153	$\{4,7,10\}$	$\{17,18,19,20,21,23,24,26,27,29,30,32,34,36,37\}$	18	0001001001000000 11111011011011010 1011

S2.27.3. A minimum six-state unital, separating family

Define $\Omega_6 = \{\sigma_1, \ldots, \sigma_6\}$ in the following order:

$$(\sigma_1, \ldots, \sigma_6) = (s_3, s_{31}, s_{75}, s_{89}, s_{113}, s_{151}). \tag{S15}$$

Table S75 prints their complete supports. Table S76 gives all 37 profiles $(\sigma_1(i), \ldots, \sigma_6(i))$ and their active state sets. Every profile is nonzero and no two coincide, so Ω_6 is unital and separating. Since five states supply at most $2^5 - 1 = 31 < 37$ nonzero binary profiles, no smaller unital, separating family can exist.

TABLE S75: Complete supports and state vectors of the minimum unital, separating family Ω_6. The full state indices are those of Table S74.

State	Full index	Complete support	Weight	37-bit vector
σ_1	s_3	$\{2,5,8,12,17,19,22,25,28,30,31,32,33,35,37\}$	15	0100100100010000 10100100100101111 0101
σ_2	s_{31}	$\{1,2,6,10,15,21,22,23,25,26,27,31,34,37\}$	14	1100010001000010 00001110111000100 1001
σ_3	s_{75}	$\{3,6,9,13,17,18,20,23,26,29,31,33,34,35,36\}$	15	0010010010001000 11010010010010111 1110
σ_4	s_{89}	$\{3,7,8,16,17,18,19,20,22,23,24,27,28,29,32,35\}$	16	0010001100000001 11110111001110010 0100
σ_5	s_{113}	$\{4,5,9,14,15,17,18,19,20,21,24,25,26,28\}$	14	0001100010000110 11111001110100000 0000
σ_6	s_{151}	$\{4,7,10,11,18,19,21,24,27,29,30,32,34,36,37\}$	15	0001001001100000 01101001001011010 1011

TABLE S76: All 37 vertex profiles on Ω_6, in two panels. Here $X_i^{(6)} = \{j : \sigma_j(i) = 1\}$, $W_6 = |X_i^{(6)}|$, and W_{153} is the weight across the full state space. The six-bit rows are nonzero and pairwise distinct.

i	Ray	$X_i^{(6)}$	Profile	W_6	W_{153}	i	Ray	$X_i^{(6)}$	Profile	W_6	W_{153}
1	z	$\{2\}$	010000	1	54	20	B_{00}	$\{3,4,5\}$	001110	3	86
2	M_{00}	$\{1,2\}$	110000	2	51	21	B_{01}	$\{2,5,6\}$	010011	3	86
3	M_{01}	$\{3,4\}$	001100	2	51	22	B_{02}	$\{1,2,4\}$	110100	3	86
4	M_{02}	$\{5,6\}$	000011	2	51	23	B_{10}	$\{2,3,4\}$	011100	3	86
5	M_{10}	$\{1,5\}$	100010	2	51	24	B_{11}	$\{4,5,6\}$	000111	3	86
6	M_{11}	$\{2,3\}$	011000	2	51	25	B_{12}	$\{1,2,5\}$	110010	3	86
7	M_{12}	$\{4,6\}$	000101	2	51	26	B_{20}	$\{2,3,5\}$	011010	3	86
8	M_{20}	$\{1,4\}$	100100	2	51	27	B_{21}	$\{2,4,6\}$	010101	3	86
9	M_{21}	$\{3,5\}$	001010	2	51	28	B_{22}	$\{1,4,5\}$	100110	3	86
10	M_{22}	$\{2,6\}$	010001	2	51	29	C_{00}	$\{3,4,6\}$	001101	3	76
11	P_0	$\{6\}$	000001	1	16	30	C_{01}	$\{1,6\}$	100001	2	76
12	P_1	$\{1\}$	100000	1	16	31	C_{02}	$\{1,2,3\}$	111000	3	76
13	P_2	$\{3\}$	001000	1	16	32	C_{10}	$\{1,4,6\}$	100101	3	76
14	Q_0	$\{5\}$	000010	1	26	33	C_{11}	$\{1,3\}$	101000	2	76
15	Q_1	$\{2,5\}$	010010	2	26	34	C_{12}	$\{2,3,6\}$	011001	3	76
16	Q_2	$\{4\}$	000100	1	26	35	C_{20}	$\{1,3,4\}$	101100	3	76
17	A_0	$\{1,3,4,5\}$	101110	4	83	36	C_{21}	$\{3,6\}$	001001	2	76
18	A_1	$\{3,4,5,6\}$	001111	4	83	37	C_{22}	$\{1,2,6\}$	110001	3	76
19	A_2	$\{1,4,5,6\}$	100111	4	83						

S2.27.4. All full 153-state vertex profiles

Table S77 prints the transpose of the complete state table: the full 153-bit profile of every vertex, in three consecutive blocks of 51 states. Thus it also enumerates the complete active state set $X_i = \{s_r : s_r(i) = 1\}$ for every vertex: a one in position r means exactly $s_r \in X_i$. The row weights are 54 for z, 51 for M, 16 for P, 26 for Q, 83 for A, 86 for B, and 76 for C.

TABLE S77: Complete 153-bit vertex profiles, with consecutive 51-bit blocks. Every row is nonzero and all 37 rows are distinct.

i	Ray	W_{153}	$s_1 \dots s_{51}$	$s_{52} \dots s_{102}$	$s_{103} \dots s_{153}$
1	z	54	100001100001100001100001100001111000011000010110000	110000110000101100001000011000011110000110000110000	110000111100001100001011000011000011000011000010000
2	M_{00}	51	111111111111111111111111111111111111111111111111111	000000000000000000000000000000000000000000000000000	000000000000000000000000000000000000000000000000000
3	M_{01}	51	000000000000000000000000000000000000000000000000000	111111111111111111111111111111111111111111111111111	000000000000000000000000000000000000000000000000000
4	M_{02}	51	000000000000000000000000000000000000000000000000000	000000000000000000000000000000000000000000000000000	111111111111111111111111111111111111111111111111111
5	M_{10}	51	111111111111111110000000000000000000000000000000000	111111111111110000000000000000000000000000000000000	111111111111111111000000000000000000000000000000000
6	M_{11}	51	000000000000000011111111111111111111000000000000000	000000000000011111111111111111000000000000000000000	000000000000000000111111111111100000000000000000000
7	M_{12}	51	000000000000000000000000000000000000111111111111111	000000000000000000000000000000011111111111111111111	000000000000000000000000000000011111111111111111111
8	M_{20}	51	111110000000000011111100000000000000111111000000000	111111000000001111110000000000111111100000000000000	111111000000000000110000000000011111100000000000000
9	M_{21}	51	000001111100000000000111111000000000000011000000000	000001111110000000011111000000000111111000000000000	000001111111100000000111111000000000111111000000000
10	M_{22}	51	000000000011111100000000000111111110000000111111000	000000000001100000000001111110000000000000111111000	000000000000111111000000011111100000000000011111000
11	P_0	16	000000000000000000000000000000000000000000000000000	000000000000000000001100001100000000001100001100000	000000000000000000000011000011000000000110001100000
12	P_1	16	011000000000110000000000000000000000011000001100000	000000000000000000000000000000000000000000000000000	001100000000000011000000000000000000011000000001010
13	P_2	16	010100011000000000110000110000000000000000000000000	001100001100000011000101000000000000000000000000000	000000000000000000000000000000000000000000000000000
14	Q_0	26	000000000000000000000000000000000000000000000000000	101010101010000000000000000000011001100101010000000	101010110011000000000000000000000010101010101000000
15	Q_1	26	000001010101010100000001010101100110000000000000000	000000000000000000000000000000000000000000000000000	000001010101010101000010101010100000000000000000000
16	Q_2	26	000000000000000010101000000010101010101000101010000	000000000000010101000001010101010100000001010100000	000000000000000000000000000000000000000000000000000
17	A_0	83	011110011110011110011110011110000111100111101001111	001111001111010011110001100001100001111000011000011	001111000011110011110100001100001100111000011000011
18	A_1	83	000110011110000110011110011110000111100001101000011	001111001111010011110111100111100001111001111001111	000011000011110000110100111001111000011001111001101
19	A_2	83	001010000110011110000110000110000111100111010011111	000011000011010000110010100111100001110011110011111	001111000011110011110100111001111000111001111011111
20	B_{00}	86	000000000000000000000000000000000000000000000000000	111111111111111111111100111001111111111110011110011	111111111111111111111110011100111111111100111001111
21	B_{01}	86	100111111111100111111111111111111111111001111110011	000000000000000000000000000000000000000000000000000	110011111111111100111111111111111111001111111110101
22	B_{02}	86	101011100111111111100111100111111111111111111111111	110011110011111100111010111111111111111111111111111	000000000000000000000000000000000000000000000000000
23	B_{10}	86	000000000000000011111111111111111111111111111111111	000000000000001111111001110011111111111110011110011	000000000000000000011100111100111111111100111001111
24	B_{11}	86	100111111111100110000000000000000000011001111110011	111111111111110000000000000000111111111111111111111	110011111111111100110000000000000011001111111110101
25	B_{12}	86	101011100111111111100111100111111111111000000000000	110011110011111100111010111111100000000000000000000	111111111111111111111111111111111111111000000000000
26	B_{20}	86	000001111111111100000011111111111111000000111111111	000000111111110000010011110011000000011001110011111	000000111111111111110011001111001100000011001110011
27	B_{21}	86	100110000011001111111000000111111111100110011011111	111111000000111111110000011111111111111000000111111	110011000000011001111000000111111100110000001011111
28	B_{22}	86	101011100110000011001111001100000001111111100000000	110011110011001100111010100000011111111111111000000	111111111111110000001111111100000011111111111100000
29	C_{00}	76	000000000000000000000000000000000000000000000000000	010101010101111111111111111111111001100110101011111	010101001100111111111111111111111101010101011111111
30	C_{01}	76	111110101010101011111110101001100111111111111111111	000000000000000000000000000000000000000000000000000	111111010101010101011101010101010111111111111111111
31	C_{02}	76	111111111111111110101011111110101010101010111010101	111111111111110101011111101010101010111111101010101	000000000000000000000000000000000000000000000000000
32	C_{10}	76	111111111111111100000000000000000000011111111111111	010101010101110000000000000000000110011010101111111	010101001100111111110000000000000010101010101111111
33	C_{11}	76	111110101010101011111110101001100110000000000000000	111111111111111111111111111111111000000000000000000	111111010101010101011101010101010100000000000000000
34	C_{12}	76	000000000000000010101111110101010101010111010101000	000000000000000010101111110101010101010111111101010	000000000000000000011111111111111111111111111111111
35	C_{20}	76	111111111110000001111111111110000000011111111000000	010101010100111111111110000000011001101010100000000	010101001100110000011111111000000010101010100000000
36	C_{21}	76	000000101010101010000000101010011001100000011111111	000000111111110000001111111111000000001111111111111	000000010101010101010001010101010100000011111111111
37	C_{22}	76	111110000001111110101010000000101010101010100010101	111111000000110101010000001010101010100000010101000	111111000000011111111100000011111111111100000011111

S2.27.5. The six-state double cover

The fractional-coloring certificate uses a second six-state family,

$$\mathcal{R}_6 = \{s_2, s_{30}, s_{73}, s_{99}, s_{109}, s_{139}\}. \tag{S16}$$

Table S78 specifies these states in the branch parametrization. Each of the 37 vertices occurs in exactly two of their supports. Thus weights $1/2$ give a fractional coloring of total weight three, and uniform weights $1/6$ reproduce the maximally mixed probabilities $1/3$ for every ray. The family Ω_6 in Eq. (S15) supplies the separating certificate; $\mathcal{R}_6$ supplies the double-cover certificate.

TABLE S78: Explicit six-state double cover $\mathcal{R}_6$. Omitted P, Q values are zero; completion values follow from the state equations.

j	State	t	ε	Active P subscripts	Active Q subscripts
1	s_2	$(0,0,0)$	0	$\{1,2\}$	$\varnothing$
2	s_{30}	$(0,1,2)$	1	$\varnothing$	$\{1,2\}$
3	s_{73}	$(1,1,1)$	0	$\{0,2\}$	$\varnothing$
4	s_{99}	$(1,2,2)$	0	$\{0\}$	$\{2\}$
5	s_{109}	$(2,0,1)$	1	$\varnothing$	$\{0,1\}$
6	s_{139}	$(2,2,0)$	0	$\{1\}$	$\{0\}$

S2.27.6. Order determination and exact verification

The companion file `sources/hesse/hesse37-canonical-evaluations.txt` records the 24 numbered contexts and all 153 binary state vectors. The parametrization in this appendix defines their canonical order independently of any external program.

Extend the canonical states to the 76 events of $L(H_{37})$ as in Sec. V A. If $X_x = \{r : s_r(x) = 1\}$, then the full profiles satisfy

$$X_x \subseteq X_y \quad \Longleftrightarrow \quad x \leq y. \tag{S17}$$

The verifier checks all $76^2 = 5{,}776$ ordered event pairs against the incidence order in Eq. (S27). Whenever $x \not\leq y$, the companion file `sources/hesse/hesse37-order-witnesses.csv` gives the least canonical index r with $s_r(x) = 1$ and $s_r(y) = 0$. Its event labels `a1`,...,`a37` denote atoms and `c1`,...,`c37` their complements; `0` and `1` denote the bounds. Every one of the 594 distinct nonorthogonal atom pairs also has a joint-one state. Thus the full family is order determining; the six-state family Ω_6 supplies the minimum unital, separating vertex representation in Eq. (S26). The distinct minima for event separation and order determination are certified in Appendix S3.3.

The verifier `hesse37-verify.py` uses exact rational arithmetic in $\mathbb{Q}(\zeta)$, with no numerical tolerance or external solver. It checks distinct rays, all 666 Hermitian inner products, the 24 bases, all graph triangles, linearity, the chromatic number and core criticality, all 153 states, the minimum separating family, both profile tables, the maximum squared ray overlap, the six-state double cover, and all event-order comparisons. It also compares the reconstructed state vectors with the archived enumeration. The accompanying machine-readable data give the same numbering and context ordering as the figure and these appendices. From the extracted joint project directory, run

```
python verification/verify-integrated.py
```

S2.28. Four-dimensional $H_{\mathrm{YO}}^{(4)}$ and H_{65}: states and profiles

This appendix provides the complete enumeration of all two-valued states on the four-dimensional examples $H_{\mathrm{YO}}^{(4)}$ and H_{65}: the 25 two-valued states of the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)}$ (Appendix S2.28.1), and the complete branch-by-branch parametrization of all $24 \times 2^{14} = 393{,}216$ two-valued states—together with the 48 canonical unital and separating states $s_{\alpha,\varepsilon}$ and the 65 vertex state-profiles—of the canonical linear hypergraph H_{65} (Appendix S2.28.2).

S2.28.1. Complete enumeration of the 25 two-valued states on $H_{\mathrm{YO}}^{(4)}$

Table S79 lists all 25 two-valued states $s_1, \ldots, s_{25} : V_{26} \to \{0,1\}$ on $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$. States $s_1, \ldots, s_{24}$ have $s_r(26) = 0$ and correspond to the 24 two-valued states of the completed Yu–Oh hypergraph H_{YO}^{+} in $\mathbb{R}^3$ (grouped into 12 pairs of states sharing the same support on the three coordinate contexts $\{1,2,3\}$, $\{4,5,6\}$, $\{7,8,9\}$), while s_{25} is the unique apex state with support $\{26\}$. Table S80 displays the transposed 26×25 vertex-by-state binary matrix, listing for each vertex $i \in \{1, \ldots, 26\}$ its state weight $\sum_{r=1}^{25} s_r(i) \in \{1, 3, 8, 13\}$, the set of states $A_i = \{s_r : s_r(i) = 1\}$, and its 25-bit profile $(s_1(i), \ldots, s_{25}(i))$.

Inspection of Table S80 highlights three structural properties of the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$:

1. *Unitality and atom separation:* Every row of Table S80 has positive weight (1, 3, 8, or 13) and all 26 binary rows are distinct ($A_i \neq A_j$ for all $1 \leq i < j \leq 26$), so $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is nonempty, unital, and separating on atoms.

2. *Exclusion of simultaneous* 1*s on the non-orthogonal triad atoms* $\{10, 11, 12, 13\}$*:* Although the four Yu–Oh triad rays $v_{10}, v_{11}, v_{12}, v_{13}$ are pairwise non-orthogonal ($v_i \cdot v_j = \pm 1 \neq 0$ for $10 \leq i < j \leq 13$), their active state sets in Table S80 are pairwise disjoint:

$$\begin{aligned} A_{10} &= \{s_{13}, s_{19}, s_{23}\}, \quad A_{11} = \{s_5, s_7, s_{15}\}, \\ A_{12} &= \{s_1, s_3, s_9\}, \qquad A_{13} = \{s_{11}, s_{17}, s_{21}\}. \end{aligned} \tag{S18}$$

Consequently, $\sum_{i=10}^{13} s(i) \leq 1$ holds for every $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$, and no two-valued state assigns $(1, 1)$ to any of the $\binom{4}{2} = 6$ non-orthogonal pairs $\{10, 11\}$, $\{10, 12\}$, $\{10, 13\}$, $\{11, 12\}$, $\{11, 13\}$, $\{12, 13\}$.

3. *Failure of order-determinacy:* Even though $A_i \neq A_j$ for all $i \neq j$, the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is not order-determining on atoms or compound propositions [4]. In particular, each triad atom $t \in \{10, 11, 12, 13\}$ (*weight* 3) is strictly dominated across all 25 states ($A_t \subsetneq A_w$, i.e., $s(t) \leq s(w)$ for all $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$) by three non-orthogonal completion atoms $w \in \{14, \ldots, 25\}$ (*weight* 13):

$$\begin{aligned} A_{10} &\subsetneq A_{16},\ A_{20},\ A_{24}, \quad A_{11} \subsetneq A_{18},\ A_{21},\ A_{25}, \\ A_{12} &\subsetneq A_{14},\ A_{22},\ A_{23}, \quad A_{13} \subsetneq A_{15},\ A_{17},\ A_{19}, \end{aligned} \tag{S19}$$

together with compound-event inclusions such as $A_{12} \subsetneq A_3 \dot{\cup} A_{14}$, $A_{11} \subsetneq A_3 \dot{\cup} A_{25}$, and their unions with the apex set $A_{26} = \{s_{25}\}$.

TABLE S79: Complete enumeration of the 25 two-valued states $s_1, \ldots, s_{25} : V_{26} \to \{0, 1\}$ on the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, showing for each state its support on the 13-ray Yu–Oh core $\{1, \ldots, 13\}$, its forced support on $\{14, \ldots, 26\}$, its total support size $|S_r|$, and its 26-bit binary vector $(s_r(1 \ldots 13)\, s_r(14 \ldots 25)\, s_r(26))$.

State	Core support	Completion/apex support	$\lvert S_r \rvert$	26-bit state vector
s_1	$\{1, 4, 8, 12\}$	$\{14, 15, 21, 22, 23\}$	9	1001000100010 1100000111000
s_2	$\{1, 4, 8\}$	$\{14, 15, 16, 17, 21, 22, 23, 25\}$	11	1001000100000 1111000111010
s_3	$\{1, 5, 7, 12\}$	$\{14, 18, 22, 23, 24\}$	9	1000101000010 1000100011100
s_4	$\{1, 5, 7\}$	$\{14, 16, 17, 18, 22, 23, 24, 25\}$	11	1000101000000 1011100011110
s_5	$\{1, 5, 9, 11\}$	$\{17, 18, 21, 22, 25\}$	9	1000100010100 0001100110010
s_6	$\{1, 5, 9\}$	$\{14, 15, 17, 18, 21, 22, 24, 25\}$	11	1000100010000 1101100110110
s_7	$\{1, 6, 8, 11\}$	$\{16, 18, 21, 23, 25\}$	9	1000010100100 0010100101010
s_8	$\{1, 6, 8\}$	$\{14, 15, 16, 18, 21, 23, 24, 25\}$	11	1000010100000 1110100101110
s_9	$\{2, 4, 7, 12\}$	$\{14, 19, 20, 22, 23\}$	9	0101001000010 1000011011000
s_{10}	$\{2, 4, 7\}$	$\{14, 16, 17, 19, 20, 22, 23, 25\}$	11	0101001000000 1011011011010
s_{11}	$\{2, 4, 9, 13\}$	$\{14, 15, 17, 19, 25\}$	9	0101000010001 1101010000010
s_{12}	$\{2, 4, 9\}$	$\{14, 15, 17, 19, 20, 21, 22, 25\}$	11	0101000010000 1101011110010
s_{13}	$\{2, 6, 7, 10\}$	$\{14, 16, 20, 24, 25\}$	9	0100011001000 1010001000110
s_{14}	$\{2, 6, 7\}$	$\{14, 16, 18, 19, 20, 23, 24, 25\}$	11	0100011000000 1010111001110
s_{15}	$\{2, 6, 9, 11\}$	$\{18, 19, 20, 21, 25\}$	9	0100010010100 0000111100010
s_{16}	$\{2, 6, 9\}$	$\{14, 15, 18, 19, 20, 21, 24, 25\}$	11	0100010010000 1100111100110
s_{17}	$\{3, 4, 8, 13\}$	$\{15, 16, 17, 19, 23\}$	9	0011000100001 0111010001000
s_{18}	$\{3, 4, 8\}$	$\{15, 16, 17, 19, 20, 21, 22, 23\}$	11	0011000100000 0111011111000
s_{19}	$\{3, 5, 7, 10\}$	$\{16, 17, 20, 22, 24\}$	9	0010101001000 0011001010100
s_{20}	$\{3, 5, 7\}$	$\{16, 17, 18, 19, 20, 22, 23, 24\}$	11	0010101000000 0011111011100
s_{21}	$\{3, 5, 9, 13\}$	$\{15, 17, 18, 19, 24\}$	9	0010100010001 0101110000100
s_{22}	$\{3, 5, 9\}$	$\{15, 17, 18, 19, 20, 21, 22, 24\}$	11	0010100010000 0101111110100
s_{23}	$\{3, 6, 8, 10\}$	$\{15, 16, 20, 21, 24\}$	9	0010010101000 0110001100100
s_{24}	$\{3, 6, 8\}$	$\{15, 16, 18, 19, 20, 21, 23, 24\}$	11	0010010100000 0110111101100
s_{25}	$\emptyset$	$\{26\}$	1	0000000000000 0000000000001

TABLE S80: Vertex state profiles across the 25 two-valued states $s_1, \dots, s_{25}$ of $H_{\mathrm{YO}}^{(4)}$, showing for each vertex $i \in \{1, \dots, 26\}$ its state weight $\sum_{r=1}^{25} s_r(i)$, the active state indices $\{r : s_r(i) = 1\}$, and the 25-bit profile $(s_{1\dots12}(i)\, s_{13\dots24}(i)\, s_{25}(i))$. All 26 rows are nonzero and pairwise distinct.

Vertex i	Weight $\sum_r s_r(i)$	Active states $\{r \in \{1, \dots, 25\} : s_r(i) = 1\}$	25-bit profile $(s_{1\dots12}(i)\, s_{13\dots24}(i)\, s_{25}(i))$
1	8	$\{1, 2, 3, 4, 5, 6, 7, 8\}$	111111110000 000000000000 0
2	8	$\{9, 10, 11, 12, 13, 14, 15, 16\}$	000000001111 111100000000 0
3	8	$\{17, 18, 19, 20, 21, 22, 23, 24\}$	000000000000 000011111111 0
4	8	$\{1, 2, 9, 10, 11, 12, 17, 18\}$	110000001111 000011000000 0
5	8	$\{3, 4, 5, 6, 19, 20, 21, 22\}$	001111000000 000000111100 0
6	8	$\{7, 8, 13, 14, 15, 16, 23, 24\}$	000000110000 111100000011 0
7	8	$\{3, 4, 9, 10, 13, 14, 19, 20\}$	001100001100 110000110000 0
8	8	$\{1, 2, 7, 8, 17, 18, 23, 24\}$	110000110000 000011000011 0
9	8	$\{5, 6, 11, 12, 15, 16, 21, 22\}$	000011000011 001100001100 0
10	3	$\{13, 19, 23\}$	000000000000 100000100010 0
11	3	$\{5, 7, 15\}$	000010100000 001000000000 0
12	3	$\{1, 3, 9\}$	101000001000 000000000000 0
13	3	$\{11, 17, 21\}$	000000000010 000010001000 0
14	13	$\{1, 2, 3, 4, 6, 8, 9, 10, 11, 12, 13, 14, 16\}$	111101011111 110100000000 0
15	13	$\{1, 2, 6, 8, 11, 12, 16, 17, 18, 21, 22, 23, 24\}$	110001010011 000111001111 0
16	13	$\{2, 4, 7, 8, 10, 13, 14, 17, 18, 19, 20, 23, 24\}$	010100110100 110011110011 0
17	13	$\{2, 4, 5, 6, 10, 11, 12, 17, 18, 19, 20, 21, 22\}$	010111000111 000011111100 0
18	13	$\{3, 4, 5, 6, 7, 8, 14, 15, 16, 20, 21, 22, 24\}$	001111110000 011100011101 0
19	13	$\{9, 10, 11, 12, 14, 15, 16, 17, 18, 20, 21, 22, 24\}$	000000001111 011111011101 0
20	13	$\{9, 10, 12, 13, 14, 15, 16, 18, 19, 20, 22, 23, 24\}$	000000001101 111101110111 0
21	13	$\{1, 2, 5, 6, 7, 8, 12, 15, 16, 18, 22, 23, 24\}$	110011110001 001101000111 0
22	13	$\{1, 2, 3, 4, 5, 6, 9, 10, 12, 18, 19, 20, 22\}$	111111001101 000001110100 0
23	13	$\{1, 2, 3, 4, 7, 8, 9, 10, 14, 17, 18, 20, 24\}$	111100111100 010011010001 0
24	13	$\{3, 4, 6, 8, 13, 14, 16, 19, 20, 21, 22, 23, 24\}$	001101010000 110100111111 0
25	13	$\{2, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16\}$	010111110111 111100000000 0
26	1	$\{25\}$	000000000000 000000000000 1

S2.28.2. Complete parametrization of all 393,216 *two-valued states and the* 48 *canonical separating states on* H_{65}

Every two-valued state $s \in \mathcal{S}(H_{65})$ on the canonical linear hypergraph $H_{65} = (V_{65}, E_{29})$ is uniquely indexed by a pair $(\alpha, \boldsymbol{\varepsilon})$ consisting of:

1. a branch index $\alpha \in \{1, \dots, 24\}$ selecting one of the 24 four-element core state supports $S_\alpha \subset V_{\mathrm{core}} = \{1, \dots, 17\}$; and
2. a 14-bit binary vector $\boldsymbol{\varepsilon} = (\varepsilon_m)_{m \in \mathcal{F}_\alpha} \in \{0,1\}^{14}$ indexed by the 14 *free completion contexts*

$$\mathcal{F}_\alpha = \big\{m \in \{1, \dots, 24\} : S_\alpha \cap \{u_m, v_m\} = \emptyset\big\},$$

whose complement is the set of 10 *hit completion contexts* $\mathcal{H}_\alpha = \{1, \dots, 24\} \setminus \mathcal{F}_\alpha$.

Explicitly, the state $s_{\alpha,\boldsymbol{\varepsilon}} : V_{65} \to \{0,1\}$ takes values

$$\begin{aligned} s_{\alpha,\boldsymbol{\varepsilon}}(i) &= \mathbf{1}_{i \in S_\alpha} \qquad (1 \le i \le 17), \\ \big(s_{\alpha,\boldsymbol{\varepsilon}}(2m+16),\, s_{\alpha,\boldsymbol{\varepsilon}}(2m+17)\big) &= \begin{cases} (0,0), & m \in \mathcal{H}_\alpha, \\ (1-\varepsilon_m,\, \varepsilon_m), & m \in \mathcal{F}_\alpha. \end{cases} \end{aligned} \tag{S20}$$

Because $|\mathcal{F}_\alpha| = 14$ for every $\alpha \in \{1, \dots, 24\}$, each branch α yields $2^{14} = 16{,}384$ states $s_{\alpha,\boldsymbol{\varepsilon}}$, giving $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$ two-valued states in total. Setting all 14 free bits equal to $\varepsilon \in \{0,1\}$ ($\varepsilon_m = \varepsilon$ for all $m \in \mathcal{F}_\alpha$) yields the 48 *canonical states* $s_{\alpha,0}, s_{\alpha,1}$ of Eq. (S29), which have support size $|S_\alpha| + 14 = 18$:

$$\begin{aligned} \mathrm{supp}(s_{\alpha,0}) &= S_\alpha \cup \big\{2m + 16 : m \in \mathcal{F}_\alpha\big\}, \\ \mathrm{supp}(s_{\alpha,1}) &= S_\alpha \cup \big\{2m + 17 : m \in \mathcal{F}_\alpha\big\}. \end{aligned} \tag{S21}$$

Table S81 enumerates all 24 branches $\alpha = 1, \ldots, 24$, specifying S_α, the 10 hit context indices $\mathcal{H}_\alpha \subset \{1, \ldots, 24\}$, the 14 free context indices $\mathcal{F}_\alpha \subset \{1, \ldots, 24\}$, and the 24-bit free-context indicator string. Table S82 lists the resulting state profiles and weights for all 65 vertices of H_{65} across both the 48 canonical states $\{s_{\alpha,\varepsilon}\}$ and the full space of 393,216 states: across the 48 canonical states $(s_{1,0}, s_{1,1}, \ldots, s_{24,0}, s_{24,1})$, each '1' in row i of Table S82 expands to '(1,1)' for core vertices $i \in \{1, \ldots, 17\}$, to '(1,0)' for first completion vertices $w_{m,1} = 2m + 16$, and to '(0,1)' for second completion vertices $w_{m,2} = 2m + 17$ (and each '0' expands to '(0,0)'), making all 65 canonical 48-bit vertex profiles nonzero and pairwise distinct by inspection.

TABLE S81: Complete branch-by-branch parametrization of the $24 \times 2^{14} = 393{,}216$ two-valued states $s_{\alpha,\boldsymbol{\varepsilon}}$ and the 48 canonical states $s_{\alpha,0}, s_{\alpha,1}$ on H_{65} via Eqs. (S20)–(S21). For each $\alpha \in \{1,\ldots,24\}$, the table lists the 4-element core support S_α, the 10 hit completion context indices $m \in \mathcal{H}_\alpha$ where $(s(2m+16), s(2m+17)) = (0,0)$ is forced on E_m, the 14 free completion context indices $m \in \mathcal{F}_\alpha$ where $(s(2m+16), s(2m+17)) = (1-\varepsilon_m, \varepsilon_m)$ with $\varepsilon_m \in \{0,1\}$ on E_m, and the 24-bit indicator string $(\mathbf{1}_{1\in\mathcal{F}_\alpha} \ldots \mathbf{1}_{12\in\mathcal{F}_\alpha}\, \mathbf{1}_{13\in\mathcal{F}_\alpha} \ldots \mathbf{1}_{24\in\mathcal{F}_\alpha})$.

Branch	Core support S_α	Hit contexts $\mathcal{H}_\alpha$	Free contexts $\mathcal{F}_\alpha$	Free indicator $m = 1 \ldots 24$
S_1	{1,5,9,16}	{3,7,9,10,11,13,19,21,22,23}	{1,2,4,5,6,8,12,14,15,16,17,18,20,24}	110111010001 011111010001
S_2	{1,5,10,14}	{3,5,6,7,11,13,17,18,19,23}	{1,2,4,8,9,10,12,14,15,16,20,21,22,24}	110100011101 011100011101
S_3	{1,6,9,16}	{3,7,9,10,11,15,17,21,22,24}	{1,2,4,5,6,8,12,13,14,16,18,19,20,23}	110111010001 110101110010
S_4	{1,6,11,12}	{1,2,3,7,11,13,14,15,17,24}	{4,5,6,8,9,10,12,16,18,19,20,21,22,23}	000111011101 000101111110
S_5	{1,7,10,14}	{3,5,6,7,11,16,17,18,20,21}	{1,2,4,8,9,10,12,13,14,15,19,22,23,24}	110100011101 111000100111
S_6	{1,7,11,12}	{1,2,3,7,11,13,14,16,20,21}	{4,5,6,8,9,10,12,15,17,18,19,22,23,24}	000111011101 001011100111
S_7	{2,5,9,16}	{1,5,9,10,12,13,19,21,22,23}	{2,3,4,6,7,8,11,14,15,16,17,18,20,24}	011101110010 011111010001
S_8	{2,5,11,13}	{1,3,4,5,12,13,15,16,19,23}	{2,6,7,8,9,10,11,14,17,18,20,21,22,24}	010001111110 010011011101
S_9	{2,6,9,16}	{1,5,9,10,12,15,17,21,22,24}	{2,3,4,6,7,8,11,13,14,16,18,19,20,23}	011101110010 110101110010
S_{10}	{2,6,10,15}	{1,5,7,8,12,15,17,19,20,24}	{2,3,4,6,9,10,11,13,14,16,18,21,22,23}	011101001110 110101001110
S_{11}	{2,8,10,15}	{1,5,7,8,12,14,18,19,20,22}	{2,3,4,6,9,10,11,13,15,16,17,21,23,24}	011101001110 101110001011
S_{12}	{2,8,11,13}	{1,3,4,5,12,14,15,16,18,22}	{2,6,7,8,9,10,11,13,17,19,20,21,23,24}	010001111110 100010111011
S_{13}	{3,5,10,14}	{2,5,6,8,9,13,17,18,19,23}	{1,3,4,7,10,11,12,14,15,16,20,21,22,24}	101100100111 011100011101
S_{14}	{3,5,11,13}	{2,3,4,8,9,13,15,16,19,23}	{1,5,6,7,10,11,12,14,17,18,20,21,22,24}	100011100111 010011011101
S_{15}	{3,7,9,17}	{2,8,9,11,12,16,20,21,23,24}	{1,3,4,5,6,7,10,13,14,15,17,18,19,22}	101111100100 111011100100
S_{16}	{3,7,10,14}	{2,5,6,8,9,16,17,18,20,21}	{1,3,4,7,10,11,12,13,14,15,19,22,23,24}	101100100111 111000100111
S_{17}	{3,8,9,17}	{2,8,9,11,12,14,18,22,23,24}	{1,3,4,5,6,7,10,13,15,16,17,19,20,21}	101111100100 101110111000
S_{18}	{3,8,11,13}	{2,3,4,8,9,14,15,16,18,22}	{1,5,6,7,10,11,12,13,17,19,20,21,23,24}	100011100111 100010111011
S_{19}	{4,6,10,15}	{4,6,7,8,10,15,17,19,20,24}	{1,2,3,5,9,11,12,13,14,16,18,21,22,23}	111010001011 110101001110
S_{20}	{4,6,11,12}	{1,2,4,6,10,13,14,15,17,24}	{3,5,7,8,9,11,12,16,18,19,20,21,22,23}	001010111011 000101111110
S_{21}	{4,7,9,17}	{4,6,10,11,12,16,20,21,23,24}	{1,2,3,5,7,8,9,13,14,15,17,18,19,22}	111010111000 111011100100
S_{22}	{4,7,11,12}	{1,2,4,6,10,13,14,16,20,21}	{3,5,7,8,9,11,12,15,17,18,19,22,23,24}	001010111011 001011100111
S_{23}	{4,8,9,17}	{4,6,10,11,12,14,18,22,23,24}	{1,2,3,5,7,8,9,13,15,16,17,19,20,21}	111010111000 101110111000
S_{24}	{4,8,10,15}	{4,6,7,8,10,14,18,19,20,22}	{1,2,3,5,9,11,12,13,15,16,17,21,23,24}	111010001011 101110001011

TABLE S82: Vertex state profiles and weights for all 65 vertices of H_{65}, showing for each core vertex $i \in \{1, \ldots, 17\}$ and each completion pair $(w_{m,1}, w_{m,2}) = (2m+16, 2m+17)$ $(m = 1, \ldots, 24)$ the number of active branches W_{24}, the weight $W_{48} = \sum_{\alpha,\varepsilon} s_{\alpha,\varepsilon}(i)$ across the 48 canonical states, the weight $W_{393,216} = \sum_{s\in\mathcal{S}(H_{65})} s(i)$ across all 393,216 states, the active branch indices $\alpha \in \{1, \ldots, 24\}$ (equal to $Z(e_m)$ for $(w_{m,1}, w_{m,2})$), and the 24-bit branch profile across $S_1, \ldots, S_{24}$ (in which each `1` expands to `11` for core vertices i, to `10` for $w_{m,1}$, and to `01` for $w_{m,2}$ across the 48 canonical states).

Vertex / pair	W_{24}	W_{48}	$W_{393,216}$	Active branches α	24-bit branch profile
Core 1	6	12	98,304	{1, 2, 3, 4, 5, 6}	11111100 00000000 00000000
Core 2	6	12	98,304	{7, 8, 9, 10, 11, 12}	00000011 11110000 00000000
Core 3	6	12	98,304	{13, 14, 15, 16, 17, 18}	00000000 00001111 11000000
Core 4	6	12	98,304	{19, 20, 21, 22, 23, 24}	00000000 00000000 00111111
Core 5	6	12	98,304	{1, 2, 7, 8, 13, 14}	11000011 00001100 00000000
Core 6	6	12	98,304	{3, 4, 9, 10, 19, 20}	00110000 11000000 00110000
Core 7	6	12	98,304	{5, 6, 15, 16, 21, 22}	00001100 00000011 00001100
Core 8	6	12	98,304	{11, 12, 17, 18, 23, 24}	00000000 00110000 11000011
Core 9	8	16	131,072	{1, 3, 7, 9, 15, 17, 21, 23}	10100010 10000010 10001010
Core 10	8	16	131,072	{2, 5, 10, 11, 13, 16, 19, 24}	01001000 01101001 00100001
Core 11	8	16	131,072	{4, 6, 8, 12, 14, 18, 20, 22}	00010101 00010100 01010100
Core 12	4	8	65,536	{4, 6, 20, 22}	00010100 00000000 00010100

Continued on the next page.

TABLE S82: Vertex profiles of H_{65} (continued).

Vertex / pair	W_{24}	W_{48}	$W_{393,216}$	Active branches α	24-bit branch profile
Core 13	4	8	65,536	$\{8, 12, 14, 18\}$	00000001 00010100 01000000
Core 14	4	8	65,536	$\{2, 5, 13, 16\}$	01001000 00001001 00000000
Core 15	4	8	65,536	$\{10, 11, 19, 24\}$	00000000 01100000 00100001
Core 16	4	8	65,536	$\{1, 3, 7, 9\}$	10100010 10000000 00000000
Core 17	4	8	65,536	$\{15, 17, 21, 23\}$	00000000 00000010 10001010
$E_1 : (18, 19)$	14	14	114,688	$\{1, 2, 3, 5, 13, 14, 15, 16, 17, 18, 19, 21, 23, 24\}$	11101000 00001111 11101011
$E_2 : (20, 21)$	14	14	114,688	$\{1, 2, 3, 5, 7, 8, 9, 10, 11, 12, 19, 21, 23, 24\}$	11101011 11110000 00101011
$E_3 : (22, 23)$	14	14	114,688	$\{7, 9, 10, 11, 13, 15, 16, 17, 19, 20, 21, 22, 23, 24\}$	00000010 11101011 10111111
$E_4 : (24, 25)$	14	14	114,688	$\{1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 13, 15, 16, 17\}$	11111110 11101011 10000000
$E_5 : (26, 27)$	14	14	114,688	$\{1, 3, 4, 6, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24\}$	10110100 00000110 11111111
$E_6 : (28, 29)$	14	14	114,688	$\{1, 3, 4, 6, 7, 8, 9, 10, 11, 12, 14, 15, 17, 18\}$	10110111 11110110 11000000
$E_7 : (30, 31)$	14	14	114,688	$\{7, 8, 9, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23\}$	00000011 10011111 11011110
$E_8 : (32, 33)$	14	14	114,688	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 20, 21, 22, 23\}$	11111111 10010000 00011110
$E_9 : (34, 35)$	14	14	114,688	$\{2, 4, 5, 6, 8, 10, 11, 12, 19, 20, 21, 22, 23, 24\}$	01011101 01110000 00111111
$E_{10} : (36, 37)$	14	14	114,688	$\{2, 4, 5, 6, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18\}$	01011101 01111111 11000000
$E_{11} : (38, 39)$	14	14	114,688	$\{7, 8, 9, 10, 11, 12, 13, 14, 16, 18, 19, 20, 22, 24\}$	00000011 11111101 01110101
$E_{12} : (40, 41)$	14	14	114,688	$\{1, 2, 3, 4, 5, 6, 13, 14, 16, 18, 19, 20, 22, 24\}$	11111100 00001101 01110101
$E_{13} : (42, 43)$	14	14	114,688	$\{3, 5, 9, 10, 11, 12, 15, 16, 17, 18, 19, 21, 23, 24\}$	00101000 11110011 11101011
$E_{14} : (44, 45)$	14	14	114,688	$\{1, 2, 3, 5, 7, 8, 9, 10, 13, 14, 15, 16, 19, 21\}$	11101011 11001111 00101000
$E_{15} : (46, 47)$	14	14	114,688	$\{1, 2, 5, 6, 7, 11, 13, 15, 16, 17, 21, 22, 23, 24\}$	11001110 00101011 10001111
$E_{16} : (48, 49)$	14	14	114,688	$\{1, 2, 3, 4, 7, 9, 10, 11, 13, 17, 19, 20, 23, 24\}$	11110010 11101000 10110011
$E_{17} : (50, 51)$	14	14	114,688	$\{1, 6, 7, 8, 11, 12, 14, 15, 17, 18, 21, 22, 23, 24\}$	10000111 00110110 11001111
$E_{18} : (52, 53)$	14	14	114,688	$\{1, 3, 4, 6, 7, 8, 9, 10, 14, 15, 19, 20, 21, 22\}$	10110111 11000110 00111100
$E_{19} : (54, 55)$	14	14	114,688	$\{3, 4, 5, 6, 9, 12, 15, 16, 17, 18, 20, 21, 22, 23\}$	00111100 10010011 11011110
$E_{20} : (56, 57)$	14	14	114,688	$\{1, 2, 3, 4, 7, 8, 9, 12, 13, 14, 17, 18, 20, 23\}$	11110011 10011100 11010010
$E_{21} : (58, 59)$	14	14	114,688	$\{2, 4, 8, 10, 11, 12, 13, 14, 17, 18, 19, 20, 23, 24\}$	01010001 01111100 11110011
$E_{22} : (60, 61)$	14	14	114,688	$\{2, 4, 5, 6, 8, 10, 13, 14, 15, 16, 19, 20, 21, 22\}$	01011101 01001111 00111100
$E_{23} : (62, 63)$	14	14	114,688	$\{3, 4, 5, 6, 9, 10, 11, 12, 16, 18, 19, 20, 22, 24\}$	00111100 11110001 01110101
$E_{24} : (64, 65)$	14	14	114,688	$\{1, 2, 5, 6, 7, 8, 11, 12, 13, 14, 16, 18, 22, 24\}$	11001111 00111101 01000101

S2.29. Parent-state and deletion certificates

The 49-ray, 87-ray, 40-ray, and 72-ray parents have no context states. The two smaller 68-ray parents are treated in Appendix S2.30.1. The first and last two have the independently verified complete bases printed in Appendix S1.16; the 87-ray parent contains the known 69-ray KS lineage. The 70-ray single-apex lift of the completed Cabello parent instead has the unique state supported at its apex.

For the real Yu–Oh parent, the intermediate 33 rays admit 192 assignments to the twenty bases alone and none after its sixteen additional pair exclusions. The sixteen missing pairs are uniquely completed by rays 34–49. The x-axis-plane extension by rays 26–29 alone has 72 basis assignments; all 24 original states extend. Including its eight additional pair exclusions leaves eight extensions, all with $s(2) = 1$. The z-axis-plane case is identical, with $s(5) = 1$. The two added bases on either axis each exclude twelve of the sixteen original axis-zero states when their further pair exclusions are included; eight original states are excluded by both, leaving none. Thus the axis forcing uses the complete orthogonality constraints, not the added bases alone.

All complete states of H_{65} are additionally supplied as `data/h65-all-states.csv.gz`. Each row gives its branch index, the fourteen-bit choice index within that branch, and its 65-bit state vector in vertex order. There are exactly $24 \times 2^{14} = 393{,}216$ data rows. This supplements the complete parametrization and the printed 48-state family; it does not assert that 48 is minimal. The 153 Hesse states, their event-order witnesses, and all original printed profiles remain in the accompanying Hesse data.

S2.30. States of the Cabello restrictions and completions

S2.30.1. States of the vertex-deleted 45-ray completion

The nine-context $C_{18,9}$, its eighteen-context faithful completion, and both 68-ray parents have no context states by their retained parity proofs. For the 45-ray construction, use exactly the local labels in Tables S31 and S32. Table S84 lists all twelve admissible core supports, and Table S85 lists the fourteen bridge pairs with their fresh labels and zero-profiles. For a support S_α, let F_α be the bridge indices missed by S_α. On every hit bridge both fresh values are zero; on each missed bridge independently choose exactly one of its two fresh vertices. This enumerates every context state once.

TABLE S83. Context-state counts after deleting each single context of the real 49-ray parent. Every count is positive. Context numbering is Table S21. Deleting a completion context 21–36 leaves its fresh ray in no context; that ray is counted as an unconstrained free vertex, which doubles the count. If the orthogonalities of the deleted context are kept as geometric exclusions, deletions 21–36 leave no state, while deletions 1–20 leave 1, 2, or 4 states.

Deleted context	States	Deleted context	States	Deleted context	States
1	12	13	2	25	2
2	4	14	4	26	2
3	4	15	4	27	2
4	32	16	16	28	2
5	2	17	4	29	2
6	2	18	4	30	2
7	4	19	4	31	2
8	4	20	4	32	2
9	12	21	2	33	2
10	4	22	2	34	2
11	4	23	2	35	2
12	2	24	2	36	2

TABLE S84: All twelve core supports; $m = |F_\alpha|$. The final column specifies extendibility under all geometric pair exclusions.

α	S_α	m	Geometric extension
1	$\{1, 6, 11, 12\}$	8	yes
2	$\{1, 10, 14, 18\}$	8	yes
3	$\{3, 7, 10, 14\}$	8	yes
4	$\{3, 8, 11, 13\}$	8	yes
5	$\{3, 13, 14, 17\}$	8	no
6	$\{2, 6, 10, 15\}$	8	yes
7	$\{6, 12, 15, 16\}$	8	no
8	$\{8, 10, 15, 18\}$	8	yes
9	$\{9, 16, 18, 19\}$	9	no
10	$\{9, 17, 18, 19\}$	9	no
11	$\{2, 11, 13, 19\}$	8	yes
12	$\{7, 11, 12, 19\}$	8	yes

TABLE S85: The fourteen bridge pairs, fresh labels, and zero-profiles; each Z_m is the set of missed-bridge branch indices.

m	Bridge	Fresh pair	Z_m
1	$\{1,2\}$	$\{21,22\}$	$\{3,4,5,7,8,9,10,12\}$
2	$\{1,13\}$	$\{23,24\}$	$\{3,6,7,8,9,10,12\}$
3	$\{1,17\}$	$\{25,26\}$	$\{3,4,6,7,8,9,11,12\}$
4	$\{2,14\}$	$\{27,28\}$	$\{1,4,7,8,9,10,12\}$
5	$\{2,17\}$	$\{29,30\}$	$\{1,2,3,4,7,8,9,12\}$
6	$\{3,16\}$	$\{31,32\}$	$\{1,2,6,8,10,11,12\}$
7	$\{6,17\}$	$\{33,34\}$	$\{2,3,4,8,9,11,12\}$
8	$\{7,8\}$	$\{35,36\}$	$\{1,2,5,6,7,9,10,11\}$
9	$\{7,15\}$	$\{37,38\}$	$\{1,2,4,5,9,10,11\}$
10	$\{7,16\}$	$\{39,40\}$	$\{1,2,4,5,6,8,10,11\}$
11	$\{8,12\}$	$\{41,42\}$	$\{2,3,5,6,9,10,11\}$
12	$\{8,16\}$	$\{43,44\}$	$\{1,2,3,5,6,10,11,12\}$
13	$\{10,19\}$	$\{45,46\}$	$\{1,4,5,7\}$
14	$\{11,18\}$	$\{47,48\}$	$\{3,5,6,7\}$

Taking the first fresh vertex on every missed bridge gives $s_{\alpha,0}$, and taking the second gives $s_{\alpha,1}$. Table S86 explicitly enumerates all 24 canonical context states. Their profiles are nonzero and pairwise distinct on all 45 vertices. The twelve branch supports and fourteen distinct nonempty zero-profiles verify this by Lemma 21(3).

TABLE S86: All 24 canonical context-state supports of the 45-ray example. Some fail geometric pair exclusions.

α	ε	Support
1	0	$\{1,6,11,12,27,29,31,35,37,39,43,45\}$
1	1	$\{1,6,11,12,28,30,32,36,38,40,44,46\}$
2	0	$\{1,10,14,18,29,31,33,35,37,39,41,43\}$
2	1	$\{1,10,14,18,30,32,34,36,38,40,42,44\}$
3	0	$\{3,7,10,14,21,23,25,29,33,41,43,47\}$
3	1	$\{3,7,10,14,22,24,26,30,34,42,44,48\}$
4	0	$\{3,8,11,13,21,25,27,29,33,37,39,45\}$
4	1	$\{3,8,11,13,22,26,28,30,34,38,40,46\}$
5	0	$\{3,13,14,17,21,35,37,39,41,43,45,47\}$
5	1	$\{3,13,14,17,22,36,38,40,42,44,46,48\}$
6	0	$\{2,6,10,15,23,25,31,35,39,41,43,47\}$
6	1	$\{2,6,10,15,24,26,32,36,40,42,44,48\}$
7	0	$\{6,12,15,16,21,23,25,27,29,35,45,47\}$
7	1	$\{6,12,15,16,22,24,26,28,30,36,46,48\}$
8	0	$\{8,10,15,18,21,23,25,27,29,31,33,39\}$
8	1	$\{8,10,15,18,22,24,26,28,30,32,34,40\}$
9	0	$\{9,16,18,19,21,23,25,27,29,33,35,37,41\}$
9	1	$\{9,16,18,19,22,24,26,28,30,34,36,38,42\}$
10	0	$\{9,17,18,19,21,23,27,31,35,37,39,41,43\}$
10	1	$\{9,17,18,19,22,24,28,32,36,38,40,42,44\}$
11	0	$\{2,11,13,19,25,31,33,35,37,39,41,43\}$
11	1	$\{2,11,13,19,26,32,34,36,38,40,42,44\}$
12	0	$\{7,11,12,19,21,23,25,27,29,31,33,43\}$
12	1	$\{7,11,12,19,22,24,26,28,30,32,34,44\}$

Ten branches miss eight pairs and two miss nine. Therefore

$$|\mathcal{S}_{\mathrm{ctx}}| = 10\,2^8 + 2\,2^9 = 3{,}584, \qquad |\mathcal{S}_{\mathrm{geom}}| = 8\,2^8 = 2048. \tag{S22}$$

The geometric branches are $1,2,3,4,6,8,11,12$; arbitrary fresh choices within each of these branches remain geometrically compatible. The other branches are excluded by the four forced pairs. Both full enumerations are supplied as compressed CSV data, in addition to this complete finite parametrization. Every geometric state vanishes on 9, 16, and 17, so this family is neither unital nor separating.

For an explicit counterexample, the context state supported at

$$\{6,12,15,16,21,23,25,27,29,35,45,47\}$$

selects both 16 and 21, although $v_{16}\cdot v_{21}=0$.

S2.30.2. Deleting one of the nine Cabello contexts

Number the nine contexts as the first nine rows of Table S29. For every deletion there are 26 context states, six geometric states, and a proper four-coloring of the remaining context graph. The color strings in Table S87 use the ordered ray list $(1, 2, 3, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20)$. The geometric states vanish precisely on the omitted context.

TABLE S87: All single-context deletion counts and four-coloring witnesses for $C_{18,9}$.

e	Omitted context	$\lvert\mathcal{S}_{\mathrm{ctx}}\rvert$	$\lvert\mathcal{S}_{\mathrm{geom}}\rvert$	Four-color string
1	$\{9, 10, 12, 13\}$	26	6	012102203020112333
2	$\{9, 11, 14, 15\}$	26	6	012021231012102333
3	$\{10, 11, 16, 17\}$	26	6	012021320012102333
4	$\{2, 3, 12, 18\}$	26	6	013103302120113322
5	$\{1, 3, 15, 19\}$	26	6	103130320102113232
6	$\{1, 2, 17, 20\}$	26	6	330130230103112222
7	$\{6, 7, 13, 18\}$	26	6	012221213030102333
8	$\{6, 8, 14, 19\}$	26	6	012202230013112333
9	$\{7, 8, 16, 20\}$	26	6	012021320012113332

For deletion of context 9, $\{7, 8, 16, 20\}$, Table S88 explicitly enumerates all 26 supports; exactly the six marked rows satisfy every geometric pair exclusion. All nine full deletion-state lists accompany the data.

TABLE S88: All 26 states after deleting context 9; geometric compatibility is marked explicitly.

j	Support	Geometric
1	$\{1, 6, 11, 12\}$	yes
2	$\{1, 7, 8, 11, 12\}$	no
3	$\{1, 7, 12, 14, 16\}$	no
4	$\{1, 8, 9, 16, 18\}$	no
5	$\{1, 10, 14, 18\}$	yes
6	$\{2, 6, 10, 15\}$	yes
7	$\{2, 7, 8, 10, 15\}$	no
8	$\{2, 8, 13, 15, 16\}$	no
9	$\{2, 7, 9, 16, 19\}$	no
10	$\{2, 11, 13, 19\}$	yes
11	$\{3, 6, 9, 17\}$	no
12	$\{3, 7, 8, 9, 17\}$	no
13	$\{3, 13, 14, 17\}$	yes
14	$\{6, 12, 15, 17\}$	no
15	$\{7, 8, 12, 15, 17\}$	no
16	$\{9, 17, 18, 19\}$	yes
17	$\{3, 6, 9, 16, 20\}$	no
18	$\{3, 7, 8, 9, 16, 20\}$	no
19	$\{3, 7, 10, 14, 20\}$	no
20	$\{3, 8, 11, 13, 20\}$	no
21	$\{3, 13, 14, 16, 20\}$	no
22	$\{6, 12, 15, 16, 20\}$	no
23	$\{7, 11, 12, 19, 20\}$	no
24	$\{7, 8, 12, 15, 16, 20\}$	no
25	$\{9, 16, 18, 19, 20\}$	no
26	$\{8, 10, 15, 18, 20\}$	no

S2.30.3. All single-ray deletions and the alternative parity subsystem

The next table records the two span failures and all forced-zero core rays after geometric pair exclusion for every possible single-ray deletion. An entry $\{u, v\} : z$ means $v_z \in \mathrm{span}(v_u, v_v)$. Every row has twelve compatible core states, 3,584 abstract completion states, and 2,048 geometric completion states, with exactly four forced extra pairs and no other extra orthogonality in the supplied realization. The complete core supports and 24 canonical states for every row are also supplied in the data.

TABLE S89: All eighteen vertex-deletion checks; the completed geometric families fail unitality and separation in every case.

Deleted ray	First span failure	Second span failure	Forced-zero core rays
1	$\{15,19\}:2$	$\{17,20\}:3$	$\{2,3,13\}$
2	$\{12,18\}:1$	$\{17,20\}:3$	$\{1,3,14\}$
3	$\{12,18\}:1$	$\{15,19\}:2$	$\{1,2,16\}$
6	$\{13,18\}:8$	$\{14,19\}:7$	$\{7,8,17\}$
7	$\{13,18\}:8$	$\{16,20\}:6$	$\{6,8,15\}$
8	$\{14,19\}:7$	$\{16,20\}:6$	$\{6,7,12\}$
9	$\{12,13\}:11$	$\{14,15\}:10$	$\{10,11,20\}$
10	$\{12,13\}:11$	$\{16,17\}:9$	$\{9,11,19\}$
11	$\{14,15\}:10$	$\{16,17\}:9$	$\{9,10,18\}$
12	$\{2,3\}:13$	$\{9,10\}:18$	$\{8,13,18\}$
13	$\{6,7\}:12$	$\{9,10\}:18$	$\{1,12,18\}$
14	$\{6,8\}:15$	$\{9,11\}:19$	$\{2,15,19\}$
15	$\{1,3\}:14$	$\{9,11\}:19$	$\{7,14,19\}$
16	$\{7,8\}:17$	$\{10,11\}:20$	$\{3,17,20\}$
17	$\{1,2\}:16$	$\{10,11\}:20$	$\{6,16,20\}$
18	$\{2,3\}:13$	$\{6,7\}:12$	$\{11,12,13\}$
19	$\{1,3\}:14$	$\{6,8\}:15$	$\{10,14,15\}$
20	$\{1,2\}:16$	$\{7,8\}:17$	$\{9,16,17\}$

For the alternative subsystem, take parent variant 2 of Table S35 and retain vertices 1–17 and 66–68. It is stateless. Each ray deletion instead leaves the positive geometric state count below; restriction of any such state shows that no eighteen-ray subset can be KS.

TABLE S90: Geometric state counts after every single-ray deletion of the twenty-ray parity subsystem.

Deleted ray	States	Deleted ray	States
1	8	2	8
3	8	4	28
5	28	6	8
7	8	8	8
9	12	10	12
11	12	12	10
13	10	14	10
15	10	16	10
17	10	66	8
67	8	68	8

S3. SUPPLEMENTARY PROOFS, REDUCTIONS, AND EXACT VERIFICATION CERTIFICATES

S3.1. Formulas used by the finite certificates

The following formulas restore the explicit conventions used by the later certificates. State indices are local to the corresponding complete table. For Hesse, the six-state vertex-separating family, its seven-state event-separating extension, and the eighteen-state order-determining family are respectively

$$\Omega_6 = \{s_3, s_{31}, s_{75}, s_{89}, s_{113}, s_{151}\}. \tag{S23}$$

$$\Omega_7 = \Omega_6 \cup \{s_{84}\} \tag{S24}$$

$$\begin{aligned}\Omega_{18} = \{&s_2, s_8, s_{20}, s_{30}, s_{40}, s_{45},\\ &s_{54}, s_{64}, s_{73}, s_{79}, s_{83}, s_{93},\\ &s_{105}, s_{109}, s_{119}, s_{123}, s_{127}, s_{150}\}.\end{aligned} \tag{S25}$$

The members of Ω_6 are denoted $\sigma_1, \dots, \sigma_6$ in the displayed order. Their atom profiles and the nontrivial atom–coatom order are

$$\kappa(i) = (\sigma_1(i), \dots, \sigma_6(i)). \tag{S26}$$

$$a_i \le a_j^{\perp} \quad \Longleftrightarrow \quad \{i, j\} \in E(\Gamma(H_{37})). \tag{S27}$$

For H_{65}, the following core coloring extends to the completed hypergraph. A base-compatible independent support S_α is extended canonically using $\varepsilon \in \{0, 1\}$. In the plane formula, η_A, η_B are an integer orthogonal basis and $n_A = \|\eta_A\|^2$, $n_B = \|\eta_B\|^2$.

$$\begin{aligned}(c(1), \dots, c(17)) = (&0, 1, 2, 3, 0, 1, 2, 4, 0, 1, 3,\\ &3, 4, 2, 1, 0, 2).\end{aligned} \tag{S28}$$

$$\begin{aligned}&\big(s_{\alpha,\varepsilon}(w_{m,1}),\ s_{\alpha,\varepsilon}(w_{m,2})\big)\\ &\quad = \begin{cases}(0, 0), & |S_\alpha \cap \{u_m, v_m\}| = 1,\\ (1-\varepsilon,\ \varepsilon), & S_\alpha \cap \{u_m, v_m\} = \emptyset.\end{cases}\end{aligned} \tag{S29}$$

$$\begin{aligned}w_{m,1}(a, b) &= a\,\eta_A + b\,\eta_B,\\ w_{m,2}(a, b) &= -n_B b\,\eta_A + n_A a\,\eta_B\end{aligned} \tag{S30}$$

The eight rays adjoined to completed Yu–Oh and the nine uncovered pairs of the Cabello subsystem are fixed by the following local labels. The last display records the completed historical Γ_1 TITS incidence.

$$\begin{aligned}&(0, 1, 2), (0, 2, -1), (0, 1, -2), (0, 2, 1),\\ &(1, 2, 0), (2, -1, 0), (1, -2, 0), (2, 1, 0).\end{aligned} \tag{S31}$$

$$\begin{aligned}&\{1, 13\}, \{2, 14\}, \{3, 16\}, \{6, 17\}, \{7, 15\},\\ &\{8, 12\}, \{9, c\}, \{10, b\}, \{11, a\}.\end{aligned} \tag{S32}$$

$$E(I_{16,9}) = E(B_{13,7}) \cup \{\{4, 11, 24\}, \{4, 10, 18\}\}. \tag{S33}$$

a. Fractional value of the faithful Cabello completion. The geometric orthogonality graph of $C_{18,9}$ has independence number four and exactly 54 independent four-sets, with each of its 18 rays belonging to twelve of them. Assigning weight 1/12 to each set gives a fractional cover of total weight 9/2, matching 18/4. Fresh pair completion preserves this value by the main completion proposition, proving $\chi_f(C_{36,18}) = 9/2$ as recorded in the main comparison. The complete 54-set certificate is in the electronic audit data.

S3.2. Snark obstructions and quantitative robustness bounds

S3.2.1. Cubic graphs, Blanuša snarks, and Isaacs dot products

For a finite simple cubic graph X, let $H(X)$ be the context hypergraph whose vertices are the edges of X and whose contexts are the three edges incident with each vertex of X. Then $\Gamma(H(X)) = L(X)$ and $\chi(H(X)) = \chi'(X)$. An orthogonal ray assignment for $H(X)$ in $\mathbb{F}^3$ is an assignment of nonzero vectors to the edges of X such that incident edges are orthogonal.

Silva and Ibiapina [5, Question 6] asked whether an orthogonal vector edge coloring in $\mathbb{R}^3$ can exist when $\chi'(X) = 4$. Theorem 11 excludes the Petersen graph P even over $\mathbb{C}$. Using the multipole forcing law of Theorem 11(1), we can extend this exclusion to the two 18-vertex Blanuša snarks [6] and the infinite family of Isaacs dot-product snarks [7].

Recall that for a cubic graph G with two nonadjacent edges $e_1 = \{x_1, y_1\}$ and $e_2 = \{x_2, y_2\}$, the Isaacs dot product $G \cdot P$ is formed by removing e_1, e_2 from G, removing two adjacent vertices u, v from the Petersen graph P (leaving the 4-pole $P \setminus \{u, v\}$ whose dual hypergraph is precisely H_8), and attaching the two dangling edges $\{2, 8\}$ of u to $\{x_1, y_1\}$ and the two dangling edges $\{13, 14\}$ of v to $\{x_2, y_2\}$; the two distinct choices of nonadjacent edges in $G = P$ produce the two Blanuša snarks B_1 and B_2 [6–8].

Corollary 1 (Obstruction for Blanuša and dot-product snarks)**.** *If a cubic graph G admits no orthogonal vector edge coloring in $\mathbb{C}^3$, then neither does any Isaacs dot product $G \cdot P$. In particular, both 18-vertex Blanuša snarks $B_1, B_2 = P \cdot P$ and all iterated dot products $X = P \cdot P \cdots P$ satisfy $\chi(H(X)) = \chi'(X) = 4 > 3 = \omega(L(X))$ and admit no orthogonal vector edge coloring in $\mathbb{C}^3$.*

Proof. In any orthogonal vector edge coloring of $G \cdot P$ in $\mathbb{C}^3$, the restriction to the embedded copy of $H_8 = H(P \setminus \{u, v\})$ satisfies $[v_2] = [v_8]$ and $[v_{13}] = [v_{14}]$ by Theorem 11(1). Assigning the common ray $[v_2] = [v_8]$ to $e_1 = \{x_1, y_1\}$ and the common ray $[v_{13}] = [v_{14}]$ to $e_2 = \{x_2, y_2\}$ preserves orthogonality at x_1, y_1, x_2, y_2 and therefore yields a valid orthogonal vector edge coloring of G in $\mathbb{C}^3$. Since P admits no such coloring by Theorem 11, induction excludes B_1, B_2 and all iterated dot products with P. □

Generalized Petersen graphs [9] and flower snarks provide further families for comparing real, complex, injective, and edge-faithful realizability.

S3.2.2. Explicit analytic lower bound and tetrahedral tight-frame upper bounds for Δ_3

Let A be the 30 edges of $\Gamma(G_{32})$ and define the minimum total squared orthogonality error in $\mathbb{C}^3$:

$$\Delta_3 = \min_{\substack{v_i \in \mathbb{C}^3 \\ \|v_i\|=1 \ (1 \le i \le 15)}} \sum_{\{i,j\} \in A} |\langle v_i, v_j \rangle|^2. \tag{S34}$$

By compactness of $(S^5)^{15}$ and Theorem 11, $\Delta_3 > 0$. We now establish both a certified rational analytic lower bound and explicit tetrahedral tight-frame upper bounds.

Proposition 2 (Quantitative bounds on Δ_3)**.** *Every assignment of unit vectors $v_1, \ldots, v_{15} \in \mathbb{C}^3$ satisfies*

$$\varepsilon := \max_{\{i,j\} \in A} |\langle v_i, v_j \rangle| > \frac{1}{120}, \tag{S35}$$

and the minimum total squared error in Eq. (S34) satisfies

$$\begin{aligned} \frac{1}{14400} < \Delta_3 &\le \min_{a^2+b^2+c^2=1} F(a, b, c) \\ &\le \frac{9275914990340646}{10001032226635921} < 0.927496 < \frac{18}{19}, \end{aligned} \tag{S36}$$

where

$$F(a,b,c) = 12a^2 + 12(ab+bc-ca)^2 + 6(b^2-a^2-c^2)^2. \tag{S37}$$

Proof. To prove the lower bound (S35), let $\varepsilon = \max_{\{i,j\}\in A} |\langle v_i, v_j\rangle|$. For any context $e = \{x,y,z\}$, the Gram matrix of $\{v_x, v_y, v_z\}$ has diagonal entries 1 and off-diagonal entries of modulus at most ε, so Gershgorin's circle theorem implies $\|I_3 - (v_x v_x^\dagger + v_y v_y^\dagger + v_z v_z^\dagger)\|_{\mathrm{op}} \le 2\varepsilon$. Hence for any unit vectors $u, w \in \mathbb{C}^3$,

$$\Big|1 - \big(|\langle u,x\rangle|^2 + |\langle u,y\rangle|^2 + |\langle u,z\rangle|^2\big)\Big| \le 2\varepsilon, \tag{S38}$$

and if $\{u,x\}, \{w,z\} \in A$, then

$$\big|\langle u,w\rangle - \langle u,y\rangle\langle y,w\rangle\big| \le 4\varepsilon. \tag{S39}$$

In $\{5,6,7\}$ and $\{9,10,11\}$, the vectors $w_7 = v_{12} - v_5\langle v_5, v_{12}\rangle$ and $w_9 = v_4 - v_{11}\langle v_{11}, v_4\rangle$ satisfy $\|w_7 - v_7\langle v_7, v_{12}\rangle\| \le 3\varepsilon$ and $\|w_9 - v_9\langle v_9, v_4\rangle\| \le 3\varepsilon$, so $|\langle w_7, w_9\rangle| \le 7\varepsilon$. Expanding,

$$\begin{aligned}\langle w_7, w_9\rangle = {} & \langle v_{12}, v_4\rangle - \langle v_{12}, v_{11}\rangle\langle v_{11}, v_4\rangle - \langle v_{12}, v_5\rangle\langle v_5, v_4\rangle \\ & + \langle v_{12}, v_5\rangle\langle v_5, v_{11}\rangle\langle v_{11}, v_4\rangle.\end{aligned}$$

The two middle terms are at most ε in modulus, since $|\langle v_{12}, v_{11}\rangle| \le \varepsilon$ and $|\langle v_5, v_4\rangle| \le \varepsilon$. Eq. (S39) on $\{1,2,3\}$ replaces each of $\langle v_{12}, v_4\rangle$, $\langle v_{12}, v_5\rangle$, $\langle v_5, v_{11}\rangle$, and $\langle v_{11}, v_4\rangle$ by its product through v_2 with error at most 4ε, and hence the triple product with error at most 12ε, since all factors have modulus at most one. The main terms combine to

$$\big(1 + |\langle v_2, v_5\rangle|^2 |\langle v_2, v_{11}\rangle|^2\big)\langle v_{12}, v_2\rangle\langle v_2, v_4\rangle,$$

whose real prefactor is at least one, so

$$\big|\langle v_{12}, v_2\rangle\langle v_2, v_4\rangle\big| \le (7+2+4+12)\varepsilon = 25\varepsilon.$$

By reflection symmetry, we may assume $|\langle v_2, v_4\rangle| \le 5\sqrt{\varepsilon}$. Applying Eq. (S38) to v_4 in $\{1,2,3\}$ and to v_2 in $\{3,4,5\}$ gives $|\langle v_1, v_4\rangle|^2 \ge 1 - 27\varepsilon - \varepsilon^2$ and $|\langle v_2, v_5\rangle|^2 \ge 1 - 27\varepsilon - \varepsilon^2$, so $\min_\theta \|v_1 - e^{i\theta} v_4\| \le \sqrt{54\varepsilon + 2\varepsilon^2}$ and $\min_\theta \|v_2 - e^{i\theta} v_5\| \le \sqrt{54\varepsilon + 2\varepsilon^2}$. Since $|\langle v_4, v_{10}\rangle| \le \varepsilon$ and $|\langle v_5, v_7\rangle| \le \varepsilon$, Eq. (S38) applied to v_1 in $\{9,10,11\}$ and to v_2 in $\{7,8,9\}$ gives

$$|\langle v_i, v_9\rangle|^2 \ge 1 - 2\varepsilon - \varepsilon^2 - \big(\varepsilon + \sqrt{54\varepsilon + 2\varepsilon^2}\big)^2 \qquad (i = 1,2).$$

Comparing the sum of these two inequalities with $|\langle v_1, v_9\rangle|^2 + |\langle v_2, v_9\rangle|^2 \le 1 + 2\varepsilon$ from $\{1,2,3\}$ gives

$$1 \le 6\varepsilon + 2\varepsilon^2 + 2\big(\varepsilon + \sqrt{54\varepsilon + 2\varepsilon^2}\big)^2.$$

The right-hand side is strictly increasing for $\varepsilon > 0$, and at $\varepsilon = 1/120$ we have $54\varepsilon + 2\varepsilon^2 = 3241/7200 < (27/40)^2$, so the right-hand side is strictly bounded above by

$$\frac{6}{120} + \frac{2}{120^2} + 2\left(\frac{1}{120} + \frac{27}{40}\right)^2 = \frac{1417}{1440} < 1,$$

certifying $\varepsilon > 1/120$ and $\Delta_3 \ge \varepsilon^2 > 1/14400$.

For the upper bounds, set $v_4 = e_1$, $v_8 = e_2$, $v_{12} = e_3$, and for any unit vector $(a,b,c) \in S^2$, define the remaining 12 unit vectors as representatives of the tetrahedral A_4-orbit of rays:

$$\begin{aligned} v_1 &= (-b,-c,-a)^T, & v_2 &= (c,-a,b)^T, \\ v_3 &= (-a,-b,c)^T, & v_5 &= (-a,b,c)^T, \\ v_6 &= (-b,c,-a)^T, & v_7 &= (c,a,b)^T, \\ v_9 &= (-c,-a,b)^T, & v_{10} &= (-a,b,-c)^T, \\ v_{11} &= (b,-c,-a)^T, & v_{13} &= (a,b,c)^T, \\ v_{14} &= (b,c,-a)^T, & v_{15} &= (-c,a,b)^T. \end{aligned} \tag{S40}$$

These 15 vectors form a tight frame $\sum_{i=1}^{15} v_i v_i^T = 5I_3$, and the sum of squared overlaps on the 30 edges of $\Gamma(G_{32})$ equals $F(a,b,c)$ in Eq. (S37). Evaluating Eq. (S40) at the normalized integer direction $(a,b,c) = (1,3,-3)/\sqrt{19}$ yields rational squared overlaps with $\Delta_3 \le 342/361 = 18/19 \approx 0.947368$ and $\max_{\{i,j\}\in A} |\langle v_i, v_j\rangle|^2 = 1/19$; evaluating at the refined normalized integer direction $(a,b,c) = (2332, 7101, -6644)/\sqrt{100005161}$ yields the certified rational bound in Eq. (S36). □

Numerical minimization of F on S^2 (as well as unrestricted local searches over $(S^5)^{15}$) finds an attained candidate minimum near 0.927495724691 at $(a,b,c) \approx (0.23319713, 0.71005263, -0.66441280)$; certifying whether the unrestricted global minimum Δ_3 coincides with $\min_{S^2} F$ remains an open optimization problem. Choosing instead the rational unit vector $(a,b,c) = \frac{1}{3}(1,2,-2)$ in Eq. (S40) makes $ab + bc - ca = 0$, so four of the ten contexts of G_{32} ($\{1,2,3\}, \{5,6,7\}, \{9,10,11\}, \{13,14,15\}$) become exact orthonormal bases of $\mathbb{R}^3$, all 15 rays are distinct, and $\Delta_3 \le 38/27$ with $\max_{\{i,j\}\in A} |\langle v_i, v_j\rangle| = 1/3$.

S3.3. Hesse H_{37}: minimum state families

All state indices in this appendix use the canonical enumeration of Appendix S2.27. The minima for unital vertex separation and event separation have the counting lower bounds $2^5 - 1 < 37$ and $2^6 < 76$. The families in Eqs. (S23) and (S24) attain those bounds: their profiles give respectively 37 distinct nonzero atom profiles and 76 distinct event profiles. The seven-state family still has 462 ordered event pairs for which inclusion of the profiles does not imply the lattice order. The family in Eq. (S25) has no such pair and hence gives the upper bound eighteen for order determination. This appendix supplies a computer-assisted proof of the matching lower bound.

S3.3.1. Reduction to families of seventeen states

Let $D = \{P_0, P_1, P_2, Q_0, Q_1, Q_2\}$. For each unordered pair $\{u, v\} \subset D$, order determination requires a state with $s(u) = s(v) = 1$: the pair is nonorthogonal, so $u \not\le v^\perp$. Every state takes value one on at most two members of D. Consequently the fifteen pairs require fifteen different states. Table S91 gives their complete witness classes. These classes are disjoint, their union T has 27 states, and a choice of one witness from each class has $1^3 2^{12} = 4{,}096$ possibilities.

TABLE S91: Complete joint-one witness classes for the fifteen pairs in D. Each entry lists all canonical indices r for which $s_r(u) = s_r(v) = 1$.

Pair $\{u, v\}$	Witness indices r
$\{P_0, P_1\}$	150
$\{P_0, P_2\}$	73
$\{P_0, Q_0\}$	93, 145
$\{P_0, Q_1\}$	127, 133
$\{P_0, Q_2\}$	79, 99
$\{P_1, P_2\}$	2
$\{P_1, Q_0\}$	105, 139
$\{P_1, Q_1\}$	14, 119
$\{P_1, Q_2\}$	40, 48
$\{P_2, Q_0\}$	54, 60
$\{P_2, Q_1\}$	8, 26
$\{P_2, Q_2\}$	20, 68
$\{Q_0, Q_1\}$	109, 113
$\{Q_0, Q_2\}$	83, 87
$\{Q_1, Q_2\}$	30, 34

The nine nonorthogonal pairs $\{z, C_{ak}\}$ require further joint-one witnesses. No state in T witnesses any of them. Indeed, if a state activates a P vertex then $s(z) = 0$. If it instead activates two Q vertices, the state equations force the selected M vertices into the remaining Q column, and all nine C values are zero. Any state outside T witnesses at most six of the nine pairs: for $s(z) = 1$, the three selected M vertices force at least three C values to vanish. Thus at least two states outside T are necessary.

It follows that an order-determining family of at most seventeen states would have exactly seventeen members: one from each of the fifteen classes in Table S91, and two outside T that together witness all nine $\{z, C_{ak}\}$ pairs. Enumerating the $\binom{126}{2}$ unordered pairs outside T leaves exactly 132 with this property. The reduction therefore leaves precisely

$$4{,}096 \cdot 132 = 540{,}672 \tag{S41}$$

candidate families. The nine pairs alone do not require three extra states; for example, s_1 and s_{72} jointly witness all nine. The remaining order requirements must also be checked.

S3.3.2. Exhaustive order certificate

For each of the 5,407 ordered event pairs $x \not\leq y$, define its nonempty witness set

$$W_{xy} = \{r \in \{1, \ldots, 153\} : s_r(x) = 1,\ s_r(y) = 0\}. \tag{S42}$$

A family is order determining exactly when its index set meets every W_{xy}. Repeated sets and proper supersets can be discarded, because meeting an inclusion-minimal set also meets every set containing it. There are exactly 81 inclusion-minimal witness sets. The exhaustive check forms each of the 540,672 families in Eq. (S41) and counts the minimal witness sets that it meets. The largest count is 75, strictly below 81. Hence no family of seventeen states is order determining. Together with Ω_{18}, this proves the minimum eighteen.

The accompanying Hesse extension checker, invoked by the joint entry point, reconstructs the contexts and all 153 states from their symbolic parametrization, checks their agreement with `hesse37-data.json`, and performs this finite search using integer bit sets and the Python standard library. No numerical tolerance, optimization solver, or unreported search cutoff is used. It also verifies the printed seven- and eighteen-state families and every entry of Table S91. From the extracted joint project directory, run the joint entry point, without Python's optimization option:

```
python verification/verify-integrated.py
```

The generated file `verification/_work/hesse/hesse37-extensions-certificate.json` records all 81 minimal witness sets, a representative event pair for each, all 132 possible extra pairs, the coverage-count distribution over the 540,672 candidates, and all event profiles for Ω_6, Ω_7, and Ω_{18}. The generated file `verification/_work/hesse/hesse37-order18-witnesses.csv` records a witness in Ω_{18} for every $x \not\leq y$. These two files use the same numbered event labels as the full-family certificate in Appendix S2.27.6.

The program additionally checks Berge girth five from the incidence graph, computes every meet and join in the 76-event order, verifies the orthomodular law, and verifies the projector sum in Eq. (30) exactly in $\mathbb{Q}(\zeta)$. The nonexistence of an injective real representation is proved analytically in Sec. V A.

S3.4. Four-dimensional $H_{\mathrm{YO}}^{(4)}$ and H_{65}: completion and verification

The coordinates in Table S17 can be reproduced by the following finite selection rule. Normalize an integer vector by dividing by the greatest common divisor of its entries and choosing the sign so that its first nonzero entry is positive. Form the set of normalized nonzero vectors obtained from $\{-3, -2, \ldots, 3\}^4$, ordered by squared norm, then maximum absolute entry, then lexicographically. Process bridge edges in the order $e_{13}, \ldots, e_{24}, e_1, \ldots, e_{12}$. For each pair, choose η_A to be the first vector in this ordered pool orthogonal to its endpoints, and η_B the first such vector also orthogonal to η_A.

For all $(a, b) \in \{-7, -6, \ldots, 7\}^2 \setminus \{(0, 0)\}$, form the pair in Eq. (S30) and normalize both vectors. Discard pairs that repeat an already placed ray or have an orthogonality to any already placed vertex other than the two endpoints. Among the remaining ordered pairs choose the lexicographically smallest score

$$\left(\max_{j,k} |(w_{m,j})_k|,\ \|w_{m,1}\|^2 + \|w_{m,2}\|^2,\ w_{m,1}, w_{m,2} \right). \tag{S43}$$

In Eq. (S43), the order on the last two entries is the usual lexicographic order on their integer coordinates. This rule yields exactly the 48 completion vectors in Table S17. It is a recipe for this specific example; the general existence assertion is established independently by Lemma 21.

The companion verifier parses the integrated coordinate and state tables, reproduces this completion rule and all 393,216 state extensions, and checks the maximal cliques, fractional-coloring certificate, atom-order failure, forced hub-completion zero, and warm-up data. It is included with the complete state file and invoked by the joint entry point in Sec. S3.7.

S3.5. Exact Cabello completion and forced-zero certificates

For each of the nine uncovered pairs in Eq. (S32), rational row reduction verifies that its two endpoint vectors span no third core ray. If two endpoint planes were orthogonal, their four endpoints would form another full basis. Exact four-clique enumeration gives only the original nine bases, all of whose pairs are already covered. This excludes such an orthogonal-plane pair. The integer vectors in Table S28 directly verify the resulting faithful completion: 36 distinct

projective rays, eighteen full bases, and 108 orthogonal pairs, with every pair covered by exactly one listed basis. The nine-context graph and the full eighteen-ray orthogonality graph both fail four-colorability and admit five-colorings, as checked by exact finite search in the companion certificate. Lemma 21(2) transfers the chromatic value to the completion.

For the vertex-deleted core with c removed, rational row reduction finds exactly the span failures

$$v_{16} \in \mathrm{span}(v_1, v_2), \qquad v_{17} \in \mathrm{span}(v_7, v_8).$$

There is no orthogonal-plane-pair obstruction. The finite-avoidance choice in Lemma 21 can therefore avoid all extra incidences except those forced by these two spans; the explicit 45-ray inventory has exactly the four pairs printed in Sec. S1.17.2. The two fresh vectors completing $\{1, 2\}$ must both be orthogonal to 16, and those completing $\{7, 8\}$ must both be orthogonal to 17.

The twelve rows of Table S84 give a short complete proof of the resulting obstruction. Whenever 16 is selected, neither 1 nor 2 is selected. The context completing $\{1, 2\}$ then needs one fresh value equal to one, while geometric exclusion with 16 requires both to be zero. Hence every geometric extension has $s(16) = 0$. The same argument with $\{7, 8\}$ gives $s(17) = 0$. These implications use the displayed core states in addition to the span incidences; the span incidences alone would not imply them. The retained basis $\{10, 11, 16, 17\}$ gives $s(10) + s(11) = 1$. If $s(9) = 1$, the other two hub bases give $s(10) = s(11) = 0$, a contradiction. Thus 9, 16, and 17 have identical zero profiles.

Conversely, each of the eight indicated core rows avoids 16 and 17, so all its fresh-pair choices satisfy the four extra exclusions. Each misses eight bridges. This proves the exact count $8\,2^8 = 2{,}048$, as well as failure of geometric unitality and separation. The two excluded core rays remain impossible for every fresh-pair completion of these fixed planes; additional accidental orthogonalities cannot restore them. Table S89 supplies the individual span checks for all eighteen deletions. Their full core tables, exact integer completions, and extension counts are included in `data/cabello18-review-checks.json`.

To audit the ancestry result, enumerate every four-clique after adjoining either three-ray choice to all 65 original rays. In both cases there are exactly 35 bases, 195 geometric pairs, and 192 context-covered pairs; the maximum basis intersection is two. The first union contains the displayed eighteen-ray parity proof. The second union contains the twenty-ray proof with multiplicities $4, 4, 2, \ldots, 2$ and eleven contexts. Each of its twenty ray deletions has a compatible assignment by Table S90. All twenty rays lie in Peres's 24-ray set. Enumerating the $\binom{24}{4} = 10{,}626$ twenty-ray subsets of that set, with all bases they contain, gives exactly 96 ray-critical KS subsets, each with eleven bases and all mutually isomorphic; the twenty-ray proof is one of them. Since Peres's set contains Kernaghan's critical twenty-ray set and 95 others [3], the twenty-ray proof is isomorphic to Kernaghan's [10]. All coordinates, full basis lists, and the preserved H_{65} restriction are checked independently of the unrestricted ancestry existence proposition.

S3.6. Transverse extension and criticality certificates

For $d = 3$, number the two copies of an n-vertex input as i and $n + i$, and the fresh transverse vertex as $2n + i$. The transverse contexts are $\{i, n + i, 2n + i\}$ and the horizontal contexts are the two numbered copies of the original list. This explicitly specifies all 45 vertices and 33 contexts for $T(H_9)$, and all 75 vertices and 57 contexts for $T(H^+_{\mathrm{YO}})$.

For each deleted original context e, a strong three-coloring of $H \setminus \{e\}$ having two equal colors and a singleton color on e gives a state after deletion of one horizontal copy of e. Use the repeated-color indicator on the deleted copy and the singleton-color indicator on the intact copy. Both have exactly one active vertex in each of their required contexts and their supports are disjoint; the remaining transverse values are forced. For deletion in the other copy, interchange the two indicators. Table S92 supplies an explicit coloring for every required context of both inputs. The color strings use original vertex order, and their labels 0,1,2 may be permuted.

TABLE S92: Explicit deletion colorings with pattern (1,1,2) and the number of states after deleting the corresponding first-copy horizontal context.

Input	Deleted original context	Color string	Extension states
H_9	$\{1,2,3\}$	212010201201122	20
H_9	$\{1,11,12\}$	201020120121221	20
H_9	$\{2,8,15\}$	021020120121220	12
H_9	$\{3,4,5\}$	012120102012212	20
H_9	$\{4,10,13\}$	012102120121000	12
H_9	$\{5,6,7\}$	012010120121220	20
H_9	$\{6,12,14\}$	012102102012202	12
H_9	$\{7,8,9\}$	012010200121222	20
H_9	$\{9,10,11\}$	012010201121222	20
H^+_{YO}	$\{1,2,3\}$	200021210110202010201122221	16
H^+_{YO}	$\{1,10,19\}$	210021201212020001011202l	8
H^+_{YO}	$\{1,13,20\}$	120210201012120011200220l	8
H^+_{YO}	$\{2,5,8\}$	021021012120201122211001 2	24
H^+_{YO}	$\{3,11,14\}$	012021102120200122210001 1	8
H^+_{YO}	$\{3,12,25\}$	021210102101222020210101 0	8
H^+_{YO}	$\{4,5,6\}$	021200210102121110220221 0	16
H^+_{YO}	$\{4,10,18\}$	021210201202121010120201 0	8
H^+_{YO}	$\{4,11,24\}$	021201012120201120211000 2	8
H^+_{YO}	$\{6,12,17\}$	021210102100222110210101 2	8
H^+_{YO}	$\{6,13,22\}$	021012102120200112210001 2	8
H^+_{YO}	$\{7,8,9\}$	021012002120101112222001 2	16
H^+_{YO}	$\{7,11,15\}$	021201012100221120211001 2	8
H^+_{YO}	$\{7,13,21\}$	021012102120100112220001 2	8
H^+_{YO}	$\{9,10,23\}$	021210201102121010220201 0	8
H^+_{YO}	$\{9,12,16\}$	021210102102222010210101 0	8

Deleting one transverse context T_v instead permits an ordered pair of original state supports S_1, S_2 exactly when $S_1 \cap S_2 \subseteq \{v\}$. In these non-three-colorable inputs disjoint pairs do not exist, so the intersection must equal $\{v\}$; the isolated fresh vertex has two choices. For H_9 this gives four states for each $v = 1, \ldots, 12$ and twelve states for each $v = 13, 14, 15$. Together with Table S92, this proves criticality of all 33 contexts. For completed Yu–Oh it gives zero states at $v = 1, \ldots, 13$ and eight states at $v = 14, \ldots, 25$. Thus horizontal criticality does not imply criticality of this full transverse construction. The corresponding G_{32} extension has 45 vertices and 35 contexts and is stateless. Its distinct six state supports have pairwise intersection size one; for H_9 the distinct eight supports have intersection size one in 21 pairs and size three in seven pairs.

S3.7. Data and verification

The companion electronic package contains coordinate and context inventories, complete state and coloring enumerations, verification programs, exact parent inventories, and finite certificates. Vertex labels and state orderings agree with the tables in this supplement. The verification programs check the common Yu–Oh numbering, the published 49-ray parent map, the parent vectors and contexts, the full H_{65} state inventory, and cross-reference targets. The finite lower-bound certificate for the eighteen-state Hesse minimum is the exhaustive 540,672-candidate argument of Sec. S3.3.

Run the companion verification entry point from the project directory:

```
python verification/verify-integrated.py
```

The exact checks use rational or $\mathbb{Q}(\zeta)$ arithmetic, with $\zeta^2 + \zeta + 1 = 0$. Reports and hashes accompany the source.

The finite checks certify the instances and certificates they explicitly test. The analytic realization and existence results also rely on their proofs in the text.

S3.8. Historical Kochen–Specker gadget certificates (1965 Fig. 1 and 1967 Γ_1)

S3.8.1. The 1965 triangles and the completed historical TITS

Identifying the repeated occurrences of vertices 1 and 2 in Kochen–Specker's 1965 Fig. 1 gives

$$\{134, 325, 2\,9\,11, 11\,1\,10, 479, 5\,8\,10, 678\}. \tag{S44}$$

These are also the triples used in the displayed function definitions on p. 182 of Ref. [11]. Their four state supports are

$$\{1,5,6,9\}, \quad \{2,4,6,10\}, \quad \{3,6,9,10\}, \quad \{4,5,6,11\}.$$

The TIFS between 1 and 2 is nonvacuous. Directly, if both were true, contexts 134, 325, 2 9 11, and 11 1 10 would make 3,4,5,9,10,11 false; contexts 479 and 5 8 10 would then make both 7 and 8 true, contradicting 678.

Every state has 6 true and 7,8 false, so no three-coloring can exist: the three color-class states of any coloring would need to be disjoint. The graph is abstract rather than an injective ray example. In a purported injective realization in $\mathbb{C}^3$, vertices 1 and 9 are both orthogonal to 4 and 11. Distinct rays 4 and 11 span a plane, whose orthogonal complement is one-dimensional; thus rays 1 and 9 coincide. This proves the obstruction even over the complex field. No claim that the 1965 authors presented this graph as a Hilbert-space realization is intended.

For the 1967 Γ_1[12, pp. 68,70], the correspondence between its original vertices and $I_{16,9}$ is

Γ_1	a_0	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9
$I_{16,9}$	10	9	1	7	3	8	2	11	4	24

The already complete triangles are $\{a_1, a_3, a_5\}$, $\{a_2, a_4, a_6\}$, and $\{a_7, a_8, a_9\}$. The six remaining edges

$$a_0a_1, \ a_0a_2, \ a_0a_8, \ a_3a_7, \ a_4a_7, \ a_5a_6$$

are completed by fresh vertices $23, 19, 18, 15, 14, 5$, respectively. The resulting nine contexts are exactly Eq. (S33). This also identifies its directed TITS $a_0 \to a_9$ with $10 \to 24$.

---

[1] M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
[2] J. Bub, Foundations of Physics **26**, 787 (1996).
[3] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
[4] K. Svozil and J. Tkadlec, Journal of Mathematical Physics **37**, 5380 (1996).
[5] A. Silva and A. Ibiapina, On orthogonal vector edge coloring (2019), arXiv:1909.01918 [cs.DM].
[6] D. Blanuša, Glasnik Matematičko-Fizički i Astronomski, Serija II **1**, 31 (1946).
[7] R. Isaacs, The American Mathematical Monthly **82**, 221 (1975).
[8] G. Brinkmann, M. Preissmann, and D. Sasaki, Discrete Mathematics & Theoretical Computer Science **17**, 369 (2015).
[9] G. Schrag, Discrete Mathematics **70**, 185 (1988).
[10] M. Kernaghan, Journal of Physics A: Mathematical and General **27**, L829 (1994).
[11] S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
[12] S. Kochen and E. P. Specker, Journal of Mathematics and Mechanics **17**, 59 (1967).