Institute for Theoretical Physics
TU Wien
Wiedner Hauptstraße 8–10/136
1040 Vienna, Austria
karl.svozil@tuwien.ac.at

November 12, 2026

The Editors
International Journal of Theoretical Physics

Re: Correction to M. Navara and K. Svozil, "Construction of Kochen–Specker Sets from Mutually Unbiased Bases," Int. J. Theor. Phys. **65**, 322 (2026), `https://doi.org/10.1007/s10773-026-06472-5`

Dear Editors,

On behalf of both authors, I ask you to publish a Correction (erratum) to the article above. After publication we rechecked all calculations in exact arithmetic and found the errors listed below.

None of them affects the main results. The 165 rays and 130 contexts, the three 69-ray, 50-context lineages with 21 common rays and 10 common contexts, the absence of two-valued states for each lineage, and the state counts 24, 36, 36 and 16 all remain correct.

Errors and corrections

1. **Fig. 2(b).** The node labeled v_{1423} carries the wrong ray. $v_{1423} = (1, -1, -1, 1)$ is not in Cabello's set and is not orthogonal to the other rays of its two contexts; for instance, $\langle v_{13}, v_{1423} \rangle = 2$. The correct ray is $(1, 1, -1, 1) = \{v_{13}, v_{23}, w_{12}\}^{\perp}$, which is Cabello's v_{47} in Table 4.
2. **Fig. 1 versus Tables 1–3.** The labels e_1, e_2, e_3 are cyclically permuted between Fig. 1 and the tables. In Fig. 1, e_i lies in the context $\{c_i, b_{jk}, e_i\}$; in the tables, e_1 is paired with c_2, e_2 with c_3 and e_3 with c_1. We relabel the tables so that $e_i = c_i \times b_{jk}$, in agreement with Fig. 1; for example, $e_{11} = (2, 1, 1)$. The rays and contexts themselves are unchanged.
3. **Table 4.** Cabello's labels v_{12}, v_{23} and v_{34} coincide with gadget labels that denote different vectors. For example, Cabello's v_{12} is $(1, 0, 0, 0)$, whereas our v_{12} is $(0, 0, 1, -1)$. We now write Cabello's labels with a superscript C, as v_{ij}^{C}, the ray shared by his contexts i and j.
4. **Sect. 4.2.2, last paragraph (p. 12).** The statement that the six pair contexts and three connector contexts "form the 10-context scaffold that enforces Proposition 2" is too strong. Proposition 2 relies on the prescribed coordinates of v_{1234}, v_{1324} and v_{1423}.
   - The incidences of these 10 contexts alone do not force u to be unbiased. For any u with nonzero coordinates they are realized with $v_{1234} \propto (s_{34}x_1, s_{34}x_2, -s_{12}x_3, -s_{12}x_4)$, where $s_{kl} = |x_k|^2 + |x_l|^2$, and analogously for v_{1324} and v_{1423}.
   - The complete 17-context hypergraph does force maximal unbiasedness by its incidences alone, through the contexts $\mathcal{E}_1$ and $\mathcal{E}_3$.
   - In $D = 5$, Proposition 3 likewise uses the prescribed coordinates of the connector rays g_i.
5. **Fig. 2 caption.** "Solid curves show imposed contexts" should read: solid curves show the 13 contexts of $\mathfrak{B}_{13}$, namely the 10 imposed contexts and the three disjoint-support contexts

$\mathcal{B}_{1234}$, $\mathcal{B}_{1324}$ and $\mathcal{B}_{1423}$.

6. **References.** Ref. [3] (Trandafir and Cabello) was published as Phys. Rev. A **111**, 022408 (2025). Ref. [24] (McNulty and Weigert) has since appeared in Quantum **10**, 2051 (2026).

Corrected version

We will post the corrected article on arXiv as version 3 of arXiv:2509.08636. It also contains clarifications that need no erratum: the 25 generation rules of Table 1 are stated explicitly, and the text now says that the 130 contexts are all orthogonal triples among the 165 rays. It also notes that Cabello's nine contexts are all the orthogonal quadruples among his 18 rays.

The corrected manuscript, with all changes marked in blue, is attached.

Sincerely,

Karl Svozil
on behalf of both authors