

Construction of Kochen–Specker Sets from Mutually Unbiased Bases

Mirko Navara^{1,*} and Karl Svozil^{2,†}

¹*Faculty of Electrical Engineering, Czech Technical University in Prague,
Technická 2, CZ-166 27 Prague 6, Czech Republic*

²*Institute for Theoretical Physics, TU Wien, Wiedner Hauptstrasse 8-10/136, 1040 Vienna, Austria*

We give a constructive analysis of Kochen–Specker (KS) contextuality at the level of complete orthogonal bases (contexts). In dimension three, three noncomputational mutually unbiased bases generate a union of 165 rays and 130 contexts. Exact provenance accounting gives a symmetric decomposition: the three 69-ray lineages share 21 rays and 10 contexts, while each contributes 48 exclusive rays and 40 lineage-specific contexts. Each 69-ray, 50-context lineage is a KS configuration, and this framework clarifies the relation among the Yu–Oh, Harding–Salinas Schmeis, and Cabello constructions. In dimensions four and five, we give parametrized orthogonality gadgets whose connector contexts force a central ray to be maximally unbiased relative to the computational basis. For the concrete four-dimensional example, the 20-ray gadget and the Cabello 18-ray set are informationally equivalent subsets of the 24-ray Peres–Mermin eigensystem, although their context hypergraphs have different two-valued-state behavior. The results show why both the rays and the complete incidence structure of their contexts are needed in assessments of KS contextuality.

Keywords: Quantum contextuality, Kochen–Specker theorem, Mutually unbiased bases, Quantum logic, Orthogonality hypergraphs, Quantum foundations

Changes made in response to the referee reports are shown in blue.

I. INTRODUCTION

The Kochen–Specker (KS) theorem shows that certain finite families of quantum propositions do not admit a context-independent assignment of classical truth values compatible with their orthogonality relations [2]. A finite KS proof is therefore specified not only by a set of rays but also by the complete bases in which those rays occur.

Definition 1 (Rays and contexts) *A ray is a one-dimensional subspace $[v] = \{\lambda v : \lambda \in \mathbb{C} \setminus \{0\}\}$. A context (block) in $\mathbb{C}^D$ is a set of D mutually orthogonal rays spanning the space. In the associated D -uniform hypergraph the rays are vertices and the contexts are hyperedges. A ray is intertwining if it belongs to at least two contexts; a ray occurring in exactly one context is nonintertwining.*

Definition 2 (Two-valued states) *For a context hypergraph (V, E) , a two-valued state is a map $s : V \rightarrow \{0, 1\}$ satisfying $\sum_{v \in C} s(v) = 1$ for every $C \in E$. A coordinatized hypergraph with no two-valued state is a KS configuration. A family of two-valued states is separating if every pair of distinct rays receives different values in at least one state; it is unital if every ray receives the value 1 in at least one state.*

These definitions distinguish the logical KS criterion from resource measures used in inequalities, chromatic

formulations, or nonlocal games [13, 14, 25]. Reduced incidence structures can be useful in those settings, but deleting singly occurring rays or omitting available contexts can change the set of two-valued states. For example, the seven-ray subconfiguration discussed in Ref. [15, Fig. 6(a)] has different classical-state behavior from its thirteen-ray completion. We therefore state explicitly which context hypergraph is being analyzed at each step.

The paper has two main contributions:

1. In $\mathbb{C}^3$, three noncomputational mutually unbiased bases (MUBs), together with a fixed family of algebraic generation maps, produce 225 labelled vectors. Projective identification reduces these to 165 rays, and an exhaustive orthogonality search gives 130 contexts. The provenance decomposition is symmetric: 21 rays and 10 contexts are common to all three 69-ray lineages, while each lineage has 48 exclusive rays and 40 lineage-specific contexts. Each lineage contains 69 rays and 50 contexts and is uncolorable.
2. In $D = 4$ and $D = 5$, connector contexts convert orthogonality into linear equations for $|x_j|^2$, forcing a central vector $u = (x_1, \dots, x_D)$ to be maximally unbiased relative to the computational basis. In the concrete $D = 4$ case, the resulting 20-ray set and the Cabello 18-ray KS set are mutually reconstructible inside the 24-ray Peres–Mermin eigensystem [16, 17, 19–21]; nevertheless, their context hypergraphs have different two-valued-state spaces.

Section II relates the putrit construction to the Yu–Oh, Harding–Salinas Schmeis, and Cabello diagrams and gives the complete ray and context inventories. Section IV separates the higher-dimensional forcing constraints from their coordinate completions and then com-

* navara@fel.cvut.cz; <https://cmp.felk.cvut.cz/~navara>

† karl.svozil@tuwien.ac.at; <http://tph.tuwien.ac.at/~svozil>

TABLE I. *Systematic generation of the vector set for contextuality analysis.* This table details the complete set of 225 vectors that form the basis of our structural analysis. A family of 25 algebraic functions (rows) is applied to nine initial vectors (columns $u_1 \dots u_9$). These initial vectors comprise all three MUBs in $\mathbb{C}^3$: the Fourier basis $\mathcal{B}_1 = \{u_1, u_2, u_3\}$, and two related MUBs, $\mathcal{B}_2 = \{u_4, u_5, u_6\}$ and its complex conjugate $\mathcal{B}_3 = \{u_7, u_8, u_9\}$; all yielding 69-50 KS sets. The final column counts the number of unique rays produced by each function, highlighting the differing projective variety of the constructions. The 165 unique rays drawn from this set form the labels for identifying orthogonal bases, hyperedges, or contexts. Labelling according to the Harding and Salinas Schmeis scheme [27]; with equivalences to the Yu–Oh numbering scheme [30].

basis	$\mathcal{B}_1$			$\mathcal{B}_2$			$\mathcal{B}_3$			#
label	u_1	u_2	u_3	u_4	u_5	u_6	u_7	u_8	u_9	
$a_1 \equiv z_1$	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	(1, 0, 0)	1
$a_2 \equiv z_2$	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	(0, 1, 0)	1
$a_3 \equiv z_3$	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	(0, 0, 1)	1
$u \equiv h_0$	(1, 1, 1)	(1, ω , ω^2)	(1, ω^2 , ω)	(1, ω , ω)	(1, ω^2 , 1)	(1, 1, ω^2)	(1, ω^2 , ω^2)	(1, ω , 1)	(1, 1, ω)	9
$b_1 \equiv y_+^1$	(0, 1, 1)	(0, ω , ω^2)	(0, ω^2 , ω)	(0, ω , ω)	(0, ω^2 , 1)	(0, 1, ω^2)	(0, ω^2 , ω^2)	(0, ω , 1)	(0, 1, ω)	3
$b_2 \equiv y_+^2$	(1, 0, 1)	(1, 0, ω^2)	(1, 0, ω)	(1, 0, ω)	(1, 0, 1)	(1, 0, ω^2)	(1, 0, ω^2)	(1, 0, 1)	(1, 0, ω)	3
$b_3 \equiv y_+^3$	(1, 1, 0)	(1, ω , 0)	(1, ω^2 , 0)	(1, ω , 0)	(1, ω^2 , 0)	(1, 1, 0)	(1, ω^2 , 0)	(1, ω , 0)	(1, 1, 0)	3
$c_1 \equiv y_-^1$	(0, 1, -1)	(0, ω , $-\omega^2$)	(0, ω^2 , $-\omega$)	(0, ω^2 , $-\omega^2$)	(0, 1, $-\omega$)	(0, ω , -1)	(0, ω , $-\omega$)	(0, 1, $-\omega^2$)	(0, ω^2 , -1)	3
$c_2 \equiv y_-^2$	(-1, 0, 1)	($-\omega$, 0, 1)	($-\omega^2$, 0, 1)	($-\omega^2$, 0, 1)	(-1, 0, 1)	($-\omega$, 0, 1)	($-\omega$, 0, 1)	(-1, 0, 1)	($-\omega^2$, 0, 1)	3
$c_3 \equiv y_-^3$	(1, -1, 0)	(ω^2 , -1, 0)	(ω , -1, 0)	(ω^2 , -1, 0)	(ω , -1, 0)	(1, -1, 0)	(ω , -1, 0)	(ω^2 , -1, 0)	(1, -1, 0)	3
d_1	(-2, 1, 1)	(-2, ω , ω^2)	(-2, ω^2 , ω)	(-2, ω , ω)	(-2, ω^2 , 1)	(-2, 1, ω^2)	(-2, ω^2 , ω^2)	(-2, ω , 1)	(-2, 1, ω)	9
d_2	(1, -2, 1)	(ω^2 , -2, ω)	(ω , -2, ω^2)	(ω^2 , -2, 1)	(ω , -2, ω)	(1, -2, ω^2)	(ω , -2, 1)	(ω^2 , -2, ω^2)	(1, -2, ω)	9
d_3	(1, 1, -2)	(ω , ω^2 , -2)	(ω^2 , ω , -2)	(ω^2 , 1, -2)	(1, ω^2 , -2)	(ω , ω , -2)	(ω , 1, -2)	(1, ω , -2)	(ω^2 , ω^2 , -2)	9
$b_{12} \equiv h_3$	(1, 1, -1)	(1, ω , $-\omega^2$)	(1, ω^2 , $-\omega$)	(ω , ω^2 , $-\omega^2$)	(ω , 1, $-\omega$)	(ω , ω , -1)	(ω^2 , ω , $-\omega$)	(ω^2 , 1, $-\omega^2$)	(ω^2 , ω^2 , -1)	9
$b_{13} \equiv h_2$	(1, -1, 1)	(1, $-\omega$, ω^2)	(1, $-\omega^2$, ω)	(ω , $-\omega^2$, ω^2)	(ω , -1, ω)	(ω , $-\omega$, 1)	(ω^2 , $-\omega$, ω)	(ω^2 , -1, ω^2)	(ω^2 , $-\omega^2$, 1)	9
$b_{23} \equiv h_1$	(-1, 1, 1)	(-1, ω , ω^2)	(-1, ω^2 , ω)	($-\omega$, ω^2 , ω^2)	($-\omega$, 1, ω)	($-\omega$, ω , 1)	($-\omega^2$, ω , ω)	($-\omega^2$, 1, ω^2)	($-\omega^2$, ω^2 , 1)	9
e_1	(1, 2, 1)	(ω^2 , 2, ω)	(ω , 2, ω^2)	(ω , $2\omega^2$, ω^2)	(1, $2\omega^2$, 1)	(ω^2 , $2\omega^2$, ω)	(ω^2 , 2 ω , ω)	(1, 2 ω , 1)	(ω , 2 ω , ω^2)	9
e_2	(1, 1, 2)	(ω , ω^2 , 2)	(ω^2 , ω , 2)	(ω , ω^2 , $2\omega^2$)	(ω^2 , ω , $2\omega^2$)	(1, 1, $2\omega^2$)	(ω^2 , ω , 2 ω)	(ω , ω^2 , 2 ω)	(1, 1, 2 ω)	9
e_3	(2, 1, 1)	(2, ω , ω^2)	(2, ω^2 , ω)	($2\omega^2$, 1, 1)	($2\omega^2$, ω , ω^2)	($2\omega^2$, ω^2 , ω)	(2 ω , 1, 1)	(2 ω , ω^2 , ω)	(2 ω , ω , ω^2)	9
b_{112}	(2, -1, 1)	(2, $-\omega$, ω^2)	(2, $-\omega^2$, ω)	(2, $-\omega$, ω)	(2, $-\omega^2$, 1)	(2, -1, ω^2)	(2, $-\omega^2$, ω^2)	(2, $-\omega$, 1)	(2, -1, ω)	9
b_{212}	(-1, 2, 1)	(-1, 2 ω , ω^2)	(-1, 2 ω^2 , ω)	(-1, 2 ω , ω)	(-1, 2 ω^2 , 1)	(-1, 2, ω^2)	(-1, 2 ω^2 , ω^2)	(-1, 2 ω , 1)	(-1, 2, ω)	9
b_{113}	(2, 1, -1)	(2, ω , $-\omega^2$)	(2, ω^2 , $-\omega$)	(2, ω , $-\omega$)	(2, ω^2 , -1)	(2, 1, $-\omega^2$)	(2, ω^2 , $-\omega^2$)	(2, ω , -1)	(2, 1, $-\omega$)	9
b_{313}	(-1, 1, 2)	(-1, ω , 2 ω^2)	(-1, ω^2 , 2 ω)	(-1, ω , 2 ω)	(-1, ω^2 , 2)	(-1, 1, 2 ω^2)	(-1, ω^2 , 2 ω^2)	(-1, ω , 2)	(-1, 1, 2 ω)	9
b_{223}	(1, 2, -1)	(1, 2 ω , $-\omega^2$)	(1, 2 ω^2 , $-\omega$)	(1, 2 ω , $-\omega$)	(1, 2 ω^2 , -1)	(1, 2, $-\omega^2$)	(1, 2 ω^2 , $-\omega^2$)	(1, 2 ω , -1)	(1, 2, $-\omega$)	9
b_{323}	(1, -1, 2)	(1, $-\omega$, 2 ω^2)	(1, $-\omega^2$, 2 ω)	(1, $-\omega$, 2 ω)	(1, $-\omega^2$, 2)	(1, -1, 2 ω^2)	(1, $-\omega^2$, 2 ω^2)	(1, $-\omega$, 2)	(1, -1, 2 ω)	9

the resulting orthogonality graph are precisely the contexts. The exhaustive search gives 50 contexts per lineage and 130 in the union. Finally, direct enumeration of maps $s : V \rightarrow \{0, 1\}$ with one 1 per context gives no two-valued state for any of the three 69-ray, 50-context lineages. $\square$

This calculation also explains why a first-occurrence labeling rule can produce a spurious 40–4–4 count: 18 of the 21 common rays were previously displayed with the color of $\mathcal{B}_1$, while the three Cartesian rays were displayed in black. Provenance must instead be assigned from the complete set of generating lineages, which restores the symmetry summarized in Proposition 1.

IV. HIGHER-DIMENSIONAL FORCING GADGETS

In $\mathcal{P}(\mathbb{C}^3)$, two independent rays determine a unique orthogonal ray. For $D \geq 4$, their orthogonal complement has dimension $D - 2$, so additional connector contexts are required to obtain a rigid finite construction.

Definition 3 (Unbiased vector) *A normalized vector u is maximally unbiased relative to an orthonormal basis $\mathcal{B}$ if*

$$|\langle e_j, u \rangle|^2 = 1/D \quad (j = 1, \dots, D).$$

For $u = (x_1, \dots, x_D)$ in the computational basis, this is equivalent to $|x_1|^2 = \dots = |x_D|^2$.

The $D = 4$ and $D = 5$ constructions have three logically separate parts: (i) a scaffold of coordinate-supported minors, (ii) connector rays whose orthogonality to the center gives equations in $|x_j|^2$, and (iii) completion vectors that turn the required orthogonal pairs into

TABLE II. The complete set of 165 projectively distinct rays, with six entries per row. The rightmost index records the seed column in Table I. The 21 rays common to all three 69-ray lineages are black. The 48 rays exclusive to $\mathcal{B}_1$, $\mathcal{B}_2$, and $\mathcal{B}_3$ are shown in red, green, and blue, respectively.

$a_{11} = (1, 0, 0)$	$a_{21} = (0, 1, 0)$	$a_{31} = (0, 0, 1)$	$u_1 = (1, 1, 1)$	$u_2 = (1, \omega, \omega^2)$	$u_3 = (1, \omega^2, \omega)$
$u_4 = (1, \omega, \omega)$	$u_5 = (1, \omega^2, 1)$	$u_6 = (1, 1, \omega^2)$	$u_7 = (1, \omega^2, \omega^2)$	$u_8 = (1, \omega, 1)$	$u_9 = (1, 1, \omega)$
$b_{11} = (0, 1, 1)$	$b_{12} = (0, \omega, \omega^2)$	$b_{13} = (0, \omega^2, \omega)$	$b_{21} = (1, 0, 1)$	$b_{22} = (1, 0, \omega^2)$	$b_{23} = (1, 0, \omega)$
$b_{31} = (1, 1, 0)$	$b_{32} = (1, \omega, 0)$	$b_{33} = (1, \omega^2, 0)$	$c_{11} = (0, 1, -1)$	$c_{12} = (0, \omega, -\omega^2)$	$c_{13} = (0, \omega^2, -\omega)$
$c_{21} = (-1, 0, 1)$	$c_{22} = (-\omega, 0, 1)$	$c_{23} = (-\omega^2, 0, 1)$	$c_{31} = (1, -1, 0)$	$c_{32} = (\omega^2, -1, 0)$	$c_{33} = (\omega, -1, 0)$
$d_{11} = (-2, 1, 1)$	$d_{12} = (-2, \omega, \omega^2)$	$d_{13} = (-2, \omega^2, \omega)$	$d_{14} = (-2, \omega, \omega)$	$d_{15} = (-2, \omega^2, 1)$	$d_{16} = (-2, 1, \omega^2)$
$d_{17} = (-2, \omega^2, \omega^2)$	$d_{18} = (-2, \omega, 1)$	$d_{19} = (-2, 1, \omega)$	$d_{21} = (1, -2, 1)$	$d_{22} = (\omega^2, -2, \omega)$	$d_{23} = (\omega, -2, \omega^2)$
$d_{24} = (\omega^2, -2, 1)$	$d_{25} = (\omega, -2, \omega)$	$d_{26} = (1, -2, \omega^2)$	$d_{27} = (\omega, -2, 1)$	$d_{28} = (\omega^2, -2, \omega^2)$	$d_{29} = (1, -2, \omega)$
$d_{31} = (1, 1, -2)$	$d_{32} = (\omega, \omega^2, -2)$	$d_{33} = (\omega^2, \omega, -2)$	$d_{34} = (\omega^2, 1, -2)$	$d_{35} = (1, \omega^2, -2)$	$d_{36} = (\omega, \omega, -2)$
$d_{37} = (\omega, 1, -2)$	$d_{38} = (1, \omega, -2)$	$d_{39} = (\omega^2, \omega^2, -2)$	$b_{121} = (1, 1, -1)$	$b_{122} = (1, \omega, -\omega^2)$	$b_{123} = (1, \omega^2, -\omega)$
$b_{124} = (\omega, \omega^2, -\omega^2)$	$b_{125} = (\omega, 1, -\omega)$	$b_{126} = (\omega, \omega, -1)$	$b_{127} = (\omega^2, \omega, -\omega)$	$b_{128} = (\omega^2, 1, -\omega^2)$	$b_{129} = (\omega^2, \omega^2, -1)$
$b_{131} = (1, -1, 1)$	$b_{132} = (1, -\omega, \omega^2)$	$b_{133} = (1, -\omega^2, \omega)$	$b_{134} = (\omega, -\omega^2, \omega^2)$	$b_{135} = (\omega, -1, \omega)$	$b_{136} = (\omega, -\omega, 1)$
$b_{137} = (\omega^2, -\omega, \omega)$	$b_{138} = (\omega^2, -1, \omega^2)$	$b_{139} = (\omega^2, -\omega^2, 1)$	$b_{231} = (-1, 1, 1)$	$b_{232} = (-1, \omega, \omega^2)$	$b_{233} = (-1, \omega^2, \omega)$
$b_{234} = (-\omega, \omega^2, \omega^2)$	$b_{235} = (-\omega, 1, \omega)$	$b_{236} = (-\omega, \omega, 1)$	$b_{237} = (-\omega^2, \omega, \omega)$	$b_{238} = (-\omega^2, 1, \omega^2)$	$b_{239} = (-\omega^2, \omega^2, 1)$
$e_{11} = (1, 2, 1)$	$e_{12} = (\omega^2, 2, \omega)$	$e_{13} = (\omega, 2, \omega^2)$	$e_{14} = (\omega, 2\omega^2, \omega^2)$	$e_{15} = (1, 2\omega^2, 1)$	$e_{16} = (\omega^2, 2\omega^2, \omega)$
$e_{17} = (\omega^2, 2\omega, \omega)$	$e_{18} = (1, 2\omega, 1)$	$e_{19} = (\omega, 2\omega, \omega^2)$	$e_{21} = (1, 1, 2)$	$e_{22} = (\omega, \omega^2, 2)$	$e_{23} = (\omega^2, \omega, 2)$
$e_{24} = (\omega, \omega^2, 2\omega^2)$	$e_{25} = (\omega^2, \omega, 2\omega^2)$	$e_{26} = (1, 1, 2\omega^2)$	$e_{27} = (\omega^2, \omega, 2\omega)$	$e_{28} = (\omega, \omega^2, 2\omega)$	$e_{29} = (1, 1, 2\omega)$
$e_{31} = (2, 1, 1)$	$e_{32} = (2, \omega, \omega^2)$	$e_{33} = (2, \omega^2, \omega)$	$e_{34} = (2\omega^2, 1, 1)$	$e_{35} = (2\omega^2, \omega, \omega^2)$	$e_{36} = (2\omega^2, \omega^2, \omega)$
$e_{37} = (2\omega, 1, 1)$	$e_{38} = (2\omega, \omega^2, \omega)$	$e_{39} = (2\omega, \omega, \omega^2)$	$b_{1121} = (2, -1, 1)$	$b_{1122} = (2, -\omega, \omega^2)$	$b_{1123} = (2, -\omega^2, \omega)$
$b_{1124} = (2, -\omega, \omega)$	$b_{1125} = (2, -\omega^2, 1)$	$b_{1126} = (2, -1, \omega^2)$	$b_{1127} = (2, -\omega^2, \omega^2)$	$b_{1128} = (2, -\omega, 1)$	$b_{1129} = (2, -1, \omega)$
$b_{2121} = (-1, 2, 1)$	$b_{2122} = (-1, 2\omega, \omega^2)$	$b_{2123} = (-1, 2\omega^2, \omega)$	$b_{2124} = (-1, 2\omega, \omega)$	$b_{2125} = (-1, 2\omega^2, 1)$	$b_{2126} = (-1, 2, \omega^2)$
$b_{2127} = (-1, 2\omega^2, \omega^2)$	$b_{2128} = (-1, 2\omega, 1)$	$b_{2129} = (-1, 2, \omega)$	$b_{1131} = (2, 1, -1)$	$b_{1132} = (2, \omega, -\omega^2)$	$b_{1133} = (2, \omega^2, -\omega)$
$b_{1134} = (2, \omega, -\omega)$	$b_{1135} = (2, \omega^2, -1)$	$b_{1136} = (2, 1, -\omega^2)$	$b_{1137} = (2, \omega^2, -\omega^2)$	$b_{1138} = (2, \omega, -1)$	$b_{1139} = (2, 1, -\omega)$
$b_{3131} = (-1, 1, 2)$	$b_{3132} = (-1, \omega, 2\omega^2)$	$b_{3133} = (-1, \omega^2, 2\omega)$	$b_{3134} = (-1, \omega, 2\omega)$	$b_{3135} = (-1, \omega^2, 2)$	$b_{3136} = (-1, 1, 2\omega^2)$
$b_{3137} = (-1, \omega^2, 2\omega^2)$	$b_{3138} = (-1, \omega, 2)$	$b_{3139} = (-1, 1, 2\omega)$	$b_{2231} = (1, 2, -1)$	$b_{2232} = (1, 2\omega, -\omega^2)$	$b_{2233} = (1, 2\omega^2, -\omega)$
$b_{2234} = (1, 2\omega, -\omega)$	$b_{2235} = (1, 2\omega^2, -1)$	$b_{2236} = (1, 2, -\omega^2)$	$b_{2237} = (1, 2\omega^2, -\omega^2)$	$b_{2238} = (1, 2\omega, -1)$	$b_{2239} = (1, 2, -\omega)$
$b_{3231} = (1, -1, 2)$	$b_{3232} = (1, -\omega, 2\omega^2)$	$b_{3233} = (1, -\omega^2, 2\omega)$	$b_{3234} = (1, -\omega, 2\omega)$	$b_{3235} = (1, -\omega^2, 2)$	$b_{3236} = (1, -1, 2\omega^2)$
$b_{3237} = (1, -\omega^2, 2\omega^2)$	$b_{3238} = (1, -\omega, 2)$	$b_{3239} = (1, -1, 2\omega)$			

full contexts. The propositions below state the forcing constraints; coordinate inventories and exact enumeration checks are given afterward.

A. Notation and conventions

Fix an unknown center vector $u = (x_1, \dots, x_D) \in \mathbb{C}^D$ with $x_j \neq 0$ for all j , and use the standard Hermitian inner product

$$\langle v, w \rangle = \sum_m \overline{v_m} w_m.$$

We use vectors to represent the one-dimensional subspaces they span and freely scale them when forming blocks, with the understanding that they must be normalized to form a literal orthonormal basis.

B. An explicit gadget for $D = 4$

First, take the computational (Cartesian) basis

$$\begin{aligned} e_1 &= (1, 0, 0, 0), & e_2 &= (0, 1, 0, 0), \\ e_3 &= (0, 0, 1, 0), & e_4 &= (0, 0, 0, 1), \end{aligned}$$

and place the context $\mathcal{B}_{\text{comp}} = \{e_1, e_2, e_3, e_4\}$ on the rim of Fig. 2(a).

In the second construction stage define, for each unordered pair $\{i, j\} \subset \{1, 2, 3, 4\}$, pair minors from a Levi-Civita contraction:

$$\begin{aligned} v_{12} &= (0, 0, \overline{x_4}, -\overline{x_3}), & v_{13} &= (0, -\overline{x_4}, 0, \overline{x_2}), \\ v_{14} &= (0, \overline{x_3}, -\overline{x_2}, 0), & v_{23} &= (\overline{x_4}, 0, 0, -\overline{x_1}), \\ v_{24} &= (-\overline{x_3}, 0, \overline{x_1}, 0), & v_{34} &= (\overline{x_2}, -\overline{x_1}, 0, 0). \end{aligned}$$

Each v_{ij} vanishes in coordinates i, j and satisfies $\langle v_{ij}, u \rangle = 0$ by direct cancellation.

In the third step, choose a vector w_{ij} with the same support as v_{ij} and orthogonal to it, that is, $\langle v_{ij}, w_{ij} \rangle = 0$:

$$\begin{aligned} w_{12} &= (0, 0, x_3, x_4), & w_{13} &= (0, x_2, 0, x_4), \\ w_{14} &= (0, x_2, x_3, 0), & w_{23} &= (x_1, 0, 0, x_4), \\ w_{24} &= (x_1, 0, x_3, 0), & w_{34} &= (x_1, x_2, 0, 0). \end{aligned}$$

TABLE III. The complete list of 130 orthogonal bases supported by the 165 rays. To avoid conflating shared rays with lineage-exclusive rays, the entries are left uncolored; the invariant provenance counts are stated in Proposition 1.

$\{a_{11}, a_{21}, a_{31}\}$	$\{a_{11}, b_{11}, c_{11}\}$	$\{a_{11}, b_{12}, c_{12}\}$	$\{a_{11}, b_{13}, c_{13}\}$	$\{a_{21}, b_{21}, c_{21}\}$	$\{a_{21}, b_{22}, c_{22}\}$
$\{a_{21}, b_{23}, c_{23}\}$	$\{a_{31}, b_{31}, c_{31}\}$	$\{a_{31}, b_{32}, c_{32}\}$	$\{a_{31}, b_{33}, c_{33}\}$	$\{u_1, u_2, u_3\}$	$\{u_1, c_{11}, d_{11}\}$
$\{u_1, c_{21}, d_{21}\}$	$\{u_1, c_{31}, d_{31}\}$	$\{u_2, c_{12}, d_{12}\}$	$\{u_2, c_{22}, d_{22}\}$	$\{u_2, c_{32}, d_{32}\}$	$\{u_3, c_{13}, d_{13}\}$
$\{u_3, c_{23}, d_{23}\}$	$\{u_3, c_{33}, d_{33}\}$	$\{u_4, u_5, u_6\}$	$\{u_4, c_{11}, d_{14}\}$	$\{u_4, c_{23}, d_{24}\}$	$\{u_4, c_{32}, d_{34}\}$
$\{u_5, c_{12}, d_{15}\}$	$\{u_5, c_{21}, d_{25}\}$	$\{u_5, c_{33}, d_{35}\}$	$\{u_6, c_{13}, d_{16}\}$	$\{u_6, c_{22}, d_{26}\}$	$\{u_6, c_{31}, d_{36}\}$
$\{u_7, u_8, u_9\}$	$\{u_7, c_{11}, d_{17}\}$	$\{u_7, c_{22}, d_{27}\}$	$\{u_7, c_{33}, d_{37}\}$	$\{u_8, c_{13}, d_{18}\}$	$\{u_8, c_{21}, d_{28}\}$
$\{u_8, c_{32}, d_{38}\}$	$\{u_9, c_{12}, d_{19}\}$	$\{u_9, c_{23}, d_{29}\}$	$\{u_9, c_{31}, d_{39}\}$	$\{b_{11}, b_{121}, b_{1121}\}$	$\{b_{11}, b_{124}, b_{1124}\}$
$\{b_{11}, b_{127}, b_{1127}\}$	$\{b_{11}, b_{131}, b_{1131}\}$	$\{b_{11}, b_{134}, b_{1134}\}$	$\{b_{11}, b_{137}, b_{1137}\}$	$\{b_{12}, b_{122}, b_{1122}\}$	$\{b_{12}, b_{125}, b_{1125}\}$
$\{b_{12}, b_{129}, b_{1129}\}$	$\{b_{12}, b_{132}, b_{1132}\}$	$\{b_{12}, b_{135}, b_{1135}\}$	$\{b_{12}, b_{139}, b_{1139}\}$	$\{b_{13}, b_{123}, b_{1123}\}$	$\{b_{13}, b_{126}, b_{1126}\}$
$\{b_{13}, b_{128}, b_{1128}\}$	$\{b_{13}, b_{133}, b_{1133}\}$	$\{b_{13}, b_{136}, b_{1136}\}$	$\{b_{13}, b_{138}, b_{1138}\}$	$\{b_{21}, b_{121}, b_{2121}\}$	$\{b_{21}, b_{125}, b_{2125}\}$
$\{b_{21}, b_{128}, b_{2128}\}$	$\{b_{21}, b_{231}, b_{2231}\}$	$\{b_{21}, b_{235}, b_{2235}\}$	$\{b_{21}, b_{238}, b_{2238}\}$	$\{b_{22}, b_{122}, b_{2122}\}$	$\{b_{22}, b_{126}, b_{2126}\}$
$\{b_{22}, b_{127}, b_{2127}\}$	$\{b_{22}, b_{232}, b_{2232}\}$	$\{b_{22}, b_{236}, b_{2236}\}$	$\{b_{22}, b_{237}, b_{2237}\}$	$\{b_{23}, b_{123}, b_{2123}\}$	$\{b_{23}, b_{124}, b_{2124}\}$
$\{b_{23}, b_{129}, b_{2129}\}$	$\{b_{23}, b_{233}, b_{2233}\}$	$\{b_{23}, b_{234}, b_{2234}\}$	$\{b_{23}, b_{239}, b_{2239}\}$	$\{b_{31}, b_{131}, b_{3131}\}$	$\{b_{31}, b_{136}, b_{3136}\}$
$\{b_{31}, b_{139}, b_{3139}\}$	$\{b_{31}, b_{231}, b_{3231}\}$	$\{b_{31}, b_{236}, b_{3236}\}$	$\{b_{31}, b_{239}, b_{3239}\}$	$\{b_{32}, b_{132}, b_{3132}\}$	$\{b_{32}, b_{134}, b_{3134}\}$
$\{b_{32}, b_{138}, b_{3138}\}$	$\{b_{32}, b_{232}, b_{3232}\}$	$\{b_{32}, b_{234}, b_{3234}\}$	$\{b_{32}, b_{238}, b_{3238}\}$	$\{b_{33}, b_{133}, b_{3133}\}$	$\{b_{33}, b_{135}, b_{3135}\}$
$\{b_{33}, b_{137}, b_{3137}\}$	$\{b_{33}, b_{233}, b_{3233}\}$	$\{b_{33}, b_{235}, b_{3235}\}$	$\{b_{33}, b_{237}, b_{3237}\}$	$\{c_{11}, b_{231}, e_{31}\}$	$\{c_{11}, b_{234}, e_{34}\}$
$\{c_{11}, b_{237}, e_{37}\}$	$\{c_{12}, b_{232}, e_{32}\}$	$\{c_{12}, b_{235}, e_{35}\}$	$\{c_{12}, b_{239}, e_{39}\}$	$\{c_{13}, b_{233}, e_{33}\}$	$\{c_{13}, b_{236}, e_{36}\}$
$\{c_{13}, b_{238}, e_{38}\}$	$\{c_{21}, b_{131}, e_{11}\}$	$\{c_{21}, b_{135}, e_{15}\}$	$\{c_{21}, b_{138}, e_{18}\}$	$\{c_{22}, b_{132}, e_{12}\}$	$\{c_{22}, b_{136}, e_{16}\}$
$\{c_{22}, b_{137}, e_{17}\}$	$\{c_{23}, b_{133}, e_{13}\}$	$\{c_{23}, b_{134}, e_{14}\}$	$\{c_{23}, b_{139}, e_{19}\}$	$\{c_{31}, b_{121}, e_{21}\}$	$\{c_{31}, b_{126}, e_{26}\}$
$\{c_{31}, b_{129}, e_{29}\}$	$\{c_{32}, b_{122}, e_{22}\}$	$\{c_{32}, b_{124}, e_{24}\}$	$\{c_{32}, b_{128}, e_{28}\}$	$\{c_{33}, b_{123}, e_{23}\}$	$\{c_{33}, b_{125}, e_{25}\}$
$\{c_{33}, b_{127}, e_{27}\}$	$\{b_{121}, b_{122}, b_{123}\}$	$\{b_{124}, b_{125}, b_{126}\}$	$\{b_{127}, b_{128}, b_{129}\}$	$\{b_{131}, b_{132}, b_{133}\}$	$\{b_{134}, b_{135}, b_{136}\}$
$\{b_{137}, b_{138}, b_{139}\}$	$\{b_{231}, b_{232}, b_{233}\}$	$\{b_{234}, b_{235}, b_{236}\}$	$\{b_{237}, b_{238}, b_{239}\}$		

For each of the $\binom{4}{2} = 6$ pairs $\{i, j\}$, the set

$$\mathcal{B}_{ij} = \{e_i, e_j, v_{ij}, w_{ij}\}$$

is an orthogonal basis (block, context).

In a fourth step, define three connector vectors

$$\begin{aligned} v_{1234} &= (x_1, x_2, -x_3, -x_4), \\ v_{1324} &= (x_1, -x_2, x_3, -x_4), \\ v_{1423} &= (x_1, -x_2, -x_3, x_4). \end{aligned}$$

Whenever u is orthogonal to one of these connector vectors, the two-dimensional complement of their span supplies completion rays h_1, h_2 . Thus the three connector contexts may be written

$$\begin{aligned} \mathcal{C}_{(a)} &= \{u, v_{1234}, h_1^{(a)}, h_2^{(a)}\}, \quad \mathcal{C}_{(b)} = \{u, v_{1324}, h_1^{(b)}, h_2^{(b)}\}, \\ \mathcal{C}_{(c)} &= \{u, v_{1423}, h_1^{(c)}, h_2^{(c)}\}. \end{aligned}$$

Proposition 2 ($D = 4$ forcing constraint)

Membership of u and the three connector rays in the contexts above forces u to be maximally unbiased relative to $\mathcal{B}_{\text{comp}}$.

Proof. The three required inner products vanish exactly when

$$|x_1|^2 + |x_2|^2 = |x_3|^2 + |x_4|^2, \quad (1)$$

$$|x_1|^2 + |x_3|^2 = |x_2|^2 + |x_4|^2, \quad (2)$$

$$|x_1|^2 + |x_4|^2 = |x_2|^2 + |x_3|^2. \quad (3)$$

Subtracting pairs of equations gives $|x_2|^2 = |x_3|^2 = |x_4|^2$; substitution gives $|x_1|^2 = |x_2|^2$. After normalization, every squared modulus is $1/4$. $\square$

1. Automatic and equality-induced contexts

Orthogonalities due only to disjoint support, such as $v_{12} \perp v_{34}$, do not constrain u . For the concrete 20-ray representation with $u = (1, 1, 1, 1)$, the computational context, six pair contexts, and three connector contexts give 10 imposed contexts. An exhaustive orthogonality search finds exactly seven further contexts. Three follow from disjoint support:

$$\begin{aligned} \mathcal{B}_{1234} &= \{v_{12}, w_{12}, v_{34}, w_{34}\}, \\ \mathcal{B}_{1324} &= \{v_{13}, w_{13}, v_{24}, w_{24}\}, \\ \mathcal{B}_{1423} &= \{v_{14}, w_{14}, v_{23}, w_{23}\}. \end{aligned}$$

The remaining four use the equal-modulus relations from Proposition 2:

$$\begin{aligned} \mathcal{C}_0 &= \{u, v_{1234}, v_{1324}, v_{1423}\}, \\ \mathcal{E}_1 &= \{w_{12}, w_{34}, v_{1324}, v_{1423}\}, \\ \mathcal{E}_2 &= \{w_{13}, w_{24}, v_{1234}, v_{1423}\}, \\ \mathcal{E}_3 &= \{w_{14}, w_{23}, v_{1234}, v_{1324}\}. \end{aligned}$$

Hence the complete orthogonality hypergraph of the concrete 20-ray set has 17 contexts, not 13. Exact enumera-

tion gives 16 two-valued states, and these states separate every pair of rays; the complete 20-ray hypergraph is therefore not a KS configuration.

2. Concrete $D = 4$ example: $u = (1, 1, 1, 1)$

Take the center vector $u = (1, 1, 1, 1)$ and the rim vectors

$$\begin{aligned} e_1 &= (1, 0, 0, 0), & e_2 &= (0, 1, 0, 0), \\ e_3 &= (0, 0, 1, 0), & e_4 &= (0, 0, 0, 1). \end{aligned}$$

The pair minors and their complementary vectors become:

$$\begin{aligned} v_{12} &= (0, 0, 1, -1), & w_{12} &= (0, 0, 1, 1), \\ v_{13} &= (0, -1, 0, 1), & w_{13} &= (0, 1, 0, 1), \\ v_{14} &= (0, 1, -1, 0), & w_{14} &= (0, 1, 1, 0), \\ v_{23} &= (1, 0, 0, -1), & w_{23} &= (1, 0, 0, 1), \\ v_{24} &= (-1, 0, 1, 0), & w_{24} &= (1, 0, 1, 0), \\ v_{34} &= (1, -1, 0, 0), & w_{34} &= (1, 1, 0, 0). \end{aligned}$$

Each pair block

$$\mathcal{B}_{ij} = \{e_i, e_j, v_{ij}, w_{ij}\}$$

forms an ONB after normalizing v_{ij} and w_{ij} .

The connector vectors are

$$\begin{aligned} v_{1234} &= (1, 1, -1, -1), \\ v_{1324} &= (1, -1, 1, -1), \\ v_{1423} &= (1, -1, -1, 1), \end{aligned}$$

and they are all orthogonal to $u = (1, 1, 1, 1)$ by inspection.

A convenient explicit completion of the connector blocks can be formed using some of the existing minor vectors:

$$\begin{aligned} C_{(a)} &= \{u, v_{1234}, v_{34}, v_{12}\}, \\ C_{(b)} &= \{u, v_{1324}, v_{24}, v_{13}\}, \\ C_{(c)} &= \{u, v_{1423}, v_{23}, v_{14}\}. \end{aligned}$$

Each of these sets is composed of four mutually orthogonal vectors. For example, in $C_{(a)}$, the vectors v_{34} and v_{12} have disjoint support, making them orthogonal. The inner products $\langle v_{1234}, v_{34} \rangle$ and $\langle v_{1234}, v_{12} \rangle$ are zero, as are all other inner products involving u . To create literal ONBs, the vectors should be normalized.

Together with $\mathcal{B}_{\text{comp}}$, the six pair contexts and three connector contexts form the 10-context scaffold that enforces Proposition 2. The equal-modulus condition also creates $\mathcal{C}_0$ and the three contexts $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ listed above. For example, $\langle w_{14}, v_{1234} \rangle = |x_2|^2 - |x_3|^2 = 0$; $\mathcal{E}_3$ is the dotted dark-green curve in Fig. 2(a).

3. Mutual reconstruction inside the Peres–Mermin eigensystem

Both ray sets are subsets of the 24-ray Peres–Mermin eigensystem [17, 18]. They share 15 rays projectively. Table IV gives rank-three orthogonal-complement constructions for the five gadget-only rays and the three Cabello-only rays. Hence each ray set generates the other under the operation “take the unique ray orthogonal to three independent rays.” This is the precise sense in which we call them informationally equivalent.

Informational equivalence does not imply equality of context hypergraphs. The complete 20-ray orthogonality hypergraph has 17 contexts and 16 separating two-valued states. The Cabello 18-ray hypergraph selects nine contexts and has no two-valued state; its parity argument makes it a KS configuration [16]. The contrast is therefore not a consequence of “more contexts allowing loopholes”—adding constraints cannot create two-valued states—but of the two constructions selecting different rays and different contexts from their common 24-ray parent.

4. Context completion and two-valued states

The context choice within the 20-ray set can itself be tracked explicitly. Let

$$\mathfrak{B}_{10} = \{\mathcal{B}_{\text{comp}}, \mathcal{B}_{ij} \ (1 \leq i < j \leq 4), \mathcal{C}_{(a)}, \mathcal{C}_{(b)}, \mathcal{C}_{(c)}\}$$

be the 10 imposed contexts, and let

$$\mathfrak{B}_{13} = \mathfrak{B}_{10} \cup \{\mathcal{B}_{1234}, \mathcal{B}_{1324}, \mathcal{B}_{1423}\}.$$

Exact enumeration shows that both hypergraphs have the same 36 two-valued states, and those states are separating. The three disjoint-support contexts therefore record exclusions already obeyed by every state of $\mathfrak{B}_{10}$; they do not remove an assignment. Adding the four equality-induced contexts gives the complete set

$$\mathfrak{B}_{17} = \mathfrak{B}_{13} \cup \{\mathcal{C}_0, \mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3\},$$

which has 16 separating two-valued states. This nested comparison makes the monotonicity transparent: adding contexts can preserve or reduce the state set, but cannot enlarge it. It also avoids attributing KS contextuality to an incomplete incidence structure merely because some available contexts were omitted.

The apparent asymmetry of Fig. 2(a) is mainly graphical and reflects the selected 20-ray subset. Both it and the Cabello 18-ray set sit inside the more symmetric 24-ray, 24-context Peres configuration [22, 23]; neither subset need inherit all symmetries of the parent.




FIG. 2. (a) Four-dimensional extension of the Harding-Salinas Schmeis hypergraph notation [27]. Solid curves show imposed contexts. The dotted dark-green curve is the equality-induced context $\mathcal{E}_3 = \{w_{14}, w_{23}, v_{1234}, v_{1324}\}$; the other contexts of the complete 20-ray orthogonality hypergraph are listed in the text. (b) The Cabello 18-ray, 9-context KS configuration [16], with labels from Ref. [13]. Shared rays are relabeled by the gadget notation; overall nonzero scalar factors are immaterial. The notation $\{v_i, v_j, v_k\}^\perp$ denotes the unique ray orthogonal to the span of three independent rays. The drawings record the stated incidence relations; no lattice-theoretic conclusion is inferred from the planar layout alone.

C. A $D = 5$ scaffold and forcing constraint

Let ε_{abcde} be totally antisymmetric with $\varepsilon_{12345} = 1$. For each triple $\{i, j, k\}$ define

$$(v_{ijk})_m = \sum_{\ell=1}^5 \varepsilon_{mijk\ell} \overline{x_\ell}.$$

The vector v_{ijk} vanishes in coordinates i, j, k and satisfies $\langle v_{ijk}, u \rangle = 0$ by antisymmetry. Choose w_{ijk} on the complementary two coordinates with $v_{ijk} \perp w_{ijk}$; then

$$\mathcal{B}_{ijk} = \{e_i, e_j, e_k, v_{ijk}, w_{ijk}\}$$

is a scaffold context.

The four connector rays are represented by

$$\begin{aligned} g_1 &= (x_1, 0, 0, 0, -x_5), & g_2 &= (0, x_2, 0, 0, -x_5), \\ g_3 &= (0, 0, x_3, 0, -x_5), & g_4 &= (0, 0, 0, x_4, -x_5). \end{aligned}$$

Whenever $u \perp g_i$, three rays spanning $\text{span}\{u, g_i\}^\perp$ complete the connector context $\mathcal{C}_i = \{u, g_i, h_1^{(i)}, h_2^{(i)}, h_3^{(i)}\}$.

Proposition 3 ($D = 5$ forcing constraint) *The four connector contexts force a normalized center vector u to be maximally unbiased relative to the computational basis.*

Proof. For $i = 1, 2, 3, 4$,

$$\langle u, g_i \rangle = |x_i|^2 - |x_5|^2.$$

Thus the four orthogonality conditions imply $|x_1|^2 = \dots = |x_5|^2$; normalization makes every squared modulus $1/5$. $\square$

V. SUMMARY AND CONCLUSION

The qutrit construction yields 165 projectively distinct rays and exactly 130 contexts. Correct provenance accounting is symmetric: the three 69-ray, 50-context KS lineages share 21 rays and 10 contexts, and each adds 48 exclusive rays and 40 lineage-specific contexts. This places the YO triple, Harding-Salinas Schmeis, and Cabello descriptions in a common MUB-based framework without assigning shared rays to an arbitrary first lineage.

In $D = 4$ and $D = 5$, the forcing mechanism is explicit at the level of constraints: connector orthogonalities yield linear equations in the squared coordinate moduli and hence maximal unbiasedness. For the concrete $D = 4$ realization, the complete 20-ray orthogonality hypergraph has 17 contexts and 16 separating two-valued states. Its ray set is mutually reconstructible with the Cabello 18-ray set inside the 24-ray Peres-Mermin eigensystem, but the Cabello nine-context hypergraph is uncolorable. Ray-set closure and context incidence are therefore distinct pieces of data.

The main methodological conclusion is that contextuality claims should specify the incidence hypergraph being tested and, when completeness is intended, enumerate every context supported by the chosen rays. Restricting attention to intertwining rays or to a selected subset of available contexts may define a legitimate reduced scenario, but its two-valued-state properties should not be transferred to a different completion without a separate check.

TABLE IV. Comparison of the 20-ray gadget and the Cabello 18-ray set [13, 16]. Both lie in the 24-ray Peres–Mermin eigensystem [17–21]. Fifteen rays are shared projectively. Each of the five gadget-only rays and each of the three Cabello-only rays is the unique orthogonal complement of a three-dimensional span generated by the other set, establishing mutual reconstructibility.

Gadget Label	Gadget Vector	Cabello Diagram Label	Vector Construction / Definition
u	$(1, 1, 1, 1)$	v_{56}	$(1, 1, 1, 1)$
e_1	$(1, 0, 0, 0)$	v_{12}	$(1, 0, 0, 0)$
e_2	$(0, 1, 0, 0)$	v_{18}	$(0, 1, 0, 0)$
e_3	$(0, 0, 1, 0)$	(Constructed)	$\{v_{12}, v_{18}, v_{28}\}^\perp$
e_4	$(0, 0, 0, 1)$	v_{28}	$(0, 0, 0, 1)$
v_{12}	$(0, 0, 1, -1)$	v_{16}	$(0, 0, 1, -1)$
v_{13}	$(0, -1, 0, 1)$	v_{45}	$(0, 1, 0, -1)$
v_{14}	$(0, 1, -1, 0)$	v_{23}	$(0, 1, -1, 0)$
v_{23}	$(1, 0, 0, -1)$	(Constructed)	$\{v_{18}, v_{23}, v_{39}\}^\perp$
v_{24}	$(-1, 0, 1, 0)$	v_{58}	$(1, 0, -1, 0)$
v_{34}	$(1, -1, 0, 0)$	v_{67}	$(1, -1, 0, 0)$
w_{12}	$(0, 0, 1, 1)$	v_{17}	$(0, 0, 1, 1)$
w_{13}	$(0, 1, 0, 1)$	(Constructed)	$\{v_{12}, v_{45}, v_{58}\}^\perp$
w_{14}	$(0, 1, 1, 0)$	v_{29}	$(0, 1, 1, 0)$
w_{23}	$(1, 0, 0, 1)$	v_{39}	$(1, 0, 0, 1)$
w_{24}	$(1, 0, 1, 0)$	v_{48}	$(1, 0, 1, 0)$
w_{34}	$(1, 1, 0, 0)$	(Constructed)	$\{v_{28}, v_{16}, v_{67}\}^\perp$
v_{1234}	$(1, 1, -1, -1)$	v_{69}	$(1, 1, -1, -1)$
v_{1324}	$(1, -1, 1, -1)$	v_{59}	$(1, -1, 1, -1)$
v_{1423}	$(1, -1, -1, 1)$	(Constructed)	$\{v_{23}, v_{17}, v_{48}\}^\perp$
(Constructed)	$\{v_{12}, v_{13}, w_{34}\}^\perp$	v_{34}	$(-1, 1, 1, 1)$
(Constructed)	$\{v_{14}, v_{24}, w_{23}\}^\perp$	v_{37}	$(1, 1, 1, -1)$
(Constructed)	$\{v_{13}, v_{23}, w_{12}\}^\perp$	v_{47}	$(1, 1, -1, 1)$

ACKNOWLEDGMENTS

We are grateful to Josef Tkadlec for providing a *Pascal* program that computes and analyses the set of two-valued states of collections of contexts. We are also grateful to Adán Cabello and Mladen Pavičić for valuable communications.

This text was partially created and revised with assistance from Gemini 2.5 Pro and OpenAI language models,

including Codex. The authors checked the calculations and text and take responsibility for the manuscript.

MN was supported by the Czech Science Foundation under Grant 25-20013L. KS was funded in whole or in part by the Austrian Science Fund (FWF) [Grant DOI:10.55776/PIN5424624].

The authors declare no conflict of interest.

-
- [1] A. M. Gleason, Measures on the closed subspaces of a Hilbert space, *Journal of Mathematics and Mechanics* (now Indiana University Mathematics Journal) **6**, 885 (1957).
 - [2] S. Kochen and E. P. Specker, The problem of hidden variables in quantum mechanics, *Journal of Mathematics and Mechanics* (now Indiana University Mathematics Journal) **17**, 59 (1967).
 - [3] E. Specker, Die Logik nicht gleichzeitig entscheidbarer Aussagen, *Dialectica* **14**, 239 (1960), english translation at <https://arxiv.org/abs/1103.4537>, arXiv:1103.4537.
 - [4] A. Bretto, *Hypergraph theory*, Mathematical Engineering (Springer, Cham, Switzerland, 2013) pp. xiv+119.
 - [5] M. H. Shekarri and K. Svozil, Noncontextual coloring of orthogonality hypergraphs, *Journal of Mathematical Physics* **63**, 032104 (2022), arXiv:2105.08520.
 - [6] K. Svozil and J. Tkadlec, Greechie diagrams, nonexistence of measures in quantum logics and Kochen–Specker type constructions, *Journal of Mathematical Physics* **37**, 5380 (1996).
 - [7] J. Bub, Schütte’s tautology and the Kochen–Specker theorem, *Foundations of Physics* **26**, 787 (1996).
 - [8] J. Tkadlec, Greechie diagrams of small quantum logics with small state spaces, *International Journal of Theo-*

- retical Physics **37**, 203 (1998).
- [9] K. Schütte, Letter to ernst specker, kiel, dated april 22nd, 1965 (1965), this letter is mentioned in Ref. [10]. A second letter to Ernst Specker from Kurt Schütte in Munich, dated November 30, 1983, hints on how Schütte had obtained this configuration: “*Zu-grunde liegt offenbar ein Schließungssatz einer vollelliptischen Ebene in einer speziellen Form, in der zwischen den Punkten und Geraden der Konfiguraton möglichst viel Polaritäten auftreten, um die Berechenbarkeit der Formel sicherzustellen und mit möglichst wenig Variablen auszukommen.*”.
 - [10] E. Clavadetscher-Seeberger, *Eine partielle Prädikatenlogik*, Ph.D. thesis, ETH-Zürich, Zürich (1983).
 - [11] S. Kochen and E. P. Specker, Logical structures arising in quantum theory, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
 - [12] A. Cabello, Reply to Pavičić’s ‘Comment on “Optimal conversion of Kochen-Specker sets into bipartite perfect quantum strategies” ’ (arxiv:2502.13787) (2025), arXiv:2503.02974.
 - [13] A. Cabello, Experimentally testable state-independent quantum contextuality, Physical Review Letters **101**, 210401 (2008), arXiv:0808.2456.
 - [14] S. Trandafir and A. Cabello, Optimal conversion of Kochen-Specker sets into bipartite perfect quantum strategies (2025), arXiv:2410.17470.
 - [15] M. Pavičić, Quantum contextual hypergraphs, operators, inequalities, and applications in higher dimensions, Entropy **27**, 54 (2025).
 - [16] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Bell-Kochen-Specker theorem: A proof with 18 vectors, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
 - [17] M. Pavičić, Kochen-Specker algorithms for qubits, AIP Conference Proceedings **734**, 195 (2004).
 - [18] K. Svozil, Converting nonlocality into contextuality, Physical Review A **110**, 012215 (2024), arXiv:2404.15793.
 - [19] A. Peres, Incompatible results of quantum measurements, Physics Letters A **151**, 107 (1990).
 - [20] D. N. Mermin, Simple unified form for the major no-hidden-variable theorems, Physical Review Letters **65**, 3373 (1990).
 - [21] A. Peres, Two simple proofs of the Kochen-Specker theorem, Journal of Physics A: Mathematical and General **24**, L175 (1991).
 - [22] M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Kochen-Specker vectors, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
 - [23] M. Pavičić, J.-P. Merlet, B. D. McKay, and N. D. Megill, Corrigendum: Kochen-Specker vectors, Journal of Physics A: Mathematical and General **38**, 3709 (2005).
 - [24] C. Budroni, A. Cabello, O. Gühne, M. Kleinmann, and J.-A. Larsson, Kochen-Specker contextuality, Reviews of Modern Physics **94**, 045007 (2022), arXiv:2102.13036.
 - [25] K. Svozil, Chromatic quantum contextuality, Entropy **27**, 387 (2025), arXiv:2501.15261.
 - [26] R. J. Greechie, Orthomodular lattices admitting no states, Journal of Combinatorial Theory. Series A **10**, 119 (1971).
 - [27] J. Harding and R. Salinas Schmeis, Remarks on orthogonality spaces, International Journal of Theoretical Physics **64**, 211 (2025), arXiv:2505.13871 [math-ph], arXiv:arXiv:2505.13871 [math-ph].
 - [28] P. Badziąg, I. Bengtsson, A. Cabello, and I. Pitowsky, Universality of state-independent violation of correlation inequalities for noncontextual theories, Physical Review Letters **103**, 050401 (2009).
 - [29] X.-D. Yu, Y.-Q. Guo, and D. M. Tong, A proof of the Kochen-Specker theorem can always be converted to a state-independent noncontextuality inequality, New Journal of Physics **17**, 093001 (2015).
 - [30] S. Yu and C. H. Oh, State-independent proof of Kochen-Specker theorem with 13 rays, Physical Review Letters **108**, 030402 (2012), arXiv:1109.4396.
 - [31] A. Cabello, The simplest Kochen-Specker set (2025), arXiv:2508.07335 [quant-ph].
 - [32] T. Durt, B.-G. Englert, I. Bengtsson, and K. Życzkowski, On mutually unbiased bases, International Journal of Quantum Information **8**, 535 (2010), arXiv:1004.3348.
 - [33] D. McNulty and S. Weigert, Mutually unbiased bases in composite dimensions – a review (2024), arXiv:2410.23997 [quant-ph], arXiv:2410.23997 [quant-ph].
 - [34] A. Cabello, A simple proof of the Kochen-Specker theorem, European Journal of Physics **15**, 179 (1994).
 - [35] J. Harding, K. Jager, and D. Smith, Group-valued measures on the lattice of closed subspaces of a Hilbert space, International Journal of Theoretical Physics **44**, 539 (2005).
 - [36] M. Navara, Mathematical questions related to non-existence of hidden variables, in *Foundations of Probability and Physics 5*, American Institute of Physics Conference Proceedings, Vol. 1101, edited by L. Accardi, G. Adenier, C. Fuchs, G. Jaeger, A. Y. Khrennikov, J.-A. Larsson, and S. Stenholm (American Institute of Physics, New York, USA, 2009) pp. 119–126.
 - [37] A. Cabello, J. R. Portillo, A. Solís, and K. Svozil, Minimal true-implies-false and true-implies-true sets of propositions in noncontextual hidden-variable theories, Physical Review A **98**, 012106 (2018), arXiv:1805.00796.
 - [38] J. Schwinger, Unitary operators bases, Proceedings of the National Academy of Sciences (PNAS) **46**, 570 (1960).
 - [39] W. K. Wootters and B. D. Fields, Optimal state-determination by mutually unbiased measurements, Annals of Physics **191**, 363 (1989).
 - [40] S. Brierley, *Mutually Unbiased Bases in Low Dimensions*, Phd thesis, University of York, York, UK (2009), eTHoS ID: uk.bl.ethos.516399, Deposited: 01 Jun 2010, Last Modified: 08 Sep 2016.
 - [41] A. Klappenecker and M. Rötteler, Constructions of mutually unbiased bases, in *Finite Fields and Applications*, Lecture Notes in Computer Science, Vol. 2948, edited by G. Mullen, A. Poli, and H. Stichtenoth (Springer, Berlin, Heidelberg, 2004) pp. 262–266, arXiv:quant-ph/0309120.
 - [42] I. D. Ivanovic, Unbiased projector basis over c^3 , Physics Letters A **228**, 329 (1997).

Appendix A: Mutually unbiased bases in $\mathbb{C}^3$

For introductions to MUBs, see Refs. [32, 33, 38–41]. Write

$$\omega = e^{2\pi i/3}, \quad \omega^3 = 1, \quad 1 + \omega + \omega^2 = 0, \quad \bar{\omega} = \omega^2.$$

In $D = 3$, a complete MUB family contains four bases [42]. Omitting the common normalization factor

$1/\sqrt{3}$ from the last three bases, we use

$$\mathcal{B}_0 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}, \quad (\text{A1})$$

$$\mathcal{B}_1 = \{(1, 1, 1), (1, \omega, \omega^2), (1, \omega^2, \omega)\}, \quad (\text{A2})$$

$$\mathcal{B}_2 = \{(1, \omega, \omega), (1, \omega^2, 1), (1, 1, \omega^2)\}, \quad (\text{A3})$$

$$\mathcal{B}_3 = \{(1, \omega^2, \omega^2), (1, \omega, 1), (1, 1, \omega)\}. \quad (\text{A4})$$

Here $\mathcal{B}_0$ is computational and $\mathcal{B}_1$ is the Fourier basis. Direct use of $1 + \omega + \omega^2 = 0$ verifies orthogonality within each basis and $|\langle b, b' \rangle|^2 = 1/3$ between normalized vectors from different bases. The ordering of $\mathcal{B}_2$ and $\mathcal{B}_3$ agrees with the seed groups in Table I.

For the higher-dimensional argument, no catalogue of a complete MUB family is required: Propositions 2 and 3 use only the criterion $|x_1|^2 = \dots = |x_D|^2$ for one vector to be maximally unbiased relative to the computational basis.