Context–Color Matrices and Latin-Square Structure in Quantum Measurements

NN

(Dated: October 30, 2026)

We study common-spectrum spectral assignments for intertwined projective quantum measurements. All worked configurations have faithful orthogonal ray representations in the stated Hilbert space, with coordinates supplied here or in the companion inventory. For n contexts in dimension d, an $n \times d$ context–color matrix places the ray labels of each basis in one row and aligns shared projectors in their assigned spectral columns. Its columns are extendable two-valued states at the exact palette; wider palettes require blanks. Real three-dimensional examples exhibit nonextendable states, forced scalar identities between distinct rays, and Born probabilities outside the convex hull of color-class states. Faithful four-dimensional extensions and the realized Yu–Oh, complex Hesse, and real 65-ray configurations illustrate spectral obstructions with surviving ordinary states. Classical Graeco–Latin pairs are distinguished from vector-valued quantum Latin squares and from general color-signature tables. Explicit orthogonal pairs organize spectral assignments on a real four-level rook net and a complex seven-level Latin net. An explicit 49-ray configuration in $\mathbb{C}^7$ realizes a cyclic Latin net as 21 projective measurements; all pairwise orthogonalities and nonorthogonalities are certified exactly.

I. QUANTUM MEASUREMENTS AND COMMON-SPECTRUM ASSIGNMENTS

A complete rank-one projective measurement in $\mathbb{F}^d$, $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, is an orthonormal basis of rays with projectors $\{P_v : v \in e\}$ satisfying $\sum_{v\in e} P_v = I_d$. Fix distinct real numbers $\Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$. Assigning every outcome a different member of Σ defines a nondegenerate observable

$$A_e = \sum_{v\in e} \lambda_{c(v)} P_v. \tag{1}$$

Every operator commuting with this nondegenerate observable is a function of it [1, 2]. We study the additional demand that a projector shared by several measurements receives the same spectral label in each. Quantum mechanics permits observables to be specified separately; the common-spectrum assignment is the extra consistency condition under examination [3–5].

Let $H = (V, E)$ list the ray labels and selected complete measurements, with $V = \bigcup E$, $E \neq \varnothing$, $d \geq 2$, and $n = |E|$. Its two-section graph $\Gamma(H)$ joins rays occurring together in a context. A *faithful orthogonal representation* (FOR) assigns distinct normalized rays $[q_v] \in \mathbb{P}(\mathbb{F}^d)$ such that $q_u \perp q_v$ exactly when u, v are adjacent. Then $P_v = q_v q_v^\dagger$, and the listed contexts are complete projective measurements. Every worked example below has this geometric foundation. The complete coordinates for the companion configurations and their orthogonality checks are supplied in Ref. [5, 6]; the small introductory example, the pentagon, the Gaussian-integer three-dimensional row–column hub grids, the four-dimensional quaternion rook net, and the seven-dimensional Latin net have coordinates printed here.

A strong d-coloring $c : V \to \{0, \ldots, d-1\}$ is injective on every context and gives Eq. (1) consistently. We write $\mathcal{C}_d(H)$ for these assignments and $\chi(H)$ for the strong chromatic number. A context contains d mutually orthogonal rays, so $\chi(H) \geq d$. The condition $\chi(H) > d$ excludes the common-spectrum labeling; it does not conflict with the existing quantum coordinates.

A two-valued context state is $s : V \to \{0, 1\}$ with $\sum_{v\in e} s(v) = 1$ for every $e \in E$. Write $\mathcal{S}(H)$ for all such states and $\mathcal{S}_{\text{ext}}(H)$ for those occurring as a color class of some $c \in \mathcal{C}_d(H)$. A state extends precisely when the residual graph $\Gamma(H)[V \setminus \operatorname{supp} s]$ admits $d-1$ colors [5]. These binary assignments differ from Born probabilities $p_v = \operatorname{Tr}(\rho P_v)$, which satisfy the same context normalization but need not be convex mixtures of color-class states.

The scope is broader than Graeco–Latin schemes alone: the seven-level net and the four-level rook net have explicit classical orthogonal Latin pairs, while the other configurations illustrate the general context–color criterion. Neither a Latin-square incidence pattern nor a Graeco–Latin pair by itself supplies quantum coordinates. Section II develops the matrix representation and distinguishes the relevant notions of Latin structure. Section III gives an exact realized Latin net for a seven-level quantum system. Sections IV and V treat coordinatized three- and four-dimensional configurations. No uncoordinatized combinatorial construction is used as a worked example.

II. CONTEXT–COLOR MATRICES FOR PROJECTIVE MEASUREMENTS

A. Ray labels as matrix entries

Order the n actual measurement contexts as rows and use the d members of the common spectrum as column labels. Each entry is a ray label, not a vector component. A projector shared by several contexts must appear in the same column throughout. The number n of measurements is independent of the Hilbert-space dimension d.

Theorem 1 (Universal $n \times d$ Context–Color Matrix Criterion)**.** *Let $H = (V, E)$ be any finite d-uniform hypergraph with $n = |E|$ contexts $E = \{C_1, \ldots, C_n\}$. Then H admits a strong d-coloring ($\chi(H) = d$) if and only if the elements*

of each context C_i can be ordered into an $n \times d$ matrix

$$M = \begin{pmatrix} M_{1,0} & M_{1,1} & \cdots & M_{1,d-1} \\ M_{2,0} & M_{2,1} & \cdots & M_{2,d-1} \\ \vdots & \vdots & \ddots & \vdots \\ M_{n,0} & M_{n,1} & \cdots & M_{n,d-1} \end{pmatrix} \in V^{n\times d}, \tag{2}$$

satisfying the following two conditions:

1. Row Bijectivity (Rainbow Contexts): *For each row $i \in \{1,\dots,n\}$, the row entries $\{M_{i,0}, M_{i,1}, \dots, M_{i,d-1}\} = C_i$ are a permutation of the d distinct vertices of context C_i.*

2. Column Alignment (Noncontextual Intertwining Consistency): *Every vertex $v \in V$ appears in* exactly one column *of M:*

$$M_{i,j} = M_{i',j'} = v \quad \Longrightarrow \quad j = j'. \tag{3}$$

When Eqs. (2)–(3) hold, setting $c(M_{i,j}) = j$ defines a strong d-coloring $c \in \mathcal{C}_d(H)$, and the distinct vertices in each column $j \in \{0,\dots,d-1\}$,

$$S_j = \{M_{i,j} : 1 \le i \le n\} \subseteq V, \tag{4}$$

form d pairwise disjoint two-valued state supports $s_j = \mathbf{1}_{S_j} \in \mathcal{S}_{\text{ext}}(H)$ partitioning V. Moreover, the vertex degrees and column support sizes obey the exact conservation laws

$$\sum_{v\in S_j} \deg_H(v) = n \qquad (0 \le j \le d-1), \tag{5}$$

$$\sum_{j=0}^{d-1} |S_j| = |V|, \tag{6}$$

$$|S_j| = n - \sum_{v\in S_j} \big(\deg_H(v) - 1\big). \tag{7}$$

Proof. Given $c \in \mathcal{C}_d(H)$, each context C_i contains a unique vertex $v \in C_i$ of color $c(v) = j$; setting $M_{i,j} = v$ satisfies Row Bijectivity, and since $c(v)$ is well-defined on V, v appears only in column $j = c(v)$, establishing Eq. (3). Conversely, given M satisfying (1) and (2), Eq. (3) assigns a unique column index $c(v) = j$ to every $v \in V = \bigcup_{i=1}^{n} C_i$, and Row Bijectivity ensures that c is injective on each C_i. Because column j contains one entry from each of the n rows and shares no vertices with any column $j' \neq j$, its underlying vertex set S_j hits every context C_i exactly once, so $\mathbf{1}_{S_j} \in \mathcal{S}_{\text{ext}}(H)$. Each vertex $v \in S_j$ appears in $\deg_H(v)$ rows of column j, summing to n entries in column j [Eq. (5)] and yielding Eq. (7); summing $|S_j|$ over the d disjoint columns partitioning V gives Eq. (6). □

Remark 2 (Incidence equations and wider palettes). Let $B \in \{0,1\}^{n\times|V|}$ be the incidence matrix and let $X_{v,a} = 1$ mean that v has color a. The criterion is equivalently

$$X\mathbf{1}_d = \mathbf{1}_{|V|}, \qquad BX = J_{n,d}, \qquad X \in \{0,1\}^{|V|\times d}. \tag{8}$$




FIG. 1. Two projective measurements in $\mathbb{R}^3$ with the coordinates (9). Each numbered ray occurs once, and each colored line is a complete context. The two incidence rings at vertex 3 mark the shared projector. The associated 2×3 matrix is Eq. (10); line colors distinguish measurements, not eigenvalue labels.

If a palette of size $k > d$ is used, the corresponding $n \times k$ vertex matrix has $k-d$ blanks in each row, and its column supports are independent sets meeting each context *at most* once. They need not be states. Algebraically $X\mathbf{1}_k = \mathbf{1}_{|V|}$ and $BX \le J_{n,k}$. The state-column interpretation applies to the exact palette $k = d$.

The matrix describes the spectral labels on already specified quantum projectors. It does not reconstruct their coordinates. If $u \sim_c v$, meaning $c(u) = c(v)$ in every $c \in \mathcal{C}_d(H)$, the corresponding rays remain distinct; it is their allowed scalar labels that coincide. Likewise, a fixed state column may fail to admit the remaining $d-1$ columns, producing a nonextendable state of the realized measurement scenario.

B. A physical two-context schematic

Take the real rays

$$\begin{aligned} &r_1 = (1,0,0), \quad r_2 = (0,1,0), \quad r_3 = (0,0,1), \\ &r_4 = (1,1,0), \quad r_5 = (1,-1,0). \end{aligned} \tag{9}$$

Set $q_v = r_v/\|r_v\|$. The contexts $C_1 = \{1,2,3\}$ and $C_2 = \{3,4,5\}$ are orthonormal bases, sharing projector P_3. Direct dot products verify faithfulness. One consistent spectral assignment is

$$M = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 3 \end{pmatrix}, \qquad \begin{aligned} S_0 = \{1,4\}, \quad S_1 = \{2,5\}, \\ S_2 = \{3\}. \end{aligned} \tag{10}$$

Thus both observables use the same spectrum, and the shared projector has eigenvalue λ_2 in both. Figure 1 shows the two actual contexts; the repeated entry 3 in Eq. (10) represents one projector. This is the minimal measurement example needed to read the larger matrices. In all hypergraph drawings, plane positions are diagram coordinates; Hilbert-space orthogonality is certified by the supplied ray vectors.

C. Graeco–Latin schemes, quantum Latin squares, and color signatures

When a coordinatized scenario is indexed by a $d \times d$ cell array and its contexts include every row and column,

a strong d-coloring is a Latin square: each spectral label occurs once per row and column. A binary state selects one cell in every row and column and is therefore a permutation transversal. A classical *Graeco–Latin square* is the superposition of two Latin squares L and G of the same order for which all d^2 ordered pairs $(L(i,j), G(i,j))$ are distinct. This orthogonality concerns scalar labels, not inner products of quantum rays.

If the actual measurement family consists of the rows, columns, and symbol classes of L, states must meet every such class once: they are transversals of L. A strong d-coloring G is then precisely an orthogonal mate of L. In this interpretation, L indexes the third family of measurements and G gives the spectral labels. The array L is not itself a valid coloring of that full measurement family, since each L-context has a constant L-label. Section III realizes this distinction with explicit quantum coordinates.

A *quantum Latin square*, by contrast, is a $d \times d$ array of normalized vectors in $\mathbb{C}^d$ whose rows and columns are orthonormal bases [7]. It need not come with any classical Graeco–Latin pair, and its definition does not require all d^2 rays to be distinct or forbid additional orthogonalities. The distinctness and precise orthogonality graph of each physical example are established by separate FOR checks. No pair of orthogonal quantum Latin squares is asserted here.

For arbitrary coordinatized contexts, two chosen colorings define

$$\sigma(v) = \big(c_1(v), c_2(v)\big). \tag{11}$$

On a cell-indexed row–column scenario, this signature is injective precisely when the two Latin colorings form a classical Graeco–Latin pair. On a general quantum scenario it is simply a table of allowed spectral labels, with no implied Latin-square row or column conditions. Several distinct projectors may have one signature; Sec. IV B gives a physical example. Such a noninjective signature table is not a Graeco–Latin square. The $n \times d$ context–color matrix requires no literal Latin-square embedding.

III. A REALIZED GRAECO–LATIN SCHEME FOR SEVEN-LEVEL MEASUREMENTS

Take the cyclic Latin square $L(i,j) = i+j \pmod 7$. We construct forty-nine distinct rank-one projectors whose rows, columns, and symbol classes are twenty-one complete measurements of a seven-level quantum system. Put $\omega = e^{2\pi i/7}$ and define the permutation f by

x	0	1	2	3	4	5	6
$f(x)$	0	2	5	1	6	4	3.

(12)

For each $(i,j) \in \mathbb{Z}_7^2$, set

$$q_{ij} = \frac{1}{\sqrt{7}}\big(\omega^{\,if(x)+jx}\big)_{x=0}^{6} \in \mathbb{C}^7. \tag{13}$$

This is the Fourier quantum-Latin construction used in Ref. [8]; the specific permutation and the full faithfulness check below are explicit finite certificates for the present example. For instance, the exponent vectors of q_{00}, q_{01}, q_{10} are respectively $(0,0,0,0,0,0,0)$, $(0,1,2,3,4,5,6)$, and $(0,2,5,1,6,4,3)$. Equation (13) specifies all forty-nine vectors without numerical approximation.

Proposition 3 (An exact 49-ray FOR in $\mathbb{C}^7$)**.** *The vectors* (13) *represent* 49 *distinct rays and satisfy*

$$q_{ij} \perp q_{k\ell} \quad \Longleftrightarrow \quad \begin{array}{c} i = k \ \text{or}\ j = \ell \\ \text{or}\ i + j = k + \ell \pmod 7, \end{array} \tag{14}$$

for distinct cells. Thus they form a FOR of the cyclic 7*-uniform* 49*-vertex,* 21*-context net.*

Proof. The inner product is $7^{-1}\sum_x \omega^{af(x)+bx}$, where $a = k-i$ and $b = \ell - j$. A sum of seven seventh roots of unity vanishes exactly when its seven exponents occur once each: writing their multiplicities as $h_0, \ldots, h_6$, the irreducibility of $1 + z + \cdots + z^6$ makes $\sum h_r \omega^r = 0$ equivalent to $h_0 = \cdots = h_6 = 1$.

If $a = 0$ and $b \neq 0$, the exponents are a permutation. If $a \neq 0$, put $m = -b/a$. Directly from Eq. (12),

$$\big(|\operatorname{im}(f - m\,\mathrm{id})|\big)_{m=0}^{6} = (7,7,4,4,4,4,4).$$

Therefore the inner product vanishes exactly for $m = 0$ or $m = 1$, namely $b = 0$ or $a + b = 0$. Together with $a = 0$, these are precisely the three listed context directions. This checks nonedges as well as edges.

All first components are $1/\sqrt{7}$, so proportionality would mean equality. Equal vectors at different cells would give $af(x) + bx = 0$ for every x. If $a \neq 0$, this would make f linear, whereas $f(1) = 2$ and $f(2) = 5 \neq 4$. If $a = 0$, then also $b = 0$. Thus all rays are distinct. □

For a density operator ρ, the probabilities $p_{ij} = \operatorname{Tr}(\rho P_{ij})$ sum to one on every row, column, and symbol context. These are three families of seven projective measurement settings sharing identical rank-one projectors at their intersections.

The classical Graeco–Latin pair used here is explicitly

$$\begin{array}{ll} L(i,j) = i + j, & G(i,j) = i + 2j \pmod 7, \\ j = G - L, & i = 2L - G \pmod 7. \end{array} \tag{15}$$

Both arrays are Latin, and the inverse formulas prove that each ordered pair occurs once. On a fixed row, fixed column, or fixed L-symbol class, G takes all seven values. Thus $c(i,j) = G(i,j)$ assigns a common nondegenerate seven-element spectrum to all twenty-one observables via Eq. (1).

Crucially, the seven G-symbol classes are *color-class state supports*, not seven additional measurement bases. Two distinct cells in one such class have differences $(a,b) = (-2b, b)$ with $b \neq 0$, so none of $a = 0$, $b = 0$, or $a + b = 0$ holds; Eq. (14) makes their rays nonorthogonal. Figure 2 therefore has exactly 21 contexts, not 28. The separately

proved FOR, rather than classical orthogonality of (L, G) alone, establishes its physical realization.

The certificate checks all $\binom{49}{2} = 1176$ pairs exactly: 441 are orthogonal and the other 735 are nonorthogonal. Among the supplied rays, the only complete seven-ray bases are the 21 listed contexts. The classical mate G in Eq. (15) gives $\chi = 7$; there are $N_c = 635$ normalized seven-colorings and 133 transversal states, and all 133 states extend. By the group-table extension theorem [9, 10], in the Cayley table $L(i, j) = ij$ of any finite group K, every transversal $T = \{(i, \pi(i)) : i \in K\}$ extends to an orthogonal mate through its left row-translates $T_g = \{(gi, \pi(i)) : i \in K\}$ $(g \in K)$, which partition $K \times K$ into $|K|$ disjoint transversals. Indeed, their symbols $gi\pi(i) = g[i\pi(i)]$ still range over K, and each cell (u, j) belongs to the unique translate $g = u\pi^{-1}(j)^{-1}$. For the present additive table this reads $T_g = \{(g + i, \pi(i)) : i \in \mathbb{Z}_7\}$. Moreover, finite-field direction bounds explain why $d = 7$ and 21 contexts are sharp for prime-field Fourier nets of the form (13): over $\mathbb{F}_3$ and $\mathbb{F}_5$ every complete mapping is affine [11], collapsing cell rays to d rays, and in general any nonlinear function $f : \mathbb{F}_p \to \mathbb{F}_p$ determines at least $(p + 3)/2$ directions [12, 13], so $f(x) - mx$ can be a permutation for at most $r \le (p - 3)/2$ slopes m [14]. Thus $p = 7$ is the minimal prime dimension admitting a faithful Fourier 3-net ($r = 2$ slopes $m \in \{0, 1\}$ plus the vertical direction $a = 0$), and 3 parallel classes (21 contexts) is the maximum achievable in $\mathbb{C}^7$ by this ansatz. This minimum-prime statement concerns the Fourier 3-net construction only; it does not exclude the different four-dimensional rook scheme below or imply a general minimum dimension for quantum Latin squares. Thus this FOR provides a geometric baseline, *not* an example of a chromatic gap or of nonextendable states. It is a complex coordinatization; no real seven-dimensional realization is asserted.

Figure 2 draws this coordinatized net with the same numbered-node and context-ring convention as the earlier figures.

IV. REALIZED QUANTUM CONFIGURATIONS WITH AN EXACT COMMON SPECTRUM

The following configurations have faithful coordinates in dimensions three or four. We use their context matrices to analyze global spectral constraints and the distinction between ordinary states and color-class states. Full coordinate inventories and geometric checks for the companion configurations are given in Ref. [5, 6]; the pentagon, the Gaussian-integer row–column hub grids $R_{21,12}$ and $R_{25,15}$, and the four-dimensional quaternion rook net $H^{(4)}_{16,8}$ have complete coordinates printed below.

A. The Pentagon P_5 (5×3 Matrix and Purely Chromatic Ternary Exclusion)

The pentagon hypergraph P_5 ($d = 3, n = 5$ contexts, $|V| = 10$, Berge girth $g = 5$, real FOR in $\mathbb{R}^3$) has contexts

$$\begin{aligned} &C_1 = \{1, 6, 2\}, \quad C_2 = \{2, 7, 3\}, \quad C_3 = \{3, 8, 4\}, \\ &C_4 = \{4, 9, 5\}, \quad C_5 = \{5, 10, 1\}, \end{aligned} \tag{16}$$

One explicit FOR uses the following unnormalized real vectors, with $q_v = r_v/\|r_v\|$:

$$\begin{aligned} r_1 &= (1, 0, 0), & r_6 &= (0, 1, -1), \\ r_2 &= (0, 1, 1), & r_7 &= (2, -1, 1), \\ r_3 &= (1, 1, -1), & r_8 &= (1, 0, 1), \\ r_4 &= (1, -2, -1), & r_9 &= (5, 2, 1), \\ r_5 &= (0, 1, -2), & r_{10} &= (0, 2, 1). \end{aligned} \tag{17}$$

Their orthogonal pairs are exactly those of the five listed contexts. Figure 3 therefore depicts five projective measurements, with each shared ray represented once.

There are five degree-two cycle hinges $\{1, 2, 3, 4, 5\}$ and five degree-one midpoint vertices $\{6, 7, 8, 9, 10\}$.

In a 5×3 Context–Color Matrix $M \in V^{5\times 3}$, each hinge $h \in \{1, 2, 3, 4, 5\}$ has $\deg(h) = 2$ and saves 1 vertex when placed in a column [Eq. (7)]. Because the 5 hinges form an odd 5-cycle, at most $\lfloor 5/2 \rfloor = 2$ nonadjacent hinges can share a single column!

- *Why* $s_{11} = \{6, 7, 8, 9, 10\}$ *is nonextendable:* The state s_{11} uses $k = 0$ hinges (size $|S| = 5$). If s_{11} is placed in Column $\mathbf{c}_0$, the remaining 5 hinges $\{1, 2, 3, 4, 5\}$ must be distributed between only 2 columns $(\mathbf{c}_1, \mathbf{c}_2)$. Since each column can hold at most 2 hinges, two columns can hold at most $2+2 = 4 < 5$ hinges—an immediate Pigeonhole obstruction!

- *Why all other* 10 *states extend (*5 *normalized* 5×3 *matrices):* Every valid 5×3 matrix M must distribute the 5 hinges across the 3 columns as $2 + 2 + 1 = 5$ hinges, yielding column support sizes $(5 - 2) + (5 - 2) + (5 - 1) = 3 + 3 + 4 = 10 = |V|$. Fixing the anchor context $C_1 = (1, 6, 2)$ in Row 1 as $(\mathbf{c}_0, \mathbf{c}_1, \mathbf{c}_2) = (0, 1, 2)$ gives the $N_c = 5$ normalized 5×3 Context–Color Matrices (corresponding to the 5 choices of the uniquely colored hinge):




FIG. 2. Twenty-one complete projective measurements on 49 distinct rays in $\mathbb{C}^7$, with coordinates (13). Vertex v_{7i+j+1} represents q_{ij} and projector P_{ij}. Horizontal, vertical, and curved contexts fix i, j, and $i+j \pmod 7$, respectively. Proposition 3 proves exact faithfulness, including nonorthogonality of every unjoined pair. Each complete seven-ray basis among these rays is listed. The mate G of Eq. (15) supplies spectral labels; its symbol classes are not drawn as contexts. The configuration admits a common seven-value spectrum and all 133 ordinary states extend. Context colors identify measurements, not eigenvalue labels.




FIG. 3. The real three-dimensional pentagon measurement scenario, with exact coordinates (17). Each colored line is an orthonormal triad. Numbered circles are the ten distinct physical rays; concentric rings show shared outcomes. Black rims identify the terminals $6, 7, 9$, which cannot all receive one spectral label in any consistent three-value assignment. The rays and contexts remain well defined independently of that spectral restriction.

$$M_1^{(P_5)} = \begin{pmatrix} \mathbf{1} & 6 & \mathbf{2} \\ \mathbf{3} & 7 & \mathbf{2} \\ \mathbf{3} & \mathbf{4} & 8 \\ 9 & \mathbf{4} & \mathbf{5} \\ \mathbf{1} & 10 & \mathbf{5} \end{pmatrix}, \quad M_2^{(P_5)} = \begin{pmatrix} \mathbf{1} & 6 & \mathbf{2} \\ \mathbf{3} & 7 & \mathbf{2} \\ \mathbf{3} & 8 & \mathbf{4} \\ 9 & \mathbf{5} & \mathbf{4} \\ \mathbf{1} & \mathbf{5} & 10 \end{pmatrix},$$

$$M_3^{(P_5)} = \begin{pmatrix} \mathbf{1} & 6 & \mathbf{2} \\ 7 & \mathbf{3} & \mathbf{2} \\ \mathbf{4} & \mathbf{3} & 8 \\ \mathbf{4} & \mathbf{5} & 9 \\ \mathbf{1} & \mathbf{5} & 10 \end{pmatrix}, \quad M_4^{(P_5)} = \begin{pmatrix} \mathbf{1} & 6 & \mathbf{2} \\ 7 & \mathbf{3} & \mathbf{2} \\ \mathbf{4} & \mathbf{3} & 8 \\ \mathbf{4} & 9 & \mathbf{5} \\ \mathbf{1} & 10 & \mathbf{5} \end{pmatrix}, \qquad (18)$$

$$M_5^{(P_5)} = \begin{pmatrix} \mathbf{1} & 6 & \mathbf{2} \\ 7 & \mathbf{3} & \mathbf{2} \\ 8 & \mathbf{3} & \mathbf{4} \\ 9 & \mathbf{5} & \mathbf{4} \\ \mathbf{1} & \mathbf{5} & 10 \end{pmatrix}.$$

Inspection of Eq. (18) immediately proves the *purely chromatic ternary exclusion* $\neg(c(6) = c(7) = c(9))$: in any 5×3 matrix where $6, 7, 9$ all sit in Column $\mathbf{c}_1$, rows C_1, C_2, C_4

force the five hinges $\{1,2,3,4,5\}$ entirely into the two remaining columns $\mathbf{c}_0$ and $\mathbf{c}_2$, which can hold at most 4 hinges!

B. Faithful Yu–Oh Restrictions $H_{17,10}$ and $H_{21,13}$ and Their Color-Signature Table

The 17-ray, 10-context hypergraph $H_{17,10}$ retains the four base contexts $C_1 = \{1,2,3\}$, $C_2 = \{7,8,9\}$, $C_3 = \{4,5,6\}$, $C_4 = \{2,8,5\}$ of completed Yu–Oh H^+_{YO} together with the six contexts of diagonal sectors $h_1 = 11$ and $h_2 = 10$:

$$\begin{aligned}\mathcal{E}_{11} &= \big\{\{3,11,14\},\{7,11,15\},\{4,11,24\}\big\},\\ \mathcal{E}_{10} &= \big\{\{9,10,23\},\{1,10,19\},\{4,10,18\}\big\}.\end{aligned} \tag{19}$$

Fixing the anchor row $C_4 = (2,8,5)$ across columns $(\mathbf{c}_0,\mathbf{c}_1,\mathbf{c}_2)$, $H_{17,10}$ admits $N_c = 4$ normalized 10×3 Context–Color Matrices (Table S41 of Ref. [6]), shown here for c_1 and c_2:

$$M_1^{(H_{17,10})} = \left(\begin{array}{ccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 \\ \hline 2 & 1 & 3 \\ 9 & 8 & 7 \\ 4 & 6 & 5 \\ 2 & 8 & 5 \\ \hline 14 & \mathbf{11} & 3 \\ 15 & \mathbf{11} & 7 \\ 4 & \mathbf{11} & \mathbf{24} \\ \hline 9 & 23 & \mathbf{10} \\ 19 & 1 & \mathbf{10} \\ 4 & \mathbf{18} & \mathbf{10} \end{array}\right),\quad M_2^{(H_{17,10})} = \left(\begin{array}{ccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 \\ \hline 2 & 1 & 3 \\ 9 & 8 & 7 \\ 6 & 4 & 5 \\ 2 & 8 & 5 \\ \hline \mathbf{11} & 14 & 3 \\ \mathbf{11} & 15 & 7 \\ \mathbf{11} & 4 & \mathbf{24} \\ \hline 9 & 23 & \mathbf{10} \\ 19 & 1 & \mathbf{10} \\ \mathbf{18} & 4 & \mathbf{10} \end{array}\right). \tag{20}$$

In every coloring of $H_{17,10}$, the Specker-bug restriction forces $c(10) \neq c(11)$ (Sec. IV C). Both hubs are also adjacent to vertex 4, through the roof contexts $\{4,11,24\}$ and $\{4,10,18\}$. Thus $4,10,11$ always have three distinct colors. The first roof forces 24 into the color of 10, and the second forces 18 into the color of 11. Hence $10 \sim_c 24$ and $11 \sim_c 18$ in all four normalized matrices.

a. A direct quantum-probability consequence. The real coordinates of these two distinct rays are $r_{10} = (1,-1,1)$ and $r_{24} = (1,-1,2)$. Prepare $\rho = P_{24} = r_{24}r_{24}^T/6$. The Born rule gives

$$p_{24} = 1, \qquad p_{10} = \frac{|r_{10}\cdot r_{24}|^2}{\|r_{10}\|^2\|r_{24}\|^2} = \frac{8}{9}. \tag{21}$$

Every color-class state has $s(10) = s(24)$, so every convex mixture of extendable states has $p_{10} = p_{24}$. This physically available preparation lies outside that convex hull. In any decomposition into ordinary context states, a nonextendable total weight η must satisfy $|p_{24} - p_{10}| \le \eta$, and hence $\eta \ge 1/9$. The companion supplies the complete Born vector and an ordinary-state mixture attaining this bound [5, 6]. A forced scalar identity therefore does not identify the two quantum projectors.

Adjoining the third diagonal sector $\mathcal{E}_{13} = \{\{1,13,20\},\{7,13,21\},\{6,13,22\}\}$ yields $H_{21,13}$ ($n = 13$ contexts, $|V| = 21$). Only two normalized 13×3 Context–Color Matrices survive ($M_1^{(H_{21,13})}$ and $M_2^{(H_{21,13})}$, corresponding to the antipodal base patterns 101 and 010):

$$M_1^{(H_{21,13})} = \left(\begin{array}{ccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 \\ \hline 2 & 1 & 3 \\ 9 & 8 & 7 \\ 4 & 6 & 5 \\ 2 & 8 & 5 \\ \hline 14 & 11 & 3 \\ 15 & 11 & 7 \\ 4 & 11 & 24 \\ \hline 9 & 23 & 10 \\ 19 & 1 & 10 \\ 4 & 18 & 10 \\ \hline 13 & 1 & 20 \\ 13 & 21 & 7 \\ 13 & 6 & 22 \end{array}\right),\quad M_2^{(H_{21,13})} = \left(\begin{array}{ccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 \\ \hline 2 & 3 & 1 \\ 7 & 8 & 9 \\ 6 & 4 & 5 \\ 2 & 8 & 5 \\ \hline 14 & 3 & 11 \\ 7 & 15 & 11 \\ 24 & 4 & 11 \\ \hline 10 & 23 & 9 \\ 10 & 19 & 1 \\ 10 & 4 & 18 \\ \hline 20 & 13 & 1 \\ 7 & 13 & 21 \\ 6 & 13 & 22 \end{array}\right). \tag{22}$$

The joint-signature criterion of Sec. II C applies as follows: each vertex $v \in V(H_{21,13})$ receives a 2-tuple chromatic signature $\sigma(v) = (c_1(v), c_2(v)) \in \{0,1,2\}^2$ recording its column index in $M_1^{(H_{21,13})}$ and $M_2^{(H_{21,13})}$. Placing the 21 vertices into the 3×3 signature grid $\{0,1,2\}\times\{0,1,2\}$ yields the 9 chromatic equivalence classes:

$c_2 \backslash c_1$	$\mathbf{c}_0 = 0$	$\mathbf{c}_1 = 1$	$\mathbf{c}_2 = 2$
0	$\{2,14\}$	$\{6\}$	$\{7,10,20,24\}$
1	$\{4,13,15,19\}$	$\{8,23\}$	$\{3\}$
2	$\{9\}$	$\{1,11,18,21\}$	$\{5,22\}$

(23)

A literal 3×3 Graeco–Latin square assigns one distinct cell to each of nine ordered symbol pairs. Equation (23) is instead a 3×3 array of signature fibers: $H_{21,13}$ has 21 vertices and only two normalized colorings, forcing multiple vertices to collide in six of the nine cells and producing $3\binom{4}{2} + 3\binom{2}{2} = 21$ chromatically inseparable pairs. Each pair consists of distinct faithfully represented quantum rays; the common signature constrains their scalar labels, not their coordinates. Consequently, this quantum configuration is not a Graeco–Latin square scheme; the square-shaped signature table is only an analysis device.

C. Three-Dimensional Logical Gadgets $B_{13,7}, D_{16,9}, I_{16,9}, T_{19,11}$

The remaining three-dimensional gadgets in the companion inventory [5] are sub-hypergraphs or roof exten-

sions of $H_{17,10}$, and their $n \times 3$ Context–Color Matrices directly exhibit how TIFS and TITS constraints operate:

1. *Completed Specker Bug $B_{13,7}$ (7×3 matrix, $N_c = 6$):* Deleting $C_3 = \{4,5,6\}$, $\{4,11,24\}$, and $\{4,10,18\}$ from $H_{17,10}$ leaves the 7 contexts of $B_{13,7}$:

   $$\{1,2,3\},\ \{7,8,9\},\ \{2,8,5\},\ \{3,11,14\},$$
   $$\{7,11,15\},\ \{9,10,23\},\ \{1,10,19\}.$$

   Suppose 10 and 11 had one common color h. Their four neighbors $1,3,7,9$ would all avoid h. If $h \neq c(2)$, the context $\{1,2,3\}$ could not contain h, contrary to its being rainbow. If $h = c(2)$, then $h \neq c(8)$ because $2,8$ share a context; now $\{7,8,9\}$ cannot contain h. Both cases contradict strong coloring. Thus $c(10) \neq c(11)$, without making a palette assumption incompatible with the fixed anchor.

2. *Completed Γ_1 Gadget $I_{16,9}$ (9×3 matrix, $N_c = 6$):* Re-adjoining the two roof contexts $\{4,11,24\}$ and $\{4,10,18\}$ to $B_{13,7}$ (without $\{4,5,6\}$) appends two rows to each of the 6 matrices of $B_{13,7}$. Since 10 and 11 always occupy distinct columns in $B_{13,7}$, the apex 4 is forced into the third column, locking 24 into the column of 10 ($10 \sim_c 24$) and 18 into the column of 11 ($11 \sim_c 18$).

3. *Purely Chromatic Binary TIFS $D_{16,9}$ (9×3 matrix, $N_c = 7$) and Two-Way TITS $T_{19,11}$ (11×3 matrix, $N_c = 7$):* Deleting $\{1,10,19\}$ from $H_{17,10}$ yields $D_{16,9}$. While the nonextendable state $s = \{1,5,7,14,18,23,24\} \in \mathcal{S}(D_{16,9})$ contains both 14 and 18, no 9×3 Context–Color Matrix can place 14 and 18 in the same column: if $14,18 \in \mathbf{c}_0$, then $\{3,11,14\},\{4,11,24\},\{4,10,18\}$ force $3,4$ into one column (say $\mathbf{c}_1$) and $10,11$ into $\mathbf{c}_2$, which forces $7,9 \notin \mathbf{c}_2 \implies 8 \in \mathbf{c}_2$, and $2,5 \notin \mathbf{c}_1 \implies 2,5 \in \mathbf{c}_0$, colliding in Row 4 ($\{2,8,5\}$)! Adjoining the two roof contexts $\{14,26,27\}$ and $\{18,26,28\}$ in $T_{19,11}$ appends two rows to the 7 normalized 9×3 matrices of $D_{16,9}$; since 14 and 18 always sit in distinct columns, their common neighbor 26 must occupy the third column, locking 28 into the column of 14 ($14 \sim_c 28$) and 27 into the column of 18 ($18 \sim_c 27$).

D. Gaussian-Integer Row–Column Hub Grids $R_{21,12}$ and $R_{25,15}$ in $\mathbb{C}^3$ (Nonextendable Permutation Transversals)

While a full 3×3 Latin 3-net cannot be faithfully realized by nine rays in $\mathbb{F}^3$ without collapsing cells to three rays, replacing the column contexts by the neighborhoods $N(P_k) = \{M_{0k}, M_{1k}, M_{2k}\}$ of three column hubs P_0, P_1, P_2 yields a faithful 21-ray, 12-context configuration $R_{21,12}$ in $\mathbb{C}^3$ whose nonextendable states are precisely the $3! = 6$ permutation transversals of its 3×3 core.

Arrange the nine core vertices $M_{ak} = 3a + k + 1$ ($a, k \in \{0,1,2\}$) as the rows $\mathcal{M}_0 = \{1,2,3\}$, $\mathcal{M}_1 = \{4,5,6\}$, $\mathcal{M}_2 = \{7,8,9\}$, let $P_0 = 10, P_1 = 11, P_2 = 12$ be the column hubs, and let $B_{ak} = 3a + k + 13 \in \{13,\dots,21\}$ complete the nine column-hub contexts $\mathcal{B}_{ak} = \{M_{ak}, P_k, B_{ak}\}$. Adjoining the apex $z = 22$ and completion rays $A_0 = 23, A_1 = 24, A_2 = 25$ in the three contexts $\mathcal{A}_k = \{z, P_k, A_k\}$ gives the 25-ray, 15-context configuration $R_{25,15}$.




FIG. 4. The Gaussian-integer row–column hub grid $R_{25,15}$ in $\mathbb{C}^3$ (25 rays, 15 contexts) and its 21-ray core $R_{21,12}$ (omitting apex 22 and $23,24,25$), with coordinates (25). The three horizontal lines are the row contexts $\mathcal{M}_0 = \{1,2,3\}$, $\mathcal{M}_1 = \{4,5,6\}$, $\mathcal{M}_2 = \{7,8,9\}$; the three three-pronged fans join the column hubs $P_0 = 10, P_1 = 11, P_2 = 12$ to the grid columns via $B_{ak} \in \{13,\dots,21\}$; the top fan joins the apex $z = 22$ to P_0, P_1, P_2 via $23,24,25$. Black rims mark the diagonal permutation transversal $\{1,5,9\}$, one of the $3! = 6$ nonextendable permutation-transversal states of $R_{21,12}$.

Fix the real hub rays $P_0 = (0,1,-1)$, $P_1 = (-1,0,1)$, $P_2 = (1,-1,0)$, all orthogonal to $z = (1,1,1)$. For any real parameter $t \in \mathbb{R}$, set $\alpha(t) = -1 + it$. The three vectors

$$v_0(t) = \big(\alpha(t),2,2\big), \qquad v_1(t) = \big(2,\alpha(t),2\big), \qquad v_2(t) = \big(2,2,\alpha(t)\big) \tag{24}$$

satisfy $\langle v_j(t), v_k(t)\rangle = 2(-1-it) + 2(-1+it) + 4 = 0$ ($j \neq k$) and $\langle v_k(t), P_k\rangle = 0$ for every $t \in \mathbb{R}$. Choosing $t = \infty$ (the standard basis) for $\mathcal{M}_0$, $t = 1$ ($\alpha = -1 + i$) for $\mathcal{M}_1$, and $t = -2$ ($\alpha = -1 - 2i$) for $\mathcal{M}_2$ gives an exact Gaussian-integer FOR $r_v \in \mathbb{Z}[i]^3$ ($q_v = r_v/\|r_v\|$) requiring

no cube roots of unity:

$$
\begin{aligned}
r_1 &= (1,0,0), & r_2 &= (0,1,0),\\
r_3 &= (0,0,1), & r_4 &= (-1+i,2,2),\\
r_5 &= (2,-1+i,2), & r_6 &= (2,2,-1+i),\\
r_7 &= (-1-2i,2,2), & r_8 &= (2,-1-2i,2),\\
r_9 &= (2,2,-1-2i), & r_{10} &= (0,1,-1),\\
r_{11} &= (-1,0,1), & r_{12} &= (1,-1,0),\\
r_{13} &= (0,1,1), & r_{14} &= (1,0,1),\\
r_{15} &= (1,1,0), & r_{16} &= (2-2i,1,1),\\
r_{17} &= (1,2-2i,1), & r_{18} &= (1,1,2-2i),\\
r_{19} &= (4,1-2i,1-2i), & r_{20} &= (1-2i,4,1-2i),\\
r_{21} &= (1-2i,1-2i,4), & r_{22} &= (1,1,1),\\
r_{23} &= (2,-1,-1), & r_{24} &= (-1,2,-1),\\
r_{25} &= (-1,-1,2).
\end{aligned}
\tag{25}
$$

All 25 rays are projectively distinct, and their orthogonal pairs are exactly those of the 12 contexts of $R_{21,12}$ (on $\{1,\dots,21\}$) and the 15 contexts of $R_{25,15}$ (on $\{1,\dots,25\}$), shown in Fig. 4.

- *Why the* $3! = 6$ *permutation transversals of the* 3×3 *core are the exact nonextendable states of* $R_{21,12}$ *(*$N_c = 32$*,* $|\mathcal{S}_{\text{ext}}|/|\mathcal{S}| = 48/54$*):* Any state $s \in \mathcal{S}(R_{21,12})$ picks one vertex M_{a,k_a} in each row $\mathcal{M}_a$ ($3^3 = 27$ choices) and sets $s(P_k) = 0$ on every used column $k \in \{k_0, k_1, k_2\}$. When k_0, k_1, k_2 are a permutation of $\{0,1,2\}$ ($3! = 6$ states), all three columns are used, forcing $s(P_0) = s(P_1) = s(P_2) = 0$. Suppose such a permutation-transversal state s were placed in Column $\mathbf{c}_0$ of a 12×3 Context–Color Matrix:

  1. Each grid column $k \in \{0,1,2\}$ has one vertex in $\mathbf{c}_0$ and two remaining vertices in $\mathbf{c}_1 \cup \mathbf{c}_2$.
  2. Since $P_k \notin \mathbf{c}_0$, the hub P_k sits in $\mathbf{c}_1$ or $\mathbf{c}_2$ and (via $\mathcal{B}_{ak}$) bars all three vertices of grid column k from $c(P_k)$, forcing *both* remaining vertices of grid column k into the single remaining matrix column $\{\mathbf{c}_1, \mathbf{c}_2\} \setminus \{c(P_k)\}$.
  3. Consequently, each grid column $k \in \{0,1,2\}$ contributes an *even* number (0 or 2) of grid vertices to matrix column $\mathbf{c}_1$, so the total number of grid vertices in $\mathbf{c}_1$ is even ($\in \{0,2,4,6\}$).
  4. Yet the three disjoint row contexts $\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$ partition the nine grid vertices and require $\mathbf{c}_1$ to contain *exactly* 3 *(odd)* grid vertices—a parity contradiction!

- *Apex-extended grid* $R_{25,15}$ *(*15×3 *matrix,* $N_c = 24$*,* $|\mathcal{S}_{\text{ext}}|/|\mathcal{S}| = 48/81$*):* Adjoining $\mathcal{A}_k = \{22, P_k, A_k\}$ bars P_0, P_1, P_2 from sharing all three colors, while the row–column structure bars $c(P_0) = c(P_1) = c(P_2)$. Thus two of P_0, P_1, P_2 share one color and the third takes the remaining non-$c(22)$ color, yielding the $N_c = 3\times 2^3 = 24$ normalized 15×3 Context–Color Matrices (realizing the forced grid pattern of Eq. (36) prior to the cyclic Latin transversal obstruction in Sec. V B). One explicit 15×3 matrix (whose top 12 rows form $M_1^{(R_{21,12})}$) is:

$$
M_1^{(R_{25,15})} = \left(\begin{array}{ccc}
\mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 \\ \hline
1 & 2 & 3 \\
4 & 5 & 6 \\
7 & 8 & 9 \\ \hline
1 & 13 & \mathbf{10} \\
14 & 2 & \mathbf{11} \\
15 & \mathbf{12} & 3 \\
4 & 16 & \mathbf{10} \\
17 & 5 & \mathbf{11} \\
18 & \mathbf{12} & 6 \\
7 & 19 & \mathbf{10} \\
20 & 8 & \mathbf{11} \\
21 & \mathbf{12} & 9 \\ \hline
\mathbf{22} & 23 & \mathbf{10} \\
\mathbf{22} & 24 & \mathbf{11} \\
\mathbf{22} & \mathbf{12} & 25
\end{array}\right).
\tag{26}
$$

E. Four-Dimensional Apex Lifts and Strictly Linear Gadgets ($n\times 4$ Matrices)

In dimension $d = 4$, the $n\times 4$ Context–Color Matrix separates into two distinct structural classes (the companion four-dimensional inventory [5]):

Proposition 4 (Common-apex lift in arbitrary dimension)**.** *Let* H *be* d_0*-uniform and adjoin* $r = d - d_0$ *common fresh vertices* A *to every context. Then*

$$\Gamma(H^A) = \Gamma(H) \vee K_r, \qquad \chi(H^A) = \chi(H) + r.$$

Its states are the old states extended by zero on A*, together with the* r *singleton apex states. If* $\chi(H) = d_0$*, its extendable states transform in the same way, so the old nonextendable states and forced color identities persist. Any injective faithful realization in* $\mathbb{F}^{d_0}$ *extends faithfully into* $\mathbb{F}^d$ *by adjoining coordinate rays.*

Proof. All new vertices are adjacent to each other and to every old vertex, proving the graph join and chromatic formula. A state either selects one apex, forcing all old vertices to zero, or selects no apex and restricts to an old state. In an exact-palette coloring the apices use r distinct colors absent from the old vertices, leaving precisely d_0 colors on H. Explicitly, embed each old vector as $(q_v, 0_r)$ and use $e_{d_0+1}, \dots, e_d$ for the apices. Every enlarged context is an orthonormal basis, and the construction preserves exactly the old orthogonalities and adds the required apex pairs. □

1. *Non-Linear Common-Apex Lifts $W^{(4)}_{11,5}$, $B^{(4)}_{14,7}$, and $D^{(4)}_{17,9}$*

Each common-apex lift adjoins a single apex vertex 0 to every 3-element context C_i of a 3-dimensional gadget H, forming $C_i^{(4)} = \{0\} \cup C_i$. In Theorem 1, since $0 \in \bigcap_{i=1}^{n} C_i^{(4)}$ has degree $\deg(0) = n$, Eq. (7) gives $|S_0| = n - (n-1) = 1$. Thus *every* $n \times 4$ Context–Color Matrix $M^{(4)}$ of an apex lift is obtained by prepending a constant Column $\mathbf{c}_0 = (0, 0, \ldots, 0)^T$ to a valid $n \times 3$ Context–Color Matrix $M^{(3)}$ of H:

$$M^{(4)} = \left(\begin{array}{c|ccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{c}_3 \\ \hline 0 & & & \\ 0 & & M^{(3)} & \\ \vdots & & & \\ 0 & & & \end{array}\right) \in V^{n\times 4}. \tag{27}$$

Equation (27) provides a one-line matrix proof of the apex-extension proposition of Ref. [5]: the normalized coloring count N_c is unchanged, the singleton column $S_0 = \{0\}$ is always an extendable state, and the remaining three columns $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$ are identical to those of $M^{(3)}$.

2. *Strictly Linear Four-Dimensional Gadgets $W^{(4)}_{16,6}$, $L^{(4)}_{18,7}$, $L^{(4)}_{22,9}$, and $H^{(4)}_{16,8}$*

To restore linearity ($|e \cap e'| \le 1$, so $g(H) \ge 3$), Ref. [5] constructs three faithful linear 4-uniform gadgets in $\mathbb{R}^4$:

- *Linear 5-Wheel $W^{(4)}_{16,6}$ (6×4 matrix, $N_c = 80$):* The 6 contexts are

$$C_1 = \{2, 0, 6, 1\}, \quad C_2 = \{2, 7, 13, 3\},$$
$$C_3 = \{3, 0, 8, 4\}, \quad C_4 = \{4, 9, 14, 5\},$$
$$C_5 = \{5, 10, 15, 1\}, \quad C_6 = \{0, 11, 12, 5\}.$$

  Here vertex 0 lies in only three contexts (C_1, C_3, C_6). In any 6×4 matrix M, placing 0 in Column $\mathbf{c}_3$ covers 0 in Rows 1, 3, 6. If the three terminals 6, 8, 11 were all placed in Column $\mathbf{c}_0$, then in Rows 1, 3, 6 the outer cycle vertices $\{1, 2, 3, 4, 5\}$ would be barred from both $\mathbf{c}_0$ and $\mathbf{c}_3$! Thus $\{1, 2, 3, 4, 5\}$ would have to be distributed across only two columns $(\mathbf{c}_1, \mathbf{c}_2)$ while pairwise sharing rows around the 5-cycle $C_1 - C_2 - C_3 - C_4 - C_5$, which is impossible. One explicit 6×4 matrix is:

$$M_1^{(W^{(4)}_{16,6})} = \begin{pmatrix} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{c}_3 \\ \hline 2 & \mathbf{0} & \mathbf{6} & 1 \\ 2 & 7 & 3 & 13 \\ \mathbf{8} & \mathbf{0} & 3 & 4 \\ 5 & 9 & 14 & 4 \\ 5 & 10 & 15 & 1 \\ 5 & \mathbf{0} & \mathbf{11} & 12 \end{pmatrix}. \tag{28}$$

- *Linear Quaternion TIFS $L^{(4)}_{18,7}$ (7×4 matrix, $N_c = 144$, $\mathcal{S}_{\text{ext}} = \mathcal{S} = 68$) and Linear TITS $L^{(4)}_{22,9}$ (9×4 matrix, $N_c = 216$):* In $L^{(4)}_{18,7}$, the 7 contexts are

$$C_1 = \{1, 2, 4, 3\}, \; C_2 = \{5, 8, 7, 6\}, \; C_3 = \{4, 9, 10, 8\},$$
$$C_4 = \{11, 1, 12, 5\}, \; C_5 = \{11, 14, 13, 2\},$$
$$C_6 = \{15, 3, 16, 7\}, \; C_7 = \{15, 17, 18, 6\}.$$

  Why can 11 and 15 never share a column in a 7×4 matrix (or even a single state column)? If $11, 15 \in \mathbf{c}_0$, then Rows 4, 5, 6, 7 bar $\{1, 2, 3\}$ and $\{5, 6, 7\}$ from $\mathbf{c}_0$; then Row 1 ($\{1, 2, 4, 3\}$) forces $4 \in \mathbf{c}_0$ and Row 2 ($\{5, 8, 7, 6\}$) forces $8 \in \mathbf{c}_0$, so 4 and 8 collide in Column $\mathbf{c}_0$ of Row 3 ($\{4, 9, 10, 8\}$)! In $L^{(4)}_{22,9}$, adjoining $C_8 = \{11, 21, 22, 19\}$ and $C_9 = \{13, 20, 15, 19\}$ whose listed context C_9 includes the pair $\{13, 15\}$ ensures that $\{11, 13, 15, 19\}$ occupy all four distinct columns, locking $20 \in C_9$ into the column of 11 ($11 \sim_c 20$). An explicit 9×4 Context–Color Matrix (with its first context fixed as $(1, 2, 3, 4)$) is:

$$M_1^{(L^{(4)}_{22,9})} = \begin{pmatrix} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{c}_3 \\ \hline 1 & 2 & 3 & 4 \\ 7 & 5 & 8 & 6 \\ 9 & 10 & 8 & 4 \\ 1 & 5 & \mathbf{11} & 12 \\ 13 & 2 & \mathbf{11} & 14 \\ 7 & 15 & 3 & 16 \\ 17 & 15 & 18 & 6 \\ 21 & 22 & \mathbf{11} & 19 \\ 13 & 15 & \mathbf{20} & 19 \end{pmatrix}. \tag{29}$$

  Notice how Column $\mathbf{c}_2$ in Eq. (29) locks **11** (in Rows 4, 5, 8) and **20** (in Row 9) into the same state column $S_2 = \{3, 8, 11, 18, 20\}$.

- *Quaternion Four-by-Four Rook Net $H^{(4)}_{16,8}$ (8×4 matrix, $N_c = 24$, $|\mathcal{S}_{\text{ext}}|/|\mathcal{S}| = 24/24$):* Identify $\mathbb{R}^4$ with the quaternion algebra $\mathbb{H}$ in the standard basis $e_0 = (1, 0, 0, 0)$, $e_1 = (0, 1, 0, 0)$, $e_2 = (0, 0, 1, 0)$, $e_3 = (0, 0, 0, 1)$. For any integer quaternion $w = (w_0, w_1, w_2, w_3) \in \mathbb{Z}^4 \setminus \{0\}$, define the 16 cell vectors $M_{a,k} = (e_a \cdot w) \cdot e_k$ $(a, k \in \{0, 1, 2, 3\})$, labeled by $v = 4a + k + 1 \in \{1, \ldots, 16\}$. Multiplication by a fixed quaternion preserves orthogonality and scales squared norms by its squared norm, so the four rows $R_a = \{4a+1, 4a+2, 4a+3, 4a+4\}$ and four columns $C_k = \{k+1, k+5, k+9, k+13\}$ are eight orthogonal bases of $\mathbb{R}^4$ (the normalized array $M_{a,k}/\|w\|$ is a real quantum Latin square [7]), while for $a \ne b$ and $k \ne \ell$,

$$\langle M_{a,k}, M_{b,\ell} \rangle = \pm R(w)_{a\oplus b,\, k\oplus \ell}, \tag{30}$$

  where $\oplus$ is the Klein 4-group addition on $\{0, 1, 2, 3\}$ and $R(w)$ is the 3×3 integer Cayley rotation matrix




FIG. 5. The four-dimensional quaternion rook net $H^{(4)}_{16,8}$ in $\mathbb{R}^4$ (16 rays, 8 contexts), with coordinates (31). Horizontal and vertical lines are the four row contexts and four column contexts. Black rims mark the diagonal transversal $S_0 = \{1, 6, 11, 16\}$ (Column $\mathbf{c}_0$ of Eq. (32)).

of w. Setting $w = (1, 1, 1, 2)$ ($\|w\|^2 = 7$) makes all nine entries of $R(w)$ nonzero ($\pm 2, \pm 3, \pm 6$), certifying exact faithfulness of all 16 rays in $\{\pm 1, \pm 2\}^4$:

$$\begin{aligned}
r_1 &= (1,1,1,2), & r_2 &= (-1,1,2,-1),\\
r_3 &= (-1,-2,1,1), & r_4 &= (-2,1,-1,1),\\
r_5 &= (-1,1,-2,1), & r_6 &= (-1,-1,1,2),\\
r_7 &= (2,-1,-1,1), & r_8 &= (-1,-2,-1,-1),\\
r_9 &= (-1,2,1,-1), & r_{10} &= (-2,-1,-1,-1),\\
r_{11} &= (-1,1,-1,2), & r_{12} &= (1,1,-2,-1),\\
r_{13} &= (-2,-1,1,1), & r_{14} &= (1,-2,1,-1),\\
r_{15} &= (-1,-1,-2,-1), & r_{16} &= (-1,1,1,-2).
\end{aligned} \tag{31}$$

As stated in Sec. II C, its $N_c = 24$ normalized 8×4 Context–Color Matrices are the 24 normalized 4×4 Latin squares, and its 24 states are the $4! = 24$ permutation transversals (Fig. 5). One explicit 8×4 matrix (corresponding to the Klein 4-group Latin square $c(a, k) = a \oplus k$) is:

$$M_1^{(H^{(4)}_{16,8})} = \left(\begin{array}{cccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{c}_3 \\ \hline 1 & 2 & 3 & 4 \\ 6 & 5 & 8 & 7 \\ 11 & 12 & 9 & 10 \\ 16 & 15 & 14 & 13 \\ \hline 1 & 5 & 9 & 13 \\ 6 & 2 & 14 & 10 \\ 11 & 15 & 3 & 7 \\ 16 & 12 & 8 & 4 \end{array}\right). \tag{32}$$

This physical rook net also supports a literal classical Graeco–Latin scheme. Write the cell indices as the Klein group $\mathbb{F}_2^2$ with addition $\oplus$, and set

$$\begin{aligned} L_4(a,k) &= a \oplus k, \\ G_4(a,k) &= a \oplus A(k), \\ \big(A(0), A(1), A(2), A(3)\big) &= (0,2,3,1). \end{aligned} \tag{33}$$

Both arrays are Latin. Their ordered pairs determine k because $L_4 \oplus G_4 = k \oplus A(k)$ takes the four distinct values $(0, 3, 1, 2)$, and then $a = L_4 \oplus k$. Hence they are orthogonal. Each separately labels every row and column measurement with the same four-element spectrum. The matrix (32) uses L_4; replacing its column assignment by G_4 gives the second observable family on the same sixteen projectors.

Here both L_4 and G_4 are colorings of the eight-context scenario. Their symbol classes are permutation-transversal state supports, whose distinct rays are nonorthogonal by the rook FOR. They therefore add no measurement contexts. This differs from Sec. III, where L indexes an actual third measurement family and G supplies one allowed spectral coloring. The two classical scalar-label arrays here are also distinct objects from the single vector-valued quantum Latin square supplied by the quaternion coordinates.

Moreover, Eq. (30) shows how to engineer selective cross-diagonal orthogonalities in $\mathbb{R}^4$: choosing $w = (1, 2, 3, 6)$ satisfies $w_1 w_2 = w_0 w_3$ ($R_{12} = 0$) while keeping the other eight entries of $R(w)$ nonzero, creating exact orthogonalities $M_{a,k} \perp M_{a\oplus 1, k\oplus 2}$ along Klein-group transversals that can be completed by two-ray bridges.

V. REALIZED QUANTUM SPECTRAL OBSTRUCTIONS

The configurations in this section have actual faithful quantum coordinates even though no exact-palette context–color matrix exists. Their spectral inconsistency is compatible with the quantum measurements themselves. Coordinates and all context inventories are supplied in Ref. [5, 6].

A. Completed Yu–Oh H^+_{YO} (16×3 Obstruction) and $H^{(4)}_{\mathrm{YO}}$

Completed Yu–Oh H^+_{YO} has $n = 16$ contexts of size $d = 3$ on $|V| = 25$ vertices: the 4×3 base block $E_{\mathrm{base}} = \{\{1, 2, 3\}, \{7, 8, 9\}, \{4, 5, 6\}, \{2, 8, 5\}\}$ and four 3×3 diagonal sector blocks $\mathcal{E}_h$ ($h \in \{10, 11, 12, 13\}$), where each $\mathcal{E}_h$ consists of three rows $\{x, h, w_{xh}\}$ for $x \in N(h)$.

In a 16×3 Context–Color Matrix M:

1. In each 3×3 sector block $\mathcal{E}_h$, the hub h appears in all three rows and must therefore occupy a *single column* $j_h \in \{\mathbf{c}_0, \mathbf{c}_1, \mathbf{c}_2\}$.

2. By Row Bijectivity, none of the three neighbors $x \in N(h)$ can occupy column j_h! That is, the three neighbors $N(h) \subset \{1,3,4,6,7,9\}$ are restricted to the *two remaining columns* $\{0,1,2\} \setminus \{j_h\}$.

3. However, in the 4×3 base block (fixing Row 4 as $(2,8,5)$), there are only $2^3 = 8$ ways to order the pairs $\{1,3\},\{7,9\},\{4,6\}$ across the two available columns in Rows 1, 2, and 3:

$$M_{\text{base}}(b_1,b_2,b_3) = \left(\begin{array}{ccc} \mathbf{c}_0 & \mathbf{c}_1 & \mathbf{c}_2 \\ \hline 2 & 1 \text{ or } 3 & 3 \text{ or } 1 \\ 9 \text{ or } 7 & 8 & 7 \text{ or } 9 \\ 6 \text{ or } 4 & 4 \text{ or } 6 & 5 \\ 2 & 8 & 5 \end{array}\right). \tag{34}$$

As shown in the base-coloring partition theorem of Ref. [5], each of the 8 choices of $(b_1,b_2,b_3) \in \{0,1\}^3$ places one of the four triples $N(10) = \{1,9,4\}$, $N(11) = \{3,7,4\}$, $N(12) = \{3,9,6\}$, or $N(13) = \{1,7,6\}$ across *all three columns* $\mathbf{c}_0, \mathbf{c}_1, \mathbf{c}_2$—leaving *zero* available columns for the corresponding hub h_j!

Consequently, no 16×3 Context–Color Matrix exists ($\chi(H^+_{\text{YO}}) = 4$). Appending the common apex 26 in $H^{(4)}_{\text{YO}}$ prepends the singleton column $\mathbf{c}_0 = (26,\ldots,26)^T$ as in Eq. (27), shifting the exact same obstruction to 16×4 ($\chi(H^{(4)}_{\text{YO}}) = 5$).

B. The 37-Ray Complex Hesse Hypergraph H_{37}: A Latin-Neighborhood Obstruction

The 37-ray, 24-context complex Hesse hypergraph H_{37} in $\mathbb{C}^3$ (the Hesse section of Ref. [5]) provides a context hypergraph whose hub neighborhoods follow Latin-square patterns. Its obstruction uses those neighborhood constraints, rather than a pair of orthogonal Latin colorings.

Consider the 9 core vertices M_{ak} $(a,k \in \mathbb{Z}_3)$, arranged as a 3×3 grid:

M_{ak}	$k=0$	$k=1$	$k=2$
$a=0$ $(\mathcal{M}_0)$	M_{00} (2)	M_{01} (3)	M_{02} (4)
$a=1$ $(\mathcal{M}_1)$	M_{10} (5)	M_{11} (6)	M_{12} (7)
$a=2$ $(\mathcal{M}_2)$	M_{20} (8)	M_{21} (9)	M_{22} (10)

(35)

The grid columns $N(P_k)$ and cyclic symbol classes $N(Q_k)$ below are neighborhoods, not contexts. They must omit a color, whereas the actual row contexts must be rainbow. Thus these patterns do not make the core into a 3-net hypergraph H_L or a literal Graeco–Latin measurement scheme.

Look at how the 24 contexts of H_{37} [the listed Hesse contexts in Ref. [5]] act on this 3×3 grid in a putative 24×3 Context–Color Matrix:

1. *Rows of the* 3×3 *Grid* ($\mathcal{M}_a = \{M_{a0}, M_{a1}, M_{a2}\}$): The first 3 contexts $\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$ are the *rows* of Eq. (35), forcing each row of Eq. (35) to contain all three colors $\{\mathbf{c}_0, \mathbf{c}_1, \mathbf{c}_2\}$.

2. *Columns of the* 3×3 *Grid* ($\mathcal{B}_{ak} = \{M_{ak}, P_k, B_{ak}\}$): For each fixed $k \in \mathbb{Z}_3$, vertex P_k appears in $\mathcal{B}_{0k}, \mathcal{B}_{1k}, \mathcal{B}_{2k}$ alongside the kth *column* $N(P_k) = \{M_{0k}, M_{1k}, M_{2k}\}$ of Eq. (35). Thus column k of the grid (35) must omit the color $c(P_k)$!

3. *Cyclic Latin-Square Transversals* ($\mathcal{C}_{ak} = \{M_{a,a+k}, Q_k, C_{ak}\}$): For each fixed $k \in \mathbb{Z}_3$, vertex Q_k appears alongside the diagonal transversal
$$N(Q_k) = \{M_{0,k},\ M_{1,k+1},\ M_{2,k+2}\},$$
which is the kth symbol class of the cyclic Latin square $L(a,\ell) = \ell - a \pmod 3$! Thus each cyclic diagonal $N(Q_k)$ must omit the color $c(Q_k)$.

4. *The Apex Constraint* ($\mathcal{A}_k = \{z, P_k, A_k\}$): Fix z in Column $\mathbf{c}_0$ ($c(z) = 0$). Then the three rows $\mathcal{A}_0, \mathcal{A}_1, \mathcal{A}_2$ bar P_0, P_1, P_2 from Column $\mathbf{c}_0$, so $c(P_k) \in \{1,2\}$.

Now watch how the Latin-square structure produces the contradiction $\chi(H_{37}) = 4$:

- If P_0, P_1, P_2 all had the same color (say 1), then by (2) no entry M_{ak} in any column of Eq. (35) could have color 1, contradicting (1) that every row $\mathcal{M}_a$ contains all three colors!

- Hence $\{c(P_0), c(P_1), c(P_2)\} = \{1,2\}$. Up to a cyclic shift of the column index $k \in \mathbb{Z}_3$ and exchanging colors 1, 2, set $(c(P_0), c(P_1), c(P_2)) = (2,1,1)$.

- Since Column 0 of Eq. (35) omits color 2, and Columns 1 and 2 omit color 1, the unique color-1 entry in each row $\mathcal{M}_a$ *must sit in Column* 0 ($c(M_{a0}) = 1$ for all $a \in \mathbb{Z}_3$), while the remaining two entries (M_{a1}, M_{a2}) in each row must be a permutation $(x_a, 2-x_a)$ of $\{0,2\}$:
$$c(M) = \begin{pmatrix} 1 & x_0 & 2-x_0 \\ 1 & x_1 & 2-x_1 \\ 1 & x_2 & 2-x_2 \end{pmatrix}, \qquad x_a \in \{0,2\}. \tag{36}$$

- Now read off the three cyclic Latin-square transversals $N(Q_k)$ from Eq. (36):
$$c(N(Q_0)) = \{1,\ x_1,\ 2-x_2\},$$
$$c(N(Q_1)) = \{x_0,\ 2-x_1,\ 1\},$$
$$c(N(Q_2)) = \{2-x_0,\ 1,\ x_2\}.$$
By (3), none of these three transversals can be rainbow ($\{0,1,2\}$), so the two non-1 entries in each transversal must be equal:
$$x_1 = 2-x_2, \qquad x_0 = 2-x_1, \qquad x_2 = 2-x_0,$$
which requires $x_0 = 2-x_1 = 2-(2-x_2) = x_2 = 2-x_0 \implies x_0 = 1 \notin \{0,2\}$, a contradiction!

Thus no 24×3 Context–Color Matrix exists, proving $\chi(H_{37}) \geq 4$. A strong four-coloring is explicit: set $c(P_k) = c(Q_k) = 3$, $c(z) = 0$, $c(A_k) = 1$, $c(M_{ak}) = k$, $c(B_{ak}) = k + 1 \pmod 3$, and $c(C_{ak}) = a + k + 1 \pmod 3$. Every listed context has three distinct colors. This supplies the upper bound and proves $\chi(H_{37}) = 4$; its four-column matrix has one blank per row.

C. The Four-Dimensional Completion H_{65} (29×4 Pigeonhole Overflow)

Finally, consider the canonical linear 4-uniform hypergraph H_{65} in $\mathbb{R}^4$ ($|V| = 65, n = 29$ contexts, the real-completion section of Ref. [5]), built from the 17-ray core $V_{\text{core}} = \{1, \ldots, 17\}$ with 5 core contexts

$$C_1 = \{1, 2, 3, 4\},\ C_2 = \{5, 6, 7, 8\},\ C_3 = \{9, 10, 12, 13\},$$
$$C_4 = \{9, 11, 14, 15\},\ C_5 = \{10, 11, 16, 17\}, \tag{37}$$

and 24 completed bridge contexts $E_m = \{u_m, v_m, 2m + 16, 2m + 17\}$ ($1 \leq m \leq 24$).

In the 29×4 Context–Color Matrix picture of Theorem 1, the non-4-colorability $\chi(H_{65}) = 5$ (despite H_{65} having $|\mathcal{S}(H_{65})| = 393{,}216$ two-valued states!) has a one-line Pigeonhole proof:

1. Every state column S_j in a 29×4 Context–Color Matrix must be an independent set of $\Gamma(H_{65})$.

2. On the 17-vertex core V_{core}, the independence number is $\alpha(G_{\text{core}}) = 4$: every state $s \in \mathcal{S}(H_{65})$ selects *at most* 4 *core vertices* (in fact $|S_j \cap V_{\text{core}}| = 4$: one from C_1, one from C_2, one hub from $\{9, 10, 11\}$, and one bridge ray from $B_1 \cup B_2 \cup B_3$).

3. Therefore, four disjoint state columns S_0, S_1, S_2, S_3 can cover at most

$$\sum_{j=0}^{3} \left|S_j \cap V_{\text{core}}\right| \leq 4 \times 4 = 16 < 17 = \left|V_{\text{core}}\right| \tag{38}$$

core vertices. At least one of the 17 core vertices is left without a column, proving $\chi(H_{65}) \geq 5$. The companion supplies a five-coloring, giving $\chi(H_{65}) = 5$. Such a five-column representation has one blank per row; its columns are independent sets, as in Remark 2.

VI. PHYSICAL INTERPRETATION AND SUMMARY

The context–color matrix organizes spectral labels on existing quantum projectors. A valid column partition gives a common nondegenerate spectrum across intertwined measurements; a column obstruction excludes that labeling. The three- and four-dimensional examples retain their quantum coordinates in either case. Table I summarizes the physically realized configurations used here.

The physical distinction is particularly sharp in $H_{17,10}$: all spectral assignments identify labels 10 and 24, while the Born preparation (21) distinguishes their probabilities and requires nonextendable ordinary-state weight. The realized Yu–Oh, Hesse, and 65-ray configurations instead obstruct the exact common spectrum altogether. These are statements about specified quantum measurement families, not deductions of geometry from coloring.

The literal Graeco–Latin applications are Eqs. (15) and (33): the former combines a measurement-indexing Latin square with its spectral-label mate, whereas the latter supplies two orthogonal spectral-label colorings of an existing rook measurement net. Neither construction obtains its FOR from classical orthogonal squares alone. The Yu–Oh signatures and Hesse neighborhoods have weaker Latin-pattern interpretations and are not Graeco–Latin square schemes. The general matrix criterion applies in every Hilbert-space dimension once the quantum ray representation has been supplied. Common-apex extensions preserve that physical foundation through their explicit coordinate embedding.

ACKNOWLEDGMENTS

This research was funded in part by the Austrian Science Fund (FWF), Grant DOI 10.55776/PIN5424624.

TABLE I. Coordinatized quantum configurations discussed in the text. N_c counts exact-palette colorings modulo global palette permutations; S_{ext}/S lists extendable and ordinary state counts. The field and dimension specify the ray realization. Complete companion coordinates are in Ref. [5, 6]; the two-context, pentagon, Gaussian-integer hub grids ($R_{21,12}$ and $R_{25,15}$), four-dimensional quaternion rook net ($H^{(4)}_{16,8}$), and cyclic-net coordinates are printed in this paper.

Configuration	Ray space	$\lvert V\rvert$	$\lvert E\rvert$	N_c	$\lvert\mathcal{S}_{\mathrm{ext}}\rvert/\lvert\mathcal{S}\rvert$
Two contexts	$\mathbb{R}^3$	5	2	2	5/5
P_5	$\mathbb{R}^3$	10	5	5	10/11
$H_{17,10}$	$\mathbb{R}^3$	17	10	4	10/18
$H_{21,13}$	$\mathbb{R}^3$	21	13	2	6/21
$B_{13,7}$	$\mathbb{R}^3$	13	7	6	12/14
$I_{16,9}$	$\mathbb{R}^3$	16	9	6	12/22
$D_{16,9}$	$\mathbb{R}^3$	16	9	7	14/21
$T_{19,11}$	$\mathbb{R}^3$	19	11	7	14/25
$R_{21,12}$	$\mathbb{C}^3$	21	12	32	48/54
$R_{25,15}$	$\mathbb{C}^3$	25	15	24	48/81
$W^{(4)}_{11,5}$	$\mathbb{R}^4$	11	5	5	11/12
$B^{(4)}_{14,7}$	$\mathbb{R}^4$	14	7	6	13/15
$D^{(4)}_{17,9}$	$\mathbb{R}^4$	17	9	7	15/22
$W^{(4)}_{16,6}$	$\mathbb{R}^4$	16	6	80	56/72
$L^{(4)}_{18,7}$	$\mathbb{R}^4$	18	7	144	68/68
$L^{(4)}_{22,9}$	$\mathbb{R}^4$	22	9	216	92/158
$H^{(4)}_{16,8}$	$\mathbb{R}^4$	16	8	24	24/24
Completed Yu–Oh	$\mathbb{R}^3$	25	16	0	0/24
$H^{(4)}_{\mathrm{YO}}$	$\mathbb{R}^4$	26	16	0	0/25
H_{37}	$\mathbb{C}^3$	37	24	0	0/153
H_{65}	$\mathbb{R}^4$	65	29	0	0/393,216
Cyclic Latin net	$\mathbb{C}^7$	49	21	635	133/133

Appendix A: Verification of the physical examples

The accompanying certificate contains only the coordinatized examples retained in this paper. Exact integer and Gaussian-integer inner products verify the full faithful orthogonality relation for Eqs. (9), (17), (25), and (31), including all nonedges and distinctness of rays. For the cyclic FOR, the checker uses exponent multiplicities modulo seven to verify all 1176 inner products exactly, with 441 zeros and 735 nonzeros, and checks that the only complete seven-ray bases are the 21 listed measurements. It also enumerates the 133 transversals and 635 normalized seven-colorings as exact covers by seven disjoint transversals. The classical pairs in Eqs. (15) and (33) are checked for Latinness, unique ordered pairs, valid spectral colorings, and nonorthogonality within their color-class supports. The phase-exponent coordinates are separately provided in `cyclic7-faithful-coordinates.json`.

For the finite context inventories, normalized coloring fixes one context's color order, giving one representative of each global palette-permutation orbit. Binary states are enumerated with exact-one propagation on the listed measurements. Color-class supports give $\mathcal{S}_{\text{ext}}$. Each displayed vertex matrix is checked against its context rows and for consistent column positions of shared projectors. Ray coordinates and geometric verification for the companion configurations are those of Ref. [5, 6]; the standalone checker does not claim to rederive those larger coordinate inventories.

The five TikZ files show only coordinatized quantum scenarios. Their trace memberships, unique vertex labels, and incidence rings are checked against the physical context inventories. Trace colors distinguish measurements, while the matrix columns distinguish spectral labels. The standalone `verify-latin-context-matrix.py` needs only Python's standard library. All active source files, coordinate certificates, and verification output are included in the accompanying package.

[1] J. von Neumann, Über Funktionen von Funktionaloperatoren, Annals of Mathematics **32**, 191 (1931).

[2] P. R. Halmos, *Finite-Dimensional Vector Spaces*, Undergraduate Texts in Mathematics (Springer, New York, 1974).

[3] K. Svozil, Classical predictions for intertwined quantum observables are contingent and thus inconclusive, Quantum Reports **2**, 278 (2020), arXiv:1808.00813.

[4] M. H. Shekarriz and K. Svozil, Noncontextual coloring of orthogonality hypergraphs, Journal of Mathematical Physics **63**, 032104 (2022), arXiv:2105.08520.

[5] K. Svozil, Chromatic constraints on quantum observables: Gaps, contextuality, and separability (2026), companion manuscript, version of 26 October 2026, with Supplemental Material.

[6] K. Svozil, Supplemental material for "Chromatic Constraints on Quantum Observables: Gaps, Contextuality, and Separability" (2026), contains all explicit unnormalized vector and context inventories, complete two-valued state and coloring enumerations, odd-cycle and separation profile certificates, and supplementary proofs and verification reductions.

[7] B. Musto and J. Vicary, Quantum Latin squares and unitary error bases, Quantum Information and Computation **16**, 1318 (2016), arXiv:1504.02715.

[8] S. Ball and R. Simoens, Large sets of mutually orthogonal quantum Latin squares (2026), version 2, 27 July 2026, arXiv:2607.12933 [math.CO].

[9] H. B. Mann, The construction of orthogonal Latin squares, The Annals of Mathematical Statistics **13**, 418 (1942).

[10] I. M. Wanless, Transversals in Latin squares: a survey, in *Surveys in Combinatorics 2011*, London Mathematical Society Lecture Note Series, Vol. 392 (Cambridge University Press, Cambridge, 2011) pp. 403–437.

[11] H. Niederreiter and K. H. Robinson, Complete mappings of finite fields, Journal of the Australian Mathematical Society (Series A) **33**, 197 (1982).

[12] L. Lovász and A. Schrijver, Remarks on a theorem of Rédei, Studia Scientiarum Mathematicarum Hungarica **16**, 449 (1981).

[13] S. Ball, The number of directions determined by a function over a finite field, Journal of Combinatorial Theory, Series A **104**, 341 (2003).

[14] R. J. Evans, J. Greene, and H. Niederreiter, Linearized polynomials and permutation polynomials of finite fields, Michigan Mathematical Journal **39**, 405 (1992).