Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments

Karl Svozil [a)]
Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria

(Dated: November 5, 2026)

Suppose intertwined complete experiments each measure a nondegenerate maximal observable with one common spectrum, and the eigenvalue carried by a shared outcome is an element of physical reality, the same in every experiment. Such noncontextual eigenvalue assignments exist exactly when the context hypergraph is colorable with as many colors as the dimension. Their failure is weaker but more widespread than the Kochen–Specker contradiction, in which no two-valued state exists. Noncontextual eigenvalues, a condition on the spectra of maximal observables, can fail even when the two-valued states separate all outcomes, so that the logic has a classical realization; completed Yu–Oh is an example. Where noncontextual eigenvalue assignments exist, they can force distinct rays to share an eigenvalue; for one quantum preparation, a third of any classical model must use truth assignments that no eigenvalue assignment produces. We also correct an earlier inference from coloring to geometry and separate eigenvalue assignments from realizability and Born probabilities.

I. INTRODUCTION

An experimental context with d rank-one outcomes admits a nondegenerate observable with d distinct eigenvalues. These eigenvalues distinguish all d outcomes and provide the maximal resolution of a complete projective experiment within that context. Such an observable is maximal in the sense that every self-adjoint observable commuting with it is a function of it, so the statistics of the latter can be obtained by processing its measurement outcomes. Noninjective functions merge outcomes and reduce this resolution. This is the finite-dimensional setting of von Neumann's functional representation of commuting observables[1] (Satz 8, p. 221); see also Halmos[2] (§84, pp. 171–172).

When complete experiments are intertwined, they share outcomes represented by the same rank-one projectors. We ask whether the eigenvalues of the maximal observables measured in such intertwined experiments can be assigned to their outcomes noncontextually, in the spirit of the elements of physical reality of Einstein, Podolsky, and Rosen[3]: whether every outcome carries a definite eigenvalue that belongs to the outcome itself, rather than to the experiment in which it is recorded. Three assumptions make this precise.

(i) *Maximality.* Each complete experiment measures a maximal observable, so its d outcomes are never merged and carry d distinct eigenvalues. A degenerate observable would discard part of what the experiment resolves.

(ii) *Uniformity.* All maximal observables share one spectrum of d eigenvalues $\Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$. The eigenvalues are nothing but labels of the d outcomes of a complete experiment. Since self-adjoint operators with a common nondegenerate spectrum are unitarily equivalent, uniformity says that every complete experiment measures one and the same quantity in a different orientation, $A_e = U_e A U_e^\dagger$.

[a)] Electronic mail: karl.svozil@tuwien.ac.at

(iii) *Noncontextuality of eigenvalues.* An outcome shared by several complete experiments carries the same eigenvalue in each of them, just as Kochen and Specker require a shared outcome to carry the same truth value[4].

No constraint is placed on how the d eigenvalues are distributed over the outcomes of different contexts; any permutation within a context is allowed. A noncontextual eigenvalue assignment in this sense labels the outcomes, the most elementary record a complete experiment produces, and assumption (iii) treats these labels as elements of physical reality. For spin one, an example is the observable $aS_x^2+bS_y^2+cS_z^2$ with distinct real a, b, c considered by Kochen and Specker[4]: every orthogonal triad of directions yields one such observable, with the same eigenvalues $b+c$, $a+c$, and $a+b$. A noncontextual eigenvalue assignment then gives each direction the same coefficient, a, b, or c, in every triad that contains it.

These assumptions concern eigenvalues rather than which outcome occurs, yet they have immediate consequences for truth assignments. A two-valued state selects one outcome in every context and retains only the distinction between that outcome and the others. A noncontextual eigenvalue assignment yields d such states at once, one for each eigenvalue λ_a: "the outcome with eigenvalue λ_a occurs in every experiment." These d truth assignments have pairwise disjoint supports covering every outcome. A noncontextual eigenvalue assignment is thus a hidden-variable model whose hidden variable is the eigenvalue itself; its d equally weighted values reproduce the maximally mixed state. Because this requirement is more demanding than Kochen–Specker noncontextuality, which asks for a single truth assignment, its failure—*chromatic contextuality*—is a weaker form of contextuality that already appears in configurations retaining many classical truth assignments. Even when a noncontextual eigenvalue assignment exists, its consistency across intertwined contexts can force relations between distinct outcomes, such as equal eigenvalues for nonorthogonal rays, that are absent from the full family of individual truth assignments. We study both the existence of noncontextual eigenvalue assignments and the event distinctions and logical relations that survive when such assignments are available[5–7].

To express these questions, let a d-uniform context hypergraph $H = (V, E)$ encode the complete experiments. An ordinary two-valued state $s : V \to \{0, 1\}$ selects exactly one outcome in every context, so $\sum_{v\in e} s(v) = 1$ for every $e \in E$. The Kochen–Specker (KS) theorem[4] establishes the nonexistence of such a global truth assignment for suitable finite families of intertwined projective contexts in dimension $d \geq 3$.

A strong d-coloring $c : V \to \{0, \dots, d-1\}$ assigns all d distinct labels bijectively within every context and independently of context at shared vertices. It is the formal counterpart of a noncontextual eigenvalue assignment: $c(i) = a$ means that outcome i carries the eigenvalue λ_a in every context containing it, so assumptions (i)–(iii) hold if and only if $\chi(H) \leq d$. For an injective ray realization $i \mapsto [v_i] \in \mathbb{P}(\mathbb{F}^d)$, with $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, rank-one projectors P_i, and a common nondegenerate real spectrum $\Sigma = \{\lambda_0, \dots, \lambda_{d-1}\}$, it defines the maximal context observables

$$A_e = \sum_{i\in e} \lambda_{c(i)} P_i. \tag{1}$$

Because the indicator functions $s_a(v) = \mathbf{1}_{\{c(v)=a\}}$ $(a \in \{0, \dots, d-1\})$ of any strong d-coloring form d two-valued states whose supports partition V, one has the strict one-way implication

$$\mathcal{S}(H) = \varnothing \quad \Longrightarrow \quad \chi(H) > d, \tag{2}$$

whose converse fails even when $\mathcal{S}(H)$ is unital and separates every pair of vertices. The earlier introduction in Ref.[6] inferred absence of global two-valued assignments from $\chi > d$. Equation (2) is the valid implication; completed Yu–Oh refutes the converse. Both the implication and the failure of its converse, for the original 13-ray Yu–Oh set, were noted by Cabello[8]. In terms of eigenvalues, if no noncontextual truth assignment exists, no noncontextual eigenvalue assignment exists either; but noncontextual eigenvalues can be impossible while truth assignments, even a unital separating family of them, survive. The

distinct Appendix B inference about geometry is corrected below. We say that H has an *exact palette*, or is *chromatically complete*, when $\chi(H) = d$, with no additional density or geometric assumption. This asymmetry separates chromatic contextuality into two distinct regimes that are unified in this work:

1. *Regime I, common-spectrum chromatic obstructions ($\chi(H) > d$, possibly with $\mathcal{S}(H) \neq \varnothing$):* Here no strong d-coloring exists ($\mathcal{C}_d(H) = \varnothing$), so the outcomes admit no noncontextual assignment of d eigenvalues: every family of context observables with one common spectrum of d eigenvalues must either merge outcomes, and so fail to be maximal, or let a shared outcome change its eigenvalue from context to context; maximal observables with noncontextual eigenvalues need more than d eigenvalues. This happens even in examples in which classical truth assignments survive. A critical foundational clarification—correcting an interpretation in Appendix B of the earlier work with Shekarriz[6]—is that the chromatic obstruction $\chi(H) > d$ is an operator spectral-value obstruction and *does not* by itself obstruct geometric ray realizability in $\mathbb{F}^d$. Indeed, for finite 3-uniform linear hypergraphs of Berge girth five with unital separating state spaces, strong 3-colorability and injective $\mathbb{C}^3$ realizability are logically independent (Proposition 10). Ruling out geometric ray assignments for Greechie's Petersen-derived G_{32}[9] requires a direct Hermitian projection-identity argument (Theorem 11) rather than $\chi(G_{32}) = 4$.

2. *Regime II, exact-palette chromatic separability and logical gadgets ($\chi(H) = d$ with $\mathcal{S}_{\text{ext}}(H) \subsetneq \mathcal{S}(H)$):* In this regime, $\mathcal{C}_d(H) \neq \varnothing$, so noncontextual eigenvalue assignments exist, but some two-valued states $s \in \mathcal{S}(H)$ cannot be completed to a strong d-coloring: they are truth assignments that no noncontextual eigenvalue assignment produces. We characterize the extendable states $\mathcal{S}_{\text{ext}}(H)$ by the residual chromatic condition $\chi(\Gamma(H)[V \setminus \operatorname{supp} s]) \leq d-1$ (Theorem 2), which in dimension $d = 3$ yields an odd-cycle nonextendability certificate. In the specified subhypergraphs—such as the faithful real 17-ray and 21-ray restrictions $H_{17,10}$ and $H_{21,13}$ of completed Yu–Oh[7,10] (Theorem 15) and our faithful 3- and 4-dimensional logical gadgets (Secs. IV C and IV D)—ordinary states separate all vertices while every admissible d-coloring forces nonadjacent rays to share a color ($u \sim_c v$, chromatic inseparability / implicit identity[11]) or obeys purely chromatic true-implies-false (TIFS) or ternary exclusions violated by nonextendable states in $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$.

These two regimes are governed by a shared structural mechanism. Lemma 3 ($\chi(H) = d$ if and only if H admits $d-1$ pairwise disjoint states) is the global unanchored form of Theorem 2. Likewise, the eight normalized 3-colorings of the four base contexts of completed Yu–Oh H_{YO}^{+} partition into four antipodal pairs that each make one of the four diagonal neighborhoods rainbow (Theorem 13): retaining all four diagonal sectors forces $\chi(H_{\text{YO}}^{+}) = 4$ (Regime I), whereas retaining two or three sectors leaves four or two normalized 3-colorings and forces two or 21 chromatically inseparable pairs in $H_{17,10}$ and $H_{21,13}$ (Regime II), which our exhaustive 65,535-subset inventory classification (Proposition 16) proves minimal among the faithful restrictions of H_{YO}^{+} with inherited coordinates.

For truth values, elements of reality fail as in the Kochen–Specker theorem: the logic has no two-valued states, or too few to separate its outcomes, and then it has no classical realization. For eigenvalues the situation is different. If the two-valued states of H are unital and separating, H can be realized classically as a partition logic[4,12,13]: outcome i becomes the set of states with $s(i) = 1$, and each context becomes a partition of this set. Noncontextual eigenvalues can still be impossible, because they require a strong d-coloring, in the classical realization as much as in the quantum one. Greechie's G_{32} is an example: it has a classical realization, no orthogonal ray assignment in $\mathbb{C}^3$ (Theorem 11), and $\chi(G_{32}) = 4$. Thus $\chi(H) > d$ is a condition on the spectra of maximal observables, not on truth values. It separates quantum from classical physics only when an experiment fixes the label of each outcome, as the d-party supersinglet of the next paragraph does; for a single system, the separation comes from the Born probabilities (Sec. VI).

Noncontextual eigenvalues and strong colorings were first related by Cabello[8]. The first version of that preprint assigned noncontextual values $-1, 0, 1$ to the spin-one components S_u along the rays u and concluded that $\chi > d$ characterizes state-independent contextuality. For an orthogonal triad u, v, w, however, the eigenvectors of S_u, S_v, S_w for the eigenvalue 1 (or -1) are not mutually orthogonal, so quantum mechanics fixes only which of them takes the value 0; the necessity of $\chi > d$ and the minimality of the 13-ray Yu–Oh set were later published in Ref.[14]. The context observables A_e of Eq. (1) are diagonal in their contexts, so assumptions (i)–(iii) are equivalent to $\chi(H) \leq d$ without such a gap. The second version of the preprint distributes the d particles of the totally antisymmetric supersinglet among d parties. If all parties measure bases containing a ray v, exactly one of them obtains v; if $d-1$ parties measure the same basis, the ray of that basis which none of them obtains is obtained with certainty by the last party in any basis containing it. A local hidden-variable model of these perfect correlations exists exactly when the hypergraph of the measured bases has a strong d-coloring, the color of a ray being the party that obtains it; for the Yu–Oh set this hypergraph is H_{YO}^{+}. Here the label of an outcome is the party that records it, and its noncontextuality, assumption (iii), follows from locality and the EPR criterion of reality[3] instead of being postulated. Quantum mechanics produces these correlations whenever H has an orthogonal ray assignment in $\mathbb{C}^d$, while local classical models need a strong d-coloring, so here $\chi(H) > d$ does separate quantum from classical physics. G_{32} has no ray assignment and gives no such separation. For a single system in the maximally mixed state the threshold is fractional: I_d/d has a noncontextual truth-assignment model exactly when $\chi_f(\Gamma(H)) = d$[15] (Proposition 6). Contrary to the claim of Ref.[8], $\chi > d$ is therefore not sufficient for the contextuality of I_d/d: the faithful complex Hesse hypergraph H_{37} has $\chi = 4$ and $\chi_f = 3$ (Theorem 19). The gap between χ and the least dimension of an orthogonal ray assignment has also been studied in connection with the quantum chromatic number[16] and, recently, as a memory advantage of quantum over classical automata[17].

We further connect these combinatorial and geometric layers to orthomodular event representations and quantum probability polytopes $\mathcal{P}_{\mathrm{ext}}(H) \subseteq \mathcal{P}(H) = \operatorname{conv} \mathcal{S}(H)$, where $\mathcal{P}_{\mathrm{ext}}(H)$ collects the probability vectors of hidden-variable models that choose a noncontextual eigenvalue assignment and one of its d eigenvalues. On $H_{17,10}$, an explicit quantum ray preparation lies in $\mathcal{P}(H_{17,10}) \setminus \mathcal{P}_{\mathrm{ext}}(H_{17,10})$ and requires nonextendable state weight exactly $1/9$; the preparation along the coordinate ray z_3 requires exactly $1/3$, the largest weight that a single chromatic identity of $H_{17,10}$ can certify (Theorem 24). In complex dimension three, the 37-ray Hesse restriction H_{37}[18] has no injective $\mathbb{R}^3$ realization, pastes into a 76-event orthomodular lattice with exact minimum state-family sizes 6, 7, and 18 for vertex separation, event separation, and order determination (Theorem 19), and demonstrates that exact event-order determination does not guarantee classical mixtures for all Born preparations (Corollary 25). In real dimension four, our two-plane completion lemma (Lemma 21) produces the canonical linear 65-ray hypergraph H_{65} completing the 17-ray core of Haynes *et al.*[19] with $\chi = 5$, $\chi_f = 17/4$, and 393,216 states that fail atom-order determination and collapse a ray with a plane (Theorem 22). Finally, we formulate and prove the existence form of the KS ancestry hypothesis (Proposition 26), give an abstract transverse chromatic-to-stateless construction (Proposition 27), establish exact inclusions into specified KS parents (Bub's 49-ray set[20], the 165-ray MUB inventory, Peres's 24-ray geometry, and 68-ray extensions containing the 18-ray parity proof $C_{18,9}$ of Cabello *et al.*[21] and a copy $C_{20,11}$ of Kernaghan's 20-ray parity proof[22]), and contrast abstract context states $\mathcal{S}_{\mathrm{ctx}}$ with geometric states $\mathcal{S}_{\mathrm{geom}}$ on the 36-ray completion and 45-ray vertex deletion of $C_{18,9}$. Complete vector, state, and coloring enumerations and finite verification certificates are provided in the supplementary material, which additionally cites Refs.[23–25].

II. UNIFIED FRAMEWORK: STATES, COLORINGS, SEPARABILITY, AND COMPLETION

A. Context hypergraphs, orthogonal realizations, and spectral regimes

All hypergraphs and graphs considered here are finite. A d-uniform *context hypergraph* $H = (V, E)$ ($d \geq 2$) consists of a nonempty collection E of d-element subsets of V, called *contexts*, with $V = \bigcup_{e \in E} e$. Its *two-section* (or *context*) *graph* $\Gamma(H)$ has vertex set V and edge set $E(\Gamma(H)) = \bigcup_{e \in E} \binom{e}{2}$. A hypergraph H is *linear* if $|e \cap e'| \leq 1$ for all distinct $e, e' \in E$. A *Berge cycle* of length $\ell \geq 2$ is an alternating sequence of distinct vertices $x_1, \ldots, x_\ell$ and distinct contexts $e_1, \ldots, e_\ell$ with $\{x_j, x_{j+1}\} \subseteq e_j$ cyclically; the least such ℓ is the *Berge girth* $g(H)$ (or ∞ if acyclic). In Greechie's pasting of Boolean blocks[9,26], linearity ($g(H) \geq 3$) with $g(H) \geq 4$ yields an *orthomodular poset*, and $g(H) \geq 5$ yields an *orthomodular lattice* (OML).

With the Hermitian inner product $\langle x, y \rangle = \sum_j \overline{x_j} y_j$ on $\mathbb{F}^d$ ($\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$), we distinguish three levels of geometric realization:

1. An *orthogonal ray assignment* $\rho : V \to \mathbb{P}(\mathbb{F}^d)$, $i \mapsto [v_i]$ ($v_i \neq 0$), requires $\langle v_i, v_j \rangle = 0$ whenever $i \sim j$ in $\Gamma(H)$; nonadjacent vertices may share a ray.
2. An *injective orthogonal representation* additionally requires $[v_i] \neq [v_j]$ for all $i \neq j$.
3. An injective representation is *edge-faithful* (abbreviated FOR) if $\langle v_i, v_j \rangle = 0 \iff i \sim j$ in $\Gamma(H)$ for all $i \neq j$[27].

A *strong k-coloring* of H is a map $c : V \to \{0, \ldots, k-1\}$ injective on every context $e \in E$, i.e., a proper k-coloring of $\Gamma(H)$; the least such k is $\chi(H) = \chi(\Gamma(H))$. We call H *chromatically complete* (equivalently, H has an exact palette) when $\chi(H) = d$ and write $\mathcal{C}_d(H)$ for the set of strong d-colorings of H. For a realized d-uniform hypergraph with rank-one projectors $P_i = v_i v_i^\dagger / \langle v_i, v_i \rangle$, three spectral labelings arise: (a) *context-dependent common spectrum*, where bijections $\ell_e : e \to \Sigma = \{\lambda_0, \ldots, \lambda_{d-1}\}$ define $A_e = \sum_{i \in e} \ell_e(i) P_i$ with shared projectors carrying context-dependent eigenvalues; (b) *context-independent vertex labeling* via a strong k-coloring c and distinct real scalars $\lambda_0, \ldots, \lambda_{k-1}$ in Eq. (1), where spectra vary across contexts whenever $|c(V)| > d$ (unavoidable when $\chi(H) > d$); and (c) *chromatic completeness* ($k = d$), which provides both context-independent eigenvalues and a single d-element spectrum Σ for all A_e. Only labeling (c) meets all three assumptions of the Introduction: labeling (a) gives up noncontextual eigenvalues, and labeling (b) keeps them at the price of more than d eigenvalues, so that different contexts have different spectra.

B. Two-valued states, color extendability, and the separation hierarchy

A *two-valued context state* is a function $s : V \to \{0, 1\}$ satisfying

$$\sum_{v \in e} s(v) = 1 \qquad (e \in E). \tag{3}$$

We write $\mathcal{S}(H) = \mathcal{S}_{\mathrm{ctx}}(H)$ for the set of context states. For a ray realization ρ, the *geometric state space*

$$\mathcal{S}_{\mathrm{geom}}(H, \rho) = \{s \in \mathcal{S}(H) : s(i) + s(j) \leq 1 \text{ whenever } i \neq j,\ v_i \perp v_j\} \tag{4}$$

additionally enforces orthogonality exclusions on uncovered pairs[28]; $\mathcal{S}_{\mathrm{geom}}(H, \rho) = \mathcal{S}(H)$ when ρ is edge-faithful. A state family $\Omega \subseteq \mathcal{S}(H)$ is *unital* if every $v \in V$ has $s(v) = 1$ for some $s \in \Omega$, and *vertex separating* if distinct $u, v \in V$ have $s(u) \neq s(v)$ for some $s \in \Omega$, inducing the partition set representation $A_i = \{s \in \Omega : s(i) = 1\}$. On an event poset $(L, \leq)$, Ω is *event separating* if distinct $x, y \in L$ have distinct profiles on Ω, and *order determining* if $x \leq y \iff (\forall s \in \Omega,\ s(x) \leq s(y))$.

Definition 1 (Chromatic separability and extendable states)**.** For chromatically complete H ($\mathcal{C}_d(H) \neq \varnothing$), vertices $u, v \in V$ are *chromatically inseparable*, written $u \sim_c v$, if

$$c(u) = c(v) \qquad \text{for every } c \in \mathcal{C}_d(H), \tag{5}$$

an *implicit identity* in graph coloring[11] (Def. 3.8). We call H *chromatically separating* if $u \sim_c v \implies u = v$. A state $s \in \mathcal{S}(H)$ is *extendable*, written $s \in \mathcal{S}_{\text{ext}}(H)$, if $\operatorname{supp} s = \{v \in V : s(v) = 1\}$ is a color class of some $c \in \mathcal{C}_d(H)$[5–7].

Physically, $u \sim_c v$ means that every noncontextual eigenvalue assignment gives u and v the same eigenvalue, and an extendable state is exactly a truth assignment of the form "the outcome with eigenvalue λ_a occurs" for some noncontextual eigenvalue assignment c and eigenvalue λ_a.

Theorem 2 (Completion criterion)**.** *For $s \in \mathcal{S}(H)$ with $S = \operatorname{supp} s$, the following are equivalent:*

1. *$s \in \mathcal{S}_{\text{ext}}(H)$;*
2. *$\chi(\Gamma(H)[V \setminus S]) \leq d - 1$ (so S is a critical independent set[11] (Def. 3.3));*
3. *s belongs to d states in $\mathcal{S}(H)$ with pairwise disjoint supports partitioning V;*
4. *s belongs to $d - 1$ states in $\mathcal{S}(H)$ with pairwise disjoint supports.*

For $d = 3$, $s \notin \mathcal{S}_{\text{ext}}(H)$ if and only if $\Gamma(H)[V \setminus S]$ contains an odd cycle.

Proof. A strong d-coloring with color class S leaves a $(d-1)$-coloring on $V \setminus S$. Conversely, any $(d-1)$-coloring of $\Gamma(H)[V \setminus S]$ combined with a dth color on S is a strong d-coloring because each context meets S once and forms a $(d-1)$-clique in $V \setminus S$; its color indicators yield (3). Conversely, d states with disjoint supports $S_1, \ldots, S_d$ partitioning V define the strong d-coloring $c(v) = k$ for $v \in S_k$, since each context meets each S_k exactly once; hence (3) $\implies$ (1). Trivially (3) $\implies$ (4), and given $d-1$ disjoint state supports $S_1, \ldots, S_{d-1}$, the complement $s_d(v) = 1 - \sum_{k=1}^{d-1} s_k(v)$ is binary and sums to $d - (d-1) = 1$ on every context, giving (3). For $d = 3$, $\chi(\Gamma(H)[V \setminus S]) \leq 2$ holds if and only if $\Gamma(H)[V \setminus S]$ is bipartite (odd-cycle free). □

Applying the equivalence (3) $\iff$ (4) of Theorem 2 globally without anchoring a state s yields the characterization of chromatic completeness:

Lemma 3 (Disjoint states and strong coloring)**.** *A finite d-uniform hypergraph H satisfies $\chi(H) = d$ if and only if $\mathcal{S}(H)$ contains $d - 1$ states with pairwise disjoint supports.*

Corollary 4 (The separation hierarchy)**.** *For chromatically complete H and $u, v \in V$,*

$$u \sim_c v \quad \iff \quad s(u) = s(v) \text{ for all } s \in \mathcal{S}_{\text{ext}}(H). \tag{6}$$

Thus inseparability by $\mathcal{S}(H)$ implies chromatic inseparability ($\sim_c$), equivalently chromatic separation implies separation by $\mathcal{S}(H)$, whereas the converse fails even for faithful real ray configurations. Moreover, $\mathcal{S}_{\text{ext}}(H)$ is always unital whenever $\mathcal{C}_d(H) \neq \varnothing$.

Proof. The indicator states $s_a(v) = \mathbf{1}_{\{c(v)=a\}}$ over all $c \in \mathcal{C}_d(H)$ and $a \in \{0, \ldots, d-1\}$ are precisely $\mathcal{S}_{\text{ext}}(H) \subseteq \mathcal{S}(H)$. For each fixed coloring, the d supports partition V. The inclusion need not be strict: for a single d-element context every state is extendable. Strict inclusion is an additional property established for the specified examples. □

Proposition 5 (Canonical chromatic quotient and closure test)**.** *Let H be chromatically complete.*

1. *The relation $\sim_c$ is an equivalence relation with no two equivalent vertices in a common context. The quotient $\overline{H} = (V/\!\sim_c, \overline{E})$ is d-uniform and chromatically separating, and pullback along $V \to V/\!\sim_c$ induces a bijection $\mathcal{C}_d(\overline{H}) \to \mathcal{C}_d(H)$.*

2. *For $u \neq v$, adjoining one context $\{u, v, w_1, \ldots, w_{d-2}\}$ with fresh vertices w_i yields chromatic number $d+1$ if and only if $u \sim_c v$, and d otherwise. Since recoloring one vertex with a new color separates any pair, the* separable chromatic number χ_{sep} *of a finite graph, the least t for which some set of proper colorings with at most t colors separates every pair of distinct vertices*[5]*, satisfies $\chi_{\mathrm{sep}} \in \{\chi, \chi+1\}$.*

Proof. (1) Adjacent vertices receive distinct colors, so each context projects injectively to $V/\!\sim_c$, and every $c \in \mathcal{C}_d(H)$ factors uniquely through $V/\!\sim_c$. (2) A coloring $c \in \mathcal{C}_d(H)$ extends to the fresh context if and only if $c(u) \neq c(v)$; if $u \sim_c v$, recoloring u with a $(d+1)$th color separates (u, v) while preserving all old context inequalities. □

The quotient preserves colorings, not an injective realization. Closing an inseparable pair is likewise abstract: faithful coordinates make the pair nonorthogonal, so the added context cannot retain those rays.

Put $\mathcal{P}_{\mathrm{ext}} = \operatorname{conv} \mathcal{S}_{\mathrm{ext}}$ and $\mathcal{P} = \operatorname{conv} \mathcal{S}$. Thus $\mathcal{P}$ is the set of probability vectors of noncontextual truth-assignment models, and $\mathcal{P}_{\mathrm{ext}}$ is the subset produced by models that choose a noncontextual eigenvalue assignment c and an eigenvalue λ_a at random and report, in every context, the outcome with eigenvalue λ_a. A binary vector cannot be a convex combination of distinct binary vectors, since each positive-weight term must agree at every coordinate. Consequently,

$$\mathcal{P}_{\mathrm{ext}} = \mathcal{P} \quad \Longleftrightarrow \quad \mathcal{S}_{\mathrm{ext}} = \mathcal{S}. \tag{7}$$

This is stronger than equality of vertex-separation kernels: the pentagon has separating colors yet a nonextendable state. Pair identities alone do not describe all completion constraints.

C. Fractional coloring and fresh-vertex context completion

Write $\alpha(G)$ for the independence number and $\chi_f(G)$ for the fractional chromatic number of $G = \Gamma(H)$.

Proposition 6 (Fractional threshold and uniform state mixture)**.** *For any finite d-uniform context hypergraph H,*

$$\chi_f(\Gamma(H)) = d \quad \Longleftrightarrow \quad \frac{1}{d}\mathbf{1} \in \mathcal{P}(H) = \operatorname{conv} \mathcal{S}(H). \tag{8}$$

Proof. A state mixture $\sum_{s \in \mathcal{S}(H)} \mu_s s = \frac{1}{d}\mathbf{1}$ assigns weight $d\mu_s$ to the independent set $\operatorname{supp} s$, covering each vertex with weight 1 at total weight d, so $\chi_f(\Gamma(H)) \leq d$; since every context is a d-clique, $\chi_f(\Gamma(H)) \geq \omega(\Gamma(H)) \geq d$. Conversely, a fractional coloring $\{w_I\}$ of total weight d obeys $d \leq \sum_{v \in e} \sum_{I \ni v} w_I = \sum_I w_I |I \cap e| \leq \sum_I w_I = d$ for each $e \in E$, forcing $|I \cap e| = 1$ whenever $w_I > 0$, so $\{w_I/d\}$ is a state mixture equal to $\frac{1}{d}\mathbf{1}$. □

Equation (8) restates, for context states, the fractional-chromatic criterion of Ramanathan and Horodecki[15] for the contextuality of the maximally mixed state.

Proposition 7 (Fresh-vertex context completion)**.** *Let G be a finite core graph with designated d-clique base contexts ($d \geq 3$). Complete selected uncovered edges $\{u, v\}$ into d-cliques using $d-2$ degree-one fresh vertices per edge, covering all core vertices and edges. For the resulting hypergraph H and $G' = \Gamma(H)$,*

$$\begin{aligned} \chi(G') &= \max\{d, \chi(G)\}, \\ \chi_f(G') &= \max\{d, \chi_f(G)\}, \\ |\mathcal{S}(H)| &= \sum_S (d-2)^{m(S)}, \end{aligned} \tag{9}$$

where S ranges over independent core supports meeting each base context once and $m(S)$ counts completed edges missed by S.

TABLE I. Principal configurations. FOR means injective and edge-faithful; g is Berge girth. Counts refer to all context states and all color-class states.

Configuration	d	$\lvert V\rvert/\lvert E\rvert$	g	χ	χ_f	$\lvert\mathcal{S}\rvert$	$\lvert\mathcal{S}_{\text{ext}}\rvert$	Geometry and principal distinction
One triad	3	3/1	∞	3	3	3	3	Real FOR; colors separate.
$H_{6,3}$	3	6/3	3	3	3	4	3	No injective complex realization.
P_5	3	10/5	5	3	3	11	10	Real FOR; colors separate; ternary exclusion.
H_8	3	14/8	5	3	3	11	7	Repeated-ray assignments; no complex injection.
H_9	3	15/9	5	4	3	8	0	No complex ray assignment; context-critical.
G_{32}	3	15/10	5	4	3	6	0	No complex ray assignment.
$H_{17,10}$	3	17/10	5	3	3	18	10	Real FOR; two color-identical pairs.
$H_{21,13}$	3	21/13	5	3	3	21	6	Real FOR; 21 color-identical pairs.
Completed Yu–Oh	3	25/16	5	4	35/11	24	0	Real FOR; context-critical.
H_{37}	3	37/24	5	4	3	153	0	Complex FOR; no real injection; OML minima 6/7/18.
$H_{\text{YO}}^{(4)}$	4	26/16	2	5	46/11	25	0	Real FOR; common-apex suspension, nonlinear.
$C_{36,18}$	4	36/18	3	5	9/2	0	0	Faithful linear KS completion.
H_{65}	4	65/29	3	5	17/4	393,216	0	Real FOR; false atom order and compound collapse.

Proof. Any k-coloring ($k \geq d$) or measurable fractional palette of size $k = \max\{d, \chi_f(G)\}$ assigns disjoint colors/sets of measure 1 to u and v, leaving $k-2 \geq d-2$ colors (or measure $k-2 \geq d-2$) for the $d-2$ fresh vertices. A core support S hitting $\{u, v\}$ forces its fresh vertices to 0, whereas a missed edge allows $d-2$ choices of a single fresh vertex set to 1. □

Table I summarizes the principal configurations studied below.

The logical situation has two distinct branches. If a hypergraph has no admissible two-valued context state, it cannot admit a strong d-coloring, because the color classes of such a coloring would supply d admissible states. The converse implication fails: a noncontextual assignment of d eigenvalues to the outcomes may be impossible while ordinary states, including a unital separating family, remain available.

When a noncontextual eigenvalue assignment does exist, its color classes provide a nonempty unital family of extendable states inside the full context-state space. A particular ordinary state belongs to this restricted family precisely when deleting its support leaves a $(d-1)$-colorable graph. Chromatic identities describe pairs having equal eigenvalues in every noncontextual eigenvalue assignment, and hence equal values in every state of that restricted family; they need not hold throughout the ordinary state space. These assignment questions have to be distinguished from the separate requirements of an injective geometric realization, a representation of event order, and a classical mixture reproducing a specified Born probability vector.

III. INDEPENDENCE OF COLORING AND COORDINATIZATION, PETERSEN OBSTRUCTIONS, AND COMPLETED YU–OH

A. Small abstract examples and four-quadrant independence

The non-linear hypergraph $H_{4,2}$ with contexts $\{1,2,3\}$ and $\{1,2,4\}$ has three states $\{1\}$, $\{2\}$, $\{3,4\}$ (all extendable), forces $3 \sim_c 4$ and $s(3) = s(4)$, and has no injective $\mathbb{C}^3$ realization because two orthogonal rays determine a unique orthogonal-complement ray.

Proposition 8 (Smallest linear chromatically inseparable hypergraph)**.** *The linear* 3*-uniform hypergraph* $H_{6,3}$ *with contexts* $\{1,2,3\}$, $\{3,4,5\}$, $\{5,6,1\}$ *of Berge girth three has four unital separating states* $\{1,4\}$, $\{2,5\}$, $\{3,6\}$, $\{2,4,6\}$*, a unique strong* 3*-coloring up to permutation with inseparable classes* $\{1,4\}$, $\{2,5\}$, $\{3,6\}$*, and three extendable states (as* $V \setminus \{2,4,6\} = \{1,3,5\}$ *is a triangle). No linear* 3*-uniform hypergraph with* $|V| < 6$ *or* $|E| < 3$ *has a chromatically inseparable pair, and* $H_{6,3}$ *admits no injective realization in* $\mathbb{C}^3$*.*

Proof. The triangle $\{1,3,5\}$ in $\Gamma(H_{6,3})$ forces three distinct colors on $1,3,5$, whence $c(4) = c(1)$, $c(2) = c(5)$, and $c(6) = c(3)$. Similarly, in $\mathbb{C}^3$, $\{v_1, v_3, v_5\}$ forms an orthonormal basis, forcing $[v_4] = [v_1]$, $[v_2] = [v_5]$, and $[v_6] = [v_3]$. With $|E| \leq 2$, independent permutations of unshared vertices separate all pairs; by inclusion–exclusion, three linear 3-contexts require $|V| \geq 6$. □

Proposition 9 (Clique number at Berge girth at least four)**.** *If* $H = (V, E)$ *is an* n*-uniform linear hypergraph* ($n \geq 3$) *of Berge girth* $g(H) \geq 4$*, every clique* $K \subseteq V$ *of size* $|K| \geq 3$ *in* $\Gamma(H)$ *is contained in a unique context* $e \in E$*. Hence* $\omega(\Gamma(H)) = n$*, and chromatic incompleteness* ($\chi(H) > n$) *is equivalent to the chromatic–clique gap* $\chi(\Gamma(H)) > \omega(\Gamma(H))$*.*

Proof. For any three vertices $u, v, w \in K$, pairwise adjacency gives contexts $e_1 \supseteq \{u, v\}$, $e_2 \supseteq \{v, w\}$, $e_3 \supseteq \{w, u\}$. Distinct e_1, e_2, e_3 would form a Berge 3-cycle, contradicting $g(H) \geq 4$. By linearity, the shared context e containing $\{u, v, w\}$ is unique for $\{u, v\}$ and thus contains all of K. □

Proposition 10 (Logical independence of coloring and coordinatization)**.** *For finite* 3*-uniform context hypergraphs, all four combinations of chromatic completeness* ($\chi(H) = 3$) *and injective coordinatizability in* $\mathbb{C}^3$ *occur even within the orthomodular-lattice admissible class of linear hypergraphs of Berge girth* $g(H) \geq 5$[9,26] *with unital separating state spaces:*

1. $\chi = 3$*, injectively realizable in* $\mathbb{R}^3$*: a single context* $C_1 = \{\{1,2,3\}\}$ *(or* P_5*,* $H_{17,10}$*,* $H_{21,13}$*);*
2. $\chi = 3$*, not injectively realizable in* $\mathbb{C}^3$*: the* 8*-context Petersen restriction* H_8 *(Theorem 11);*
3. $\chi = 4$*, edge-faithfully realizable in* $\mathbb{R}^3$*: completed Yu–Oh* H^{+}_{YO} *(Theorem 13);*
4. $\chi = 4$*, admitting no orthogonal ray assignment in* $\mathbb{C}^3$*: Greechie's* G_{32} *and* H_9 *(Theorem 11).*

Injectivity is essential to quadrant (2): any strong n-coloring c yields a noninjective ray assignment $i \mapsto [e_{c(i)+1}] \in \mathbb{P}(\mathbb{F}^n)$. At Berge girth four (an orthomodular poset that is not a lattice), $E_{\square} = \{\{1,2,3\}, \{3,4,5\}, \{5,6,7\}, \{7,8,1\}\}$ also satisfies $\chi = 3$ while forcing $[v_3] = [v_7] \perp \mathrm{span}(v_1, v_5)$ whenever $[v_1] \neq [v_5]$.

B. Greechie's G_{32}, multipole forcing, and correction of the JMP spectral argument

Greechie's G_{32}[9] (Fig. 6) has 15 vertices and ten contexts

$$\begin{aligned} E(G_{32}) = \{&123, 345, 567, 789, 9\,10\,11,\\ &11\,12\,1, 4\,10\,13, 6\,12\,14,\\ &8\,2\,15, 13\,14\,15\}. \end{aligned} \tag{10}$$

Each vertex lies in two contexts, so G_{32} is dual to the Petersen graph P, $\Gamma(G_{32}) = L(P)$, $\mathrm{Aut}(G_{32}) \cong S_5$, and $\chi(G_{32}) = \chi'(P) = 4$. Its six two-valued state supports satisfy

$$|\mathcal{S}(G_{32})| = 6, \qquad |S_\alpha| = 5, \qquad |S_\alpha \cap S_\beta| = 1 \ (\alpha \neq \beta). \tag{11}$$

These supports yield the $\binom{6}{2} = 15$ two-element vertex profiles $B_i = \{\alpha : i \in S_\alpha\}$. Their nonempty pairwise intersections exclude even two disjoint states, confirming $\chi(G_{32}) = 4$ by Lemma 3.

In Appendix B of Shekarriz and Svozil[6], the chromatic obstruction $\chi(G_{32}) = 4 > 3$ was invoked to argue against a 3-dimensional realization of G_{32}. As Proposition 10 demonstrates via H^+_{YO} ($\chi = 4$, yet faithfully realizable in $\mathbb{R}^3$), $\chi(H) > 3$ rules out only a context-independent 3-value spectral assignment—a noncontextual assignment of the eigenvalues of uniform maximal observables to their outcomes—*not* geometric rays. A genuine geometric exclusion requires the projection identity: for an orthonormal basis $\{x, y, z\} \subset \mathbb{C}^3$ with $u \perp x$ and $w \perp z$,

$$\langle u, w\rangle = \langle u, y\rangle\langle y, w\rangle. \tag{12}$$

Theorem 11 (Multipole forcing and minimal complex obstructions)**.** *Let* $H_8 = G_{32} \setminus \{\{8, 2, 15\}, \{13, 14, 15\}\}$ *(*14 *vertices,* 8 *contexts) and* $H_9 = G_{32} \setminus \{\{13, 14, 15\}\}$ *(*15 *vertices,* 9 *contexts).*

1. *In every orthogonal ray assignment of* H_8 *in* $\mathbb{C}^3$ *(allowing repeated rays), the terminal rays satisfy*

$$[v_2] = [v_8] \qquad \textit{and} \qquad [v_{13}] = [v_{14}], \tag{13}$$

and either $[v_1] = [v_4] = [v_9]$ *or* $[v_3] = [v_7] = [v_{12}]$*. Thus* H_8 *has no injective realization in* $\mathbb{C}^3$*.*

2. H_8 *is chromatically complete (*$\chi(H_8) = 3$*) with* 11 *unital separating states, of which* 7 *are extendable (the* 4 *nonextendable states leave odd cycles in* $\Gamma(H_8)$*; see Table S38). Its* 3 *normalized* 3*-colorings (fixing* $(c(1), c(2), c(3)) = (0, 1, 2)$*) are listed in Table S39, forcing the* 6 *chromatically inseparable pairs* $(1, 9)$*,* $(2, 8)$*,* $(3, 7)$*,* $(4, 6)$*,* $(10, 12)$*,* $(13, 14)$*; deleting any single context of* H_8 *removes all forced color identities (leaving* $6, 20, 20, 6, 20, 20, 6, 6$ *normalized colorings on the remaining active vertices, up to global color permutation). Isolated vertices are dropped: retaining the isolated vertex created by deleting* $\{1, 2, 3\}$*,* $\{7, 8, 9\}$*,* $\{4, 10, 13\}$*, or* $\{6, 12, 14\}$ *multiplies the corresponding count by three, from* 6 *to* 18*.*

3. *Neither* H_9*,* G_{32}*, nor the* 12*-vertex,* 21*-edge subgraph* Γ_0 *of* $\Gamma(H_9)$ *on* $\{1, \ldots, 12\}$ *admits any orthogonal ray assignment in* $\mathbb{C}^3$*. Moreover,* H_9 *is context-minimal and* Γ_0 *is edge-minimal: deleting any context of* H_9 *or edge of* Γ_0 *yields a* 3*-colorable structure realizable in* $\mathbb{R}^3$ *by standard basis rays.*

Proof. (1) Let $v_1, \ldots, v_{14} \in \mathbb{C}^3$ be unit vectors for H_8. In $\{v_1, v_2, v_3\}$, $v_{11}, v_{12} \perp v_1$ and $v_4, v_5 \perp v_3$, so Eq. (12) gives $\langle x, y\rangle = \langle x, v_2\rangle\langle v_2, y\rangle$ for $x \in \{v_{11}, v_{12}\}, y \in \{v_4, v_5\}$. Expanding v_{12} in $\{v_5, v_6, v_7\}$ ($v_{12} \perp v_6$) and v_4 in $\{v_9, v_{10}, v_{11}\}$ ($v_4 \perp v_{10}$) yields $v_7\langle v_7, v_{12}\rangle = v_{12} - v_5\langle v_5, v_{12}\rangle$ and $v_9\langle v_9, v_4\rangle = v_4 - v_{11}\langle v_{11}, v_4\rangle$. Since $v_7 \perp v_9$, $v_{12} \perp v_{11}$, and $v_5 \perp v_4$, their inner product vanishes:

$$\begin{aligned} 0 &= \langle v_{12}, v_4\rangle + \langle v_{12}, v_5\rangle\langle v_5, v_{11}\rangle\langle v_{11}, v_4\rangle \\ &= \langle v_{12}, v_2\rangle\langle v_2, v_4\rangle\big(1 + |\langle v_2, v_5\rangle|^2|\langle v_2, v_{11}\rangle|^2\big). \end{aligned} \tag{14}$$

Thus $\langle v_2, v_4\rangle = 0$ or $\langle v_{12}, v_2\rangle = 0$. If $\langle v_2, v_4\rangle = 0$, set $(v_1, v_3, v_2) = (e_1, e_2, e_3)$; then $v_4 \perp v_2, v_3 \implies [v_4] = [e_1]$, $\{3, 4, 5\} \implies [v_5] = [e_3] = [v_2]$, $\{9, 10, 11\}$ with $v_{10}, v_{11} \perp e_1$ forces $[v_9] = [e_1]$, $\{5, 6, 7\}$ and $\{7, 8, 9\}$ force $[v_7] = [e_2], [v_6] = [e_1], [v_8] = [e_3] = [v_2]$, and $\{4, 10, 13\}$, $\{9, 10, 11\}$, $\{11, 12, 1\}$, $\{6, 12, 14\}$ in $\mathrm{span}(e_2, e_3)$ force $[v_{13}] = [v_{11}] = [v_{14}]$ and $[v_1] = [v_4] = [v_9]$. Reflection symmetry gives the second branch when $\langle v_{12}, v_2\rangle = 0$.

(2) In H_8, fixing $(c(1), c(2), c(3)) = (0, 1, 2)$ leaves 16 choices along $345, 567, 789, 9\,10\,11$, of which 13 fail on $11\,12\,1, 4\,10\,13, 6\,12\,14$, leaving c_1, c_2, c_3 in Table S39. Equal columns identify the 6 pairs.

(3) Both H_9 (hence G_{32}) and Γ_0 add the orthogonality $v_2 \perp v_8$ to H_8, contradicting $[v_2] = [v_8]$. For minimality, deleting $\{8, 2, 15\}$ from H_9 leaves H_8 ($\chi = 3$); deleting $\{4, 10, 13\}$ or $\{6, 12, 14\}$ is isomorphic by D_6 symmetry to deleting $\{8, 2, 15\}$; and deleting any ring context (say $\{1, 2, 3\}$) allows the 3-coloring $(c(1), \ldots, c(15)) = (2, 1, 2, 0, 1, 0, 2, 0, 1, 2, 0, 1, 1, 2, 2)$. On Γ_0, the following deletions cover all 21 edges up to D_6 symmetry: $\{1, 2\}$ (colored by 001020120121), $\{2, 3\}$ (011020120121), $\{1, 3\}$ (010120102012), and $\{4, 10\}$ (012010201021), each yielding a standard-basis $\mathbb{R}^3$ assignment $i \mapsto [e_{c(i)+1}]$. For the minimum Δ_3 of $\sum |\langle v_i, v_j \rangle|^2$ over the 30 edges $\{i, j\}$ of $\Gamma(G_{32})$ and all unit vectors $v_1, \ldots, v_{15} \in \mathbb{C}^3$, the robustness bounds $1/14400 < \Delta_3 < 0.927496$ are proved in Sec. S3 of the supplementary material. □

In his proof that state-independent contextuality in $\mathbb{C}^3$ requires 13 rays, Cabello[8] found Γ_0 to be the only square-free graph with fewer than 13 vertices and chromatic number above 3, and excluded it by showing, in real coordinates, that ten of its vertices (in our labels, all but 2 and 8) cannot be distinct rays. Part (1) proves this over $\mathbb{C}^3$: the rays $2, 8, 13, 14$ of H_8 are then fixed as orthogonal completions, and the ten rays satisfy $[v_1] = [v_4] = [v_9]$ or $[v_3] = [v_7] = [v_{12}]$. Part (3) shows more: Γ_0 admits no orthogonal ray assignment in $\mathbb{C}^3$ at all, even with repeated rays.

The complete normalized coloring lists for H_8 and $H_{17,10}$ are in the supplement.

Corollary 12 (Blanuša and Isaacs dot-product snarks)**.** *If a cubic graph G admits no orthogonal vector edge coloring in $\mathbb{C}^3$, neither does any Isaacs dot product $G \cdot P$[29], obtained by deleting two nonadjacent edges of G and two adjacent vertices u, v of the Petersen graph P, and joining the two dangling edges of u to the ends of one deleted edge and those of v to the ends of the other. In particular, both 18-vertex Blanuša snarks $B_1, B_2 = P \cdot P$[30] and all iterated dot products $X = P \cdot P \cdots P$ satisfy $\chi(H(X)) = 4 > 3 = \omega(L(X))$ and admit no orthogonal ray assignment in $\mathbb{C}^3$.*

Proof. In $G \cdot P$, the embedded copy of $P \setminus \{u, v\}$ has dual hypergraph H_8, so Theorem 11(1) forces $[v_2] = [v_8]$ and $[v_{13}] = [v_{14}]$ on the two pairs of cut edges, inducing a valid orthogonal ray assignment on G. □

Silva and Ibiapina[23] (Question 6) asked whether a cubic graph with chromatic index 4 can have an orthogonal vector edge coloring in $\mathbb{R}^3$. For the Petersen graph, the Blanuša snarks, and these dot products the answer is no, even in $\mathbb{C}^3$.

C. Completed Yu–Oh H^+_{YO}, base-coloring partition, and context criticality

Completing each of the 12 uncovered orthogonal pairs of the 13-ray Yu–Oh set[10] ($r_1, \ldots, r_{13} \in \mathbb{R}^3$, consisting of $z_1 = r_2, z_2 = r_8, z_3 = r_5$, $y_1^{\pm} = r_{1,3}, y_2^{\pm} = r_{7,9}, y_3^{\pm} = r_{4,6}$, and diagonals $h_0 = r_{12}, h_1 = r_{11}, h_2 = r_{10}, h_3 = r_{13}$) by cross-product rays $r_{14}, \ldots, r_{25}$ (Table S2) yields the 25-ray, 16-context linear hypergraph H^+_{YO} of Berge girth five[7,31]:

$$\begin{aligned} E^+_{\mathrm{YO}} &= \{123, 789, 456, 285\} \cup \bigcup_{h=10}^{13} \mathcal{E}_h, \\ \mathcal{E}_h &= \{\{x, h, w_{xh}\} : x \in N(h)\}, \end{aligned} \tag{15}$$

All 25 vectors, 16 contexts, and restriction-membership flags are tabulated in the supplement. Here the neighborhoods are $N(10) = \{1, 9, 4\}$, $N(11) = \{3, 7, 4\}$, $N(12) = \{3, 9, 6\}$ and $N(13) = \{1, 7, 6\}$, and each w_{xh} is the unique completing ray. It has a real FOR in $\mathbb{R}^3$ (48 orthogonal pairs) and 24 unital separating states (each of the 12 binary assignments on the four base contexts leaves uniquely one diagonal h_j with all three neighbors zero, allowing $s(h_j) \in \{0, 1\}$ and hence $12 + 4 \times 3 = 24$ states). Completed Yu–Oh is therefore a faithfully realized real configuration in which noncontextual truth assignments survive, enough of them to separate every pair of rays, while by the following theorem no noncontextual assignment of three eigenvalues to its outcomes exists.

Theorem 13 (Base-coloring partition and context criticality of H^+_{YO})**.** *The completed Yu–Oh hypergraph H^+_{YO} satisfies $\omega(\Gamma(H^+_{\mathrm{YO}})) = 3$, $\chi_f(\Gamma(H^+_{\mathrm{YO}})) = 35/11 > 3 = \chi_f(\Gamma(G_{32}))$, and $\chi(H^+_{\mathrm{YO}}) = 4$, and is context-critical for $\chi > 3$:*

1. *Fixing $(c(2), c(8), c(5)) = (0, 1, 2)$, the four base contexts $\{1, 2, 3\}$, $\{7, 8, 9\}$, $\{4, 5, 6\}$, $\{2, 8, 5\}$ admit eight strong 3-colorings indexed by $(b_1, b_2, b_3) \in \{0, 1\}^3$:*

$$\begin{aligned} (c(1), c(3)) &= (2 - b_1, 1 + b_1), \\ (c(7), c(9)) &= (2 - 2b_2, 2b_2), \\ (c(4), c(6)) &= (1 - b_3, b_3). \end{aligned} \tag{16}$$

   Each antipodal pair of (b_1, b_2, b_3) makes the neighborhood $N(h_j)$ of uniquely one *diagonal ray $h_j \in \{10, 11, 12, 13\}$ rainbow ($\{0, 1, 2\}$) while the other three neighborhoods are 2-colored:*

Antipodal binary patterns	*Blocked diagonal*
$101, 010$	12
$110, 001$	11
$000, 111$	10
$011, 100$	13

(17)

2. *Consequently, $\chi(H^+_{\mathrm{YO}}) = 4$, whereas deleting any single context $e \in E^+_{\mathrm{YO}}$ yields $\chi(H^+_{\mathrm{YO}} \setminus \{e\}) = 3$.*

Proof. Direct check of Eqs. (16)–(17) shows that every base 3-coloring blocks one h_j, so $\chi(H^+_{\mathrm{YO}}) \geq 4$, matched by the four-coloring

$$(c(1), \ldots, c(25)) = (2, 0, 1, 1, 2, 0, 2, 1, 0, 3, 0, 2, 1, 2, 1, 1, 1, 0, 0, 0, 0, 2, 1, 2, 0),$$

in which only $h_2 = 10$ receives the fourth color. Deleting a sector context $\{x, h_j, w\}$ frees h_j in its antipodal base colorings by setting $c(h_j) = c(x)$; deleting $\{2, 8, 5\}$ allows $c(2) = c(8) = c(5) = 0$; deleting $\{1, 2, 3\}$ allows $(c(1), c(2), c(3)) = (0, 0, 1)$, $(c(7), c(8), c(9)) = (2, 1, 0)$, $(c(4), c(5), c(6)) = (1, 2, 0)$, and $(c(10), c(11), c(12), c(13)) = (2, 0, 2, 1)$; the cases $\{7, 8, 9\}$ and $\{4, 5, 6\}$ follow by the threefold cyclic coordinate symmetry of the Yu–Oh set. For χ_f, weight $1/2$ on the 6 states of G_{32} gives $\chi_f(G_{32}) = 3$, while Proposition 7 preserves the 13-ray value $\chi_f = 35/11$[14]. □

IV. EXACT-PALETTE CHROMATIC SEPARABILITY AND LOGICAL GADGETS IN DIMENSIONS THREE AND FOUR

From now on we assume that noncontextual assignments of d eigenvalues exist ($\chi(H) = d$) and ask what they imply for the relations between distinct outcomes. Such assignments may force outcomes to share or to avoid an eigenvalue although ordinary truth assignments permit otherwise; these forced relations are their logical content.

A. Kochen–Specker antecedents and palette-invariant implications

Three historical Kochen–Specker constructions clarify the hierarchy of logical gadgets:

1. *1965 abstract TIFS precursor:* Identifying repeated labels v_1 and v_2 in Kochen and Specker's 1965 Fig. 1[32] (p. 182) yields an 11-vertex, 7-context hypergraph $\{134, 325, 2\,9\,11, 11\,1\,10, 479, 5\,8\,10, 678\}$ implementing $s(1) = 1 \implies s(2) = 0$. It has only four states ($\{1, 5, 6, 9\}$, $\{2, 4, 6, 10\}$, $\{3, 6, 9, 10\}$, $\{4, 5, 6, 11\}$), all with $s(6) = 1$ and $s(7) = s(8) = 0$, so $\chi > 3$; moreover, 1 and 9 are both orthogonal to 4 and 11, so the distinct rays 4 and 11 of an injective realization in $\mathbb{C}^3$ would force $[v_1] = [v_9]$; no injective realization exists (Sec. S3 of the supplementary material).

2. *1967 directed TITS* Γ_1*:* The 10-vertex graph Γ_1 of Lemma 1 in Ref.[4] (pp. 68, 70) implements the one-way implication $s(a_0) = 1 \implies s(a_9) = 1$. Completing its six uncompleted edges by fresh rays $23, 19, 18, 15, 14, 5$ yields the 16-ray, 9-context configuration $I_{16,9}$ in Fig. 2(d) (explicit label map in Sec. S3 of the supplementary material).

3. *1967 state-inseparable* Γ_3*:* In Ref.[4] (p. 70), Γ_3 forces $s(a) = s(b)$ in *every* two-valued state (a, b are nonseparated by $\mathcal{S}(H)$). By contrast, our chromatically inseparable examples below have $\mathcal{S}(H)$ separating all vertices, with $c(u) = c(v)$ forced only on $\mathcal{S}_{\text{ext}}(H)$.

For distinct nonadjacent terminals $u, v \in V$, an ordinary TIFS[33] obeys $s(u) = 1 \implies s(v) = 0$ (equivalently $s(u) + s(v) \leq 1$, which is always bidirectional), and a directed ordinary TITS obeys $s(u) = 1 \implies s(v) = 1$. When $\mathcal{C}_d(H) \neq \varnothing$, an *unanchored chromatic TIFS* obeys $c(u) = 0 \implies c(v) \in \{1, \ldots, d-1\}$, and an *unanchored chromatic TITS* obeys $c(u) = 0 \implies c(v) = 0$.

Proposition 14 (Palette symmetry and purely chromatic gadgets)**.** *For chromatically complete H and nonadjacent $u, v \in V$, the unanchored chromatic TIFS is equivalent to $c(u) \neq c(v)$ for all $c \in \mathcal{C}_d(H)$ (an* implicit edge[11] *(Def. 3.1)), and the unanchored chromatic TITS is equivalent to $u \sim_c v$ (an* implicit identity*). Both chromatic relations are bidirectional. Ordinary TIFS and directed ordinary TITS entail their chromatic counterparts, and with only $c(u) = 0$ fixed, the allowed output set $\{c(v) : c \in \mathcal{C}_d(H),\ c(u) = 0\}$ is one of $\{0\}$, $\{1, \ldots, d-1\}$, or $\{0, \ldots, d-1\}$.*

Proof. Global palette permutations act transitively on $\{0, \ldots, d-1\}$ and, stabilizing 0, transitively on $\{1, \ldots, d-1\}$. Since every color class $s_a(x) = \mathbf{1}_{\{c(x)=a\}}$ is in $\mathcal{S}_{\text{ext}}(H) \subseteq \mathcal{S}(H)$, any ordinary TIFS or one-way TITS forces $c(u) \neq c(v)$ or $c(u) = c(v)$, respectively. □

We call a chromatic TIFS *purely chromatic* if some $s \in \mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$ satisfies $s(u) = s(v) = 1$, and a chromatic identity $u \sim_c v$ *two-way purely chromatic* if ordinary states in $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$ realize both $(s(u), s(v)) = (1, 0)$ and $(0, 1)$, so neither directed ordinary TITS holds. Since a state with $s(u) = s(v) = 1$ cannot be a color class when $c(u) \neq c(v)$ for all c, and a state with $s(u) \neq s(v)$ cannot be one when $u \sim_c v$, the qualifier $\mathcal{S}(H) \setminus \mathcal{S}_{\text{ext}}(H)$ is automatic: a chromatic TIFS is purely chromatic exactly when the ordinary TIFS fails, and a chromatic identity is two-way purely chromatic exactly when both directed ordinary TITS fail.

B. Faithful Yu–Oh restrictions $H_{17,10}$ and $H_{21,13}$ and exhaustive inventory search

The base-coloring partition of H^+_{YO} in Theorem 13 directly produces chromatically inseparable faithful restrictions when diagonal sectors are omitted. Retain the four base contexts $\{1, 2, 3\}$, $\{7, 8, 9\}$, $\{4, 5, 6\}$, $\{2, 8, 5\}$ of Eq. (15). Keeping the two diagonal sectors at 10 and 11 defines $H_{17,10}$ on 17 rays and 10 contexts [Fig. 1(a)]; keeping the three sectors at $10, 11, 13$ defines $H_{21,13}$ on 21 rays and 13 contexts [Fig. 1(b)]. Deleting $\{4, 5, 6\}$, $\{4, 11, 24\}$, $\{4, 10, 18\}$ from $H_{17,10}$ recovers the 13-ray, 7-context completed Specker bug $B_{13,7}$ [Fig. 2(b)].

Theorem 15 (Chromatic inseparability in $H_{17,10}$ and $H_{21,13}$)**.** *Both $H_{17,10}$ and $H_{21,13}$ are linear of Berge girth five and inherit injective edge-faithful realizations in $\mathbb{R}^3$ from H^+_{YO}:*

1. *$H_{17,10}$ has* 18 *unital separating states,* 4 *normalized strong* 3*-colorings (Table S41), and* 10 *extendable states. Its only distinct chromatically inseparable pairs are* $(10, 24)$ *and* $(11, 18)$*. The ordinary implications* $10 \Rightarrow 24$ *and* $11 \Rightarrow 18$ *already hold, but their converses fail; these identities are not two-way purely chromatic. Relative to the anchor* $(c(2), c(8), c(5)) = (0, 1, 2)$*, its four colorings restrict on* $(9, 10, 11)$ *to* $(0, 2, 1)$*,* $(0, 2, 0)$*,* $(2, 1, 2)$*,* $(2, 0, 2)$*, yielding the anchored implications*

$$\begin{aligned} c(9) = 0 &\implies c(11) \in \{0, 1\}, \\ c(10) = 0 &\implies c(11) = 2. \end{aligned} \tag{18}$$

(a) $H_{17,10}$ (17 rays, 10 contexts, 2 inseparable pairs)

(b) $H_{21,13}$ (21 rays, 13 contexts, 21 inseparable pairs)




FIG. 1. Faithful three-dimensional Yu–Oh restrictions in $\mathbb{R}^3$, drawn in the common vertex layout of completed Yu–Oh. (a) $H_{17,10}$, omitting diagonal sectors $h_0 = 12$ and $h_3 = 13$, with four normalized strong 3-colorings and chromatically inseparable pairs $10 \sim_c 24$ and $11 \sim_c 18$; (b) $H_{21,13}$, omitting only $h_0 = 12$, with two normalized strong 3-colorings and 21 chromatically inseparable pairs across nine chromatic signature classes.

2. *$H_{21,13}$ has* 21 *unital separating states,* 2 *normalized strong* 3*-colorings, and* 6 *extendable states. Its nine chromatic signature classes* $(c_1(v), c_2(v)) \in \{0,1,2\}^2$ *are listed in Table S44, giving* $3\binom{4}{2} + 3\binom{2}{2} = 21$ *chromatically inseparable pairs. Exactly* 15 *are two-way purely chromatic identities (Table S70). The remaining six pairs obey the directed ordinary implications*

$$11 \Rightarrow 18, \quad 11 \Rightarrow 21, \quad 13 \Rightarrow 15,$$
$$13 \Rightarrow 19, \quad 10 \Rightarrow 20, \quad 10 \Rightarrow 24,$$

each with a false converse. This classification exhausts all 21 *pairs; no distinct vertices are inseparable under all ordinary states.*

Proof. By Theorem 13(1), among the 8 base colorings in Eq. (16), sector 11 excludes $(1,1,0)$, $(0,0,1)$ and sector 10 excludes $(0,0,0)$, $(1,1,1)$. The 4 surviving (b_1,b_2,b_3) extend uniquely to $H_{17,10}$ (Table S41): in each, $N(10) = \{1,9,4\}$ and $N(11) = \{3,7,4\}$ are 2-colored with $c(10) \neq c(4)$ and $c(11) \neq c(4)$, and $c(10) \neq c(11)$ by the restrictions to $(9,10,11)$ listed in (1), forcing $c(24) = c(10)$ in $\{4,11,24\}$ and $c(18) = c(11)$ in $\{4,10,18\}$. Adding sector 13 in $H_{21,13}$ additionally excludes $(0,1,1)$, $(1,0,0)$, leaving only $(1,0,1)$ and $(0,1,0)$, whose column signatures are listed in Table S44. Each retained diagonal adds 3 states to the 12 base states, giving $12 + 2 \times 3 = 18$ and $12 + 3 \times 3 = 21$ states (Sec. S2 of the supplementary material). Testing both directed truth patterns on these 21 states gives the exhaustive 15-versus-6 classification; Table S70 of the supplementary material supplies witnesses for all 15 two-way pairs, together with a reproducible enumeration procedure. □

The full chromatic signatures and both-direction truth witnesses are in the supplement. The witnesses are generated from the complete state families and checked against every context.

Physically, every noncontextual eigenvalue assignment on $H_{17,10}$ gives the nonorthogonal rays 10 and 24, and likewise 11 and 18, the same eigenvalue, although ordinary truth assignments distinguish them. In $H_{21,13}$, fifteen of the twenty-one forced identities are not even shadowed by a one-way truth implication: they exist only at the level of noncontextual eigenvalues.




FIG. 2. Faithful real three-dimensional logical gadgets in $\mathbb{R}^3$. (a) Pentagon P_5 with purely chromatic ternary exclusion $\neg(c(6) = c(7) = c(9))$; (b) completed Specker bug $B_{13,7}$ (ordinary TIFS $c(10) \neq c(11)$); (c) purely chromatic binary TIFS $D_{16,9}$ ($c(14) \neq c(18)$); (d) completed historical Γ_1 TITS $I_{16,9}$ ($10 \sim_c 24, 11 \sim_c 18$); (e) two-way purely chromatic identity $T_{19,11}$ ($14 \sim_c 28, 18 \sim_c 27$); (f) $H_{21,13}$ showing three of its fifteen two-way purely chromatic identities: $2 \sim_c 14, 5 \sim_c 22, 8 \sim_c 23$. Red triangles forbid monochromatic triples; red dashed arrows denote $c(u) \neq c(v)$; blue dotted arrows denote $u \sim_c v$.

Proposition 16 (Complete inventory classification)**.** *Among all* $2^{16} - 1 = 65{,}535$ *nonempty context subsets* $E' \subseteq E(H_{\mathrm{YO}}^{+})$ *with inherited coordinates from Tables S1–S3, exactly* 539 *are connected and edge-faithful, and among these* precisely ten *are* 3*-colorable with a nontrivial chromatic inseparability class: the* $\binom{4}{2} = 6$ *two-sector restrictions (*17 *rays,* 10 *contexts,* 18 *states,* 4 *normalized colorings,* 2 *inseparable pairs) and the* $\binom{4}{3} = 4$ *three-sector restrictions (*21 *rays,* 13 *contexts,* 21 *states,* 2 *normalized colorings,* 21 *inseparable pairs). Thus* 17 *rays and* 10 *contexts are minimal for chromatic inseparability within the faithful restrictions of* H_{YO}^{+} *with inherited coordinates.*

Proof. Since H_{YO}^{+} is linear and edge-faithful, a subset E' is edge-faithful if and only if no omitted context contains two vertices of $V' = \bigcup E'$. Exact enumeration by two independent coloring algorithms over all 539 connected faithful subsets certifies the ten solutions listed in Table S71. Disconnected restrictions separate distinct components by independent palette permutations. □

The ten retained-sector cases and binary search masks are enumerated in the supplement. The restriction to inherited coordinates matters: $I_{16,9}$ of Sec. IV C is $H_{17,10}$ without the context $\{4, 5, 6\}$; it has 16 rays and 9 contexts and keeps both identities, but it is edge-faithful only with refitted coordinates.

C. Faithful three-dimensional TIFS, TITS, and purely chromatic gadgets

Figure 2 and Table II synthesize our faithful real 3-dimensional gadgets (coordinates and state/coloring tables in Secs. S1–S2 of the supplementary material):

- *Pentagon P_5 [Fig. 2(a)]:* Contexts $\{1,6,2\}$, $\{2,7,3\}$, $\{3,8,4\}$, $\{4,9,5\}$, $\{5,10,1\}$ have 5 normalized 3-colorings, 11 states, and 10 extendable states, separating all vertex pairs (showing that chromatic separation does not require $\mathcal{S}_{\text{ext}} = \mathcal{S}$). Its degree-one vertices obey the purely chromatic ternary exclusion $\neg(c(6) = c(7) = c(9))$: assigning color 0 to $6,7,9$ forces $1,2,3,4,5$ into $\{1,2\}$ with $c(1) = c(3)$ and $c(4) = c(5)$, contradicting $\{4,9,5\}$, whereas the nonextendable state $\{6,7,8,9,10\}$ (whose complement is the 5-cycle 12345) assigns 1 to all three terminals.

- *Specker bug $B_{13,7}$ and completed Γ_1 $I_{16,9}$ [Fig. 2(b),(d)]:* $B_{13,7}$ is an ordinary and chromatic TIFS ($c(10) \neq c(11)$, overlap 1/9). Adjoining $\{4,11,24\}$ and $\{4,10,18\}$ gives $I_{16,9}$ (deleting $\{4,5,6\}$ from $H_{17,10}$ with refitted faithful coordinates), where $c(10) \neq c(11)$ forces $c(4)$ to be the third color, so $c(24) = c(10)$ and $c(18) = c(11)$ via the one-way ordinary TITS $s(10) = 1 \implies s(24) = 1$ (overlap 25/27).

Proposition 17 (Purely chromatic binary TIFS $D_{16,9}$ and two-way identity $T_{19,11}$)**.** *Let $D_{16,9} = H_{17,10} \setminus \{\{1,10,19\}\}$ (with orphan 19 removed) and $T_{19,11} = D_{16,9} \cup \{\{14,26,27\}, \{18,26,28\}\}$, realized faithfully in $\mathbb{R}^3$ [Fig. 2(c),(e)]:*

1. *Every $c \in \mathcal{C}_3(D_{16,9})$ satisfies $c(14) \neq c(18)$ (7 normalized colorings, 14 extendable states), whereas the nonextendable state $\{1,5,7,14,18,23,24\} \in \mathcal{S}(D_{16,9})$ assigns $s(14) = s(18) = 1$ (squared terminal overlap 16/2025).*

2. *Every $c \in \mathcal{C}_3(T_{19,11})$ satisfies $14 \sim_c 28$ and $18 \sim_c 27$ (7 normalized colorings, 25 states, 14 extendable), while for both pairs, nonextendable states $\{1,5,7,14,18,23,24\}$ and $\{1,6,8,10,11,27,28\}$ realize $(1,0)$ and $(0,1)$, violating both directed ordinary TITS implications (squared overlap 2009/2025; Table S70).*

Proof. (1) If $c(14) = c(18) = 0$, set $c(3) = 1, c(11) = 2$ by $\{3,11,14\}$. Then $\{4,11,24\}$ and $\{4,10,18\}$ force $c(4) = 1$ and $c(10) = 2$. In $\{7,8,9\}$, $7 \perp 11$ and $9 \perp 10$ avoid color 2, forcing $c(8) = 2$. Then $\{1,2,3\}$ and $\{4,5,6\}$ make 2 and 5 avoid color 1, so $\{2,8,5\}$ forces $c(2) = c(5) = 0$, a contradiction. (2) In $T_{19,11}$, $c(14) \neq c(18)$ forces their common neighbor 26 ($r_{26} \parallel r_{14} \times r_{18}$) to take the third color, whence $c(27) = c(18)$ and $c(28) = c(14)$. □

D. Apex extensions and faithful four-dimensional linear gadgets

Proposition 18 (Apex extension)**.** *Embedding any 3-uniform hypergraph H realized in $\mathbb{F}^3$ into $\mathbb{F}^d$ ($d \geq 3$) and appending $d-3$ mutually orthogonal coordinate apices to every context preserves injectivity, edge-faithfulness, $\chi(H^{(d)}) = \chi(H) + d - 3$, normalized coloring count, and old chromatic inseparability classes (with apices as singletons), while adding $d-3$ apex-only states (all extendable when $\chi(H) = 3$) to the zero-extended states of H.*

Proof. In every coloring of $H^{(d)}$, the $d-3$ apices occupy $d-3$ distinct colors across all contexts, leaving a 3-coloring of H; similarly, any state either sets one apex to 1 (forcing all old vertices to 0) or sets all apices to 0 and restricts to $\mathcal{S}(H)$. □

Applying Proposition 18 at $d = 4$ to P_5, $B_{13,7}$, and $D_{16,9}$ yields the non-linear ($g = 2$) four-dimensional gadgets $W^{(4)}_{11,5}$, $B^{(4)}_{14,7}$, and $D^{(4)}_{17,9}$ in Fig. 3(a),(c),(e). To achieve *linearity* ($|e \cap e'| \leq 1$) in $\mathbb{R}^4$, we construct three faithful linear gadgets ($g = 3$) in Fig. 3(b),(d),(f) (Table II):

TABLE II. Faithful geometric gadgets. $N_c = |\mathcal{C}_d|/d!$ fixes one context palette. Terminal relations describe colors unless explicitly identified as ordinary truth implications.

Gadget	d	$\lvert V\rvert/\lvert E\rvert$	N_c	$\lvert\mathcal{S}\rvert$	$\lvert\mathcal{S}_{\text{ext}}\rvert$	Relation and mechanism
P_5	3	10/5	5	11	10	Ternary exclusion on $6, 7, 9$; purely chromatic.
$B_{13,7}$	3	13/7	6	14	12	$c(10) \neq c(11)$; ordinary TIFS.
$D_{16,9}$	3	16/9	7	21	14	$c(14) \neq c(18)$; purely chromatic TIFS.
$I_{16,9}$	3	16/9	6	22	12	$10 \sim_c 24$, $11 \sim_c 18$; directed ordinary TITS.
$T_{19,11}$	3	19/11	7	25	14	$14 \sim_c 28$, $18 \sim_c 27$; both truth directions fail.
$H_{21,13}$	3	21/13	2	21	6	21 identical pairs: 15 two-way purely chromatic; six directed ordinary TITS.
$H_{17,10}$	3	17/10	4	18	10	Two color-identical pairs from directed ordinary TITS.
$W^{(4)}_{11,5}$	4	11/5	5	12	11	Apex ternary exclusion.
$W^{(4)}_{16,6}$	4	16/6	80	72	56	Linear ternary exclusion on $6, 8, 11$.
$B^{(4)}_{14,7}$	4	14/7	6	15	13	Apex ordinary TIFS.
$L^{(4)}_{18,7}$	4	18/7	144	68	68	Linear ordinary TIFS $c(11) \neq c(15)$.
$D^{(4)}_{17,9}$	4	17/9	7	22	15	Apex purely chromatic TIFS.
$L^{(4)}_{22,9}$	4	22/9	216	158	92	Linear identity $11 \sim_c 20$ from directed ordinary TITS.

1. *Linear purely chromatic ternary gadget $W^{(4)}_{16,6}$ [Fig. 3(b)]:* Contexts $\{2, 0, 6, 1\}$, $\{2, 7, 13, 3\}$, $\{3, 0, 8, 4\}$, $\{4, 9, 14, 5\}$, $\{5, 10, 15, 1\}$, $\{0, 11, 12, 5\}$ satisfy $\neg(c(6) = c(8) = c(11))$ (if $c(6) = c(8) = c(11) = 0$ and $c(0) = 3$, then $1, 2, 3, 4, 5$ must 2-color a 5-cycle), while $\{6, 7, 8, 9, 10, 11\} \in \mathcal{S}(W^{(4)}_{16,6})$ sets all three terminals to 1.

2. *Linear ordinary TIFS $L^{(4)}_{18,7}$ [Fig. 3(d)]:* Contexts $\{1, 2, 4, 3\}$, $\{5, 8, 7, 6\}$, $\{4, 9, 10, 8\}$, $\{11, 1, 12, 5\}$, $\{11, 14, 13, 2\}$, $\{15, 3, 16, 7\}$, $\{15, 17, 18, 6\}$ enforce $s(11) = 1 \implies s(15) = 0$ and $c(11) \neq c(15)$ (if $c(11) = c(15) = a$, then $1, 2, 3$ and $5, 6, 7$ avoid a, forcing $c(4) = c(8) = a$ in $\{4, 9, 10, 8\}$). Using $r_{13} = (1, 0, 1, 1), r_{14} = (2, 0, -1, -1)$ avoids an unlisted orthogonality $13 \perp 15$; all 68 states are extendable (terminal overlap 1/42).

3. *Linear directed TITS and color identity $L^{(4)}_{22,9}$ [Fig. 3(f)]:* Adjoining $\{11, 21, 22, 19\}$ and $\{13, 20, 15, 19\}$ to $L^{(4)}_{18,7}$ (with $r_{13} = (1, 0, 4, 4), r_{14} = (8, 0, -1, -1)$ so $r_{13} \perp r_{15}$) makes $\{11, 13, 15, 19\}$ pairwise differently colored, forcing $11 \sim_c 20$ and $s(11) = 1 \implies s(20) = 1$ (while $\{1, 6, 9, 14, 16, 20, 21\}$ has $s(11) = 0, s(20) = 1$; terminal overlap 41/42).

V. THE 37-RAY COMPLEX HESSE HYPERGRAPH AND RATIONAL FOUR-DIMENSIONAL COMPLETIONS

A. The 37-ray complex Hesse hypergraph H_{37} and its 76-event orthomodular lattice

With $\zeta = \exp(2\pi i/3)$ ($\zeta^2 + \zeta + 1 = 0$) and $a, k \in \mathbb{Z}_3$, define 37 rays in $\mathbb{C}^3$ (core rays z, M_{ak}, P_k, Q_k and unique cross-product completion rays A_k, B_{ak}, C_{ak}, numbered $1, \ldots, 37$ via $z \leftrightarrow 1, M_{ak} \leftrightarrow 2{+}3a{+}k, P_k \leftrightarrow 11{+}k, Q_k \leftrightarrow 14{+}k, A_k \leftrightarrow 17{+}k, B_{ak} \leftrightarrow 20{+}3a{+}k, C_{ak} \leftrightarrow$




FIG. 3. Faithful real four-dimensional gadgets in $\mathbb{R}^4$. Panels (a),(c),(e) are common-apex lifts ($g(H) = 2$); panels (b),(d),(f) are linear ($g(H) = 3$). (a),(b) Purely chromatic ternary exclusions; (c),(d) ordinary TIFS; (e) purely chromatic binary TIFS $c(14) \neq c(18)$; (f) linear color identity $11 \sim_c 20$ from directed ordinary TITS $s(11) = 1 \implies s(20) = 1$.

$29 + 3a + k$):

$$\begin{aligned} z &= (1,0,0), & M_{ak} &= (1, \zeta^{a+k}, \zeta^{a+2k}), \\ P_k &= (0,1,-\zeta^k), & Q_k &= (-\zeta^k, 0, 1), \\ A_k &= (0,1,\zeta^k), & B_{ak} &= (-2, \zeta^{a+k}, \zeta^{a+2k}), \\ C_{ak} &= (1, -2\zeta^{2a+k}, \zeta^{2k}), \end{aligned} \tag{19}$$

and the 24 complete orthogonal triad contexts (Fig. 4):

$$\begin{aligned} \mathcal{M}_a &= \{M_{a0}, M_{a1}, M_{a2}\}, & \mathcal{A}_k &= \{z, P_k, A_k\}, \\ \mathcal{B}_{ak} &= \{M_{ak}, P_k, B_{ak}\}, & \mathcal{C}_{ak} &= \{M_{a,a+k}, Q_k, C_{ak}\}. \end{aligned} \tag{20}$$

Theorem 19 (Properties of H_{37})**.** *The hypergraph H_{37} defined by Eqs. (19)–(20) satisfies:*

1. *It is linear with Berge girth five,* 72 *edges, and $\omega(\Gamma(H_{37})) = 3$; its triangles are precisely its* 24 *contexts.*

2. *Equation (19) defines an injective edge-faithful representation in $\mathbb{C}^3$, whereas H_{37} admits* no *injective orthogonal representation in $\mathbb{R}^3$.*

3. *Its chromatic numbers are $\chi(H_{37}) = 4 > 3 = \chi_f(\Gamma(H_{37})) = \omega(\Gamma(H_{37}))$.*

4. *Its* 24 *Boolean blocks paste into a* 76*-event orthomodular lattice $L(H_{37}) = \{0,1\} \cup \{a_i, a_i^{\perp} : 1 \le i \le 37\}$. Its full state space has $|\mathcal{S}(H_{37})| = 153$ states and is order determining on $L(H_{37})$, with exact minimum state-family sizes* 6 *for unital vertex separation,* 7 *for separation of all* 76 *events, and* 18 *for order determination.*




FIG. 4. Complete 3-uniform hypergraph diagram of H_{37} (37 rays, 24 orthogonal triads, 72 edges in $\mathbb{C}^3$). Bold circles denote e_1, e_2, e_3 $(\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2)$; the other 21 smooth traces join core pairs through their unique completion rays $17, \ldots, 37$. Concentric rings record incident context colors.

Proof. (1)–(2) Zero patterns, magnitudes, and ζ-phases distinguish all 37 rays. Algebraic inner products confirm that $\langle v_i, v_j \rangle = 0$ holds strictly for the $24\binom{3}{2} = 72$ context pairs: $\langle M_{ak}, M_{c\ell} \rangle = 1 + \zeta^{c-a}(\zeta^{\ell-k} + \zeta^{2(\ell-k)}) = 0 \iff (c = a, \ell \neq k)$; $P_j \perp M_{ak} \iff j = k$; $Q_j \perp M_{ak} \iff k = a + j$; $\langle M_{c\ell}, B_{ak} \rangle = -2 + \zeta^{a-c}(\zeta^{k-\ell} + \zeta^{2(k-\ell)}) = 0 \iff (c, \ell) = (a, k)$; $\langle M_{c\ell}, C_{ak} \rangle = 1 - 2\zeta^{2a+k-c-\ell} + \zeta^{2k-c-2\ell} = 0 \iff (c = a, \ell = a + k)$; and unequal term magnitudes exclude orthogonalities among A, B, C. Uniqueness of cross-product completions in $\mathbb{C}^3$ excludes Berge 3- and 4-cycles, while $(\mathcal{A}_0, \mathcal{B}_{00}, \mathcal{M}_0, \mathcal{B}_{01}, \mathcal{A}_1)$ is a Berge 5-cycle.

To exclude any injective representation in $\mathbb{R}^3$, suppose unit $z \in \mathbb{R}^3$ and real $p_k, m_{ak} \in \mathbb{R}^3$ realize $\mathcal{A}_k, \mathcal{B}_{ak}, \mathcal{M}_a$. If $m_{ak} \cdot z = 0$, then $\{z, p_k, m_{ak}\}$ is an orthogonal basis of $\mathbb{R}^3$, and for $\ell \neq k$, $p_\ell \perp z$ and $m_{a\ell} \perp m_{ak}$ with $p_\ell \perp m_{a\ell}$ force $[p_\ell] = [m_{ak}]$ or $[m_{a\ell}] = [z]$. Hence $m_{ak} \cdot z \neq 0$. Rescaling so $m_{ak} \cdot z = 1$ and choosing unit $u_k \in z^\perp \cap p_k^\perp$ gives $m_{ak} = z + t_{ak} u_k$ $(t_{ak} \in \mathbb{R} \backslash \{0\})$. Orthogonality of $\mathcal{M}_a$ yields $1 + t_{ak} t_{a\ell}(u_k \cdot u_\ell) = 0$ $(k \neq \ell)$. Setting $q_k = t_{ak}/t_{0k}$ gives $q_0 q_1 = q_0 q_2 = q_1 q_2 = 1$, whose only real solutions are $(q_0, q_1, q_2) = \pm(1, 1, 1)$. Thus at most two distinct bases $\mathcal{M}_a$ exist in $\mathbb{R}^3$, contradicting injectivity on the three bases $\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_2$. The argument uses only the 13 rays z, P_k, M_{ak} and the orthogonalities among them, so already the 13-vertex subgraph of $\Gamma(H_{37})$ induced by these rays has no injective orthogonal representation in $\mathbb{R}^3$.

(3) In any 3-coloring with $c(z) = 0$, $P_0, P_1, P_2 \in \{1, 2\}$ cannot be monochromatic (as

$\mathcal{M}_a$ uses all three colors and $M_{ak} \sim P_k$). Up to symmetry $(c(P_0), c(P_1), c(P_2)) = (2, 1, 1)$, forcing $c(M_{a0}) = 1$ and $(c(M_{a1}), c(M_{a2})) = (x_a, 2 - x_a)$ with $x_a \in \{0, 2\}$. Since $N(Q_k) = \{M_{0k}, M_{1,k+1}, M_{2,k+2}\}$ avoids a color, $x_1 = 2 - x_2$, $x_0 = 2 - x_1$, $x_2 = 2 - x_0$, impossible on a 3-cycle. Thus $\chi(H_{37}) = 4$ (attained by $c(z) = 0, c(M_{ak}) = k, c(P_k) = c(Q_k) = 3$). The 6-state double cover $\mathcal{R}_6 = \{s_2, s_{30}, s_{73}, s_{99}, s_{109}, s_{139}\}$ covers every vertex twice, giving $\chi_f(\Gamma(H_{37})) = 3$.

(4) Each state is fixed by $t = (t_0, t_1, t_2) \in \mathbb{Z}_3^3$ with $s(M_{ak}) = \mathbf{1}_{k=t_a}$, $s(z) = \varepsilon \in \{0, 1\}$, and independent bits on $U(t) = \mathbb{Z}_3 \setminus \{t_0, t_1, t_2\}$ (for $\varepsilon = 0$) and $W(t) = \mathbb{Z}_3 \setminus \{t_a - a\}$, yielding $(1 + 2^{|U(t)|})2^{|W(t)|}$ states per t:

Class of $t \in \mathbb{Z}_3^3$	$\lvert\#t\rvert$	$(\lvert U(t)\rvert, \lvert W(t)\rvert)$	$(1 + 2^{\lvert U\rvert})2^{\lvert W\rvert}$
(c, c, c)	3	$(2, 0)$	$5 \times 1 = 5$
$(c, c + 1, c + 2)$	3	$(0, 2)$	$2 \times 4 = 8$
$(c, c + 2, c + 1)$	3	$(0, 0)$	$2 \times 1 = 2$
Two equal entries	18	$(1, 1)$	$3 \times 2 = 6$

(21)

and hence $|\mathcal{S}(H_{37})| = 3 \cdot 5 + 3 \cdot 8 + 3 \cdot 2 + 18 \cdot 6 = 153$. Since $2^5 - 1 = 31 < 37$ and $2^6 = 64 < 76$, the explicit families Ω_6 and Ω_7 given in the supplement prove the minima 6 and 7. For order determination ($a_i \le a_j^\perp \iff i \sim j$), the 15 nonorthogonal pairs in $D = \{P_0, P_1, P_2, Q_0, Q_1, Q_2\}$ require 15 disjoint joint-one witnesses in $T \subset \mathcal{S}(H_{37})$ ($|T| = 27$), and the 9 pairs $\{z, C_{ak}\}$ require at least 2 states outside T. Exhaustive bit-set verification over all $4{,}096 \times 132 = 540{,}672$ candidate 17-state families shows each meets at most 75 of the 81 minimal witness sets W_{xy}, whereas the 18-state family Ω_{18} in the supplement meets all 81 (Sec. S3 of the supplementary material). □

Theorem 19(2) is not the first finite separation of complex from real three-dimensional orthogonality: Harding and Salinas Schmeis[34] (Thm. 2.6) construct, from mutually unbiased bases, a two-gadget configuration that is realizable in $\mathbb{C}^3$ but not in $\mathbb{R}^3$ (48 rays, 32 bases, 189 context states, and 180 geometric states). H_{37} is smaller, linear of Berge girth five, and edge-faithful in $\mathbb{C}^3$, and its real obstruction already resides in the 13-ray subgraph on z, P_k, M_{ak}.

By Proposition 18, appending $d - 3$ common apices to H_{37} in $\mathbb{C}^d$ ($d \ge 3$) gives faithful d-uniform hypergraphs with $|V| = d + 34$, $|E| = 24$, $\omega = d$, $\chi = d + 1$, and $|\mathcal{S}| = d + 150$.

An order-determining family represents the specified 76-event order, orthocomplements, and Boolean operations within listed contexts. It does not preserve arbitrary meets as set intersections: nonorthogonal atoms have meet zero although some states make both true. It is therefore not a Boolean-sublattice embedding of the full lattice, nor a claim of closure under every Hilbert operation. In particular, the minimum six-state separating family and the six-state uniform double cover solve different problems.

B. Hyperplane obstruction, two-plane completion lemma, and canonical linear H_{65} in $\mathbb{R}^4$

Appending a single apex $v_{26} = (0, 0, 0, 1)^T \in \mathbb{R}^4$ to H_{YO}^+ gives the 26-ray, 16-context non-linear suspension $H_{\mathrm{YO}}^{(4)}$ (Fig. 5) with $\omega = 4, \chi = 5$, and 25 unital separating states (Proposition 18), but $|e \cap e'| = 2$ ($g = 2$). Restricted to the 13 Yu–Oh rays and the apex, its orthogonality graph is the Yu–Oh graph joined to one common vertex, the example that Cabello, Kleinmann, and Budroni[14] (Thm. 1) used to show that $\chi_f > d$ is necessary but not sufficient for state-independent contextuality; H_{65} below is a linear, apex-free example of the same phenomenon.

Proposition 20 (Hyperplane obstruction to linearity in $\mathbb{F}^4$)**.** *In any injective realization in $\mathbb{P}(\mathbb{F}^4)$, if two contexts e, e' each have three rays in a common 3-dimensional hyperplane $W = z^\perp$, both must contain $[z]$; hence if $|e_W \cap e'_W| \ge 1$, then $|e \cap e'| \ge 2$, violating linearity.*




FIG. 5. The completed 25-ray, 16-context Yu–Oh hypergraph H_{YO}^{+} in $\mathbb{R}^3$ ($\omega = 3, \chi = 4$) and its non-linear 26-ray, 16-context apex suspension $H_{\mathrm{YO}}^{(4)}$ in $\mathbb{R}^4$ ($\omega = 4, \chi = 5, g = 2$) obtained by adjoining the common orthogonal apex $v_{26} = (0, 0, 0, 1)^T$ to every context.

Proof. Three mutually orthogonal rays in W span W, forcing the fourth orthogonal ray of both e and e' to equal $[z] = W^{\perp}$. □

Lemma 21 (Simultaneous two-plane completion in $\mathbb{R}^4$)**.** *Let $V_{\mathrm{core}} \subset \mathbb{Q}^4 \setminus \{0\}$ be a finite set of non-collinear rational vectors with 4-element orthogonal base contexts E_{base} ($|C \cap C'| \leq 1$) and let all remaining orthogonal pairs be the bridge pairs $E_{\mathrm{bridge}} = \{e_1, \ldots, e_M\}$ ($e_m = \{u_m, v_m\}$, $L_m = \mathrm{span}(v_{u_m}, v_{v_m})$, $P_m = L_m^{\perp}$) such that every core vertex lies in a base context or is a bridge endpoint.*

1. *The pairs e_m can be simultaneously completed to 4-bases $E_m = \{u_m, v_m, w_{m,1}, w_{m,2}\}$ by fresh integer vectors $w_{m,1}, w_{m,2} \in P_m \cap \mathbb{Z}^4$ forming a linear edge-faithful hypergraph H if and only if*

$$\begin{aligned} v_z &\notin L_m \quad (z \in V_{\mathrm{core}} \setminus e_m), \\ P_\ell &\neq L_m \quad (\ell \neq m). \end{aligned} \tag{22}$$

2. *$\omega(\Gamma(H)) = 4$ and $\chi(H) = \max\{4, \chi(G_{\mathrm{core}})\}$; here G_{core} is automatically K_5-free, since no five nonzero vectors of $\mathbb{R}^4$ are mutually orthogonal.*

3. *If all G_{core}-independent base supports $S_1, \ldots, S_N$ form a unital separating family on the core, and the zero-profiles $Z(e_m) = \{\alpha : S_\alpha \cap e_m = \varnothing\}$ are nonempty and pairwise distinct, the $\leq 2N$ canonical extensions $s_{\alpha,\varepsilon}$ ($\varepsilon \in \{0, 1\}$, setting $(s_{\alpha,\varepsilon}(w_{m,1}), s_{\alpha,\varepsilon}(w_{m,2})) = (1 - \varepsilon, \varepsilon)$ when $\alpha \in Z(e_m)$ and $(0, 0)$ otherwise) are unital and separating on $V(H)$.*

Proof. If $v_z \in L_m$ then $P_m \subset v_z^{\perp}$, and if $P_\ell = L_m$ then $P_\ell \perp P_m$, forcing extra orthogonalities. Conversely, given Eq. (22), an integer orthogonal basis $\{\eta_A, \eta_B\} \subset P_m \cap \mathbb{Z}^4$ defines the infinite family $w_{m,1}(a, b) = a\eta_A + b\eta_B$, $w_{m,2}(a, b) = -\|\eta_B\|^2 b\eta_A + \|\eta_A\|^2 a\eta_B$, avoiding inductively the finitely many 1-dimensional intersections $z^{\perp} \cap P_m$ and $\mathbb{R}z \cap P_m$, where z ranges over the core rays *and* the fresh rays placed at earlier steps, together with $P_m \cap L_r$ ($r > m$). The last condition keeps every fresh ray outside each later L_r, so that $z^{\perp} \cap P_r$ remains 1-dimensional for fresh z as well. Linearity, coloring extension, and distinctness of canonical vertex profiles follow from Proposition 7 and distinctness of $Z(e_m)$. □




FIG. 6. Canonical linear 4-uniform hypergraph H_{65} in $\mathbb{R}^4$ (65 rays, 29 contexts, 174 edges, $g = 3, \omega = 4, \chi = 5$). Thick colored curves denote the five core contexts $C_1, \ldots, C_5$; the 24 thinner lines denote the completed bridge contexts $E_1, \ldots, E_{24}$ carrying degree-one completion pairs $\{18, 19\}, \ldots, \{64, 65\}$.

To construct a *linear* 4-uniform hypergraph in $\mathbb{R}^4$, we take the 17-ray core $V_{\text{core}} = \{1, \ldots, 17\} \subset \{-1, 0, 1\}^4$, which is orthogonally equivalent to the 17-ray Peres restriction of Haynes *et al.*[19] (pp. 8–9) via the orthogonal matrix

$$T = \frac{1}{2}\begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & -1 & -1 & -1 \end{pmatrix}, \qquad T^T T = I_4. \tag{23}$$

The core consists of the coordinate basis C_1, an even-parity Hadamard basis C_2, three odd-parity hub rays $9, 10, 11$, and the three binary bridge pairs $B_1 = \{12, 13\}$, $B_2 = \{14, 15\}$, $B_3 = \{16, 17\}$. The remaining base contexts are $C_3 = \{9, 10, 12, 13\}$, $C_4 = \{9, 11, 14, 15\}$ and $C_5 = \{10, 11, 16, 17\}$. Complete coordinates and all compatible supports are in Tables S17–S18 and S81 of the supplementary material. Completing its 24 uncovered bridge edges $e_m \in (C_1 \cup C_2) \times (B_1 \cup B_2 \cup B_3)$ by fresh integer pairs $\{2m+16, 2m+17\} \subset \{18, \ldots, 65\}$ via Lemma 21 yields $H_{65} = (V_{65}, E_{29})$ (Fig. 6; Tables S17–S18 of the supplementary material):

$$\begin{aligned} E_{29} &= \{C_1, \ldots, C_5\} \cup \{E_m : 1 \le m \le 24\}, \\ E_m &= \{u_m, v_m, 2m + 16, 2m + 17\}. \end{aligned} \tag{24}$$

Theorem 22 (Properties of H_{65})**.** *The* 65*-ray,* 29*-context hypergraph* H_{65} *satisfies:*

1. *It is linear* ($g(H_{65}) = 3$) *and edge-faithful in* $\mathbb{R}^4$ (174 *edges) with* $\omega(\Gamma(H_{65})) = 4$*, and nonconformal (*13 *uncovered maximal triangles:* 12 *bridge triangles plus* $\{9, 10, 11\}$*).*

2. *It has* $\alpha(G_{\text{core}}) = 4$*,* $\chi_f(\Gamma(H_{65})) = 17/4 = 4.25 > 4$*, and* $\chi(H_{65}) = 5$*.*

TABLE III. Bridge neighbors in the two core bases. The complete 24 state supports are supplementary.

b	$N(b)\cap C_1$	$N(b)\cap C_2$
12	$\{2,3\}$	$\{5,8\}$
13	$\{1,4\}$	$\{6,7\}$
14	$\{2,4\}$	$\{6,8\}$
15	$\{1,3\}$	$\{5,7\}$
16	$\{3,4\}$	$\{7,8\}$
17	$\{1,2\}$	$\{5,6\}$

3. *Its core has* 24 *compatible state supports* $S_\alpha=\{i_1,i_2,h,b\}$ *(Table S81), each missing* 14 *bridges and yielding* $|\mathcal{S}(H_{65})|=24\times 2^{14}=393{,}216$ *states and* 48 *canonical separating states.*

4. *Every* $s\in\mathcal{S}(H_{65})$ *obeys* $s(9)+s(10)+s(11)=1$*, so adjoining the fourth hub ray* $q=(1,1,1,-1)^T$ *forces* $\widetilde{s}(q)=0$*; moreover,* $A_9=A_{16}\dot\cup A_{17}$ *on all of* $\mathcal{S}(H_{65})$*, forcing the false atom-order implication* $s(16)\le s(9)$ *and collapsing the* 1*-ray* $[v_9]=[v_{16}-v_{17}]$ *with the* 2*-plane* $\mathrm{span}(v_{16},v_{17})$*.*

Proof. (1)–(2) No K_4 in G_{core} contains a bridge edge $\{u_m,v_m\}$, since $u_m\in C_a$ and $v_m\in B_k$ have exactly one common neighbor in G_{core}, namely the second vertex of $N(v_m)\cap C_a$; this verifies $P_\ell\neq L_m$, and support-size / magnitude checks verify $v_z\notin L_m$ in Lemma 21. For any independent $I\subseteq V_{\mathrm{core}}$, $|I\cap C_1|\le 1$ and $|I\cap C_2|\le 1$. If $I\cap\{9,10,11\}\neq\varnothing$, $|I\cap(C_3\cup C_4\cup C_5)|\le 2$ so $|I|\le 4$; if $I\cap\{9,10,11\}=\varnothing$ and I picks $(b_1,b_2,b_3)\in B_1\times B_2\times B_3$, even-parity triples cover C_2 and odd-parity triples cover C_1 (Table III), so again $|I|\le 3{+}1=4$. Thus $\alpha(G_{\mathrm{core}})=4$, $\chi_f\ge 17/4$ (matched by weights $1/8$ on the 24 core states and $1/16$ on the 20 remaining independent 4-sets), and $\chi(H_{65})=5$. (3)–(4) By parity, no state can avoid $\{9,10,11\}$, giving $s(9)+s(10)+s(11)=1$, $3\times 2\times 2\times 2=24$ core supports (Table S81), $24-10=14$ missed bridges per support, and $A_9=A_{16}\dot\cup A_{17}$. □

VI. EVENT ORDER, QUANTUM PROBABILITIES, AND NONEXTENDABLE WEIGHT

A. Extendable state polytopes and nonextendable weight on $H_{17,10}$

For any context hypergraph H, the classical state polytope $\mathcal{P}(H)=\mathrm{conv}\,\mathcal{S}(H)$ and the extendable state polytope $\mathcal{P}_{\mathrm{ext}}(H)=\mathrm{conv}\,\mathcal{S}_{\mathrm{ext}}(H)$ satisfy $\mathcal{P}_{\mathrm{ext}}(H)\subseteq\mathcal{P}(H)$, with $p_u=p_v$ for all $p\in\mathcal{P}_{\mathrm{ext}}(H)$ whenever $u\sim_c v$ by Corollary 4. For $p\in\mathcal{P}(H)$, define its *minimum nonextendable weight*

$$\begin{aligned}\eta_H(p)=\min_{\mu}\quad & \sum_{s\notin\mathcal{S}_{\mathrm{ext}}}\mu_s,\\ \text{subject to}\quad & p=\sum_{s\in\mathcal{S}}\mu_s s,\\ & \mu_s\ge 0,\qquad \sum_s\mu_s=1.\end{aligned}\tag{25}$$

Proposition 23 (Nonextendable weight lower bound)**.** *For any chromatically inseparable pair* $u\sim_c v$ *in* H *and any* $p\in\mathcal{P}(H)$*,* $\eta_H(p)\ge|p_u-p_v|$*.*

Proof. Every $s\in\mathcal{S}_{\mathrm{ext}}(H)$ satisfies $s(u)-s(v)=0$, while every $s\notin\mathcal{S}_{\mathrm{ext}}(H)$ satisfies $|s(u)-s(v)|\le 1$, so $|p_u-p_v|\le\sum_{s\notin\mathcal{S}_{\mathrm{ext}}(H)}\mu_s|s(u)-s(v)|\le\sum_{s\notin\mathcal{S}_{\mathrm{ext}}(H)}\mu_s$. □

Theorem 24 (Quantum preparations in $\mathcal{P}(H_{17,10}) \setminus \mathcal{P}_{\text{ext}}(H_{17,10})$)**.** *In the real FOR of $H_{17,10}$ (Table S2 of the supplementary material):*

1. *The pure quantum preparation along $r_{24} = (1, -1, 2)^T$,*

$$\rho_\star = \frac{1}{6}\begin{pmatrix} 1 & -1 & 2 \\ -1 & 1 & -2 \\ 2 & -2 & 4 \end{pmatrix}, \qquad p_i = \mathrm{Tr}(\rho_\star P_i) = \frac{(r_{24} \cdot r_i)^2}{6\|r_i\|^2}, \tag{26}$$

belongs to $\mathcal{P}(H_{17,10})$ via the exact five-state convex combination (state indices in Table S40 of the supplementary material; see also Tables S46–S47)

$$p = \frac{1}{12}s_4 + \frac{5}{36}s_9 + \frac{1}{36}s_{10} + \frac{7}{12}s_{14} + \frac{1}{6}s_{17}, \tag{27}$$

yet satisfies $p_{24} = 1$ and $p_{10} = 8/9$ for the chromatically inseparable pair $(10, 24)$, so

$$p \notin \mathcal{P}_{\text{ext}}(H_{17,10}) \qquad \text{and} \qquad \eta_{H_{17,10}}(p) = \frac{1}{9}. \tag{28}$$

2. *The pure preparation along the coordinate ray $z_3 = r_5 = (0, 0, 1)^T$, with Born vector $p'_i = (r_5 \cdot r_i)^2/\|r_i\|^2$, has $p'_{24} = p'_{18} = 2/3$ and $p'_{10} = p'_{11} = 1/3$, belongs to $\mathcal{P}(H_{17,10})$ via the four-state convex combination $p' = \frac{1}{3}s_3 + \frac{1}{6}s_4 + \frac{1}{3}s_{14} + \frac{1}{6}s_{15}$, and satisfies $\eta_{H_{17,10}}(p') = 1/3$.*

3. *For every density operator ρ and both inseparable pairs $(u, v) \in \{(10, 24), (11, 18)\}$, $|p_u(\rho) - p_v(\rho)| \le \|P_u - P_v\| = 1/3$. Hence Proposition 23 certifies at most $1/3$ on $H_{17,10}$, and the preparation in (2) attains this value.*

Proof. (1) Direct evaluation of Eq. (27) on all 17 vertices matches p_i in Eq. (26) (Table S46), with $r_{10} = (1, -1, 1)^T$ giving $p_{10} = 4^2/(6 \cdot 3) = 8/9$. Since $10 \sim_c 24$, Proposition 23 gives $\eta_{H_{17,10}}(p) \ge |1 - 8/9| = 1/9$, which is attained in Eq. (27) because among $\{s_4, s_9, s_{10}, s_{14}, s_{17}\}$ precisely s_4 and s_{10} are nonextendable, with total weight $1/12 + 1/36 = 1/9$. (2) Direct evaluation of the four-state mixture on all 17 vertices reproduces p' (Tables S46–S47). Proposition 23 gives $\eta_{H_{17,10}}(p') \ge |2/3 - 1/3| = 1/3$, attained because s_3, s_{14} are extendable and s_4, s_{15} are not, with total weight $1/6 + 1/6 = 1/3$. (3) For unit vectors with squared overlap κ, $P_u - P_v$ has eigenvalues $\pm\sqrt{1 - \kappa}$ and 0. Both pairs have $\kappa = (r_u \cdot r_v)^2/(\|r_u\|^2\|r_v\|^2) = 16/18 = 8/9$, so $|\,\mathrm{Tr}\,\rho(P_u - P_v)| \le \|P_u - P_v\| = 1/3$. □

The exact support-and-weight certificates and the complete Born probability vectors are given in the supplement. These tables are regenerated from the checked state inventory, rather than manually transcribed.

These preparations exclude only models built from noncontextual eigenvalue assignments: their ordinary-state models are exhibited explicitly. A state is extendable exactly when some noncontextual eigenvalue assignment c makes it the indicator of a single eigenvalue λ_a in every context observable A_e of Eq. (1). For a model built from noncontextual eigenvalue assignments, every chromatic identity $u \sim_c v$ entails the equality $p_u = p_v$; quantum mechanics violates it, and Proposition 23 converts the violation into a lower bound on η. The weight η therefore measures how far the quantum predictions force any noncontextual truth-assignment model away from noncontextual eigenvalues: on $H_{17,10}$, at least one third of the weight in the preparation along z_3. Since $\mathcal{P}_{\text{ext}}(H_{17,10})$ lies in the hyperplane $p_{10} = p_{24}$, a Born vector in $\mathcal{P}(H_{17,10})$ generically lies outside $\mathcal{P}_{\text{ext}}(H_{17,10})$; the content of Theorem 24 is the size of the required weight. Conversely, averaging the d color classes of any admissible coloring gives the maximally mixed Born vector $1/d$, so the restricted obstruction is preparation dependent. The $H_{17,10}$ ordinary implication $s(10) \le s(24)$ also prevents order

determination for distinct ray atoms; vertex separation and order representation must not be conflated.

For a single system, nothing forces a classical model to use only extendable states. The supersinglet game of Sec. I does. A local hidden-variable model that reproduces its perfect correlations gives the d parties the d color classes of a strong d-coloring, so each party follows an extendable state. If $d-1$ parties measure the same basis and obtain all of its rays but v, the last particle is prepared along v, and in such a model its outcome probabilities lie in $\mathcal{P}_{\mathrm{ext}}(H)$. On $H_{17,10}$ the two preparations of Theorem 24 arise in this way from the bases $\{4,11,24\}$ and $\{2,8,5\}$. Both lie outside $\mathcal{P}_{\mathrm{ext}}(H_{17,10})$, so quantum mechanics contradicts every such local model, although $\chi(H_{17,10})=3$. More directly, if $u \sim_c v$, such a model never lets two different parties obtain u and v. Quantum mechanics gives this event the probability $(1-\kappa)/[d(d-1)]$, where κ is the squared overlap of the two rays, because the reduced state of two particles of the supersinglet is the normalized projector onto the antisymmetric subspace. For the pairs $(10,24)$ and $(11,18)$ of $H_{17,10}$ this probability is $1/54$.

B. Order determination versus Born probabilities on H_{37}, $H_{\mathrm{YO}}^{(4)}$, and H_{65}

When $\chi(H) > d$ (Regime I), no noncontextual eigenvalue assignment exists, $\mathcal{P}_{\mathrm{ext}}(H)=\varnothing$, and the relevant probability question is whether Born assignments $p_i(\rho)=\mathrm{Tr}(\rho\Pi_i)$ lie in $\mathcal{P}(H)=\mathrm{conv}\,\mathcal{S}(H)$ (Table IV).

Corollary 25 (Order determination does not imply classical probability models)**.** *The* 76*-event orthomodular lattice $L(H_{37})$ admits an order-determining state space $\mathcal{S}(H_{37})$ and a classical mixture for the maximally mixed state $I_3/3$, yet every $p \in \mathcal{P}(H_{37})$ satisfies the tight inequality*

$$\mathcal{C}_{37}(p) = \sum_{k=0}^{2} \big[p(P_k) + p(Q_k)\big] \le 2, \tag{29}$$

whereas the complex FOR of H_{37} satisfies

$$\sum_{k=0}^{2} (\Pi_{P_k} + \Pi_{Q_k}) = \mathrm{diag}(3/2, 3/2, 3) = \frac{3}{2} I_3 + \frac{3}{2} |e_3\rangle\langle e_3|, \tag{30}$$

yielding $\mathcal{C}_{37}(\rho) = \frac{3}{2}(1+\rho_{33}) > 2$ whenever $\rho_{33} > 1/3$ (and $\mathcal{C}_{37}(|e_3\rangle\langle e_3|) = 3$).

Proof. In the state parametrization of Theorem 19(4), every $t \in \mathbb{Z}_3^3$ has $|U(t)| + |W(t)| \le 2$, establishing Eq. (29), while summing the normalized projectors of P_k, Q_k in Eq. (19) using $1+\zeta+\zeta^2=0$ gives Eq. (30). Uniform weights $1/6$ on $\mathcal{R}_6$ reproduce $p_i(I_3/3) = 1/3$. □

In dimension four, the non-linear suspension $H_{\mathrm{YO}}^{(4)}$ obeys $\mathcal{Y}(p) = p_{26} + \sum_{i=10}^{13} p_i \le 1$ on $\mathcal{P}(H_{\mathrm{YO}}^{(4)})$, whereas $\sum_{i=10}^{13} \Pi_i = \frac{4}{3}(I_4 - \Pi_{26})$ gives $\mathcal{Y}(\rho) = 1 + (1 - p_{26}(\rho))/3 > 1$ for every density operator $\rho \neq \Pi_{26}$[14]. For the canonical linear H_{65}, every $s \in \mathcal{S}(H_{65})$ has 4 active core vertices, so every $p \in \mathcal{P}(H_{65})$ obeys $\sum_{i=1}^{17} p_i = 4$. However, with $q = (1,1,1,-1)^T$ and $Q = qq^T/4$, direct summation gives

$$\sum_{i=1}^{17} \Pi_i = 4I_4 + Q \quad \Longrightarrow \quad \sum_{i=1}^{17} p_i(\rho) = 4 + \mathrm{Tr}(\rho Q), \tag{31}$$

violating the classical bound 4 whenever $\mathrm{Tr}(\rho Q) > 0$ (including $I_4/4$, where the sum is $17/4 = \chi_f$, and $\rho = Q$, where it is 5). Nevertheless, on the orthogonal subspace $q^{\perp}$, the density operator $\rho_\star = (I_4 - Q)/3$ has core Born probabilities $1/4$ on $C_1 \cup C_2$, $1/3$ on $\{9,10,11\}$, and $1/6$ on $B_1 \cup B_2 \cup B_3$, and completion probabilities $r_{m,1} + r_{m,2} = 7/12$. Choosing $\alpha \in \{1,\ldots,24\}$ uniformly and $\lambda \in [0,1]$ uniformly and setting $\varepsilon_m = \mathbf{1}_{\{\lambda < \tau_m\}}$ with

TABLE IV. Event order and preparation-dependent probability models for the fixed realizations. Membership statements concern Born vectors $p(\rho)$; the restricted model is $\mathcal{P}_{\text{ext}}$.

Configuration	Order determination	An ordinary-state model	A failed model or witness
$H_{17,10}$	No: $s(10) \leq s(24)$ is a false ray inclusion.	P_{24} and P_{z_3} have ordinary mixtures.	Neither lies in $\mathcal{P}_{\text{ext}}$; $\eta = 1/9$ and $1/3$ (maximal pair certificate).
Completed Yu–Oh	No.	No Born preparation lies in $\mathcal{P}$.	Every preparation gives $\sum_{10}^{13} p_i = 4/3 > 1$.
H_{37}	Yes on its specified 76-event lattice; 18 states minimal.	$I_3/3$ has a six-state mixture.	$\rho_{33} > 1/3$ violates the P, Q bound.
$H_{\text{YO}}^{(4)}$	No.	Pure apex P_{26} has its singleton-state model.	Every $\rho \neq P_{26}$ violates the displayed bound.
H_{65}	No: $s(16) \leq s(9)$ is a false ray inclusion.	$(I_4 - Q)/3$ has a model for all 65 rays.	$\text{Tr}(\rho Q) > 0$ violates the core sum bound.

$\tau_m = \frac{12}{7} r_{m,2} \in \{4/9, 5/11, 6/11, 5/9\}$ on the 14 missed bridges of S_α gives $\mathbb{E}[s(w_{m,2})] = \frac{14}{24}\tau_m = r_{m,2}$, expressing $p(\rho_\star) \in \mathcal{P}(H_{65})$ as a convex mixture of at most 120 states. Rotating V_{core} by $D_0 = I_4, D_1 = \text{diag}(1,1,-1,-1), D_2 = \text{diag}(1,-1,1,-1), D_3 = \text{diag}(1,-1,-1,1)$ gives a fixed enlarged family with

$$\sum_{a,i} \Pi_i^{(a)} = 17 I_4, \qquad \sum_{a,i} p_i^{(a)} \leq 16 \text{ classically.} \tag{32}$$

Each copy admits at most four true rays under geometric pair exclusions; repeated projectors are counted with multiplicity. The 24 distinct rays form the Peres set[35]. This is an enlargement, not a state-independent test of the original 65 rays.

VII. KS ANCESTRY, SPECIFIED PARENTS, AND ABSTRACT VERSUS GEOMETRIC STATES

A. The KS ancestry hypothesis and abstract transverse extensions

Motivated by the recurring parent structures of our examples, we formulate the *KS ancestry hypothesis*: informally, every noncolorable ($\chi(H) > d$) configuration has a larger KS ancestor. For realized ray configurations, its existence form follows from an elementary union argument, even without the chromatic premise; the substantive open question is whether the retained rays can be made essential to the contradiction (Sec. VIII):

Proposition 26 (Same-dimensional finite KS ancestry)**.** *Every finite ray set $R \subset \mathbb{P}(\mathbb{F}^d)$ ($\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$, $d \geq 3$) is a proper subset of a finite KS ray set $P \subset \mathbb{P}(\mathbb{F}^d)$ whose orthogonality graph and full-basis hypergraph induce those on R, with every ray of P in a complete context. Consequently, a finite graph G has an injective edge-faithful realization in $\mathbb{F}^d$ if and only if G is an induced subgraph of a finite KS orthogonality graph in $\mathbb{F}^d$.*

Proof. Choose a finite full-basis KS set $K \subset \mathbb{P}(\mathbb{F}^d)$[4,36], orient K by $U \in \text{U}(d)$ (or $\text{O}(d)$) so $UK \not\subseteq R$, and complete any uncovered rays of $R \cup UK$ to bases to obtain P. Any state on P would restrict to a state on UK, a contradiction, and keeping the rays of R fixed preserves all induced orthogonalities and full bases on R. □

By Theorem 11, H_8 has no injective realization, and H_9, G_{32} have no ray assignment; none can embed into any 3-dimensional geometric KS parent. For abstract hypergraphs, Lemma 3 yields a direct chromatic-to-stateless construction:

Proposition 27 (Transverse chromatic-to-stateless extension)**.** *For any finite d-uniform hypergraph $H = (V, E)$ ($d \geq 3$), let $T(H)$ take $d-1$ disjoint copies $H^{(1)}, \ldots, H^{(d-1)}$ and adjoin for each $v \in V$ a fresh vertex w_v and the transverse context $T_v = \{v^{(1)}, \ldots, v^{(d-1)}, w_v\}$. Then $T(H)$ has $d|V|$ vertices, $(d-1)|E| + |V|$ contexts, is linear if H is linear, and its two-valued states correspond bijectively to labeled strong d-colorings of H; in particular,*

$$\mathcal{S}(T(H)) = \varnothing \quad \Longleftrightarrow \quad \chi(H) > d. \tag{33}$$

Proof. A state $s \in \mathcal{S}(T(H))$ restricts to $d-1$ states on $H^{(k)}$ that are pairwise disjoint by T_v (forcing $s(w_v) = 1 - \sum_{k=1}^{d-1} s(v^{(k)})$), which by Lemma 3 is equivalent to a strong d-coloring of H. □

In dimension $d = 3$, any edge $u \sim v$ of H creates a Berge 4-loop $(u^{(1)}, v^{(1)}, v^{(2)}, u^{(2)})$ in $T(H)$, excluding an injective realization in $\mathbb{C}^3$ by the common orthogonal-complement argument; thus $T(H)$ is an abstract construction for a single system. It describes, however, the supersinglet game of Ref.[8] (Sec. I): the copies $H^{(k)}$ are the bases measured by $d-1$ of the parties, the transverse context T_v states that exactly one party obtains v, and the fresh vertex w_v stands for the last party, whose truth assignment is fixed by Lemma 3. Equation (33) is thus the combinatorial content of the statement that this game has no local hidden-variable model exactly when $\chi(H) > d$. For H_9, $T(H_9)$ (45 vertices, 33 contexts) is context-critical for statelessness on all 33 contexts; for H_{YO}^{+}, $T(H_{\mathrm{YO}}^{+})$ (75 vertices, 57 contexts) is critical on all 32 horizontal contexts and 12 transverse contexts $v \in \{14, \ldots, 25\}$, but remains stateless upon deleting T_v for $v \in \{1, \ldots, 13\}$ (Sec. S3 of the supplementary material).

B. Explicit inclusions in known KS parents and Cabello completions

Beyond unrestricted existence, the following explicit unions use known KS components; the unions are not asserted to be previously named, smallest, or critical parents (explicit coordinate dictionaries and certificates in Secs. S1–S3 of the supplementary material):

1. *Bub's 49-ray context-critical real parent of H_{YO}^{+}:*

   Proposition 28 (Context-critical real parent of H_{YO}^{+})**.** *Adjoining to H_{YO}^{+} the eight rays $(0, 1, \pm 2)$, $(0, 2, \pm 1)$, $(1, \pm 2, 0)$, $(2, \pm 1, 0)$ yields 33 rays with 20 bases and 16 uncovered orthogonal pairs that eliminate all 24 states of H_{YO}^{+}. Completing those 16 pairs yields the 49-ray, 36-context full-context Schütte–Bub KS configuration[20,37], containing H_{YO}^{+} as an induced restriction, in which every one of the 36 contexts is critical for $\mathcal{S}_{\mathrm{ctx}} = \varnothing$ (Table S83 of the supplementary material). If the orthogonalities of a deleted context are retained as geometric exclusions, only the 20 bases of the 33-ray stage remain critical; deleting any of the 16 completion contexts leaves $\mathcal{S}_{\mathrm{geom}} = \varnothing$.*

2. *MUB-inventory parents of H_{37}:* In the 165-ray, 130-context MUB inventory[18] formed by three 69-ray, 50-context KS lineages L_0, L_1, L_2 (where each L_a completes the 36 uncovered pairs of Cabello's 33-ray, 14-basis KS set[38]; Pavičić[39] points out that this 33-ray set already occurs as a subset of a 69-ray, 50-basis set in Ref.[40]), H_{37} embeds as an induced restriction via the seed-index matrix $j(a,k) = \left(\begin{smallmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 9 & 8 \end{smallmatrix}\right)$ mapping $z \leftrightarrow a_{1,1}$, $M_{ak} \leftrightarrow u_{j(a,k)}$, $P_k \leftrightarrow c_{1,k+1}$, $Q_k \leftrightarrow c_{2,k+1}$, $A_k \leftrightarrow b_{1,k+1}$, $B_{ak} \leftrightarrow d_{1,j(a,k)}$, $C_{ak} \leftrightarrow d_{2,j(a,a+k)}$. It lies in the 57-ray, 40-context intermediate spoke hypergraph (297 states) and in each 87-ray, 64-context KS union $H_{37} \cup L_a$.

3. *Four-dimensional Peres and Cabello parents of $H_{\mathrm{YO}}^{(4)}$ and H_{65}:* Taking the union with Peres's 24-ray set P_{24}[35] gives a 40-ray, 36-context KS parent of $H_{\mathrm{YO}}^{(4)}$ ($|H_{\mathrm{YO}}^{(4)} \cap P_{24}| = 10$; note that a single-apex lift of a 3-dimensional KS set always retains the unique apex-one state) and a 72-ray, 48-context KS parent of H_{65} ($|H_{65} \cap P_{24}| = 17$). More economically:

Proposition 29 (Three-ray KS extensions of H_{65}). *Adjoining* $a = (1,0,0,1), b = (0,1,0,1), c = (0,0,1,1)$ *to* H_{65} *yields a* 68*-ray,* 35*-basis KS parent containing* H_{65} *as an induced restriction and Cabello's* 18*-ray,* 9*-context parity proof* $C_{18,9}$[21] *(*C_3, C_4, C_5 *plus* $D_1 = \{2,3,12,a\}, D_2 = \{1,3,15,b\}, D_3 = \{1,2,17,c\}, D_4 = \{6,7,13,a\}, D_5 = \{6,8,14,b\}, D_6 = \{7,8,16,c\}$*). Adjoining* $(0,1,-1,0)$, $(1,0,-1,0)$, $(1,-1,0,0)$ *instead yields a* 68*-ray,* 35*-basis KS parent containing a ray-critical* 20*-ray,* 11*-context parity proof* $C_{20,11}$ *(*$11 = 4s(4)+4s(5)+2\sum_{i\neq 4,5} s(i)$*). All* 20 *rays of* $C_{20,11}$ *belong to Peres's* 24*-ray set, and* $C_{20,11}$ *is one of its* 96 *critical* 20*-ray KS subsets, which are mutually isomorphic and include Kernaghan's set*[21,22].

Proof. In $C_{18,9}$, summing s over C_3, C_4, C_5 gives $3 = 2(s(9)+s(10)+s(11))+\sum_{k=12}^{17} s(k)$, so $s(12)+\cdots+s(17)$ is odd, whereas $D_1 \oplus D_4, D_2 \oplus D_5, D_3 \oplus D_6$ give $s(12)+s(13) \equiv s(2)+s(3)+s(6)+s(7)$, $s(14)+s(15) \equiv s(1)+s(3)+s(6)+s(8)$, $s(16)+s(17) \equiv s(1)+s(2)+s(7)+s(8)$ (mod 2), summing to 0 (mod 2). In $C_{20,11}$ (comprising $C_1, \ldots, C_5$ and six contexts on $\{1,\ldots,8,12,\ldots,17\}$ with $(0,1,-1,0)$, $(1,0,-1,0)$, $(1,-1,0,0)$), vertices 4, 5 have degree 4 and the other 18 vertices have degree 2, so $11 = 4s(4)+4s(5)+2\sum_{i\neq 4,5} s(i)$, an odd-equals-even contradiction; explicit states after deleting any single ray are certified in Sec. S3 of the supplementary material. □

Both 68-ray presentations have 35 bases covering 192 of 195 geometric orthogonal pairs and context intersections of size up to two. They are not edge-faithful in the listed bases. The 72-ray Peres union covers all pairs; no size minimum or criticality of either full parent is asserted.

4. *Faithful completion* $C_{36,18}$ *and abstract versus geometric states on* $C_{45,21}$*:*

Corollary 30 (Faithful Cabello completion and vertex deletion). *The* 9 *uncovered orthogonal pairs of* $C_{18,9}$ *satisfy Eq.* (22)*, so Lemma 21 yields a linear edge-faithful* 36*-ray,* 18*-context KS hypergraph* $C_{36,18}$ *in* $\mathbb{R}^4$ *with* $\omega = 4, \chi = 5$*, and* $\mathcal{S} = \varnothing$*. Deleting one vertex (say* c*) and its two contexts from* $C_{18,9}$ *leaves* 17 *rays,* 7 *bases,* 14 *uncovered pairs, and* 12 *core states whose abstract completion* $C_{45,21}$ *has* $\chi = 5$ *and* $|\mathcal{S}_{\mathrm{ctx}}(C_{45,21})| = 3{,}584$ *unital separating context states (*24 *canonical). However,* $v_{16} \in \mathrm{span}(v_1, v_2)$ *and* $v_{17} \in \mathrm{span}(v_7, v_8)$ *violate Eq.* (22)*, forcing four extra orthogonalities that reduce the geometric state space to* $|\mathcal{S}_{\mathrm{geom}}(C_{45,21})| = 2{,}048$ *with* $s(9) = s(16) = s(17) = 0$*, destroying unitality and separation (Table V).*

Proof. For $C_{18,9}$, the 9 uncovered pairs are the 6 core bridges $\{1,13\}$, $\{2,14\}$, $\{3,16\}$, $\{6,17\}$, $\{7,15\}$, $\{8,12\}$ from H_{65} plus $\{11,a\}$, $\{10,b\}$, $\{9,c\}$. They satisfy Eq. (22) and are completed in Tables S28–S29 of the supplementary material. Deleting c, D_3, D_6 leaves 12 core supports: 10 miss 8 of the 14 uncovered pairs and 2 miss 9 (Table S84), giving $10 \cdot 2^8 + 2 \cdot 2^9 = 3{,}584$ context states. Since $v_{16} = v_1 + v_2$ and $v_{17} = \frac{1}{2}(v_7 - v_8)$, completing $\{1,2\}$ by $\{21,22\}$ and $\{7,8\}$ by $\{35,36\}$ forces $v_{16} \perp v_{21}, v_{22}$ and $v_{17} \perp v_{35}, v_{36}$ in the local labels of Tables S31–S32. If $s(16) = 1$, neither 1 nor 2 is selected in any admissible core support, so $s(21)+s(22) = 1$, contradicting geometric exclusion; the same argument gives $s(17) = 0$. The retained hub bases then force $s(9) = 0$. Exactly 8 core supports satisfy $s(9) = s(16) = s(17) = 0$, each missing 8 uncovered pairs. All their fresh-pair choices obey the four extra exclusions, yielding $8 \cdot 2^8 = 2{,}048$ geometric states. □

VIII. DEPENDENCIES, COUNTEREXAMPLES, AND REMAINING QUESTIONS

The common dependency structure is incidence → states and colorings, followed by independent tests of geometry, event order, and quantum probabilities. A chromatic argument can be made on an abstract hypergraph; its operational interpretation requires

TABLE V. Context and geometric state spaces in the four-dimensional examples. Linearity of the listed incidence structure does not imply edge-faithfulness.

Configuration	$\lvert V\rvert/\lvert E\rvert$	$\lvert\mathcal{S}_{\mathrm{ctx}}\rvert$	$\lvert\mathcal{S}_{\mathrm{geom}}\rvert$	Scope
$C_{18,9}$	18/9	0	0	Nine uncovered geometric pairs; parity KS proof.
$C_{36,18}$	36/18	0	0	Faithful linear completion.
$C_{45,21}$	45/21	3,584	2,048	Four additional pairs destroy geometric unitality and separation.
H_{65}	65/29	393,216	393,216	Edge-faithful; all context states respect the geometry.

TABLE VI. Logical boundaries of the integrated results. Each row prevents transfer of a conclusion between different assignment or representation problems.

Tempting unrestricted inference	Counterexample or necessary qualification
$\chi > d$ excludes states or ray realizations	Completed Yu–Oh has faithful real rays and 24 unital separating states.
$\chi > d$ makes I_d/d contextual[8]	One system needs $\chi_f > d$[15]: H_{37} has $\chi = 4$, $\chi_f = 3$, and a six-state mixture for $I_3/3$.
Ordinary states separate ⇒ colors separate	Faithful $H_{17,10}$ and $H_{21,13}$ have forced color identities.
Colors separate ⇒ every state extends	The pentagon separates all vertex pairs but has one nonextendable state.
Vertex separation ⇒ event-order determination	$H_{17,10}$ has 10 ⇒ 24; H_{65} has 16 ⇒ 9 and ray–plane collapse.
Order determination ⇒ all Born mixtures	The specified event order of H_{37} is represented, while $\rho_{33} > 1/3$ violates the state inequality.
$\chi_f > d$ establishes fixed-measurement state independence	Ref.[14] (Thm. 1) (Yu–Oh plus apex); linear and apex-free: H_{65} admits the Born model of $(I_4 - Q)/3$ on all 65 rays.
Abstract stateless extension ⇒ geometric KS parent	Closing a color-identical pair changes orthogonality; transverse four-loops exclude injective triadic geometry.
Generic KS ancestry explains the original chromatic gap	Unrestricted extension holds even for colorable sets and need not use the old rays essentially.

a ray realization. Conversely, a realization does not select an ontic state family or a classical probability mixture. The exact-palette assumption, that the outcomes admit a noncontextual assignment of d eigenvalues, adds a precise completion test at the level of whole truth assignments: a truth assignment is compatible with noncontextual eigenvalues exactly when the outcomes it leaves false can carry the remaining $d - 1$ eigenvalues.

Table VI also explains why the probability and gadget examples carry different force. A one-way ordinary TITS already gives a symmetric color identity; a two-way purely chromatic identity requires counterstates in both truth directions. A ternary color exclusion may survive even when every pair can be equal. A strict inclusion $\mathcal{P}_{\mathrm{ext}} \subsetneq \mathcal{P}$ therefore need not be detectable through pair equalities alone.

Open problems concern global minimum faithful examples beyond the specified inventory, economical rational completions with fewer degree-one rays, residual-coloring certificates in dimensions above three, and geometric ancestry in which the retained rays are essential to the contradiction. The current statements establish existence and the recorded finite classifications, without asserting these stronger minima or essential-ancestry properties. All complete enumerations and the detailed verification scope remain in the supplement and electronic records.

IX. CONCLUSION

We started from a simple physical requirement on intertwined complete experiments: that each is the measurement of a maximal, nondegenerate observable with one common spectrum of d eigenvalues, and that the eigenvalue carried by an outcome is an element of physical reality in the sense of Einstein, Podolsky, and Rosen[3], the same in every experiment in which the outcome occurs. This requirement of noncontextual eigenvalues is exactly strong d-colorability of the context hypergraph. It asks for more than Kochen–Specker noncontextuality: not one classical truth assignment, but d of them with disjoint supports covering every outcome. Its failure, $\chi(H) > d$, is therefore a weaker but more widespread form of contextuality. Completed Yu–Oh has 24 truth assignments separating all of its rays, yet its outcomes cannot carry three eigenvalues noncontextually. These truth assignments also give a classical realization, which fails in the same way, so $\chi(H) > d$ restricts the spectra of maximal observables rather than truth values.

By unifying these common-spectrum chromatic obstructions ($\chi(H) > d$) with exact-palette chromatic separability constraints ($\chi(H) = d$, $\mathcal{S}_{\text{ext}}(H) \subsetneq \mathcal{S}(H)$), we have distinguished the requirements across noncontextual eigenvalue assignments, geometric ray realizations, orthomodular event representations, and quantum probability models. At the foundation, $\chi(H) > d$ excludes noncontextual eigenvalues for uniform maximal context observables, whereas the exclusion of an injective realization for H_8 and of every ray assignment for H_9, G_{32} requires direct Hermitian projection forcing, correcting the earlier spectral argument for G_{32}[6] and establishing four-quadrant independence even at Berge girth five. Within chromatically complete hypergraphs ($\chi(H) = d$), where noncontextual eigenvalue assignments exist, our completion criterion ($\chi(\Gamma(H)[V \setminus \operatorname{supp} s]) \leq d - 1$) characterizes which truth assignments arise from noncontextual eigenvalue assignments and explains the specified examples of chromatic identities ($u \sim_c v$) and purely chromatic TIFS/TITS/ternary constraints ($H_{17,10}, H_{21,13}, D_{16,9}, T_{19,11}, W^{(4)}_{16,6}$) and yields measurable nonextendable weights 1/9 and 1/3 for quantum ray preparations on $H_{17,10}$, the latter being the largest value a single chromatic identity can certify there. In higher complexity, the Hesse hypergraph H_{37} (6/7/18 OML minima) and our rational four-dimensional completion H_{65} (393,216 states) demonstrate that neither exact order determination nor a fractional chromatic gap decides classicality for all Born preparations, while the elementary KS ancestry result, the 68-ray KS parents of H_{65}, and the completions $C_{36,18}$ and $C_{45,21}$ of $C_{18,9}$ delineate unrestricted embedding from specified critical parents and geometric pair exclusions.

The common thread is that a classical account of quantum outcomes can succeed at the level of truth values and still fail at the level of eigenvalues. For the preparation along z_3, the quantum predictions on $H_{17,10}$ admit noncontextual truth-assignment models, but every such model must place at least one third of its weight on truth assignments that no noncontextual eigenvalue assignment produces.

SUPPLEMENTARY MATERIAL

See the supplementary material for the explicit vector and context inventories, the complete two-valued state and coloring enumerations, the odd-cycle and separation-profile certificates, and the supplementary proofs and verification reductions.

ACKNOWLEDGMENTS

I thank Mohammad Hadi Shekarriz for valuable comments and suggestions. This research was funded in part by the Austrian Science Fund (FWF), Grant DOI 10.55776/PIN5424624. The author acknowledges FWF and TU Wien Bibliothek for support through the Open Access Funding Programme. Generative AI tools were used for editorial work, including language editing and the LaTeX layout of figures and tables, and for independent checks of

finite computations. The author reviewed all such output and takes full responsibility for the content of this article.

AUTHOR DECLARATIONS

Conflict of Interest

The author has no conflicts to disclose.

DATA AVAILABILITY

The data that support the findings of this study are available within the article and its supplementary material. The verification programs and machine-readable certificates are available from the author upon reasonable request.

[1] J. von Neumann, Annals of Mathematics **32**, 191 (1931).
[2] P. R. Halmos, *Finite-Dimensional Vector Spaces*, Undergraduate Texts in Mathematics (Springer, New York, 1974).
[3] A. Einstein, B. Podolsky, and N. Rosen, Physical Review **47**, 777 (1935).
[4] S. Kochen and E. P. Specker, Journal of Mathematics and Mechanics **17**, 59 (1967).
[5] K. Svozil, Quantum Reports **2**, 278 (2020), arXiv:1808.00813.
[6] M. H. Shekarriz and K. Svozil, Journal of Mathematical Physics **63**, 032104 (2022), arXiv:2105.08520.
[7] K. Svozil, Entropy **27**, 387 (2025), arXiv:2501.15261.
[8] A. Cabello, "State-independent quantum contextuality and maximum nonlocality," (2012), version 1 (2011): "Simple unified proof of state-independent contextuality", arXiv:1112.5149 [quant-ph].
[9] R. J. Greechie, Journal of Combinatorial Theory. Series A **10**, 119 (1971).
[10] S. Yu and C. H. Oh, Physical Review Letters **108**, 030402 (2012), arXiv:1109.4396.
[11] J. A. Martin H., "Graph-chromatic implicit relations," (2009), arXiv:0901.1255 [math.CO].
[12] A. Dvurečenskij, S. Pulmannová, and K. Svozil, Helvetica Physica Acta **68**, 407 (1995), arXiv:1806.04271.
[13] K. Svozil, International Journal of Theoretical Physics **44**, 745 (2005), arXiv:quant-ph/0209136.
[14] A. Cabello, M. Kleinmann, and C. Budroni, Physical Review Letters **114**, 250402 (2015), arXiv:1501.03432.
[15] R. Ramanathan and P. Horodecki, Physical Review Letters **112**, 040404 (2014), arXiv:1212.5933.
[16] G. Scarpa and S. Severini, IEEE Transactions on Information Theory **58**, 2524 (2012), arXiv:1106.0712.
[17] S. Prakash, "Quantum memory advantage from contextuality," (2026), arXiv:2607.00507 [quant-ph].
[18] M. Navara and K. Svozil, "Construction of Kochen-Specker sets from mutually unbiased bases," (2025), arXiv:2509.08636 [quant-ph].
[19] G. Haynes, C. Park, A. Schaeffer, J. Webster, and L. H. Mitchell, The Electronic Journal of Combinatorics **17**, R55 (2010).
[20] J. Bub, Foundations of Physics **26**, 787 (1996).
[21] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
[22] M. Kernaghan, Journal of Physics A: Mathematical and General **27**, L829 (1994).
[23] A. Silva and A. Ibiapina, "On orthogonal vector edge coloring," (2019), arXiv:1909.01918 [cs.DM].
[24] G. Brinkmann, M. Preissmann, and D. Sasaki, Discrete Mathematics & Theoretical Computer Science **17**, 369 (2015).
[25] G. Schrag, Discrete Mathematics **70**, 185 (1988).
[26] K. Svozil and J. Tkadlec, Journal of Mathematical Physics **37**, 5380 (1996).
[27] L. Hogben, K. F. Palmowski, D. E. Roberson, and S. Severini, Electronic Journal of Linear Algebra **32**, 98 (2017).
[28] M. Waegell and P. K. Aravind, Foundations of Physics **41**, 1786 (2011), arXiv:1103.6058 [quant-ph].
[29] R. Isaacs, The American Mathematical Monthly **82**, 221 (1975).
[30] D. Blanuša, Glasnik Matematičko-Fizički i Astronomski, Serija II **1**, 31 (1946).
[31] M. Pavičić, Entropy **21**, 1107 (2019).
[32] S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
[33] A. Cabello, J. R. Portillo, A. Solís, and K. Svozil, Physical Review A **98**, 012106 (2018), arXiv:1805.00796.
[34] J. Harding and R. Salinas Schmeis, International Journal of Theoretical Physics **64**, 211 (2025), arXiv:2505.13871 [math-ph].
[35] A. Peres, Journal of Physics A: Mathematical and General **24**, L175 (1991).
[36] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **339**, 425 (2005).

[37]M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
[38]A. Cabello, Physical Review Letters **135**, 190203 (2025), arXiv:2508.07335 [quant-ph].
[39]M. Pavičić, "On simplest Kochen-Specker sets," (2025), arXiv:2512.10483 [quant-ph].
[40]M. Pavičić, Quantum **7**, 953 (2023).

Supplementary Material for "Strong Colorings of Orthogonality Hypergraphs as Noncontextual Eigenvalue Assignments"

Karl Svozil
Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, 1040 Vienna, Austria

(Dated: November 5, 2026)

This supplementary material provides the complete vector and context enumerations, two-valued state tables, normalized strong coloring tables, odd-cycle nonextendability certificates, chromatic signature and separation profile tables, parent-embedding dictionaries, and supplementary proofs and finite verification certificates supporting the main text. Section S1 lists the explicit unnormalized coordinates and context tables for the completed Yu–Oh hypergraph H_{YO}^{+} and its faithful restrictions $H_{17,10}$ and $H_{21,13}$, the three- and four-dimensional logical gadgets, the 37-ray complex Hesse hypergraph H_{37}, the 26-ray suspension $H_{\mathrm{YO}}^{(4)}$ and 65-ray canonical linear completion H_{65}, the Cabello completions $C_{36,18}$ and $C_{45,21}$, and the specified finite KS parents (49-ray Schütte–Bub, 87-ray MUB-inventory union, 40- and 72-ray Peres unions, and 68-ray three-ray extensions). Section S2 gives the complete two-valued state tables, extendable-state classifications, normalized coloring tables, odd-cycle certificates, and separation profiles across both regimes. Section S3 presents the supplementary proofs and exact reductions, including Isaacs dot-product snarks and quantitative Δ_3 robustness bounds for G_{32}, the 540,672-candidate reduction for the minimum 18-state order-determining family on $L(H_{37})$, the canonical completion selection rule for H_{65}, the forced-zero and ray-criticality certificates for the four-dimensional Cabello completions, transverse criticality deletion colorings, and historical Kochen–Specker gadget certificates.

READING GUIDE AND VERIFICATION SCOPE

This supplement collects complete vector and context inventories (S1), state and coloring enumerations (S2), and extended proofs and certificates (S3). Labels are local to each configuration. "Exact palette" means $\chi(H) = d$, with no additional density or geometric assumption: the outcomes admit a noncontextual assignment of d eigenvalues, that is, every complete experiment can be measured by a maximal observable with one common spectrum of d eigenvalues, with each outcome carrying the same eigenvalue in every context. Colorings are such eigenvalue assignments, and extendable states are the truth assignments "the outcome with eigenvalue λ_a occurs" that they produce.

The five-state and four-state $H_{17,10}$ mixtures and both-direction gadget witnesses use the complete state families. Every selected support satisfies every context equation, and rational arithmetic verifies every Born marginal and the attained nonextendable weights $1/9$ and $1/3$. The ordinary implication $10 \Rightarrow 24$ on $H_{17,10}$ gives a false inclusion between distinct rays and hence a failure of order determination.

The companion electronic data include all coordinates, state enumerations (including the compressed 393,216-state H_{65} inventory), verification programs, and the exhaustive certificates for the eighteen-state Hesse minimum and the 65,535-subset Yu–Oh classification. Checking an eighteen-state family establishes an upper bound; the matching lower bound requires the exhaustive certificate in Sec. S3.3. The directories `part-a/data`, `part-a/verification`, `part-b/data`, and `part-b/verification` contain the underlying data and verifiers. Reports and source-integrity hashes are in `audit/`; Sec. S3.7 describes the joint verification entry point and its scope.

S1. EXPLICIT UNNORMALIZED VECTORS AND CONTEXTS

All numerical vertex labels in this section are local to the named configuration and match the figures and theorems of the main text.

S1.1. Completed Yu–Oh and the Petersen-derived contexts

The three-dimensional vectors below give all projective rays used in Sec. III C; signs are chosen to make the first nonzero integer coordinate positive. The 26-ray suspension appends zero and adds e_4. The abstract G_{32}, H_9, H_8 context lists in Eq. (10) and Theorem 11 specify their incidences completely. The geometric obstructions exclude a corresponding injective three-dimensional vector enumeration for these abstract examples.

TABLE S1: All 25 unnormalized completed Yu–Oh vectors.

i	r_i	i	r_i
1	$(0,1,1)$	14	$(2,1,1)$
2	$(1,0,0)$	15	$(1,2,-1)$
3	$(0,1,-1)$	16	$(1,-2,1)$
4	$(1,1,0)$	17	$(1,1,-2)$
5	$(0,0,1)$	18	$(1,-1,-2)$
6	$(1,-1,0)$	19	$(2,1,-1)$
7	$(1,0,1)$	20	$(2,-1,1)$
8	$(0,1,0)$	21	$(1,-2,-1)$
9	$(1,0,-1)$	22	$(1,1,2)$
10	$(1,-1,1)$	23	$(1,2,1)$
11	$(1,-1,-1)$	24	$(1,-1,2)$
12	$(1,1,1)$	25	$(2,-1,-1)$
13	$(1,1,-1)$		

S1.2. Faithful Yu–Oh restrictions $H_{17,10}$ and $H_{21,13}$ and logical gadgets in dimensions three and four

Tables S2 and S3 record the restriction membership of the 25 parent rays and 16 parent contexts of H^{+}_{YO} in $H_{17,10}$ and $H_{21,13}$. Tables S4–S14 provide the explicit unnormalized integer coordinates and context lists for the faithful real 3-dimensional gadgets P_5, $B_{13,7}$, $D_{16,9}$, $I_{16,9}$, and $T_{19,11}$ and the faithful real 4-dimensional gadgets $W^{(4)}_{11,5}$, $W^{(4)}_{16,6}$, $B^{(4)}_{14,7}$, $L^{(4)}_{18,7}$, $D^{(4)}_{17,9}$, and $L^{(4)}_{22,9}$. Note that for the standalone gadgets P_5, $D_{16,9}$, $I_{16,9}$, and $T_{19,11}$, the vertex numbering is local to each gadget panel; in $L^{(4)}_{18,7}$ [Fig. 3(d)], $r_{13} = (1,0,1,1)$ and $r_{14} = (2,0,-1,-1)$ are chosen so that $r_{13} \not\perp r_{15}$, whereas in $L^{(4)}_{22,9}$ [Fig. 3(f)], $r_{13} = (1,0,4,4)$ and $r_{14} = (8,0,-1,-1)$ are used so that $r_{13} \perp r_{15}$ in the added context $\{13,20,15,19\}$.

TABLE S2: All 25 unnormalized parent vectors; membership of the two geometric examples.

i	r_i	$H_{17,10}$	$H_{21,13}$	i	r_i	$H_{17,10}$	$H_{21,13}$
1	$(0,1,1)$	yes	yes	14	$(2,1,1)$	yes	yes
2	$(1,0,0)$	yes	yes	15	$(1,2,-1)$	yes	yes
3	$(0,1,-1)$	yes	yes	16	$(1,-2,1)$	–	–
4	$(1,1,0)$	yes	yes	17	$(1,1,-2)$	–	–
5	$(0,0,1)$	yes	yes	18	$(1,-1,-2)$	yes	yes
6	$(1,-1,0)$	yes	yes	19	$(2,1,-1)$	yes	yes
7	$(1,0,1)$	yes	yes	20	$(2,-1,1)$	–	yes
8	$(0,1,0)$	yes	yes	21	$(1,-2,-1)$	–	yes
9	$(1,0,-1)$	yes	yes	22	$(1,1,2)$	–	yes
10	$(1,-1,1)$	yes	yes	23	$(1,2,1)$	yes	yes
11	$(1,-1,-1)$	yes	yes	24	$(1,-1,2)$	yes	yes
12	$(1,1,1)$	–	–	25	$(2,-1,-1)$	–	–
13	$(1,1,-1)$	–	yes				

TABLE S3: The 16 parent contexts and the two selected restrictions.

j	context	$H_{17,10}$	$H_{21,13}$
1	$\{1,2,3\}$	yes	yes
2	$\{7,8,9\}$	yes	yes
3	$\{4,5,6\}$	yes	yes
4	$\{2,8,5\}$	yes	yes
5	$\{3,12,25\}$	–	–
6	$\{9,12,16\}$	–	–
7	$\{6,12,17\}$	–	–
8	$\{3,11,14\}$	yes	yes
9	$\{7,11,15\}$	yes	yes
10	$\{4,11,24\}$	yes	yes
11	$\{9,10,23\}$	yes	yes
12	$\{1,10,19\}$	yes	yes
13	$\{4,10,18\}$	yes	yes
14	$\{1,13,20\}$	–	yes
15	$\{7,13,21\}$	–	yes
16	$\{6,13,22\}$	–	yes

S1.3. 3D panel (a): P_5

TABLE S4: Unnormalized vectors and all contexts of P_5.

i	r_i	j	Context
1	$(1,0,0)$	1	$\{1,6,2\}$
2	$(0,1,1)$	2	$\{2,7,3\}$
3	$(1,1,-1)$	3	$\{3,8,4\}$
4	$(1,-2,-1)$	4	$\{4,9,5\}$
5	$(0,1,-2)$	5	$\{5,10,1\}$
6	$(0,1,-1)$		
7	$(2,-1,1)$		
8	$(1,0,1)$		
9	$(5,2,1)$		
10	$(0,2,1)$		

S1.4. 3D panel (b): $B_{13,7}$

TABLE S5: Unnormalized vectors and all contexts of $B_{13,7}$.

i	r_i	j	Context
1	$(0,1,1)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,1,-1)$	3	$\{2,8,5\}$
5	$(0,0,1)$	4	$\{3,11,14\}$
7	$(1,0,1)$	5	$\{7,11,15\}$
8	$(0,1,0)$	6	$\{9,10,23\}$
9	$(1,0,-1)$	7	$\{1,10,19\}$
10	$(1,-1,1)$		
11	$(1,-1,-1)$		
14	$(2,1,1)$		
15	$(1,2,-1)$		
19	$(2,1,-1)$		
23	$(1,2,1)$		

S1.5. 3D panel (c): $D_{16,9}$

TABLE S6: Unnormalized vectors and all contexts of $D_{16,9}$.

i	r_i	j	Context
1	$(0,1,2)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,2,-1)$	3	$\{4,5,6\}$
4	$(1,2,0)$	4	$\{2,8,5\}$
5	$(0,0,1)$	5	$\{3,11,14\}$
6	$(2,-1,0)$	6	$\{7,11,15\}$
7	$(1,0,1)$	7	$\{4,11,24\}$
8	$(0,1,0)$	8	$\{9,10,23\}$
9	$(1,0,-1)$	9	$\{4,10,18\}$
10	$(2,-1,2)$		
11	$(2,-1,-2)$		
14	$(5,2,4)$		
15	$(1,4,-1)$		
18	$(4,-2,-5)$		
23	$(1,4,1)$		
24	$(4,-2,5)$		

S1.6. 3D panel (d): $I_{16,9}$

TABLE S7: Unnormalized vectors and all contexts of $I_{16,9}$.

i	r_i	j	Context
1	$(0,1,1)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,1,-1)$	3	$\{2,8,5\}$
4	$(4,5,3)$	4	$\{3,11,14\}$
5	$(0,0,1)$	5	$\{7,11,15\}$
7	$(1,0,2)$	6	$\{4,11,24\}$
8	$(0,1,0)$	7	$\{9,10,23\}$
9	$(2,0,-1)$	8	$\{1,10,19\}$
10	$(1,-2,2)$	9	$\{4,10,18\}$
11	$(2,-1,-1)$		
14	$(1,1,1)$		
15	$(2,5,-1)$		
18	$(16,-5,-13)$		
19	$(4,1,-1)$		
23	$(2,5,4)$		
24	$(1,-5,7)$		

S1.7. 3D panel (e): $T_{19,11}$

TABLE S8: Unnormalized vectors and all contexts of $T_{19,11}$.

i	r_i	j	Context
1	$(0,1,2)$	1	$\{1,2,3\}$
2	$(1,0,0)$	2	$\{7,8,9\}$
3	$(0,2,-1)$	3	$\{4,5,6\}$
4	$(1,2,0)$	4	$\{2,8,5\}$
5	$(0,0,1)$	5	$\{3,11,14\}$
6	$(2,-1,0)$	6	$\{7,11,15\}$
7	$(1,0,1)$	7	$\{4,11,24\}$
8	$(0,1,0)$	8	$\{9,10,23\}$
9	$(1,0,-1)$	9	$\{4,10,18\}$
10	$(2,-1,2)$	10	$\{14,26,27\}$
11	$(2,-1,-2)$	11	$\{18,26,28\}$
14	$(5,2,4)$		
15	$(1,4,-1)$		
18	$(4,-2,-5)$		
23	$(1,4,1)$		
24	$(4,-2,5)$		
26	$(2,-41,18)$		
27	$(200,-82,-209)$		
28	$(241,82,160)$		

S1.8. 4D panel (a): $W^{(4)}_{11,5}$

TABLE S9: Unnormalized vectors and all contexts of $W^{(4)}_{11,5}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{0,1,6,2\}$
1	$(1,0,0,0)$	2	$\{0,2,7,3\}$
2	$(0,1,1,0)$	3	$\{0,3,8,4\}$
3	$(1,1,-1,0)$	4	$\{0,4,9,5\}$
4	$(1,-2,-1,0)$	5	$\{0,5,10,1\}$
5	$(0,1,-2,0)$		
6	$(0,1,-1,0)$		
7	$(2,-1,1,0)$		
8	$(1,0,1,0)$		
9	$(5,2,1,0)$		
10	$(0,2,1,0)$		

S1.9. 4D panel (b): $W^{(4)}_{16,6}$

TABLE S10: Unnormalized vectors and all contexts of $W^{(4)}_{16,6}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{2,0,6,1\}$
1	$(1,0,0,0)$	2	$\{2,7,13,3\}$
2	$(0,1,1,0)$	3	$\{3,0,8,4\}$
3	$(1,1,-1,0)$	4	$\{4,9,14,5\}$
4	$(1,-2,-1,0)$	5	$\{5,10,15,1\}$
5	$(0,1,-2,0)$	6	$\{0,11,12,5\}$
6	$(0,1,-1,0)$		
7	$(2,-1,1,1)$		
8	$(1,0,1,0)$		
9	$(5,2,1,1)$		
10	$(0,4,2,1)$		
11	$(1,2,1,0)$		
12	$(5,-2,-1,0)$		
13	$(2,-1,1,-6)$		
14	$(5,2,1,-30)$		
15	$(0,2,1,-10)$		

S1.10. 4D panel (c): $B^{(4)}_{14,7}$

TABLE S11: Unnormalized vectors and all contexts of $B^{(4)}_{14,7}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{0,1,2,3\}$
1	$(0,1,1,0)$	2	$\{0,7,8,9\}$
2	$(1,0,0,0)$	3	$\{0,2,8,5\}$
3	$(0,1,-1,0)$	4	$\{0,3,11,14\}$
5	$(0,0,1,0)$	5	$\{0,7,11,15\}$
7	$(1,0,1,0)$	6	$\{0,1,10,19\}$
8	$(0,1,0,0)$	7	$\{0,9,10,23\}$
9	$(1,0,-1,0)$		
10	$(1,-1,1,0)$		
11	$(1,-1,-1,0)$		
14	$(2,1,1,0)$		
15	$(1,2,-1,0)$		
19	$(2,1,-1,0)$		
23	$(1,2,1,0)$		

S1.11. 4D panel (d): $L^{(4)}_{18,7}$

TABLE S12: Unnormalized vectors and all contexts of $L^{(4)}_{18,7}$.

i	r_i	j	Context
1	$(1,0,0,0)$	1	$\{1,2,4,3\}$
2	$(0,1,0,0)$	2	$\{5,8,7,6\}$
3	$(0,0,1,0)$	3	$\{4,9,10,8\}$
4	$(0,0,0,1)$	4	$\{11,1,12,5\}$
5	$(0,1,-1,-1)$	5	$\{11,14,13,2\}$
6	$(4,-5,-11,6)$	6	$\{15,3,16,7\}$
7	$(1,-4,0,-4)$	7	$\{15,17,18,6\}$
8	$(4,1,1,0)$		
9	$(1,-5,1,0)$		
10	$(2,-1,-7,0)$		
11	$(0,0,1,-1)$		
12	$(0,2,1,1)$		
13	$(1,0,1,1)$		
14	$(2,0,-1,-1)$		
15	$(4,2,0,-1)$		
16	$(4,-5,0,6)$		
17	$(5,-9,7,2)$		
18	$(2,113,77,234)$		

S1.12. 4D panel (e): $D^{(4)}_{17,9}$

TABLE S13: Unnormalized vectors and all contexts of $D^{(4)}_{17,9}$.

i	r_i	j	Context
0	$(0,0,0,1)$	1	$\{0,1,2,3\}$
1	$(0,1,2,0)$	2	$\{0,7,8,9\}$
2	$(1,0,0,0)$	3	$\{0,4,5,6\}$
3	$(0,2,-1,0)$	4	$\{0,2,8,5\}$
4	$(1,2,0,0)$	5	$\{0,3,11,14\}$
5	$(0,0,1,0)$	6	$\{0,7,11,15\}$
6	$(2,-1,0,0)$	7	$\{0,9,10,23\}$
7	$(1,0,1,0)$	8	$\{0,4,11,24\}$
8	$(0,1,0,0)$	9	$\{0,4,10,18\}$
9	$(1,0,-1,0)$		
10	$(2,-1,2,0)$		
11	$(2,-1,-2,0)$		
14	$(5,2,4,0)$		
15	$(1,4,-1,0)$		
18	$(4,-2,-5,0)$		
23	$(1,4,1,0)$		
24	$(4,-2,5,0)$		

S1.13. 4D panel (f): $L^{(4)}_{22,9}$

TABLE S14: Unnormalized vectors and all contexts of $L^{(4)}_{22,9}$.

i	r_i	j	Context
1	$(1,0,0,0)$	1	$\{1,2,4,3\}$
2	$(0,1,0,0)$	2	$\{5,8,7,6\}$
3	$(0,0,1,0)$	3	$\{4,9,10,8\}$
4	$(0,0,0,1)$	4	$\{11,1,12,5\}$
5	$(0,1,-1,-1)$	5	$\{11,14,13,2\}$
6	$(4,-5,-11,6)$	6	$\{15,3,16,7\}$
7	$(1,-4,0,-4)$	7	$\{15,17,18,6\}$
8	$(4,1,1,0)$	8	$\{11,21,22,19\}$
9	$(1,-5,1,0)$	9	$\{13,20,15,19\}$
10	$(2,-1,-7,0)$		
11	$(0,0,1,-1)$		
12	$(0,2,1,1)$		
13	$(1,0,4,4)$		
14	$(8,0,-1,-1)$		
15	$(4,2,0,-1)$		
16	$(4,-5,0,6)$		
17	$(5,-9,7,2)$		
18	$(2,113,77,234)$		
19	$(16,-33,-2,-2)$		
20	$(4,2,-21,20)$		
21	$(22,12,-11,-11)$		
22	$(129,44,153,153)$		

S1.14. Hesse H_{37}: vectors, contexts, and non-collinearity

This appendix uses exactly the vertex numbers and context numbers of Fig. 4. All indices a, k in the family notation are in $\mathbb{Z}_3$. The vectors are unnormalized and the inner product is Hermitian. Table S15 lists every vector, squared norm, color, and incident context. Table S16 lists all 24 contexts with their actual vectors. Together they specify the faithful orthogonal representation completely.

S1.14.1. All 37 vectors, their colors, and incident contexts

TABLE S15: Complete enumeration of the 37 vectors of H_{37}. The color $c(i) \in \{0,1,2,3\}$ is the explicit four-coloring in the main text. Context indices refer to Table S16.

i	Ray	v_i	$\|v_i\|^2$	$c(i)$	Incident contexts
1	z	$(1,0,0)$	1	0	$\{4,5,6\}$
2	M_{00}	$(1,1,1)$	3	0	$\{1,7,16\}$
3	M_{01}	$(1,\zeta,\zeta^2)$	3	1	$\{1,8,17\}$
4	M_{02}	$(1,\zeta^2,\zeta)$	3	2	$\{1,9,18\}$
5	M_{10}	$(1,\zeta,\zeta)$	3	0	$\{2,10,21\}$
6	M_{11}	$(1,\zeta^2,1)$	3	1	$\{2,11,19\}$
7	M_{12}	$(1,1,\zeta^2)$	3	2	$\{2,12,20\}$
8	M_{20}	$(1,\zeta^2,\zeta^2)$	3	0	$\{3,13,23\}$
9	M_{21}	$(1,1,\zeta)$	3	1	$\{3,14,24\}$
10	M_{22}	$(1,\zeta,1)$	3	2	$\{3,15,22\}$
11	P_0	$(0,1,-1)$	2	3	$\{4,7,10,13\}$
12	P_1	$(0,1,-\zeta)$	2	3	$\{5,8,11,14\}$
13	P_2	$(0,1,-\zeta^2)$	2	3	$\{6,9,12,15\}$
14	Q_0	$(-1,0,1)$	2	3	$\{16,19,22\}$
15	Q_1	$(-\zeta,0,1)$	2	3	$\{17,20,23\}$
16	Q_2	$(-\zeta^2,0,1)$	2	3	$\{18,21,24\}$
17	A_0	$(0,1,1)$	2	1	$\{4\}$

Continued on the next page.

TABLE S15: Complete enumeration of the 37 vectors of H_{37}. The color $c(i) \in \{0,1,2,3\}$ is the explicit four-coloring in the main text. Context indices refer to Table S16. (continued).

i	Ray	v_i	$\|v_i\|^2$	$c(i)$	Incident contexts
18	A_1	$(0,1,\zeta)$	2	1	{5}
19	A_2	$(0,1,\zeta^2)$	2	1	{6}
20	B_{00}	$(-2,1,1)$	6	1	{7}
21	B_{01}	$(-2,\zeta,\zeta^2)$	6	2	{8}
22	B_{02}	$(-2,\zeta^2,\zeta)$	6	0	{9}
23	B_{10}	$(-2,\zeta,\zeta)$	6	1	{10}
24	B_{11}	$(-2,\zeta^2,1)$	6	2	{11}
25	B_{12}	$(-2,1,\zeta^2)$	6	0	{12}
26	B_{20}	$(-2,\zeta^2,\zeta^2)$	6	1	{13}
27	B_{21}	$(-2,1,\zeta)$	6	2	{14}
28	B_{22}	$(-2,\zeta,1)$	6	0	{15}
29	C_{00}	$(1,-2,1)$	6	1	{16}
30	C_{01}	$(1,-2\zeta,\zeta^2)$	6	2	{17}
31	C_{02}	$(1,-2\zeta^2,\zeta)$	6	0	{18}
32	C_{10}	$(1,-2\zeta^2,1)$	6	2	{19}
33	C_{11}	$(1,-2,\zeta^2)$	6	0	{20}
34	C_{12}	$(1,-2\zeta,\zeta)$	6	1	{21}
35	C_{20}	$(1,-2\zeta,1)$	6	0	{22}
36	C_{21}	$(1,-2\zeta^2,\zeta^2)$	6	1	{23}
37	C_{22}	$(1,-2,\zeta)$	6	2	{24}

S1.14.2. All 24 contexts and their orthogonal vectors

TABLE S16: All 24 orthogonal triads, numbered in the same order as the context colors in Fig. 4. Vectors are given in increasing order of their vertex numbers. Each triad contributes three orthogonality edges; the 72 resulting pairs are all the orthogonal pairs.

Context	Family	Vertices	Unnormalized vectors
e_1	$\mathcal{M}_0$	$\{2,3,4\}$	$\{(1,1,1),\ (1,\zeta,\zeta^2),\ (1,\zeta^2,\zeta)\}$
e_2	$\mathcal{M}_1$	$\{5,6,7\}$	$\{(1,\zeta,\zeta),\ (1,\zeta^2,1),\ (1,1,\zeta^2)\}$
e_3	$\mathcal{M}_2$	$\{8,9,10\}$	$\{(1,\zeta^2,\zeta^2),\ (1,1,\zeta),\ (1,\zeta,1)\}$
e_4	$\mathcal{A}_0$	$\{1,11,17\}$	$\{(1,0,0),\ (0,1,-1),\ (0,1,1)\}$
e_5	$\mathcal{A}_1$	$\{1,12,18\}$	$\{(1,0,0),\ (0,1,-\zeta),\ (0,1,\zeta)\}$
e_6	$\mathcal{A}_2$	$\{1,13,19\}$	$\{(1,0,0),\ (0,1,-\zeta^2),\ (0,1,\zeta^2)\}$
e_7	$\mathcal{B}_{00}$	$\{2,11,20\}$	$\{(1,1,1),\ (0,1,-1),\ (-2,1,1)\}$
e_8	$\mathcal{B}_{01}$	$\{3,12,21\}$	$\{(1,\zeta,\zeta^2),\ (0,1,-\zeta),\ (-2,\zeta,\zeta^2)\}$
e_9	$\mathcal{B}_{02}$	$\{4,13,22\}$	$\{(1,\zeta^2,\zeta),\ (0,1,-\zeta^2),\ (-2,\zeta^2,\zeta)\}$
e_{10}	$\mathcal{B}_{10}$	$\{5,11,23\}$	$\{(1,\zeta,\zeta),\ (0,1,-1),\ (-2,\zeta,\zeta)\}$
e_{11}	$\mathcal{B}_{11}$	$\{6,12,24\}$	$\{(1,\zeta^2,1),\ (0,1,-\zeta),\ (-2,\zeta^2,1)\}$
e_{12}	$\mathcal{B}_{12}$	$\{7,13,25\}$	$\{(1,1,\zeta^2),\ (0,1,-\zeta^2),\ (-2,1,\zeta^2)\}$
e_{13}	$\mathcal{B}_{20}$	$\{8,11,26\}$	$\{(1,\zeta^2,\zeta^2),\ (0,1,-1),\ (-2,\zeta^2,\zeta^2)\}$
e_{14}	$\mathcal{B}_{21}$	$\{9,12,27\}$	$\{(1,1,\zeta),\ (0,1,-\zeta),\ (-2,1,\zeta)\}$
e_{15}	$\mathcal{B}_{22}$	$\{10,13,28\}$	$\{(1,\zeta,1),\ (0,1,-\zeta^2),\ (-2,\zeta,1)\}$
e_{16}	$\mathcal{C}_{00}$	$\{2,14,29\}$	$\{(1,1,1),\ (-1,0,1),\ (1,-2,1)\}$

Continued on the next page.

TABLE S16: All 24 orthogonal triads, numbered in the same order as the context colors in Fig. 4. Vectors are given in increasing order of their vertex numbers. Each triad contributes three orthogonality edges; the 72 resulting pairs are all the orthogonal pairs. (continued).

Context	Family	Vertices	Unnormalized vectors
e_{17}	$\mathcal{C}_{01}$	$\{3,15,30\}$	$\{(1,\zeta,\zeta^2),\ (-\zeta,0,1),\ (1,-2\zeta,\zeta^2)\}$
e_{18}	$\mathcal{C}_{02}$	$\{4,16,31\}$	$\{(1,\zeta^2,\zeta),\ (-\zeta^2,0,1),\ (1,-2\zeta^2,\zeta)\}$
e_{19}	$\mathcal{C}_{10}$	$\{6,14,32\}$	$\{(1,\zeta^2,1),\ (-1,0,1),\ (1,-2\zeta^2,1)\}$
e_{20}	$\mathcal{C}_{11}$	$\{7,15,33\}$	$\{(1,1,\zeta^2),\ (-\zeta,0,1),\ (1,-2,\zeta^2)\}$
e_{21}	$\mathcal{C}_{12}$	$\{5,16,34\}$	$\{(1,\zeta,\zeta),\ (-\zeta^2,0,1),\ (1,-2\zeta,\zeta)\}$
e_{22}	$\mathcal{C}_{20}$	$\{10,14,35\}$	$\{(1,\zeta,1),\ (-1,0,1),\ (1,-2\zeta,1)\}$
e_{23}	$\mathcal{C}_{21}$	$\{8,15,36\}$	$\{(1,\zeta^2,\zeta^2),\ (-\zeta,0,1),\ (1,-2\zeta^2,\zeta^2)\}$
e_{24}	$\mathcal{C}_{22}$	$\{9,16,37\}$	$\{(1,1,\zeta),\ (-\zeta^2,0,1),\ (1,-2,\zeta)\}$

S1.14.3. Norm shells and a non-collinearity certificate

The four squared-norm shells have sizes

$$\#\{i : \|v_i\|^2 = 1\} = 1, \quad \#\{i : \|v_i\|^2 = 2\} = 9, \quad \#\{i : \|v_i\|^2 = 3\} = 9, \quad \#\{i : \|v_i\|^2 = 6\} = 18. \tag{S1}$$

Exact evaluation over the $\binom{37}{2} = 666$ unordered pairs gives

$$\max_{i<j} \frac{|\langle v_i, v_j\rangle|^2}{\|v_i\|^2\|v_j\|^2} = \frac{3}{4} < 1. \tag{S2}$$

For example, the maximum is attained by $v_{11} = P_0 = (0,1,-1)$ and $v_{29} = C_{00} = (1,-2,1)$:

$$\frac{|\langle v_{11}, v_{29}\rangle|^2}{\|v_{11}\|^2\|v_{29}\|^2} = \frac{(-3)^2}{2\cdot 6} = \frac{3}{4}. \tag{S3}$$

Thus no two printed vectors represent the same ray. The maximum and all 72 zero inner products are checked exactly by the accompanying verifier; the family-wise argument in the main text explains why there are no further orthogonalities.

S1.15. Four-dimensional $H^{(4)}_{\mathrm{YO}}$ and H_{65}: vectors and contexts

For direct human and computational verification, this appendix enumerates all unnormalized integer vectors in $\mathbb{Z}^4$ and all four-element contexts for the two four-dimensional examples H_{65} and $H^{(4)}_{\mathrm{YO}}$:

1. The *canonical linear hypergraph* $H_{65} = (V_{65}, E_{29})$ (Theorem 22, Appendix S1.15.1), which has exactly 65 **pairwise non-collinear integer vectors** (17 core vectors $v_1, \ldots, v_{17}$ and 48 degree-1 completion vectors $v_{18}, \ldots, v_{65}$), 29 **four-element contexts** (5 core contexts $C_1, \ldots, C_5$ and 24 bridge-completion contexts $E_1, \ldots, E_{24}$), and 174 **orthogonality edges** ($29 \times \binom{4}{2} = 174$).

2. The *non-linear warm-up apex suspension* $H^{(4)}_{\mathrm{YO}} = (V_{26}, E^{(4)}_{16})$ (Proposition 20, Appendix S1.15.2), which has exactly 26 **pairwise non-collinear integer vectors** (13 Yu–Oh rays $v_1, \ldots, v_{13}$, 12 completion rays $v_{14}, \ldots, v_{25}$ in $\mathbb{R}^3 \times \{0\}$, and 1 apex ray v_{26}), 16 **four-element contexts**, and 73 **orthogonality edges** ($16 \times \binom{3}{2} = 48$ edges in $\mathbb{R}^3 \times \{0\}$ plus 25 apex edges).

Canonical primitive normal form and human-verifiable non-collinearity. Every vector $v_i = (x_{i,1}, x_{i,2}, x_{i,3}, x_{i,4})^T \in \mathbb{Z}^4$ listed below in both H_{65} and $H_{\mathrm{YO}}^{(4)}$ is written in *canonical primitive projective form*:

$$\begin{gathered} \gcd\big(|x_{i,1}|, |x_{i,2}|, |x_{i,3}|, |x_{i,4}|\big) = 1, \\ \text{first nonzero entry of } v_i > 0. \end{gathered} \tag{S4}$$

Because every projective ray $[v] \in \mathbb{P}(\mathbb{Q}^4)$ possesses a *unique* representative in $\mathbb{Z}^4$ satisfying Eq. (S4), two vectors v_i, v_j in canonical primitive form are collinear (proportional over $\mathbb{R}$ or $\mathbb{C}$) *if and only if they are identical* 4*-tuples* ($v_i = v_j$) and, in particular, have the same squared Euclidean norm $\|v_i\|^2 = \|v_j\|^2$.

S1.15.1. The canonical linear 65*-vertex,* 29*-context hypergraph* H_{65}

Table S17 enumerates all 65 unnormalized integer vectors $v_1, \ldots, v_{65} \in \mathbb{Z}^4$ of H_{65}, together with their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0, 1, 2, 3, 4\}$ in the strong 5-coloring extending Eq. (S28), and the contexts in E_{29} to which each vertex belongs. Table S18 enumerates all 29 contexts $C_1, \ldots, C_5, E_1, \ldots, E_{24} \in E_{29}$, displaying side by side both the vertex indices $\{i, j, k, \ell\}$ and the four mutually orthogonal integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

To verify by hand that all 65 vectors in Table S17 are pairwise non-collinear across all $\binom{65}{2} = 2{,}080$ pairs, observe that all 65 tuples satisfy Eq. (S4) and partition by squared norm $\|v_i\|^2$ into seven small, disjoint norm shells ($4 + 6 + 7 + 12 + 12 + 12 + 12 = 65$):

- $\|v_i\|^2 = 1$ (4 vectors): indices $\{1, 2, 3, 4\}$, the 4 standard coordinate basis vectors;
- $\|v_i\|^2 = 2$ (6 vectors): indices $\{12, \ldots, 17\}$, one vector for each of the $\binom{4}{2} = 6$ coordinate-pair supports;
- $\|v_i\|^2 = 4$ (7 vectors): indices $\{5, \ldots, 11\}$, seven distinct ± 1 vectors with $x_{i,1} = +1$;
- $\|v_i\|^2 = 11$ (12 vectors): indices $\{18, 20, 22, 25, 26, 28, 30, 33, 34, 36, 39, 41\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct sign/entry patterns of $\{0, \pm 1, \pm 1, \pm 3\}$;
- $\|v_i\|^2 = 18$ (12 vectors): indices $\{43, 45, 46, 49, 51, 53, 54, 56, 59, 61, 62, 65\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 2, \pm 2, \pm 3\}$;
- $\|v_i\|^2 = 22$ (12 vectors): indices $\{19, 21, 23, 24, 27, 29, 31, 32, 35, 37, 38, 40\}$, each having a single zero coordinate (three each with $x_1 = 0$, $x_2 = 0$, $x_3 = 0$, $x_4 = 0$) and distinct patterns of $\{0, \pm 2, \pm 3, \pm 3\}$;
- $\|v_i\|^2 = 36$ (12 vectors): indices $\{42, 44, 47, 48, 50, 52, 55, 57, 58, 60, 63, 64\}$, twelve distinct permutations and sign patterns of $\{\pm 1, \pm 1, \pm 3, \pm 5\}$.

Because no two 4-tuples within any of the seven norm shells coincide, no two of the 65 vectors are collinear; quantitatively, the maximum squared cosine over all $\binom{65}{2} = 2{,}080$ pairs is $\max_{1 \le i < j \le 65} (v_i \cdot v_j)^2 / (\|v_i\|^2 \|v_j\|^2) = 19^2/(11 \times 36) = 361/396 \approx 0.9116 < 1$, attained by (v_{41}, v_{47}), whose squared norms are 11 and 36 and whose inner product is 19.

TABLE S17: Complete enumeration of the 65 unnormalized integer vectors $v_i \in \mathbb{Z}^4$ of H_{65}, their squared norms $\|v_i\|^2$, their colors $c(i) \in \{0,1,2,3,4\}$, and the contexts in E_{29} containing vertex i.

i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i	i	$v_i \in \mathbb{Z}^4$	$\|v_i\|^2$	$c(i)$	Contexts containing i
1	$(1,0,0,0)$	1	0	C_1, E_3, E_7, E_{11}	34	$(1,-1,0,3)$	11	1	$E_9 = \{3,16,34,35\}$
2	$(0,1,0,0)$	1	1	C_1, E_1, E_5, E_{12}	35	$(3,-3,0,-2)$	22	3	$E_9 = \{3,16,34,35\}$
3	$(0,0,1,0)$	1	2	C_1, E_2, E_8, E_9	36	$(1,-1,3,0)$	11	1	$E_{10} = \{4,16,36,37\}$
4	$(0,0,0,1)$	1	3	C_1, E_4, E_6, E_{10}	37	$(3,-3,-2,0)$	22	2	$E_{10} = \{4,16,36,37\}$
5	$(1,1,1,1)$	4	0	$C_2, E_{13}, E_{19}, E_{23}$	38	$(0,2,3,3)$	22	1	$E_{11} = \{1,17,38,39\}$
6	$(1,1,-1,-1)$	4	1	$C_2, E_{15}, E_{17}, E_{24}$	39	$(0,3,-1,-1)$	11	3	$E_{11} = \{1,17,38,39\}$
7	$(1,-1,1,-1)$	4	2	$C_2, E_{16}, E_{20}, E_{21}$	40	$(2,0,-3,-3)$	22	0	$E_{12} = \{2,17,40,41\}$
8	$(1,-1,-1,1)$	4	4	$C_2, E_{14}, E_{18}, E_{22}$	41	$(3,0,1,1)$	11	3	$E_{12} = \{2,17,40,41\}$
9	$(1,1,-1,1)$	4	0	C_3, C_4	42	$(1,-5,3,1)$	36	1	$E_{13} = \{5,12,42,43\}$
10	$(1,-1,1,1)$	4	1	C_3, C_5	43	$(2,-1,-3,2)$	18	2	$E_{13} = \{5,12,42,43\}$
11	$(1,-1,-1,-1)$	4	3	C_4, C_5	44	$(1,-3,5,1)$	36	0	$E_{14} = \{8,12,44,45\}$
12	$(1,0,0,-1)$	2	3	$C_3, E_1, E_2, E_{13}, E_{14}$	45	$(2,3,1,2)$	18	1	$E_{14} = \{8,12,44,45\}$
13	$(0,1,1,0)$	2	4	$C_3, E_3, E_4, E_{15}, E_{16}$	46	$(1,-2,2,-3)$	18	0	$E_{15} = \{6,13,46,47\}$
14	$(1,0,1,0)$	2	2	$C_4, E_5, E_6, E_{17}, E_{18}$	47	$(5,-1,1,3)$	36	2	$E_{15} = \{6,13,46,47\}$
15	$(0,1,0,-1)$	2	1	$C_4, E_7, E_8, E_{19}, E_{20}$	48	$(3,-1,1,5)$	36	0	$E_{16} = \{7,13,48,49\}$
16	$(1,1,0,0)$	2	0	$C_5, E_9, E_{10}, E_{21}, E_{22}$	49	$(3,2,-2,-1)$	18	1	$E_{16} = \{7,13,48,49\}$
17	$(0,0,1,-1)$	2	2	$C_5, E_{11}, E_{12}, E_{23}, E_{24}$	50	$(1,3,-1,5)$	36	0	$E_{17} = \{6,14,50,51\}$
18	$(1,0,-3,1)$	11	0	$E_1 = \{2,12,18,19\}$	51	$(2,-3,-2,1)$	18	3	$E_{17} = \{6,14,50,51\}$
19	$(3,0,2,3)$	22	2	$E_1 = \{2,12,18,19\}$	52	$(1,5,-1,3)$	36	0	$E_{18} = \{8,14,52,53\}$
20	$(1,3,0,1)$	11	0	$E_2 = \{3,12,20,21\}$	53	$(2,1,-2,-3)$	18	1	$E_{18} = \{8,14,52,53\}$
21	$(3,-2,0,3)$	22	1	$E_2 = \{3,12,20,21\}$	54	$(3,-2,1,-2)$	18	2	$E_{19} = \{5,15,54,55\}$
22	$(0,1,-1,3)$	11	1	$E_3 = \{1,13,22,23\}$	55	$(3,1,-5,1)$	36	3	$E_{19} = \{5,15,54,55\}$
23	$(0,3,-3,-2)$	22	2	$E_3 = \{1,13,22,23\}$	56	$(1,2,3,2)$	18	0	$E_{20} = \{7,15,56,57\}$
24	$(2,-3,3,0)$	22	0	$E_4 = \{4,13,24,25\}$	57	$(5,1,-3,1)$	36	3	$E_{20} = \{7,15,56,57\}$
25	$(3,1,-1,0)$	11	1	$E_4 = \{4,13,24,25\}$	58	$(1,-1,-5,-3)$	36	1	$E_{21} = \{7,16,58,59\}$
26	$(1,0,-1,-3)$	11	0	$E_5 = \{2,14,26,27\}$	59	$(2,-2,-1,3)$	18	3	$E_{21} = \{7,16,58,59\}$
27	$(3,0,-3,2)$	22	3	$E_5 = \{2,14,26,27\}$	60	$(1,-1,-3,-5)$	36	1	$E_{22} = \{8,16,60,61\}$
28	$(1,-3,-1,0)$	11	0	$E_6 = \{4,14,28,29\}$	61	$(2,-2,3,-1)$	18	2	$E_{22} = \{8,16,60,61\}$
29	$(3,2,-3,0)$	22	1	$E_6 = \{4,14,28,29\}$	62	$(1,3,-2,-2)$	18	1	$E_{23} = \{5,17,62,63\}$
30	$(0,1,3,1)$	11	2	$E_7 = \{1,15,30,31\}$	63	$(5,-3,-1,-1)$	36	3	$E_{23} = \{5,17,62,63\}$
31	$(0,3,-2,3)$	22	3	$E_7 = \{1,15,30,31\}$	64	$(3,-5,-1,-1)$	36	0	$E_{24} = \{6,17,64,65\}$
32	$(2,3,0,3)$	22	0	$E_8 = \{3,15,32,33\}$	65	$(3,1,2,2)$	18	3	$E_{24} = \{6,17,64,65\}$
33	$(3,-1,0,-1)$	11	3	$E_8 = \{3,15,32,33\}$					

TABLE S18: Complete enumeration of the 29 four-element contexts of $H_{65} = (V_{65}, E_{29})$, listing for each context both its vertex indices $\{i,j,k,\ell\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_\ell\} \subset \mathbb{Z}^4$.

Context	Vertex set	v_i	v_j	v_k	v_ℓ
Five core 4-contexts $C_1, \ldots, C_5$ on $V_{\text{core}} = \{1,\ldots,17\}$:					
C_1	$\{1,2,3,4\}$	$v_1 = (1,0,0,0)$	$v_2 = (0,1,0,0)$	$v_3 = (0,0,1,0)$	$v_4 = (0,0,0,1)$
C_2	$\{5,6,7,8\}$	$v_5 = (1,1,1,1)$	$v_6 = (1,1,-1,-1)$	$v_7 = (1,-1,1,-1)$	$v_8 = (1,-1,-1,1)$
C_3	$\{9,10,12,13\}$	$v_9 = (1,1,-1,1)$	$v_{10} = (1,-1,1,1)$	$v_{12} = (1,0,0,-1)$	$v_{13} = (0,1,1,0)$
C_4	$\{9,11,14,15\}$	$v_9 = (1,1,-1,1)$	$v_{11} = (1,-1,-1,-1)$	$v_{14} = (1,0,1,0)$	$v_{15} = (0,1,0,-1)$
C_5	$\{10,11,16,17\}$	$v_{10} = (1,-1,1,1)$	$v_{11} = (1,-1,-1,-1)$	$v_{16} = (1,1,0,0)$	$v_{17} = (0,0,1,-1)$
Twelve coordinate-to-bridge completion contexts $E_1, \ldots, E_{12}$ (between C_1 and $B_1 \cup B_2 \cup B_3$):					
E_1	$\{2,12,18,19\}$	$v_2 = (0,1,0,0)$	$v_{12} = (1,0,0,-1)$	$v_{18} = (1,0,-3,1)$	$v_{19} = (3,0,2,3)$
E_2	$\{3,12,20,21\}$	$v_3 = (0,0,1,0)$	$v_{12} = (1,0,0,-1)$	$v_{20} = (1,3,0,1)$	$v_{21} = (3,-2,0,3)$
E_3	$\{1,13,22,23\}$	$v_1 = (1,0,0,0)$	$v_{13} = (0,1,1,0)$	$v_{22} = (0,1,-1,3)$	$v_{23} = (0,3,-3,-2)$
E_4	$\{4,13,24,25\}$	$v_4 = (0,0,0,1)$	$v_{13} = (0,1,1,0)$	$v_{24} = (2,-3,3,0)$	$v_{25} = (3,1,-1,0)$
E_5	$\{2,14,26,27\}$	$v_2 = (0,1,0,0)$	$v_{14} = (1,0,1,0)$	$v_{26} = (1,0,-1,-3)$	$v_{27} = (3,0,-3,2)$
E_6	$\{4,14,28,29\}$	$v_4 = (0,0,0,1)$	$v_{14} = (1,0,1,0)$	$v_{28} = (1,-3,-1,0)$	$v_{29} = (3,2,-3,0)$
E_7	$\{1,15,30,31\}$	$v_1 = (1,0,0,0)$	$v_{15} = (0,1,0,-1)$	$v_{30} = (0,1,3,1)$	$v_{31} = (0,3,-2,3)$
E_8	$\{3,15,32,33\}$	$v_3 = (0,0,1,0)$	$v_{15} = (0,1,0,-1)$	$v_{32} = (2,3,0,3)$	$v_{33} = (3,-1,0,-1)$
E_9	$\{3,16,34,35\}$	$v_3 = (0,0,1,0)$	$v_{16} = (1,1,0,0)$	$v_{34} = (1,-1,0,3)$	$v_{35} = (3,-3,0,-2)$
E_{10}	$\{4,16,36,37\}$	$v_4 = (0,0,0,1)$	$v_{16} = (1,1,0,0)$	$v_{36} = (1,-1,3,0)$	$v_{37} = (3,-3,-2,0)$
E_{11}	$\{1,17,38,39\}$	$v_1 = (1,0,0,0)$	$v_{17} = (0,0,1,-1)$	$v_{38} = (0,2,3,3)$	$v_{39} = (0,3,-1,-1)$
E_{12}	$\{2,17,40,41\}$	$v_2 = (0,1,0,0)$	$v_{17} = (0,0,1,-1)$	$v_{40} = (2,0,-3,-3)$	$v_{41} = (3,0,1,1)$
Twelve Hadamard-to-bridge completion contexts $E_{13}, \ldots, E_{24}$ (between C_2 and $B_1 \cup B_2 \cup B_3$):					
E_{13}	$\{5,12,42,43\}$	$v_5 = (1,1,1,1)$	$v_{12} = (1,0,0,-1)$	$v_{42} = (1,-5,3,1)$	$v_{43} = (2,-1,-3,2)$
E_{14}	$\{8,12,44,45\}$	$v_8 = (1,-1,-1,1)$	$v_{12} = (1,0,0,-1)$	$v_{44} = (1,-3,5,1)$	$v_{45} = (2,3,1,2)$

Continued on the next page.

TABLE S18: All 29 contexts of H_{65} (continued).

Context	Vertex set	v_i	v_j	v_k	v_ℓ
E_{15}	$\{6,13,46,47\}$	$v_6=(1,1,-1,-1)$	$v_{13}=(0,1,1,0)$	$v_{46}=(1,-2,2,-3)$	$v_{47}=(5,-1,1,3)$
E_{16}	$\{7,13,48,49\}$	$v_7=(1,-1,1,-1)$	$v_{13}=(0,1,1,0)$	$v_{48}=(3,-1,1,5)$	$v_{49}=(3,2,-2,-1)$
E_{17}	$\{6,14,50,51\}$	$v_6=(1,1,-1,-1)$	$v_{14}=(1,0,1,0)$	$v_{50}=(1,3,-1,5)$	$v_{51}=(2,-3,-2,1)$
E_{18}	$\{8,14,52,53\}$	$v_8=(1,-1,-1,1)$	$v_{14}=(1,0,1,0)$	$v_{52}=(1,5,-1,3)$	$v_{53}=(2,1,-2,-3)$
E_{19}	$\{5,15,54,55\}$	$v_5=(1,1,1,1)$	$v_{15}=(0,1,0,-1)$	$v_{54}=(3,-2,1,-2)$	$v_{55}=(3,1,-5,1)$
E_{20}	$\{7,15,56,57\}$	$v_7=(1,-1,1,-1)$	$v_{15}=(0,1,0,-1)$	$v_{56}=(1,2,3,2)$	$v_{57}=(5,1,-3,1)$
E_{21}	$\{7,16,58,59\}$	$v_7=(1,-1,1,-1)$	$v_{16}=(1,1,0,0)$	$v_{58}=(1,-1,-5,-3)$	$v_{59}=(2,-2,-1,3)$
E_{22}	$\{8,16,60,61\}$	$v_8=(1,-1,-1,1)$	$v_{16}=(1,1,0,0)$	$v_{60}=(1,-1,-3,-5)$	$v_{61}=(2,-2,3,-1)$
E_{23}	$\{5,17,62,63\}$	$v_5=(1,1,1,1)$	$v_{17}=(0,0,1,-1)$	$v_{62}=(1,3,-2,-2)$	$v_{63}=(5,-3,-1,-1)$
E_{24}	$\{6,17,64,65\}$	$v_6=(1,1,-1,-1)$	$v_{17}=(0,0,1,-1)$	$v_{64}=(3,-5,-1,-1)$	$v_{65}=(3,1,2,2)$

S1.15.2. The non-linear warm-up 26-vertex, 16-context apex suspension $H^{(4)}_{\mathrm{YO}}$

For completeness, the 26 unnormalized integer vectors $v_1,\dots,v_{26}\in\mathbb{Z}^4$ of the non-linear warm-up hypergraph $H^{(4)}_{\mathrm{YO}}=(V_{26},E^{(4)}_{16})$ in Proposition 20 (written in canonical primitive form satisfying Eq. (S4)) are:

$$\begin{aligned}
v_1&=(0,1,1,0), & v_2&=(1,0,0,0),\\
v_3&=(0,1,-1,0), & v_4&=(1,1,0,0),\\
v_5&=(0,0,1,0), & v_6&=(1,-1,0,0),\\
v_7&=(1,0,1,0), & v_8&=(0,1,0,0),\\
v_9&=(1,0,-1,0), & v_{10}&=(1,-1,1,0),\\
v_{11}&=(1,-1,-1,0), & v_{12}&=(1,1,1,0),\\
v_{13}&=(1,1,-1,0), & v_{14}&=(2,1,1,0),\\
v_{15}&=(1,2,-1,0), & v_{16}&=(1,-2,1,0),\\
v_{17}&=(1,1,-2,0), & v_{18}&=(1,-1,-2,0),\\
v_{19}&=(2,1,-1,0), & v_{20}&=(2,-1,1,0),\\
v_{21}&=(1,-2,-1,0), & v_{22}&=(1,1,2,0),\\
v_{23}&=(1,2,1,0), & v_{24}&=(1,-1,2,0),\\
v_{25}&=(2,-1,-1,0), & v_{26}&=(0,0,0,1).
\end{aligned}\tag{S5}$$

These 26 canonical primitive vectors partition by squared norm $\|v_i\|^2$ into four disjoint norm shells $(4+6+4+12=26)$:

- $\|v_i\|^2=1$ (4 vectors): $\{v_2,v_5,v_8,v_{26}\}$, the four coordinate basis vectors of $\mathbb{R}^4$;
- $\|v_i\|^2=2$ (6 vectors): $\{v_1,v_3,v_4,v_6,v_7,v_9\}$, the six binary rays in $\mathbb{R}^3\times\{0\}$;
- $\|v_i\|^2=3$ (4 vectors): $\{v_{10},v_{11},v_{12},v_{13}\}$, the four triad rays $(1,\pm1,\pm1,0)^T$;
- $\|v_i\|^2=6$ (12 vectors): $\{v_{14},\dots,v_{25}\}$, all $3\times4=12$ canonical primitive permutations and sign patterns of $(\pm2,\pm1,\pm1,0)^T$.

Since all 26 canonical 4-tuples are distinct, all $\binom{26}{2}=325$ pairs of vectors are strictly non-collinear, with maximum squared cosine $\max_{1\le i<j\le26}(v_i\cdot v_j)^2/(\|v_i\|^2\|v_j\|^2)=4^2/(3\times6)=8/9\approx0.8889<1$. Table S19 lists all 16 four-element contexts of $E^{(4)}_{16}$ side by side with their four mutually orthogonal integer vectors.

TABLE S19: Complete enumeration of the 16 four-element contexts of the non-linear warm-up apex suspension $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, listing for each context both its vertex indices $\{i, j, k, 26\}$ and the four mutually orthogonal unnormalized integer vectors $\{v_i, v_j, v_k, v_{26}\} \subset \mathbb{Z}^4$.

Context	Vertex set	v_i	v_j	v_k	v_{26}
$e_1^{(4)}$	$\{1,2,3,26\}$	$v_1 = (0,1,1,0)$	$v_2 = (1,0,0,0)$	$v_3 = (0,1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_2^{(4)}$	$\{7,8,9,26\}$	$v_7 = (1,0,1,0)$	$v_8 = (0,1,0,0)$	$v_9 = (1,0,-1,0)$	$v_{26} = (0,0,0,1)$
$e_3^{(4)}$	$\{4,5,6,26\}$	$v_4 = (1,1,0,0)$	$v_5 = (0,0,1,0)$	$v_6 = (1,-1,0,0)$	$v_{26} = (0,0,0,1)$
$e_4^{(4)}$	$\{2,5,8,26\}$	$v_2 = (1,0,0,0)$	$v_5 = (0,0,1,0)$	$v_8 = (0,1,0,0)$	$v_{26} = (0,0,0,1)$
$e_5^{(4)}$	$\{3,12,25,26\}$	$v_3 = (0,1,-1,0)$	$v_{12} = (1,1,1,0)$	$v_{25} = (2,-1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_6^{(4)}$	$\{9,12,16,26\}$	$v_9 = (1,0,-1,0)$	$v_{12} = (1,1,1,0)$	$v_{16} = (1,-2,1,0)$	$v_{26} = (0,0,0,1)$
$e_7^{(4)}$	$\{6,12,17,26\}$	$v_6 = (1,-1,0,0)$	$v_{12} = (1,1,1,0)$	$v_{17} = (1,1,-2,0)$	$v_{26} = (0,0,0,1)$
$e_8^{(4)}$	$\{3,11,14,26\}$	$v_3 = (0,1,-1,0)$	$v_{11} = (1,-1,-1,0)$	$v_{14} = (2,1,1,0)$	$v_{26} = (0,0,0,1)$
$e_9^{(4)}$	$\{7,11,15,26\}$	$v_7 = (1,0,1,0)$	$v_{11} = (1,-1,-1,0)$	$v_{15} = (1,2,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{10}^{(4)}$	$\{4,11,24,26\}$	$v_4 = (1,1,0,0)$	$v_{11} = (1,-1,-1,0)$	$v_{24} = (1,-1,2,0)$	$v_{26} = (0,0,0,1)$
$e_{11}^{(4)}$	$\{9,10,23,26\}$	$v_9 = (1,0,-1,0)$	$v_{10} = (1,-1,1,0)$	$v_{23} = (1,2,1,0)$	$v_{26} = (0,0,0,1)$
$e_{12}^{(4)}$	$\{1,10,19,26\}$	$v_1 = (0,1,1,0)$	$v_{10} = (1,-1,1,0)$	$v_{19} = (2,1,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{13}^{(4)}$	$\{4,10,18,26\}$	$v_4 = (1,1,0,0)$	$v_{10} = (1,-1,1,0)$	$v_{18} = (1,-1,-2,0)$	$v_{26} = (0,0,0,1)$
$e_{14}^{(4)}$	$\{1,13,20,26\}$	$v_1 = (0,1,1,0)$	$v_{13} = (1,1,-1,0)$	$v_{20} = (2,-1,1,0)$	$v_{26} = (0,0,0,1)$
$e_{15}^{(4)}$	$\{7,13,21,26\}$	$v_7 = (1,0,1,0)$	$v_{13} = (1,1,-1,0)$	$v_{21} = (1,-2,-1,0)$	$v_{26} = (0,0,0,1)$
$e_{16}^{(4)}$	$\{6,13,22,26\}$	$v_6 = (1,-1,0,0)$	$v_{13} = (1,1,-1,0)$	$v_{22} = (1,1,2,0)$	$v_{26} = (0,0,0,1)$

S1.16. Explicit parent configurations and inclusion maps

S1.16.1. The 49-ray real Yu–Oh parent and published Bub labels

Table S20 lists all rays, including the original 25. The final column gives the exact labels in the published full 49-ray encoding[1] (p. 14, Sec. 5(xi)) of Bub's set[2]. Under this map the 36 bases in Table S21 equal all 36 published contexts; their 108 pairs equal all geometric orthogonalities. In this labeling 26–33 are Eq. (S31), and 34–49 complete the sixteen remaining orthogonal pairs. The first 25 vectors are those of Table S1.

TABLE S20: All unnormalized vectors of the real 49-ray parent, with the verified published Bub labels.

i	v_i	Published label
1	$(0,1,1)$	`G`
2	$(1,0,0)$	`1`
3	$(0,1,-1)$	`P`
4	$(1,1,0)$	`H`
5	$(0,0,1)$	`3`
6	$(1,-1,0)$	`U`
7	$(1,0,1)$	`Q`
8	$(0,1,0)$	`2`
9	$(1,0,-1)$	`e`
10	$(1,-1,1)$	`D`
11	$(1,-1,-1)$	`h`
12	$(1,1,1)$	`S`
13	$(1,1,-1)$	`a`
14	$(2,1,1)$	`g`
15	$(1,2,-1)$	`n`
16	$(1,-2,1)$	`d`
17	$(1,1,-2)$	`T`
18	$(1,-1,-2)$	`F`
19	$(2,1,-1)$	`E`
20	$(2,-1,1)$	`I`
21	$(1,-2,-1)$	`k`
22	$(1,1,2)$	`b`
23	$(1,2,1)$	`A`

i	v_i	Published label
24	$(1,-1,2)$	N
25	$(2,-1,-1)$	R
26	$(0,1,2)$	i
27	$(0,2,-1)$	O
28	$(0,1,-2)$	6
29	$(0,2,1)$	7
30	$(1,2,0)$	5
31	$(2,-1,0)$	4
32	$(1,-2,0)$	V
33	$(2,1,0)$	j
34	$(2,1,-5)$	Z
35	$(5,-2,1)$	m
36	$(1,2,5)$	W
37	$(5,2,-1)$	c
38	$(1,-2,-5)$	f
39	$(5,-1,2)$	L
40	$(5,1,2)$	9
41	$(2,1,5)$	8
42	$(2,-1,-5)$	J
43	$(5,2,1)$	X
44	$(1,-2,5)$	l
45	$(5,-1,-2)$	Y
46	$(5,-2,-1)$	B
47	$(1,2,-5)$	C
48	$(5,1,-2)$	M
49	$(2,-1,5)$	K

TABLE S21: All 36 complete bases of the real parent.

e	Context	e	Context	e	Context
1	$\{1,2,3\}$	2	$\{1,10,19\}$	3	$\{1,13,20\}$
4	$\{2,5,8\}$	5	$\{2,26,27\}$	6	$\{2,28,29\}$
7	$\{3,11,14\}$	8	$\{3,12,25\}$	9	$\{4,5,6\}$
10	$\{4,10,18\}$	11	$\{4,11,24\}$	12	$\{5,30,31\}$
13	$\{5,32,33\}$	14	$\{6,12,17\}$	15	$\{6,13,22\}$
16	$\{7,8,9\}$	17	$\{7,11,15\}$	18	$\{7,13,21\}$
19	$\{9,10,23\}$	20	$\{9,12,16\}$	21	$\{14,32,34\}$
22	$\{15,26,35\}$	23	$\{15,31,36\}$	24	$\{16,26,37\}$
25	$\{16,33,38\}$	26	$\{17,29,39\}$	27	$\{18,27,40\}$
28	$\{19,32,41\}$	29	$\{20,30,42\}$	30	$\{21,28,43\}$
31	$\{21,33,44\}$	32	$\{22,27,45\}$	33	$\{23,28,46\}$
34	$\{23,31,47\}$	35	$\{24,29,48\}$	36	$\{25,30,49\}$

S1.16.2. The 87-ray Hesse parent

Use $\zeta = \exp(2\pi i/3)$ and $\zeta^2 + \zeta + 1 = 0$. The first 37 rays retain exactly the Hesse numbering; projective representatives have their first nonzero coordinate one. The inventory labels use the generation convention of Sec. VII B. Tables S22 and S23 list all 87 rays and all 64 complete bases. The six additional orthogonal pairs outside the 64 complete bases are listed separately: this larger parent's full orthogonality graph need not equal its context two-section. The retained Hesse restriction has all its original orthogonalities covered by contexts.

TABLE S22: All 87 unnormalized rays of the first Hesse/lineage union.

i	v_i	Inventory label
1	$(1,0,0)$	a_{11}
2	$(1,1,1)$	u_1
3	$(1,\zeta,-1-\zeta)$	u_2
4	$(1,-1-\zeta,\zeta)$	u_3
5	$(1,\zeta,\zeta)$	u_4
6	$(1,-1-\zeta,1)$	u_5
7	$(1,1,-1-\zeta)$	u_6
8	$(1,-1-\zeta,-1-\zeta)$	u_7
9	$(1,1,\zeta)$	u_9
10	$(1,\zeta,1)$	u_8
11	$(0,1,-1)$	c_{11}
12	$(0,1,-\zeta)$	c_{12}
13	$(0,1,1+\zeta)$	c_{13}
14	$(1,0,-1)$	c_{21}
15	$(1,0,1+\zeta)$	c_{22}
16	$(1,0,-\zeta)$	c_{23}
17	$(0,1,1)$	b_{11}
18	$(0,1,\zeta)$	b_{12}
19	$(0,1,-1-\zeta)$	b_{13}
20	$(1,\frac{-1}{2},\frac{-1}{2})$	d_{11}
21	$(1,-\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$	d_{12}
22	$(1,\frac{1}{2}+\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$	d_{13}
23	$(1,-\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$	d_{14}
24	$(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{-1}{2})$	d_{15}
25	$(1,\frac{-1}{2},\frac{1}{2}+\frac{1}{2}\zeta)$	d_{16}
26	$(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$	d_{17}
27	$(1,\frac{-1}{2},-\frac{1}{2}\zeta)$	d_{19}
28	$(1,-\frac{1}{2}\zeta,\frac{-1}{2})$	d_{18}
29	$(1,-2,1)$	d_{21}
30	$(1,-2\zeta,-1-\zeta)$	d_{22}
31	$(1,2+2\zeta,\zeta)$	d_{23}
32	$(1,2+2\zeta,1)$	d_{25}
33	$(1,-2,-1-\zeta)$	d_{26}
34	$(1,-2\zeta,\zeta)$	d_{24}
35	$(1,-2\zeta,1)$	d_{28}
36	$(1,2+2\zeta,-1-\zeta)$	d_{27}
37	$(1,-2,\zeta)$	d_{29}
38	$(0,1,0)$	a_{21}
39	$(0,0,1)$	a_{31}
40	$(1,\frac{-1}{2},\frac{1}{2})$	b_{1121}
41	$(1,-\frac{1}{2}\zeta,\frac{-1}{2}-\frac{1}{2}\zeta)$	b_{1122}
42	$(1,\frac{1}{2}+\frac{1}{2}\zeta,\frac{1}{2}\zeta)$	b_{1123}
43	$(1,\frac{1}{2},\frac{-1}{2})$	b_{1131}
44	$(1,\frac{1}{2}\zeta,\frac{1}{2}+\frac{1}{2}\zeta)$	b_{1132}
45	$(1,\frac{-1}{2}-\frac{1}{2}\zeta,-\frac{1}{2}\zeta)$	b_{1133}
46	$(1,1,-1)$	b_{121}
47	$(1,\zeta,1+\zeta)$	b_{122}
48	$(1,-1-\zeta,-\zeta)$	b_{123}
49	$(1,-1,1)$	b_{131}
50	$(1,-\zeta,-1-\zeta)$	b_{132}
51	$(1,1+\zeta,\zeta)$	b_{133}
52	$(1,0,1)$	b_{21}
53	$(1,-2,-1)$	b_{2121}
54	$(1,-2\zeta,1+\zeta)$	b_{2122}
55	$(1,2+2\zeta,-\zeta)$	b_{2123}
56	$(1,0,-1-\zeta)$	b_{22}
57	$(1,2,-1)$	b_{2231}
58	$(1,2\zeta,1+\zeta)$	b_{2232}
59	$(1,-2-2\zeta,-\zeta)$	b_{2233}
60	$(1,0,\zeta)$	b_{23}
61	$(1,-1,-1)$	b_{231}
62	$(1,-\zeta,1+\zeta)$	b_{232}
63	$(1,1+\zeta,-\zeta)$	b_{233}
64	$(1,1,0)$	b_{31}
65	$(1,-1,-2)$	b_{3131}

i	v_i	Inventory label
66	$(1, -\zeta, 2+2\zeta)$	b_{3132}
67	$(1, 1+\zeta, -2\zeta)$	b_{3133}
68	$(1, \zeta, 0)$	b_{32}
69	$(1, -1, 2)$	b_{3231}
70	$(1, -\zeta, -2-2\zeta)$	b_{3232}
71	$(1, 1+\zeta, 2\zeta)$	b_{3233}
72	$(1, -1-\zeta, 0)$	b_{33}
73	$(1, -1, 0)$	c_{31}
74	$(1, -\zeta, 0)$	c_{32}
75	$(1, 1+\zeta, 0)$	c_{33}
76	$(1, 1, -2)$	d_{31}
77	$(1, \zeta, 2+2\zeta)$	d_{32}
78	$(1, -1-\zeta, -2\zeta)$	d_{33}
79	$(1, 2, 1)$	e_{11}
80	$(1, 2\zeta, -1-\zeta)$	e_{12}
81	$(1, -2-2\zeta, \zeta)$	e_{13}
82	$(1, 1, 2)$	e_{21}
83	$(1, \zeta, -2-2\zeta)$	e_{22}
84	$(1, -1-\zeta, 2\zeta)$	e_{23}
85	$(1, \frac{1}{2}, \frac{1}{2})$	e_{31}
86	$(1, \frac{1}{2}\zeta, \frac{-1}{2} - \frac{1}{2}\zeta)$	e_{32}
87	$(1, \frac{-1}{2} - \frac{1}{2}\zeta, \frac{1}{2}\zeta)$	e_{33}

TABLE S23: All 64 complete bases of the first 87-ray Hesse parent.

e	Context	e	Context	e	Context
1	$\{1, 11, 17\}$	2	$\{1, 12, 18\}$	3	$\{1, 13, 19\}$
4	$\{1, 38, 39\}$	5	$\{2, 3, 4\}$	6	$\{2, 11, 20\}$
7	$\{2, 14, 29\}$	8	$\{2, 73, 76\}$	9	$\{3, 12, 21\}$
10	$\{3, 15, 30\}$	11	$\{3, 74, 77\}$	12	$\{4, 13, 22\}$
13	$\{4, 16, 31\}$	14	$\{4, 75, 78\}$	15	$\{5, 6, 7\}$
16	$\{5, 11, 23\}$	17	$\{5, 16, 34\}$	18	$\{6, 12, 24\}$
19	$\{6, 14, 32\}$	20	$\{7, 13, 25\}$	21	$\{7, 15, 33\}$
22	$\{8, 9, 10\}$	23	$\{8, 11, 26\}$	24	$\{8, 15, 36\}$
25	$\{9, 12, 27\}$	26	$\{9, 16, 37\}$	27	$\{10, 13, 28\}$
28	$\{10, 14, 35\}$	29	$\{11, 61, 85\}$	30	$\{12, 62, 86\}$
31	$\{13, 63, 87\}$	32	$\{14, 38, 52\}$	33	$\{14, 49, 79\}$
34	$\{15, 38, 56\}$	35	$\{15, 50, 80\}$	36	$\{16, 38, 60\}$
37	$\{16, 51, 81\}$	38	$\{17, 40, 46\}$	39	$\{17, 43, 49\}$
40	$\{18, 41, 47\}$	41	$\{18, 44, 50\}$	42	$\{19, 42, 48\}$
43	$\{19, 45, 51\}$	44	$\{39, 64, 73\}$	45	$\{39, 68, 74\}$
46	$\{39, 72, 75\}$	47	$\{46, 47, 48\}$	48	$\{46, 52, 53\}$
49	$\{46, 73, 82\}$	50	$\{47, 54, 56\}$	51	$\{47, 74, 83\}$
52	$\{48, 55, 60\}$	53	$\{48, 75, 84\}$	54	$\{49, 50, 51\}$
55	$\{49, 64, 65\}$	56	$\{50, 66, 68\}$	57	$\{51, 67, 72\}$
58	$\{52, 57, 61\}$	59	$\{56, 58, 62\}$	60	$\{59, 60, 63\}$
61	$\{61, 62, 63\}$	62	$\{61, 64, 69\}$	63	$\{62, 68, 70\}$
64	$\{63, 71, 72\}$				

Its six further orthogonal pairs are $\{5, 74\}$, $\{6, 75\}$, $\{7, 73\}$, $\{8, 75\}$, $\{9, 73\}$, $\{10, 74\}$.

S1.16.3. Peres24 and the complete four-dimensional extensions

The coordinates of H_{65} and $H_{\mathrm{YO}}^{(4)}$ are those already listed above. Table S24 explicitly enumerates the known Peres rays. In each union, retain the original local labels and append the missing rays in the order displayed in Table S25. Tables S26 and S27 list every available complete basis. Full orthogonality-pair lists accompany the coordinates in the data; the 40-ray union has additional pairs outside its complete contexts.

TABLE S24: All 24 primitive unnormalized Peres rays, in the certificate order.

i	p_i
1	$(1,0,0,0)$
2	$(0,1,0,0)$
3	$(0,0,1,0)$
4	$(0,0,0,1)$
5	$(1,-1,0,0)$
6	$(1,1,0,0)$
7	$(1,0,-1,0)$
8	$(1,0,1,0)$
9	$(1,0,0,-1)$
10	$(1,0,0,1)$
11	$(0,1,-1,0)$
12	$(0,1,1,0)$
13	$(0,1,0,-1)$
14	$(0,1,0,1)$
15	$(0,0,1,-1)$
16	$(0,0,1,1)$
17	$(1,-1,-1,-1)$
18	$(1,-1,-1,1)$
19	$(1,-1,1,-1)$
20	$(1,-1,1,1)$
21	$(1,1,-1,-1)$
22	$(1,1,-1,1)$
23	$(1,1,1,-1)$
24	$(1,1,1,1)$

TABLE S25: Every ray added to the original four-dimensional examples to obtain the stated parents.

Original	New label	Added vector
$H_{\mathrm{YO}}^{(4)}$	27	$(1,0,0,-1)$
$H_{\mathrm{YO}}^{(4)}$	28	$(1,0,0,1)$
$H_{\mathrm{YO}}^{(4)}$	29	$(0,1,0,-1)$
$H_{\mathrm{YO}}^{(4)}$	30	$(0,1,0,1)$
$H_{\mathrm{YO}}^{(4)}$	31	$(0,0,1,-1)$
$H_{\mathrm{YO}}^{(4)}$	32	$(0,0,1,1)$
$H_{\mathrm{YO}}^{(4)}$	33	$(1,-1,-1,-1)$
$H_{\mathrm{YO}}^{(4)}$	34	$(1,-1,-1,1)$
$H_{\mathrm{YO}}^{(4)}$	35	$(1,-1,1,-1)$
$H_{\mathrm{YO}}^{(4)}$	36	$(1,-1,1,1)$
$H_{\mathrm{YO}}^{(4)}$	37	$(1,1,-1,-1)$
$H_{\mathrm{YO}}^{(4)}$	38	$(1,1,-1,1)$
$H_{\mathrm{YO}}^{(4)}$	39	$(1,1,1,-1)$
$H_{\mathrm{YO}}^{(4)}$	40	$(1,1,1,1)$
H_{65}	66	$(1,-1,0,0)$
H_{65}	67	$(1,0,-1,0)$
H_{65}	68	$(1,0,0,1)$
H_{65}	69	$(0,1,-1,0)$
H_{65}	70	$(0,1,0,1)$
H_{65}	71	$(0,0,1,1)$
H_{65}	72	$(1,1,1,-1)$

TABLE S26: All 36 complete bases of the 40-ray parent.

e	Context	e	Context
1	$\{1,2,3,26\}$	2	$\{1,3,27,28\}$
3	$\{1,10,19,26\}$	4	$\{1,13,20,26\}$
5	$\{1,27,36,38\}$	6	$\{1,28,35,37\}$
7	$\{2,5,8,26\}$	8	$\{2,5,29,30\}$
9	$\{2,8,31,32\}$	10	$\{3,11,14,26\}$
11	$\{3,12,25,26\}$	12	$\{3,27,34,40\}$
13	$\{3,28,33,39\}$	14	$\{4,5,6,26\}$
15	$\{4,6,31,32\}$	16	$\{4,10,18,26\}$
17	$\{4,11,24,26\}$	18	$\{4,31,33,36\}$
19	$\{4,32,34,35\}$	20	$\{5,8,27,28\}$
21	$\{6,12,17,26\}$	22	$\{6,13,22,26\}$
23	$\{6,31,37,40\}$	24	$\{6,32,38,39\}$
25	$\{7,8,9,26\}$	26	$\{7,9,29,30\}$
27	$\{7,11,15,26\}$	28	$\{7,13,21,26\}$
29	$\{7,29,33,38\}$	30	$\{7,30,34,37\}$
31	$\{9,10,23,26\}$	32	$\{9,12,16,26\}$
33	$\{9,29,35,40\}$	34	$\{9,30,36,39\}$
35	$\{33,36,38,39\}$	36	$\{34,35,37,40\}$

TABLE S27: All 48 complete bases of the 72-ray parent.

e	Context	e	Context
1	$\{1,2,3,4\}$	2	$\{1,2,17,71\}$
3	$\{1,3,15,70\}$	4	$\{1,4,13,69\}$
5	$\{1,13,22,23\}$	6	$\{1,15,30,31\}$
7	$\{1,17,38,39\}$	8	$\{2,3,12,68\}$
9	$\{2,4,14,67\}$	10	$\{2,12,18,19\}$
11	$\{2,14,26,27\}$	12	$\{2,17,40,41\}$
13	$\{3,4,16,66\}$	14	$\{3,12,20,21\}$
15	$\{3,15,32,33\}$	16	$\{3,16,34,35\}$
17	$\{4,13,24,25\}$	18	$\{4,14,28,29\}$
19	$\{4,16,36,37\}$	20	$\{5,6,7,8\}$
21	$\{5,6,17,66\}$	22	$\{5,7,15,67\}$
23	$\{5,8,12,69\}$	24	$\{5,12,42,43\}$
25	$\{5,15,54,55\}$	26	$\{5,17,62,63\}$
27	$\{6,7,13,68\}$	28	$\{6,8,14,70\}$
29	$\{6,13,46,47\}$	30	$\{6,14,50,51\}$
31	$\{6,17,64,65\}$	32	$\{7,8,16,71\}$
33	$\{7,13,48,49\}$	34	$\{7,15,56,57\}$
35	$\{7,16,58,59\}$	36	$\{8,12,44,45\}$
37	$\{8,14,52,53\}$	38	$\{8,16,60,61\}$
39	$\{9,10,11,72\}$	40	$\{9,10,12,13\}$
41	$\{9,11,14,15\}$	42	$\{9,66,71,72\}$
43	$\{10,11,16,17\}$	44	$\{10,67,70,72\}$
45	$\{11,68,69,72\}$	46	$\{12,13,68,69\}$
47	$\{14,15,67,70\}$	48	$\{16,17,66,71\}$

S1.17. Cabello completions and the two 68-ray parents

S1.17.1. The eighteen-ray input and faithful 36-ray completion

All labels in this subsection are local. In $C_{18,9}$ and $C_{36,18}$, $18 = a$, $19 = b$, and $20 = c$; labels 4 and 5 are absent. They do not denote the completion rays of H_{65}. The core is the first eighteen rows of Table S28; labels 21–38 are fresh. The contexts of $C_{18,9}$ are the first nine rows of Table S29.

TABLE S28: All 36 unnormalized vectors of the faithful Cabello completion.

i	v_i
1	$(1,0,0,0)$
2	$(0,1,0,0)$
3	$(0,0,1,0)$
6	$(1,1,-1,-1)$
7	$(1,-1,1,-1)$
8	$(1,-1,-1,1)$
9	$(1,1,-1,1)$
10	$(1,-1,1,1)$
11	$(1,-1,-1,-1)$
12	$(1,0,0,-1)$
13	$(0,1,1,0)$
14	$(1,0,1,0)$
15	$(0,1,0,-1)$
16	$(1,1,0,0)$
17	$(0,0,1,-1)$
18	$(1,0,0,1)$
19	$(0,1,0,1)$
20	$(0,0,1,1)$
21	$(0,1,-1,3)$
22	$(0,3,-3,-2)$
23	$(1,0,-1,3)$
24	$(3,0,-3,-2)$
25	$(1,-1,0,3)$
26	$(3,-3,0,-2)$
27	$(3,1,2,2)$
28	$(3,-5,-1,-1)$
29	$(3,2,1,2)$
30	$(3,-1,-5,-1)$
31	$(2,3,1,2)$
32	$(1,-3,5,1)$
33	$(4,2,3,-3)$
34	$(5,-7,-1,1)$
35	$(4,3,2,-3)$
36	$(5,-1,-7,1)$
37	$(3,4,2,-3)$
38	$(1,-5,7,-1)$

TABLE S29: All eighteen full bases of the faithful Cabello completion.

e	Context
1	$\{9,10,12,13\}$
2	$\{9,11,14,15\}$
3	$\{10,11,16,17\}$
4	$\{2,3,12,18\}$
5	$\{1,3,15,19\}$
6	$\{1,2,17,20\}$
7	$\{6,7,13,18\}$
8	$\{6,8,14,19\}$
9	$\{7,8,16,20\}$
10	$\{1,13,21,22\}$
11	$\{2,14,23,24\}$
12	$\{3,16,25,26\}$
13	$\{6,17,27,28\}$
14	$\{7,15,29,30\}$
15	$\{8,12,31,32\}$
16	$\{9,20,33,34\}$
17	$\{10,19,35,36\}$
18	$\{11,18,37,38\}$

The orthogonal matrix $Q = M/\sqrt{2}$, with

$$M = \begin{pmatrix} 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \qquad M^T M = 2I_4, \tag{S6}$$

maps the published eighteen-ray set and all nine bases of Ref.[3] (Eqs. (1)–(9)) onto this realization. Table S30 gives the complete ray map; the input representatives are unnormalized and signs are immaterial.

TABLE S30: Exact ray map from the originally published Cabello coordinates.

Published vector u	Local label of $[Qu]$
$(0,0,0,1)$	12
$(0,0,1,0)$	13
$(0,0,1,1)$	11
$(0,1,-1,0)$	8
$(0,1,0,-1)$	1
$(0,1,0,0)$	18
$(1,-1,-1,-1)$	19
$(1,-1,-1,1)$	16
$(1,-1,0,0)$	9
$(1,-1,1,-1)$	17
$(1,0,-1,0)$	2
$(1,0,0,-1)$	7
$(1,0,0,1)$	6
$(1,0,1,0)$	3
$(1,1,-1,1)$	15
$(1,1,0,0)$	10
$(1,1,1,-1)$	14
$(1,1,1,1)$	20

S1.17.2. An injective 45-ray completion after deleting c

Delete local ray 20 and contexts 6 and 9 from the first nine contexts above. The seven remaining bases are followed by fourteen fresh-pair contexts in Table S32. The next table prints all 45 vectors; 4, 5, and 20 are absent and 21–48 are fresh. Its only uncovered orthogonal pairs are $\{16,21\}, \{16,22\}, \{17,35\}, \{17,36\}$.

TABLE S31: All 45 unnormalized vectors of the vertex-deleted completion.

i	v_i
1	$(1,0,0,0)$
2	$(0,1,0,0)$
3	$(0,0,1,0)$
6	$(1,1,-1,-1)$
7	$(1,-1,1,-1)$
8	$(1,-1,-1,1)$
9	$(1,1,-1,1)$
10	$(1,-1,1,1)$
11	$(1,-1,-1,-1)$
12	$(1,0,0,-1)$
13	$(0,1,1,0)$
14	$(1,0,1,0)$
15	$(0,1,0,-1)$
16	$(1,1,0,0)$
17	$(0,0,1,-1)$
18	$(1,0,0,1)$
19	$(0,1,0,1)$
21	$(0,0,1,2)$
22	$(0,0,2,-1)$
23	$(0,1,-1,3)$
24	$(0,3,-3,-2)$
25	$(0,1,2,2)$
26	$(0,4,-1,-1)$
27	$(1,0,-1,3)$
28	$(3,0,-3,-2)$
29	$(1,0,2,2)$
30	$(4,0,-1,-1)$
31	$(1,-1,0,3)$
32	$(3,-3,0,-2)$
33	$(5,3,4,4)$
34	$(7,-9,-1,-1)$
35	$(1,1,3,3)$
36	$(3,3,-1,-1)$
37	$(5,4,3,4)$
38	$(7,-1,-9,-1)$
39	$(2,-2,-1,3)$
40	$(1,-1,-5,-3)$
41	$(4,5,3,4)$
42	$(1,-7,9,1)$
43	$(4,-4,-1,-9)$
44	$(5,-5,13,3)$
45	$(6,5,4,-5)$
46	$(9,-1,-11,1)$
47	$(3,4,2,-3)$
48	$(1,-5,7,-1)$

TABLE S32: All 21 full bases of the vertex-deleted completion.

e	Context
1	$\{9, 10, 12, 13\}$
2	$\{9, 11, 14, 15\}$
3	$\{10, 11, 16, 17\}$
4	$\{2, 3, 12, 18\}$
5	$\{1, 3, 15, 19\}$
6	$\{6, 7, 13, 18\}$
7	$\{6, 8, 14, 19\}$
8	$\{1, 2, 21, 22\}$
9	$\{1, 13, 23, 24\}$
10	$\{1, 17, 25, 26\}$
11	$\{2, 14, 27, 28\}$
12	$\{2, 17, 29, 30\}$
13	$\{3, 16, 31, 32\}$
14	$\{6, 17, 33, 34\}$
15	$\{7, 8, 35, 36\}$
16	$\{7, 15, 37, 38\}$
17	$\{7, 16, 39, 40\}$
18	$\{8, 12, 41, 42\}$
19	$\{8, 16, 43, 44\}$
20	$\{10, 19, 45, 46\}$
21	$\{11, 18, 47, 48\}$

S1.17.3. All vectors and bases of the two 68-ray parents

In each parent retain all 65 vectors of Appendix S1.15.1, with their original labels, and add the three vectors in the corresponding rows of Table S33; Tables S34 and S35 list all 35 bases of each parent. Thus the six rows specify every added ray; the two parent inventories use distinct local meanings for labels 66–68. These labels also differ from the 72-ray parent numbering in Appendix S1.16.

TABLE S33: The three added rays in each local 68-ray parent inventory.

Parent variant	i	v_i
18–9 subsystem	66	$(1, 0, 0, 1)$
18–9 subsystem	67	$(0, 1, 0, 1)$
18–9 subsystem	68	$(0, 0, 1, 1)$
20–11 subsystem	66	$(0, 1, -1, 0)$
20–11 subsystem	67	$(1, 0, -1, 0)$
20–11 subsystem	68	$(1, -1, 0, 0)$

TABLE S34: All 35 full bases of 68-ray parent variant 1.

e	Context	e	Context
1	$\{1,2,3,4\}$	2	$\{1,2,17,68\}$
3	$\{1,3,15,67\}$	4	$\{1,13,22,23\}$
5	$\{1,15,30,31\}$	6	$\{1,17,38,39\}$
7	$\{2,3,12,66\}$	8	$\{2,12,18,19\}$
9	$\{2,14,26,27\}$	10	$\{2,17,40,41\}$
11	$\{3,12,20,21\}$	12	$\{3,15,32,33\}$
13	$\{3,16,34,35\}$	14	$\{4,13,24,25\}$
15	$\{4,14,28,29\}$	16	$\{4,16,36,37\}$
17	$\{5,6,7,8\}$	18	$\{5,12,42,43\}$
19	$\{5,15,54,55\}$	20	$\{5,17,62,63\}$
21	$\{6,7,13,66\}$	22	$\{6,8,14,67\}$
23	$\{6,13,46,47\}$	24	$\{6,14,50,51\}$
25	$\{6,17,64,65\}$	26	$\{7,8,16,68\}$
27	$\{7,13,48,49\}$	28	$\{7,15,56,57\}$
29	$\{7,16,58,59\}$	30	$\{8,12,44,45\}$
31	$\{8,14,52,53\}$	32	$\{8,16,60,61\}$
33	$\{9,10,12,13\}$	34	$\{9,11,14,15\}$
35	$\{10,11,16,17\}$		

TABLE S35: All 35 full bases of 68-ray parent variant 2.

e	Context	e	Context
1	$\{1,2,3,4\}$	2	$\{1,4,13,66\}$
3	$\{1,13,22,23\}$	4	$\{1,15,30,31\}$
5	$\{1,17,38,39\}$	6	$\{2,4,14,67\}$
7	$\{2,12,18,19\}$	8	$\{2,14,26,27\}$
9	$\{2,17,40,41\}$	10	$\{3,4,16,68\}$
11	$\{3,12,20,21\}$	12	$\{3,15,32,33\}$
13	$\{3,16,34,35\}$	14	$\{4,13,24,25\}$
15	$\{4,14,28,29\}$	16	$\{4,16,36,37\}$
17	$\{5,6,7,8\}$	18	$\{5,6,17,68\}$
19	$\{5,7,15,67\}$	20	$\{5,8,12,66\}$
21	$\{5,12,42,43\}$	22	$\{5,15,54,55\}$
23	$\{5,17,62,63\}$	24	$\{6,13,46,47\}$
25	$\{6,14,50,51\}$	26	$\{6,17,64,65\}$
27	$\{7,13,48,49\}$	28	$\{7,15,56,57\}$
29	$\{7,16,58,59\}$	30	$\{8,12,44,45\}$
31	$\{8,14,52,53\}$	32	$\{8,16,60,61\}$
33	$\{9,10,12,13\}$	34	$\{9,11,14,15\}$
35	$\{10,11,16,17\}$		

For variant 1 the nine-context parity subsystem consists of exactly $C_3, C_4, C_5, D_1, \ldots, D_6$ with $a = 66, b = 67, c = 68$. For variant 2 the twenty-ray subsystem consists of all rows whose labels lie in $\{1, \ldots, 17, 66, 67, 68\}$. The machine-readable inventories repeat every coordinate, basis, and inclusion map.

S2. TWO-VALUED STATES, EXTENDABILITY CERTIFICATES, COLORINGS, AND SEPARATION PROFILES

S2.1. State, coloring, odd-cycle certificate, and profile tables for chromatically complete configurations ($\chi(H) = d$)

This subsection provides the complete state and coloring enumerations and odd-cycle nonextendability certificates for $H_{6,3}$ (Table S36), P_5 (Table S37), H_8 (Tables S38 and S39), $H_{17,10}$ (Tables S40 and S41), $H_{21,13}$ (Tables S42, S43, and S44), the vertex separation profiles and Born probability decomposition on $H_{17,10}$ and $H_{21,13}$ (Tables S45–S47), the complete state and normalized coloring tables for all faithful 3- and 4-dimensional logical gadgets (Tables S48–S69 and S70), and the ten chromatically inseparable faithful restrictions of H_{YO}^{+} (Table S71).

TABLE S36: All two-valued states of $H_{6,3}$. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4\}$	yes	–
2	$\{2,4,6\}$	no	$(3,1,5)$
3	$\{2,5\}$	yes	–
4	$\{3,6\}$	yes	–

TABLE S37: All two-valued states of P_5. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,3,9\}$	yes	–
2	$\{1,4,7\}$	yes	–
3	$\{1,7,8,9\}$	yes	–
4	$\{2,4,10\}$	yes	–
5	$\{2,5,8\}$	yes	–
6	$\{2,8,9,10\}$	yes	–
7	$\{3,5,6\}$	yes	–
8	$\{3,6,9,10\}$	yes	–
9	$\{4,6,7,10\}$	yes	–
10	$\{5,6,7,8\}$	yes	–
11	$\{6,7,8,9,10\}$	no	$(3,2,1,5,4)$

TABLE S38: All two-valued states of H_8. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4,6,9\}$	yes	–
2	$\{1,5,8,10,14\}$	no	$(9,7,6,12,11)$
3	$\{1,5,9,13,14\}$	yes	–
4	$\{2,4,6,8,11\}$	yes	–
5	$\{2,4,7,11,14\}$	no	$(5,3,1,12,6)$
6	$\{2,5,8,10,12\}$	yes	–
7	$\{2,5,8,11,13,14\}$	yes	–
8	$\{2,5,9,12,13\}$	no	$(4,3,1,11,10)$
9	$\{3,6,8,11,13\}$	no	$(7,5,4,10,9)$
10	$\{3,7,10,12\}$	yes	–
11	$\{3,7,11,13,14\}$	yes	–

TABLE S39. All strong three-colorings of H_8 up to color permutation. The anchor is $(1,2,3) = (0,1,2)$.

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14
c_1	0	1	2	0	1	0	2	1	0	1	2	1	2	2
c_2	0	1	2	0	1	0	2	1	0	2	1	2	1	1
c_3	0	1	2	1	0	1	2	1	0	2	1	2	0	0

TABLE S40: All two-valued states of $H_{17,10}$. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4,8,14,15,23\}$	yes	–
2	$\{1,5,7,14,18,23,24\}$	no	$(4,11,3,2,8,9,10)$
3	$\{1,5,9,11,18\}$	yes	–
4	$\{1,5,9,14,15,18,24\}$	no	$(11,3,2,8,7)$
5	$\{1,6,8,11,18,23\}$	yes	–
6	$\{1,6,8,14,15,18,23,24\}$	no	$(11,3,2,5,4)$
7	$\{2,4,7,14,19,23\}$	yes	–
8	$\{2,4,9,14,15,19\}$	yes	–
9	$\{2,6,7,10,14,24\}$	yes	–
10	$\{2,6,7,14,18,19,23,24\}$	no	$(11,3,1,10,4)$
11	$\{2,6,9,11,18,19\}$	yes	–
12	$\{2,6,9,14,15,18,19,24\}$	no	$(11,3,1,10,4)$
13	$\{3,4,8,15,19,23\}$	yes	–
14	$\{3,5,7,10,24\}$	yes	–
15	$\{3,5,7,18,19,23,24\}$	no	$(8,2,1,10,9)$
16	$\{3,5,9,15,18,19,24\}$	no	$(7,8,2,1,10,4,11)$
17	$\{3,6,8,10,15,24\}$	yes	–
18	$\{3,6,8,15,18,19,23,24\}$	no	$(5,2,1,10,4)$

TABLE S41. All strong three-colorings of $H_{17,10}$ up to color permutation. The anchor is $(2,8,5) = (0,1,2)$.

i	1	2	3	4	5	6	7	8	9	10	11	14	15	18	19	23	24
c_1	1	0	2	0	2	1	2	1	0	2	1	0	0	1	0	1	2
c_2	1	0	2	1	2	0	2	1	0	2	0	1	1	0	0	1	2
c_3	2	0	1	0	2	1	0	1	2	1	2	0	1	2	0	0	1
c_4	2	0	1	1	2	0	0	1	2	0	2	0	1	2	1	1	0

TABLE S42: All two-valued states of $H_{21,13}$. The last column gives an odd cycle in the complement of each nonextendable support.

j	support of s_j	extendable	odd cycle
1	$\{1,4,8,14,15,21,22,23\}$	no	$(7,11,3,2,5,6,13)$
2	$\{1,5,7,14,18,22,23,24\}$	no	$(4,11,3,2,8,9,10)$
3	$\{1,5,9,11,18,21,22\}$	yes	–
4	$\{1,5,9,14,15,18,21,22,24\}$	no	$(11,3,2,8,7)$
5	$\{1,6,8,11,18,21,23\}$	yes	–
6	$\{1,6,8,14,15,18,21,23,24\}$	no	$(11,3,2,5,4)$
7	$\{2,4,7,14,19,20,22,23\}$	no	$(8,9,10,1,13,6,5)$
8	$\{2,4,9,13,14,15,19\}$	yes	–
9	$\{2,4,9,14,15,19,20,21,22\}$	no	$(11,3,1,13,7)$
10	$\{2,6,7,10,14,20,24\}$	yes	–
11	$\{2,6,7,14,18,19,20,23,24\}$	no	$(11,3,1,10,4)$
12	$\{2,6,9,11,18,19,20,21\}$	no	$(5,4,10,1,13,7,8)$
13	$\{2,6,9,14,15,18,19,20,21,24\}$	no	$(11,3,1,10,4)$
14	$\{3,4,8,13,15,19,23\}$	yes	–
15	$\{3,4,8,15,19,20,21,22,23\}$	no	$(5,2,1,13,6)$
16	$\{3,5,7,10,20,22,24\}$	yes	–
17	$\{3,5,7,18,19,20,22,23,24\}$	no	$(8,2,1,10,9)$
18	$\{3,5,9,13,15,18,19,24\}$	no	$(7,8,2,1,10,4,11)$
19	$\{3,5,9,15,18,19,20,21,22,24\}$	no	$(8,2,1,13,7)$
20	$\{3,6,8,10,15,20,21,24\}$	no	$(4,5,2,1,13,7,11)$
21	$\{3,6,8,15,18,19,20,21,23,24\}$	no	$(5,2,1,10,4)$

TABLE S43. All strong three-colorings of $H_{21,13}$ up to color permutation. The anchor is $(2,8,5) = (0,1,2)$.

i	1	2	3	4	5	6	7	8	9	10	11	13	14	15	18	19	20	21	22	23	24
c_1	1	0	2	0	2	1	2	1	0	2	1	0	0	0	1	0	2	1	2	1	2
c_2	2	0	1	1	2	0	0	1	2	0	2	1	0	1	2	1	0	2	2	1	0

TABLE S44. The nine chromatic signatures of $H_{21,13}$. Columns give c_1 and rows give c_2 for the two normalized colorings in Table S43. Each cell is one inseparability class: its vertices have the same color in every admissible coloring. Ordinary two-valued states distinguish every pair, including vertices in the same cell.

$c_2 \backslash c_1$	0	1	2
2	$\{9\}$	$\{1,11,18,21\}$	$\{5,22\}$
1	$\{4,13,15,19\}$	$\{8,23\}$	$\{3\}$
0	$\{2,14\}$	$\{6\}$	$\{7,10,20,24\}$

TABLE S46. Valid states and exact weights reproducing two $H_{17,10}$ preparations: the five-state mixture for P_{24} (upper block, Eq. (27)) and the four-state mixture for P_{z_3}, the preparation along $z_3 = r_5$ (lower block). State numbering is that of Table S40. Every context equation and every Born marginal are checked with rational arithmetic.

State	Weight	Support	Extendable?
Preparation P_{24}, nonextendable weight $\frac{1}{9}$			
s_4	$\frac{1}{12}$	$\{1, 5, 9, 14, 15, 18, 24\}$	no
s_9	$\frac{5}{36}$	$\{2, 6, 7, 10, 14, 24\}$	yes
s_{10}	$\frac{1}{36}$	$\{2, 6, 7, 14, 18, 19, 23, 24\}$	no
s_{14}	$\frac{7}{12}$	$\{3, 5, 7, 10, 24\}$	yes
s_{17}	$\frac{1}{6}$	$\{3, 6, 8, 10, 15, 24\}$	yes
Preparation P_{z_3}, nonextendable weight $\frac{1}{3}$			
s_3	$\frac{1}{3}$	$\{1, 5, 9, 11, 18\}$	yes
s_4	$\frac{1}{6}$	$\{1, 5, 9, 14, 15, 18, 24\}$	no
s_{14}	$\frac{1}{3}$	$\{3, 5, 7, 10, 24\}$	yes
s_{15}	$\frac{1}{6}$	$\{3, 5, 7, 18, 19, 23, 24\}$	no

TABLE S45: Complete state profiles in hexadecimal, and all Born probabilities for the 17-ray preparation in Eq. (26). The probabilities are also reproduced by Eq. (27).

i	$H_{17,10}$ profile	$H_{21,13}$ profile	p_i on $H_{17,10}$
1	3F	3F	1/12
2	FC0	1FC0	1/6
3	3F000	1FE000	3/4
4	10C1	61C1	0
5	E00E	7800E	2/3
6	30F30	181E30	1/3
7	6342	18642	3/4
8	31031	186031	1/6
9	8C8C	6198C	1/12
10	12100	88200	8/9
11	414	814	0
13	-	22080	–
14	BEB	17EB	1/4
15	398A9	1E71A9	1/4
18	2CE3E	171C3E	1/9
19	2DEC0	177DC0	1/36
20	-	1DDF40	–
21	-	1C593D	–
22	-	5C14F	–
23	25273	116473	1/36
24	3EB2A	1F962A	1

TABLE S47. Exact Born probabilities $p_i = (r_{24} \cdot r_i)^2/(6\|r_i\|^2)$ for P_{24} and $p'_i = (r_5 \cdot r_i)^2/\|r_i\|^2$ for P_{z_3}, also reproduced by the two mixtures of Table S46.

i	p_i	p'_i	i	p_i	p'_i
1	$\frac{1}{12}$	$\frac{1}{2}$	10	$\frac{8}{9}$	$\frac{1}{3}$
2	$\frac{1}{6}$	0	11	0	$\frac{1}{3}$
3	$\frac{3}{4}$	$\frac{1}{2}$	14	$\frac{1}{4}$	$\frac{1}{6}$
4	0	0	15	$\frac{1}{4}$	$\frac{1}{6}$
5	$\frac{2}{3}$	1	18	$\frac{1}{9}$	$\frac{2}{3}$
6	$\frac{1}{3}$	0	19	$\frac{1}{36}$	$\frac{1}{6}$
7	$\frac{3}{4}$	$\frac{1}{2}$	23	$\frac{1}{36}$	$\frac{1}{6}$
8	$\frac{1}{6}$	0	24	1	$\frac{2}{3}$
9	$\frac{1}{12}$	$\frac{1}{2}$			

S2.2. State conventions for the additional gadgets

The state tables below are local to each labeled configuration. Supports are in lexicographic order; E is yes precisely for color-class supports. The three-dimensional pentagon and $H_{21,13}$ use the complete earlier tables and are not repeated. The four-dimensional apex-only states are included explicitly. All ordinary state families listed here are unital and separating.

S2.3. All states of $B_{13,7}$

TABLE S48: All two-valued states of $B_{13,7}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,5,7,14,23\}$	yes	8	$\{2,9,11,19\}$	yes
2	$\{1,5,9,11\}$	yes	9	$\{2,9,14,15,19\}$	yes
3	$\{1,5,9,14,15\}$	no	10	$\{3,5,7,10\}$	yes
4	$\{1,8,11,23\}$	yes	11	$\{3,5,7,19,23\}$	no
5	$\{1,8,14,15,23\}$	yes	12	$\{3,5,9,15,19\}$	yes
6	$\{2,7,10,14\}$	yes	13	$\{3,8,10,15\}$	yes
7	$\{2,7,14,19,23\}$	yes	14	$\{3,8,15,19,23\}$	yes

S2.4. All states of $D_{16,9}$

TABLE S49: All two-valued states of $D_{16,9}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,4,8,14,15,23\}$	yes	12	$\{2,6,7,10,14,24\}$	yes
2	$\{1,5,7,10,14,24\}$	yes	13	$\{2,6,7,14,18,23,24\}$	no
3	$\{1,5,7,14,18,23,24\}$	no	14	$\{2,6,9,11,18\}$	yes
4	$\{1,5,9,11,18\}$	yes	15	$\{2,6,9,14,15,18,24\}$	no
5	$\{1,5,9,14,15,18,24\}$	no	16	$\{3,4,8,15,23\}$	yes
6	$\{1,6,8,10,11\}$	yes	17	$\{3,5,7,10,24\}$	yes
7	$\{1,6,8,10,14,15,24\}$	no	18	$\{3,5,7,18,23,24\}$	yes
8	$\{1,6,8,11,18,23\}$	yes	19	$\{3,5,9,15,18,24\}$	yes
9	$\{1,6,8,14,15,18,23,24\}$	no	20	$\{3,6,8,10,15,24\}$	yes
10	$\{2,4,7,14,23\}$	yes	21	$\{3,6,8,15,18,23,24\}$	no
11	$\{2,4,9,14,15\}$	yes			

S2.5. All states of $I_{16,9}$

TABLE S50: All two-valued states of $I_{16,9}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,4,5,7,14,23\}$	yes	12	$\{2,7,14,18,19,23,24\}$	no
2	$\{1,4,5,9,14,15\}$	no	13	$\{2,9,11,18,19\}$	yes
3	$\{1,4,8,14,15,23\}$	yes	14	$\{2,9,14,15,18,19,24\}$	no
4	$\{1,5,7,14,18,23,24\}$	no	15	$\{3,4,5,7,19,23\}$	no
5	$\{1,5,9,11,18\}$	yes	16	$\{3,4,5,9,15,19\}$	yes
6	$\{1,5,9,14,15,18,24\}$	no	17	$\{3,4,8,15,19,23\}$	yes
7	$\{1,8,11,18,23\}$	yes	18	$\{3,5,7,10,24\}$	yes
8	$\{1,8,14,15,18,23,24\}$	no	19	$\{3,5,7,18,19,23,24\}$	no
9	$\{2,4,7,14,19,23\}$	yes	20	$\{3,5,9,15,18,19,24\}$	no
10	$\{2,4,9,14,15,19\}$	yes	21	$\{3,8,10,15,24\}$	yes
11	$\{2,7,10,14,24\}$	yes	22	$\{3,8,15,18,19,23,24\}$	no

S2.6. All states of $T_{19,11}$

TABLE S51: All two-valued states of $T_{19,11}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,4,8,14,15,23,28\}$	yes	14	$\{2,6,7,14,18,23,24\}$	no
2	$\{1,5,7,10,14,24,28\}$	yes	15	$\{2,6,9,11,18,27\}$	yes
3	$\{1,5,7,14,18,23,24\}$	no	16	$\{2,6,9,14,15,18,24\}$	no
4	$\{1,5,9,11,18,27\}$	yes	17	$\{3,4,8,15,23,26\}$	yes
5	$\{1,5,9,14,15,18,24\}$	no	18	$\{3,4,8,15,23,27,28\}$	no
6	$\{1,6,8,10,11,26\}$	yes	19	$\{3,5,7,10,24,26\}$	yes
7	$\{1,6,8,10,11,27,28\}$	no	20	$\{3,5,7,10,24,27,28\}$	no
8	$\{1,6,8,10,14,15,24,28\}$	no	21	$\{3,5,7,18,23,24,27\}$	yes
9	$\{1,6,8,11,18,23,27\}$	yes	22	$\{3,5,9,15,18,24,27\}$	yes
10	$\{1,6,8,14,15,18,23,24\}$	no	23	$\{3,6,8,10,15,24,26\}$	yes
11	$\{2,4,7,14,23,28\}$	yes	24	$\{3,6,8,10,15,24,27,28\}$	no
12	$\{2,4,9,14,15,28\}$	yes	25	$\{3,6,8,15,18,23,24,27\}$	no
13	$\{2,6,7,10,14,24,28\}$	yes			

S2.7. All states of $W^{(4)}_{11,5}$

TABLE S52: All two-valued states of $W^{(4)}_{11,5}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0\}$	yes	7	$\{2,8,9,10\}$	yes
2	$\{1,3,9\}$	yes	8	$\{3,5,6\}$	yes
3	$\{1,4,7\}$	yes	9	$\{3,6,9,10\}$	yes
4	$\{1,7,8,9\}$	yes	10	$\{4,6,7,10\}$	yes
5	$\{2,4,10\}$	yes	11	$\{5,6,7,8\}$	yes
6	$\{2,5,8\}$	yes	12	$\{6,7,8,9,10\}$	no

S2.8. All states of $W^{(4)}_{16,6}$

TABLE S53: All two-valued states of $W^{(4)}_{16,6}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0,7,9,10\}$	yes	37	$\{2,8,12,14,15\}$	yes
2	$\{0,7,9,15\}$	yes	38	$\{3,5,6\}$	yes
3	$\{0,7,10,14\}$	yes	39	$\{3,6,9,10,11\}$	yes
4	$\{0,7,14,15\}$	yes	40	$\{3,6,9,10,12\}$	yes
5	$\{0,9,10,13\}$	yes	41	$\{3,6,9,11,15\}$	yes
6	$\{0,9,13,15\}$	yes	42	$\{3,6,9,12,15\}$	yes
7	$\{0,10,13,14\}$	yes	43	$\{3,6,10,11,14\}$	yes
8	$\{0,13,14,15\}$	yes	44	$\{3,6,10,12,14\}$	yes
9	$\{1,3,9,11\}$	yes	45	$\{3,6,11,14,15\}$	yes
10	$\{1,3,9,12\}$	yes	46	$\{3,6,12,14,15\}$	yes
11	$\{1,3,11,14\}$	yes	47	$\{4,6,7,10,11\}$	yes
12	$\{1,3,12,14\}$	yes	48	$\{4,6,7,10,12\}$	yes
13	$\{1,4,7,11\}$	yes	49	$\{4,6,7,11,15\}$	yes
14	$\{1,4,7,12\}$	yes	50	$\{4,6,7,12,15\}$	yes
15	$\{1,4,11,13\}$	yes	51	$\{4,6,10,11,13\}$	yes
16	$\{1,4,12,13\}$	yes	52	$\{4,6,10,12,13\}$	yes
17	$\{1,7,8,9,11\}$	yes	53	$\{4,6,11,13,15\}$	yes
18	$\{1,7,8,9,12\}$	yes	54	$\{4,6,12,13,15\}$	yes
19	$\{1,7,8,11,14\}$	yes	55	$\{5,6,7,8\}$	yes
20	$\{1,7,8,12,14\}$	yes	56	$\{5,6,8,13\}$	yes
21	$\{1,8,9,11,13\}$	yes	57	$\{6,7,8,9,10,11\}$	no
22	$\{1,8,9,12,13\}$	yes	58	$\{6,7,8,9,10,12\}$	no
23	$\{1,8,11,13,14\}$	yes	59	$\{6,7,8,9,11,15\}$	no
24	$\{1,8,12,13,14\}$	yes	60	$\{6,7,8,9,12,15\}$	no
25	$\{2,4,10,11\}$	yes	61	$\{6,7,8,10,11,14\}$	no
26	$\{2,4,10,12\}$	yes	62	$\{6,7,8,10,12,14\}$	no
27	$\{2,4,11,15\}$	yes	63	$\{6,7,8,11,14,15\}$	no
28	$\{2,4,12,15\}$	yes	64	$\{6,7,8,12,14,15\}$	no
29	$\{2,5,8\}$	yes	65	$\{6,8,9,10,11,13\}$	no
30	$\{2,8,9,10,11\}$	yes	66	$\{6,8,9,10,12,13\}$	no
31	$\{2,8,9,10,12\}$	yes	67	$\{6,8,9,11,13,15\}$	no
32	$\{2,8,9,11,15\}$	yes	68	$\{6,8,9,12,13,15\}$	no
33	$\{2,8,9,12,15\}$	yes	69	$\{6,8,10,11,13,14\}$	no
34	$\{2,8,10,11,14\}$	yes	70	$\{6,8,10,12,13,14\}$	no
35	$\{2,8,10,12,14\}$	yes	71	$\{6,8,11,13,14,15\}$	no
36	$\{2,8,11,14,15\}$	yes	72	$\{6,8,12,13,14,15\}$	no

S2.9. All states of $B_{14,7}^{(4)}$

TABLE S54: All two-valued states of $B_{14,7}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0\}$	yes	9	$\{2,9,11,19\}$	yes
2	$\{1,5,7,14,23\}$	yes	10	$\{2,9,14,15,19\}$	yes
3	$\{1,5,9,11\}$	yes	11	$\{3,5,7,10\}$	yes
4	$\{1,5,9,14,15\}$	no	12	$\{3,5,7,19,23\}$	no
5	$\{1,8,11,23\}$	yes	13	$\{3,5,9,15,19\}$	yes
6	$\{1,8,14,15,23\}$	yes	14	$\{3,8,10,15\}$	yes
7	$\{2,7,10,14\}$	yes	15	$\{3,8,15,19,23\}$	yes
8	$\{2,7,14,19,23\}$	yes			

S2.10. All states of $L_{18,7}^{(4)}$

TABLE S55: All two-valued states of $L_{18,7}^{(4)}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,6,9,13,16\}$	yes	35	$\{3,5,9,13,18\}$	yes
2	$\{1,6,9,14,16\}$	yes	36	$\{3,5,9,14,17\}$	yes
3	$\{1,6,10,13,16\}$	yes	37	$\{3,5,9,14,18\}$	yes
4	$\{1,6,10,14,16\}$	yes	38	$\{3,5,10,13,17\}$	yes
5	$\{1,7,9,13,17\}$	yes	39	$\{3,5,10,13,18\}$	yes
6	$\{1,7,9,13,18\}$	yes	40	$\{3,5,10,14,17\}$	yes
7	$\{1,7,9,14,17\}$	yes	41	$\{3,5,10,14,18\}$	yes
8	$\{1,7,9,14,18\}$	yes	42	$\{3,6,9,11\}$	yes
9	$\{1,7,10,13,17\}$	yes	43	$\{3,6,9,12,13\}$	yes
10	$\{1,7,10,13,18\}$	yes	44	$\{3,6,9,12,14\}$	yes
11	$\{1,7,10,14,17\}$	yes	45	$\{3,6,10,11\}$	yes
12	$\{1,7,10,14,18\}$	yes	46	$\{3,6,10,12,13\}$	yes
13	$\{1,8,13,15\}$	yes	47	$\{3,6,10,12,14\}$	yes
14	$\{1,8,13,16,17\}$	yes	48	$\{3,8,11,17\}$	yes
15	$\{1,8,13,16,18\}$	yes	49	$\{3,8,11,18\}$	yes
16	$\{1,8,14,15\}$	yes	50	$\{3,8,12,13,17\}$	yes
17	$\{1,8,14,16,17\}$	yes	51	$\{3,8,12,13,18\}$	yes
18	$\{1,8,14,16,18\}$	yes	52	$\{3,8,12,14,17\}$	yes
19	$\{2,5,9,15\}$	yes	53	$\{3,8,12,14,18\}$	yes
20	$\{2,5,9,16,17\}$	yes	54	$\{4,5,13,15\}$	yes
21	$\{2,5,9,16,18\}$	yes	55	$\{4,5,13,16,17\}$	yes
22	$\{2,5,10,15\}$	yes	56	$\{4,5,13,16,18\}$	yes
23	$\{2,5,10,16,17\}$	yes	57	$\{4,5,14,15\}$	yes
24	$\{2,5,10,16,18\}$	yes	58	$\{4,5,14,16,17\}$	yes
25	$\{2,6,9,12,16\}$	yes	59	$\{4,5,14,16,18\}$	yes
26	$\{2,6,10,12,16\}$	yes	60	$\{4,6,11,16\}$	yes
27	$\{2,7,9,12,17\}$	yes	61	$\{4,6,12,13,16\}$	yes
28	$\{2,7,9,12,18\}$	yes	62	$\{4,6,12,14,16\}$	yes
29	$\{2,7,10,12,17\}$	yes	63	$\{4,7,11,17\}$	yes
30	$\{2,7,10,12,18\}$	yes	64	$\{4,7,11,18\}$	yes
31	$\{2,8,12,15\}$	yes	65	$\{4,7,12,13,17\}$	yes
32	$\{2,8,12,16,17\}$	yes	66	$\{4,7,12,13,18\}$	yes
33	$\{2,8,12,16,18\}$	yes	67	$\{4,7,12,14,17\}$	yes
34	$\{3,5,9,13,17\}$	yes	68	$\{4,7,12,14,18\}$	yes

S2.11. All states of $D^{(4)}_{17,9}$

TABLE S56: All two-valued states of $D^{(4)}_{17,9}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{0\}$	yes	12	$\{2,4,9,14,15\}$	yes
2	$\{1,4,8,14,15,23\}$	yes	13	$\{2,6,7,10,14,24\}$	yes
3	$\{1,5,7,10,14,24\}$	yes	14	$\{2,6,7,14,18,23,24\}$	no
4	$\{1,5,7,14,18,23,24\}$	no	15	$\{2,6,9,11,18\}$	yes
5	$\{1,5,9,11,18\}$	yes	16	$\{2,6,9,14,15,18,24\}$	no
6	$\{1,5,9,14,15,18,24\}$	no	17	$\{3,4,8,15,23\}$	yes
7	$\{1,6,8,10,11\}$	yes	18	$\{3,5,7,10,24\}$	yes
8	$\{1,6,8,10,14,15,24\}$	no	19	$\{3,5,7,18,23,24\}$	yes
9	$\{1,6,8,11,18,23\}$	yes	20	$\{3,5,9,15,18,24\}$	yes
10	$\{1,6,8,14,15,18,23,24\}$	no	21	$\{3,6,8,10,15,24\}$	yes
11	$\{2,4,7,14,23\}$	yes	22	$\{3,6,8,15,18,23,24\}$	no

S2.12. All states of $L^{(4)}_{22,9}$

TABLE S57: All two-valued states of $L^{(4)}_{22,9}$. E indicates extendability.

j	Support	E	j	Support	E
1	$\{1,6,9,13,16,21\}$	yes	80	$\{2,8,12,16,17,20,21\}$	no
2	$\{1,6,9,13,16,22\}$	yes	81	$\{2,8,12,16,17,20,22\}$	no
3	$\{1,6,9,14,16,19\}$	yes	82	$\{2,8,12,16,18,19\}$	yes
4	$\{1,6,9,14,16,20,21\}$	no	83	$\{2,8,12,16,18,20,21\}$	no
5	$\{1,6,9,14,16,20,22\}$	no	84	$\{2,8,12,16,18,20,22\}$	no
6	$\{1,6,10,13,16,21\}$	yes	85	$\{3,5,9,13,17,21\}$	yes
7	$\{1,6,10,13,16,22\}$	yes	86	$\{3,5,9,13,17,22\}$	yes
8	$\{1,6,10,14,16,19\}$	yes	87	$\{3,5,9,13,18,21\}$	yes
9	$\{1,6,10,14,16,20,21\}$	no	88	$\{3,5,9,13,18,22\}$	yes
10	$\{1,6,10,14,16,20,22\}$	no	89	$\{3,5,9,14,17,19\}$	yes
11	$\{1,7,9,13,17,21\}$	yes	90	$\{3,5,9,14,17,20,21\}$	no
12	$\{1,7,9,13,17,22\}$	yes	91	$\{3,5,9,14,17,20,22\}$	no
13	$\{1,7,9,13,18,21\}$	yes	92	$\{3,5,9,14,18,19\}$	yes
14	$\{1,7,9,13,18,22\}$	yes	93	$\{3,5,9,14,18,20,21\}$	no
15	$\{1,7,9,14,17,19\}$	yes	94	$\{3,5,9,14,18,20,22\}$	no
16	$\{1,7,9,14,17,20,21\}$	no	95	$\{3,5,10,13,17,21\}$	yes
17	$\{1,7,9,14,17,20,22\}$	no	96	$\{3,5,10,13,17,22\}$	yes
18	$\{1,7,9,14,18,19\}$	yes	97	$\{3,5,10,13,18,21\}$	yes
19	$\{1,7,9,14,18,20,21\}$	no	98	$\{3,5,10,13,18,22\}$	yes
20	$\{1,7,9,14,18,20,22\}$	no	99	$\{3,5,10,14,17,19\}$	yes
21	$\{1,7,10,13,17,21\}$	yes	100	$\{3,5,10,14,17,20,21\}$	no
22	$\{1,7,10,13,17,22\}$	yes	101	$\{3,5,10,14,17,20,22\}$	no
23	$\{1,7,10,13,18,21\}$	yes	102	$\{3,5,10,14,18,19\}$	yes
24	$\{1,7,10,13,18,22\}$	yes	103	$\{3,5,10,14,18,20,21\}$	no
25	$\{1,7,10,14,17,19\}$	yes	104	$\{3,5,10,14,18,20,22\}$	no
26	$\{1,7,10,14,17,20,21\}$	no	105	$\{3,6,9,11,20\}$	yes
27	$\{1,7,10,14,17,20,22\}$	no	106	$\{3,6,9,12,13,21\}$	yes
28	$\{1,7,10,14,18,19\}$	yes	107	$\{3,6,9,12,13,22\}$	yes
29	$\{1,7,10,14,18,20,21\}$	no	108	$\{3,6,9,12,14,19\}$	yes
30	$\{1,7,10,14,18,20,22\}$	no	109	$\{3,6,9,12,14,20,21\}$	no
31	$\{1,8,13,16,17,21\}$	yes	110	$\{3,6,9,12,14,20,22\}$	no
32	$\{1,8,13,16,17,22\}$	yes	111	$\{3,6,10,11,20\}$	yes
33	$\{1,8,13,16,18,21\}$	yes	112	$\{3,6,10,12,13,21\}$	yes
34	$\{1,8,13,16,18,22\}$	yes	113	$\{3,6,10,12,13,22\}$	yes
35	$\{1,8,14,15,21\}$	yes	114	$\{3,6,10,12,14,19\}$	yes
36	$\{1,8,14,15,22\}$	yes	115	$\{3,6,10,12,14,20,21\}$	no
37	$\{1,8,14,16,17,19\}$	yes	116	$\{3,6,10,12,14,20,22\}$	no
38	$\{1,8,14,16,17,20,21\}$	no	117	$\{3,8,11,17,20\}$	yes
39	$\{1,8,14,16,17,20,22\}$	no	118	$\{3,8,11,18,20\}$	yes
40	$\{1,8,14,16,18,19\}$	yes	119	$\{3,8,12,13,17,21\}$	yes
41	$\{1,8,14,16,18,20,21\}$	no	120	$\{3,8,12,13,17,22\}$	yes
42	$\{1,8,14,16,18,20,22\}$	no	121	$\{3,8,12,13,18,21\}$	yes
43	$\{2,5,9,15,21\}$	yes	122	$\{3,8,12,13,18,22\}$	yes
44	$\{2,5,9,15,22\}$	yes	123	$\{3,8,12,14,17,19\}$	yes

j	Support	E	j	Support	E
45	$\{2,5,9,16,17,19\}$	yes	124	$\{3,8,12,14,17,20,21\}$	no
46	$\{2,5,9,16,17,20,21\}$	no	125	$\{3,8,12,14,17,20,22\}$	no
47	$\{2,5,9,16,17,20,22\}$	no	126	$\{3,8,12,14,18,19\}$	yes
48	$\{2,5,9,16,18,19\}$	yes	127	$\{3,8,12,14,18,20,21\}$	no
49	$\{2,5,9,16,18,20,21\}$	no	128	$\{3,8,12,14,18,20,22\}$	no
50	$\{2,5,9,16,18,20,22\}$	no	129	$\{4,5,13,16,17,21\}$	yes
51	$\{2,5,10,15,21\}$	yes	130	$\{4,5,13,16,17,22\}$	yes
52	$\{2,5,10,15,22\}$	yes	131	$\{4,5,13,16,18,21\}$	yes
53	$\{2,5,10,16,17,19\}$	yes	132	$\{4,5,13,16,18,22\}$	yes
54	$\{2,5,10,16,17,20,21\}$	no	133	$\{4,5,14,15,21\}$	yes
55	$\{2,5,10,16,17,20,22\}$	no	134	$\{4,5,14,15,22\}$	yes
56	$\{2,5,10,16,18,19\}$	yes	135	$\{4,5,14,16,17,19\}$	yes
57	$\{2,5,10,16,18,20,21\}$	no	136	$\{4,5,14,16,17,20,21\}$	no
58	$\{2,5,10,16,18,20,22\}$	no	137	$\{4,5,14,16,17,20,22\}$	no
59	$\{2,6,9,12,16,19\}$	yes	138	$\{4,5,14,16,18,19\}$	yes
60	$\{2,6,9,12,16,20,21\}$	no	139	$\{4,5,14,16,18,20,21\}$	no
61	$\{2,6,9,12,16,20,22\}$	no	140	$\{4,5,14,16,18,20,22\}$	no
62	$\{2,6,10,12,16,19\}$	yes	141	$\{4,6,11,16,20\}$	yes
63	$\{2,6,10,12,16,20,21\}$	no	142	$\{4,6,12,13,16,21\}$	yes
64	$\{2,6,10,12,16,20,22\}$	no	143	$\{4,6,12,13,16,22\}$	yes
65	$\{2,7,9,12,17,19\}$	yes	144	$\{4,6,12,14,16,19\}$	yes
66	$\{2,7,9,12,17,20,21\}$	no	145	$\{4,6,12,14,16,20,21\}$	no
67	$\{2,7,9,12,17,20,22\}$	no	146	$\{4,6,12,14,16,20,22\}$	no
68	$\{2,7,9,12,18,19\}$	yes	147	$\{4,7,11,17,20\}$	yes
69	$\{2,7,9,12,18,20,21\}$	no	148	$\{4,7,11,18,20\}$	yes
70	$\{2,7,9,12,18,20,22\}$	no	149	$\{4,7,12,13,17,21\}$	yes
71	$\{2,7,10,12,17,19\}$	yes	150	$\{4,7,12,13,17,22\}$	yes
72	$\{2,7,10,12,17,20,21\}$	no	151	$\{4,7,12,13,18,21\}$	yes
73	$\{2,7,10,12,17,20,22\}$	no	152	$\{4,7,12,13,18,22\}$	yes
74	$\{2,7,10,12,18,19\}$	yes	153	$\{4,7,12,14,17,19\}$	yes
75	$\{2,7,10,12,18,20,21\}$	no	154	$\{4,7,12,14,17,20,21\}$	no
76	$\{2,7,10,12,18,20,22\}$	no	155	$\{4,7,12,14,17,20,22\}$	no
77	$\{2,8,12,15,21\}$	yes	156	$\{4,7,12,14,18,19\}$	yes
78	$\{2,8,12,15,22\}$	yes	157	$\{4,7,12,14,18,20,21\}$	no
79	$\{2,8,12,16,17,19\}$	yes	158	$\{4,7,12,14,18,20,22\}$	no

S2.13. Complete normalized colorings of the gadget panels

Each color word gives one representative under global color permutation; the vertices and the anchor convention are specified separately for each table. The full labeled family is recovered by all palette permutations. Equal color labels across different colorings carry no extra constraint.

S2.14. Colorings of P_5

The columns of each word correspond, in order, to vertices $1,2,3,4,5,6,7,8,9,10$. The anchor context $(1,6,2)$ has colors $(0,\dots,2)$.

TABLE S58: Every normalized coloring of P_5.

j	Color word	j	Color word
1	0201211201	4	0210210211
2	0202111102	5	0212110002
3	0210110222		

S2.15. Colorings of $B_{13,7}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 5, 7, 8, 9, 10, 11, 14, 15, 19, 23$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S59: Every normalized coloring of $B_{13,7}$.

j	Color word	j	Color word
1	1022012010221	4	2012012020111
2	1022210201101	5	2012012120100
3	1022210210001	6	2012210102102

S2.16. Colorings of $D_{16,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S60: Every normalized coloring of $D_{16,9}$.

j	Color word	j	Color word
1	1020210121102202	5	2010210121201201
2	1020212101100222	6	2011200120201210
3	1020212102100112	7	2011202102021012
4	1021202102011012		

S2.17. Colorings of $I_{16,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 14, 15, 18, 19, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S61: Every normalized coloring of $I_{16,9}$.

j	Color word	j	Color word
1	1020221021001012	4	2010201212012001
2	1021221020110012	5	2011201202012110
3	1022201201021210	6	2012221010210021

S2.18. Colorings of $T_{19,11}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24, 26, 27, 28$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S62: Every normalized coloring of $T_{19,11}$.

j	Color word	j	Color word
1	1020210121102202120	5	2010210121201201120
2	1020212101100222120	6	2011200120201210120
3	1020212102100112210	7	2011202102021012102
4	1021202102011012201		

S2.19. Colorings of $H_{21,13}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 18, 19, 20, 21, 22, 23, 24$. The anchor context $(2, 8, 5)$ has colors $(0, \ldots, 2)$.

TABLE S63: Every normalized coloring of $H_{21,13}$.

j	Color word	j	Color word
1	102021210210001021212	2	201120012021012102210

S2.20. Colorings of $W^{(4)}_{11,5}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10$. The anchor context $(0, 1, 6, 2)$ has colors $(0, \ldots, 3)$.

TABLE S64: Every normalized coloring of $W^{(4)}_{11,5}$.

j	Color word	j	Color word
1	01312322312	4	01321321322
2	01313222213	5	01323221113
3	01321221333		

S2.21. Colorings of $W^{(4)}_{16,6}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15$. The anchor context $(2, 0, 6, 1)$ has colors $(0, \ldots, 3)$.

TABLE S65: Every normalized coloring of $W^{(4)}_{16,6}$.

j	Color word	j	Color word
1	1302302101123322	41	1302022331003131
2	1302302101132322	42	1302022331030131
3	1302302101223321	43	1302022331103130
4	1302302101232321	44	1302022331130130
5	1302302102123312	45	1302022333003111
6	1302302102132312	46	1302022333030111
7	1302302102223311	47	1302022333103110
8	1302302102232311	48	1302022333130110
9	1302302301123122	49	1302322100003311
10	1302302301132122	50	1302322100030311
11	1302302301223121	51	1302322100103310
12	1302302301232121	52	1302322100130310
13	1302302302123112	53	1302322101003301
14	1302302302132112	54	1302322101030301
15	1302302302223111	55	1302322101103300
16	1302302302232111	56	1302322101130300
17	1303202101123232	57	1302322300003111
18	1303202101132232	58	1302322300030111
19	1303202101223231	59	1302322300103110
20	1303202101232231	60	1302322300130110
21	1303202103123212	61	1302322301003101

j	Color word	j	Color word
22	1303202103132212	62	1302322301030101
23	1303202103223211	63	1302322301103100
24	1303202103232211	64	1302322301130100
25	1303202201123132	65	1303022121003231
26	1303202201132132	66	1303022121030231
27	1303202201223131	67	1303022121103230
28	1303202201232131	68	1303022121130230
29	1303202203123112	69	1303022123003211
30	1303202203132112	70	1303022123030211
31	1303202203223111	71	1303022123103210
32	1303202203232111	72	1303022123130210
33	1302022131003331	73	1303022221003131
34	1302022131030331	74	1303022221030131
35	1302022131103330	75	1303022221103130
36	1302022131130330	76	1303022221130130
37	1302022133003311	77	1303022223003111
38	1302022133030311	78	1303022223030111
39	1302022133103310	79	1303022223103110
40	1302022133130310	80	1303022223130110

S2.22. Colorings of $B^{(4)}_{14,7}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 5, 7, 8, 9, 10, 11, 14, 15, 19, 23$. The anchor context $(0, 1, 2, 3)$ has colors $(0, \ldots, 3)$.

TABLE S66: Every normalized coloring of $B^{(4)}_{14,7}$.

j	Color word	j	Color word
1	01233213212331	4	01231132321321
2	01233312312221	5	01231231212333
3	01233312321121	6	01231231312322

S2.23. Colorings of $L^{(4)}_{18,7}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18$. The anchor context $(1, 2, 4, 3)$ has colors $(0, \ldots, 3)$.

TABLE S67: Every normalized coloring of $L^{(4)}_{18,7}$.

j	Color word	j	Color word
1	013212030123031203	73	013210230132021023
2	013212030123031230	74	013210230132021032
3	013212030123301203	75	013210230132201023
4	013212030123301230	76	013210230132201032
5	013212031023031203	77	013210231032021023
6	013212031023031230	78	013210231032021032
7	013212031023301203	79	013210231032201023
8	013212031023301230	80	013210231032201032

j	Color word	j	Color word
9	013210230123031023	81	013213201332020112
10	013210230123031032	82	013213201332020121
11	013210230123301023	83	013213201332200112
12	013210230123301032	84	013213201332200121
13	013210231023031023	85	013213203132020112
14	013210231023031032	86	013213203132020121
15	013210231023301023	87	013213203132200112
16	013210231023301032	88	013213203132200121
17	013213201323030112	89	013213201332021002
18	013213201323030121	90	013213201332021020
19	013213201323300112	91	013213201332201002
20	013213201323300121	92	013213201332201020
21	013213203123030112	93	013213203132021002
22	013213203123030121	94	013213203132021020
23	013213203123300112	95	013213203132201002
24	013213203123300121	96	013213203132201020
25	013213201323031002	97	013221030131022103
26	013213201323031020	98	013221030131022130
27	013213201323301002	99	013221030131202103
28	013213201323301020	100	013221030131202130
29	013213203123031002	101	013221031031022103
30	013213203123031020	102	013221031031022130
31	013213203123301002	103	013221031031202103
32	013213203123301020	104	013221031031202130
33	013231201321030123	105	013223010331021202
34	013231201321030132	106	013223010331021220
35	013231201321300123	107	013223010331201202
36	013231201321300132	108	013223010331201220
37	013231203121030123	109	013223013031021202
38	013231203121030132	110	013223013031021220
39	013231203121300123	111	013223013031201202
40	013231203121300132	112	013223013031201220
41	013232101321030213	113	013223010331022101
42	013232101321030231	114	013223010331022110
43	013232101321300213	115	013223010331202101
44	013232101321300231	116	013223010331202110
45	013232103121030213	117	013223013031022101
46	013232103121030231	118	013223013031022110
47	013232103121300213	119	013223013031202101
48	013232103121300231	120	013223013031202110
49	013230210321031023	121	013220130131022013
50	013230210321031032	122	013220130131022031
51	013230210321301023	123	013220130131202013
52	013230210321301032	124	013220130131202031
53	013230213021031023	125	013220131031022013
54	013230213021031032	126	013220131031022031
55	013230213021301023	127	013220131031202013
56	013230213021301032	128	013220131031202031

j	Color word	j	Color word
57	013232010321031203	129	013223101331020212
58	013232010321031230	130	013223101331020221
59	013232010321301203	131	013223101331200212
60	013232010321301230	132	013223101331200221
61	013232013021031203	133	013223103131020212
62	013232013021031230	134	013223103131020221
63	013232013021301203	135	013223103131200212
64	013232013021301230	136	013223103131200221
65	013212030132021203	137	013223101331022001
66	013212030132021230	138	013223101331022010
67	013212030132201203	139	013223101331202001
68	013212030132201230	140	013223101331202010
69	013212031032021203	141	013223103131022001
70	013212031032021230	142	013223103131022010
71	013212031032201203	143	013223103131202001
72	013212031032201230	144	013223103131202010

S2.24. Colorings of $D^{(4)}_{17,9}$

The columns of each word correspond, in order, to vertices $0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 23, 24$. The anchor context $(0, 1, 2, 3)$ has colors $(0, \ldots, 3)$.

TABLE S68: Every normalized coloring of $D^{(4)}_{17,9}$.

j	Color word	j	Color word
1	01232132313123123	5	01233122312123132
2	01232312131123323	6	01231323123211213
3	01232313121122333	7	01233121321213231
4	01232313123122113		

S2.25. Colorings of $L^{(4)}_{22,9}$

The columns of each word correspond, in order, to vertices $1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22$. The anchor context $(1, 2, 4, 3)$ has colors $(0, \ldots, 3)$.

TABLE S69: Every normalized coloring of $L^{(4)}_{22,9}$.

j	Color word	j	Color word
1	0132120301230312033201	109	0132120310320212032301
2	0132120301230312033210	110	0132120310320212032310
3	0132120301230312303201	111	0132120310320212302301
4	0132120301230312303210	112	0132120310320212302310
5	0132120310230312033201	113	0132120301322012030312
6	0132120310230312033210	114	0132120301322012030321
7	0132120310230312303201	115	0132120301322012300312
8	0132120310230312303210	116	0132120301322012300321
9	0132120301233012030213	117	0132120310322012030312
10	0132120301233012030231	118	0132120310322012030321

j	Color word	j	Color word
11	013212030l233012300213	119	01321203l0322012300312
12	013212030l233012300231	120	01321203l0322012300321
13	013212031023301203021 3	121	013213201332200112130 2
14	013212031023301203023 1	122	013213201332200112132 0
15	013212031023301230021 3	123	013213201332200121130 2
16	013212031023301230023 1	124	013213201332200121132 0
17	013213201323300112120 3	125	013213203132200112130 2
18	013213201323300112123 0	126	013213203132200112132 0
19	013213201323300121120 3	127	013213203132200121130 2
20	013213201323300121123 0	128	013213203132200121132 0
21	013213203123300112120 3	129	013210230132021023230 1
22	013213203123300112123 0	130	013210230132021023231 0
23	013213203123300121120 3	131	013210230132021032230 1
24	013213203123300121123 0	132	013210230132021032231 0
25	013210230123031023320 1	133	013210231032021023230 1
26	013210230123031023321 0	134	013210231032021023231 0
27	013210230123031032320 1	135	013210231032021032230 1
28	013210230123031032321 0	136	013210231032021032231 0
29	013210231023031023320 1	137	013210230132201023031 2
30	013210231023031023321 0	138	013210230132201023032 1
31	013210231023031032320 1	139	013210230132201032031 2
32	013210231023031032321 0	140	013210230132201032032 1
33	013210230123301023021 3	141	013210231032201023031 2
34	013210230123301023023 1	142	013210231032201023032 1
35	013210230123301032021 3	143	013210231032201032031 2
36	013210230123301032023 1	144	013210231032201032032 1
37	013210231023301023021 3	145	013213201332021002230 1
38	013210231023301023023 1	146	013213201332021002231 0
39	013210231023301032021 3	147	013213201332021020230 1
40	013210231023301032023 1	148	013213201332021020231 0
41	013213201323031002320 1	149	013213203132021002230 1
42	013213201323031002321 0	150	013213203132021002231 0
43	013213201323031020320 1	151	013213203132021020230 1
44	013213201323031020321 0	152	013213203132021020231 0
45	013213203123031002320 1	153	013213201332201002031 2
46	013213203123031002321 0	154	013213201332201002032 1
47	013213203123031020320 1	155	013213201332201020031 2
48	013213203123031020321 0	156	013213201332201020032 1
49	013213201323301002021 3	157	013213203132201002031 2
50	013213201323301002023 1	158	013213203132201002032 1
51	013213201323301020021 3	159	013213203132201020031 2
52	013213201323301020023 1	160	013213203132201020032 1
53	013213203123301002021 3	161	013223010331021202230 1
54	013213203123301002023 1	162	013223010331021202231 0
55	013213203123301020021 3	163	013223010331021220230 1
56	013213203123301020023 1	164	013223010331021220231 0
57	013231201321300123120 3	165	013223013031021202230 1
58	013231201321300123123 0	166	013223013031021202231 0

j	Color word	j	Color word
59	013231201321300132 1203	167	013223013031021220 2301
60	013231201321300132 1230	168	013223013031021220 2310
61	013231203121300123 1203	169	013223010331201202 0312
62	013231203121300123 1230	170	013223010331201202 0321
63	013231203121300132 1203	171	013223010331201220 0312
64	013231203121300132 1230	172	013223010331201220 0321
65	013232101321300213 1203	173	013223013031201202 0312
66	013232101321300213 1230	174	013223013031201202 0321
67	013232101321300231 1203	175	013223013031201220 0312
68	013232101321300231 1230	176	013223013031201220 0321
69	013232103121300213 1203	177	013221030131022103 1302
70	013232103121300213 1230	178	013221030131022103 1320
71	013232103121300231 1203	179	013221030131022130 1302
72	013232103121300231 1230	180	013221030131022130 1320
73	013230210321031023 3201	181	013221031031022103 1302
74	013230210321031023 3210	182	013221031031022103 1320
75	013230210321031032 3201	183	013221031031022130 1302
76	013230210321031032 3210	184	013221031031022130 1320
77	013230213021031023 3201	185	013223010331022101 1302
78	013230213021031023 3210	186	013223010331022101 1320
79	013230213021031032 3201	187	013223010331022110 1302
80	013230213021031032 3210	188	013223010331022110 1320
81	013230210321301023 0213	189	013223013031022101 1302
82	013230210321301023 0231	190	013223013031022101 1320
83	013230210321301032 0213	191	013223013031022110 1302
84	013230210321301032 0231	192	013223013031022110 1320
85	013230213021301023 0213	193	013223101331200212 1302
86	013230213021301023 0231	194	013223101331200212 1320
87	013230213021301032 0213	195	013223101331200221 1302
88	013230213021301032 0231	196	013223101331200221 1320
89	013232010321031203 3201	197	013223103131200212 1302
90	013232010321031203 3210	198	013223103131200212 1320
91	013232010321031230 3201	199	013223103131200221 1302
92	013232010321031230 3210	200	013223103131200221 1320
93	013232013021031203 3201	201	013220130131022013 1302
94	013232013021031203 3210	202	013220130131022013 1320
95	013232013021031230 3201	203	013220130131022031 1302
96	013232013021031230 3210	204	013220130131022031 1320
97	013232010321301203 0213	205	013220131031022013 1302
98	013232010321301203 0231	206	013220131031022013 1320
99	013232010321301230 0213	207	013220131031022031 1302
100	013232010321301230 0231	208	013220131031022031 1320
101	013232013021301203 0213	209	013223101331022001 1302
102	013232013021301203 0231	210	013223101331022001 1320
103	013232013021301230 0213	211	013223101331022010 1302
104	013232013021301230 0231	212	013223101331022010 1320
105	013212030132021203 2301	213	013223103131022001 1302
106	013212030132021203 2310	214	013223103131022001 1320

j	Color word	j	Color word
107	0132120301320212302301	215	0132231031310220101302
108	0132120301320212302310	216	0132231031310220101320

Exhaustive classification of chromatic identities. For a chromatically inseparable pair (u, v), form the two directed witness sets

$$W_{10}(u,v) = \{j : s_j(u) = 1,\ s_j(v) = 0\},$$
$$W_{01}(u,v) = \{j : s_j(u) = 0,\ s_j(v) = 1\}.$$

Both sets are nonempty exactly for a two-way purely chromatic identity. If only W_{10} is empty, $u \Rightarrow v$ holds and its converse fails; if only W_{01} is empty, the implication is reversed. If both are empty, the pair is inseparable under all ordinary states. Any member of either witness set is automatically nonextendable, since every color-class state has equal values on a chromatically inseparable pair.

The following Python standard-library procedure reconstructs $H_{21,13}$ directly from its 13 contexts in Table S3, enumerates all binary assignments, and reconstructs normalized strong colorings as partitions into three disjoint state supports. Lexicographic support order agrees with Table S42. It prints both complete directed witness sets for every chromatic identity, rather than only the representatives in Table S70.

```
from itertools import combinations
E = [(1,2,3),(7,8,9),(4,5,6),(2,8,5),
     (3,11,14),(7,11,15),(4,11,24),
     (9,10,23),(1,10,19),(4,10,18),
     (1,13,20),(7,13,21),(6,13,22)]
V = sorted(set().union(*map(set, E)))
M = [sum(1 << V.index(v) for v in e) for e in E]
S = sorted(tuple(v for i,v in enumerate(V) if b >> i & 1)
           for b in range(1 << len(V))
           if all((b & m).bit_count() == 1 for m in M))
S = list(map(frozenset, S))
C = []
for blocks in combinations(S, 3):
    if any(a & b for a,b in combinations(blocks, 2)):
        continue
    blocks = [next(s for s in blocks if a in s) for a in (2,8,5)]
    C.append({v:k for k,s in enumerate(blocks) for v in s})
assert (len(S), len(C)) == (21, 2)
counts = [0, 0, 0]
for u,v in combinations(V, 2):
    if not all(c[u] == c[v] for c in C):
        continue
    W10 = [j for j,s in enumerate(S, 1) if u in s and v not in s]
    W01 = [j for j,s in enumerate(S, 1) if v in s and u not in s]
    k = 0 if W10 and W01 else 1 if W10 or W01 else 2
    counts[k] += 1
    print((u,v), W10, W01)
assert counts == [15, 6, 0]
```

TABLE S70. Both directed truth-pattern witnesses for all 15 two-way purely chromatic pairs of $H_{21,13}$ and both displayed pairs of $T_{19,11}$. For $H_{21,13}$, s_j is the state in Table S42; for $T_{19,11}$, entries give supports. Every witness is nonextendable and satisfies every context equation.

Configuration	Pair (u,v)	$(s(u),s(v))=(1,0)$	$(s(u),s(v))=(0,1)$
$T_{19,11}$	$(14,28)$	$\{1,5,7,14,18,23,24\}$	$\{1,6,8,10,11,27,28\}$
$T_{19,11}$	$(18,27)$	$\{1,5,7,14,18,23,24\}$	$\{1,6,8,10,11,27,28\}$
$H_{21,13}$	$(1,11)$	s_1	s_{12}
$H_{21,13}$	$(1,18)$	s_1	s_{11}
$H_{21,13}$	$(1,21)$	s_2	s_9
$H_{21,13}$	$(2,14)$	s_{12}	s_1
$H_{21,13}$	$(4,13)$	s_1	s_{18}
$H_{21,13}$	$(4,15)$	s_7	s_4
$H_{21,13}$	$(4,19)$	s_1	s_{11}
$H_{21,13}$	$(5,22)$	s_{18}	s_1
$H_{21,13}$	$(7,10)$	s_2	s_{20}
$H_{21,13}$	$(7,20)$	s_2	s_9
$H_{21,13}$	$(7,24)$	s_7	s_4
$H_{21,13}$	$(8,23)$	s_{20}	s_2
$H_{21,13}$	$(15,19)$	s_1	s_7
$H_{21,13}$	$(18,21)$	s_2	s_1
$H_{21,13}$	$(20,24)$	s_7	s_2

TABLE S71: All ten connected, edge-faithful, colorable restrictions with a forced color identity in the specified inventory.

retained diagonals	rays	contexts	colorings	inseparable pairs
$\{10,11\}$	17	10	4	$(10,24)$, $(11,18)$
$\{10,12\}$	17	10	4	$(10,16)$, $(12,23)$
$\{10,13\}$	17	10	4	$(10,20)$, $(13,19)$
$\{11,12\}$	17	10	4	$(11,25)$, $(12,14)$
$\{11,13\}$	17	10	4	$(11,21)$, $(13,15)$
$\{12,13\}$	17	10	4	$(12,22)$, $(13,17)$
$\{10,11,12\}$	21	13	2	21 pairs
$\{10,11,13\}$	21	13	2	21 pairs
$\{10,12,13\}$	21	13	2	21 pairs
$\{11,12,13\}$	21	13	2	21 pairs

S2.26. An explicit state certificate for the completed Yu–Oh hypergraph

The first four contexts of Eq. (15) force exactly one of vertices 2, 5, and 8 to have value one. In the triad containing that coordinate vertex, both other vertices have value zero. In each of the other two triads, one of the two remaining vertices has value one. There are therefore $3 \cdot 2^2 = 12$ possible supports T_k on vertices $1,\dots,9$, listed in Table S72.

For each such support, exactly one of vertices $10,11,12,13$ has no orthogonal neighbor in T_k; call it η_k. The other three are forced to zero, while $s(\eta_k)$ may be either zero or one. This gives precisely 24 assignments on the original 13 vertices. Every completion vertex w belongs to one context $\{i,j,w\}$ and has the uniquely determined value

$$s(w) = 1 - s(i) - s(j). \tag{S7}$$

The listed assignments have $s(i)+s(j) \le 1$ on every original orthogonal pair, so these

extensions are valid two-valued states. Conversely, every state of the completed hypergraph restricts to one of these assignments.

For a compact separation certificate, encode the profile of vertex i as the nonnegative integer

$$m_i = \sum_{k=1}^{12} \sum_{b=0}^{1} s_{k,b}(i)\, 2^{2(k-1)+b}.$$

In hexadecimal notation these 25 integers are

i	1	2	3	4	5
m_i	0000FF	00FF00	FF0000	030F03	3C003C
i	6	7	8	9	10
m_i	C0F0C0	0C330C	C300C3	30CC30	882000
i	11	12	13	14	15
m_i	0080A0	00020A	220800	007F5F	F34C53
i	16	17	18	19	20
m_i	CF31C5	3F0D35	74D0FC	77DF00	DDF700
i	21	22	23	24	25
m_i	D1C4F3	1D073F	4713CF	FC705C	00FDF5

(S8)

All 25 integers are nonzero and pairwise distinct, certifying unitality and separation. Notably, the Hamming weights (number of states in which $s(i) = 1$) partition the vertices into three classes: 8 for $i \in \{1, \ldots, 9\}$, 3 for $i \in \{10, \ldots, 13\}$, and 13 for the completion vertices $i \in \{14, \ldots, 25\}$ (with $8+3+13 = 24$ in each completion context), in contrast to the uniform state weight 2 across all 15 vertices of G_{32} and consistent with the fractional chromatic–clique gap $\chi_f(\Gamma(H^+_{\mathrm{YO}})) = 35/11 > 3 = \omega(\Gamma(H^+_{\mathrm{YO}}))$.

TABLE S72. The 24 states are indexed by (k, b), where $1 \le k \le 12$ and $b \in \{0, 1\}$. Their support on vertices $1, \ldots, 13$ is T_k when $b = 0$ and $T_k \cup \{\eta_k\}$ when $b = 1$. Equation (S7) determines all remaining values.

k	T_k	η_k
1	$\{1, 4, 8\}$	12
2	$\{1, 5, 7\}$	12
3	$\{1, 5, 9\}$	11
4	$\{1, 6, 8\}$	11
5	$\{2, 4, 7\}$	12
6	$\{2, 4, 9\}$	13
7	$\{2, 6, 7\}$	10
8	$\{2, 6, 9\}$	11
9	$\{3, 4, 8\}$	13
10	$\{3, 5, 7\}$	10
11	$\{3, 5, 9\}$	13
12	$\{3, 6, 8\}$	10

S2.27. Hesse H_{37}: all states and separating profiles

The canonical states $s_1, \dots, s_{153}$ are listed in Table S74. A two-valued coloring is a state with exactly one value one per context. Vertex numbers are those of Appendix S1.14 and Fig. 4: vertices $1, \dots, 16$ form the core, and $17, \dots, 37$ are completions. We list every state, a minimum separating family, and every vertex profile.

S2.27.1. State parametrization and all 27 branches

Choose $t = (t_0, t_1, t_2) \in \mathbb{Z}_3^3$ and $\varepsilon \in \{0, 1\}$. Put

$$s(M_{ak}) = \mathbf{1}_{k=t_a}, \qquad s(z) = \varepsilon, \tag{S9}$$

$$U(t) = \mathbb{Z}_3 \setminus \{t_0, t_1, t_2\}, \qquad W(t) = \mathbb{Z}_3 \setminus \{t_a - a : a \in \mathbb{Z}_3\}. \tag{S10}$$

For $\varepsilon = 0$, choose the values of P_k freely for $k \in U(t)$, with other P values zero. For $\varepsilon = 1$, all P values are zero. Independently, choose the values of Q_k freely for $k \in W(t)$, with other Q values zero. All completion values are forced:

$$s(A_k) = 1 - \varepsilon - s(P_k), \tag{S11}$$

$$s(B_{ak}) = 1 - s(M_{ak}) - s(P_k), \tag{S12}$$

$$s(C_{ak}) = 1 - s(M_{a,a+k}) - s(Q_k). \tag{S13}$$

Enumerate t lexicographically with $0 < 1 < 2$, then $\varepsilon = 1, 0$. Within each branch, list the available P bits in increasing k, followed by the available Q bits in increasing k, and order the resulting bit strings with 1 before 0. This rule fixes all state indices used in the text, tables, and verification data. Table S73 gives the index ranges of all 27 branches.

TABLE S73: All 27 state branches. The sets $U(t), W(t)$ identify the free core bits. The two choices of ε give $N_0 = 2^{|U|+|W|}$ and $N_1 = 2^{|W|}$ states; their index ranges refer to Table S74.

t	$U(t)$	$W(t)$	N_0	r for $\varepsilon = 0$	N_1	r for $\varepsilon = 1$
$(0,0,0)$	$\{1,2\}$	$\varnothing$	4	2–5	1	1
$(0,0,1)$	$\{2\}$	$\{1\}$	4	8–11	2	6–7
$(0,0,2)$	$\{1\}$	$\{1\}$	4	14–17	2	12–13
$(0,1,0)$	$\{2\}$	$\{2\}$	4	20–23	2	18–19
$(0,1,1)$	$\{2\}$	$\{1\}$	4	26–29	2	24–25
$(0,1,2)$	$\varnothing$	$\{1,2\}$	4	34–37	4	30–33
$(0,2,0)$	$\{1\}$	$\{2\}$	4	40–43	2	38–39
$(0,2,1)$	$\varnothing$	$\varnothing$	1	45	1	44
$(0,2,2)$	$\{1\}$	$\{2\}$	4	48–51	2	46–47
$(1,0,0)$	$\{2\}$	$\{0\}$	4	54–57	2	52–53
$(1,0,1)$	$\{2\}$	$\{0\}$	4	60–63	2	58–59
$(1,0,2)$	$\varnothing$	$\varnothing$	1	65	1	64
$(1,1,0)$	$\{2\}$	$\{2\}$	4	68–71	2	66–67
$(1,1,1)$	$\{0,2\}$	$\varnothing$	4	73–76	1	72
$(1,1,2)$	$\{0\}$	$\{2\}$	4	79–82	2	77–78
$(1,2,0)$	$\varnothing$	$\{0,2\}$	4	87–90	4	83–86
$(1,2,1)$	$\{0\}$	$\{0\}$	4	93–96	2	91–92
$(1,2,2)$	$\{0\}$	$\{2\}$	4	99–102	2	97–98
$(2,0,0)$	$\{1\}$	$\{0\}$	4	105–108	2	103–104
$(2,0,1)$	$\varnothing$	$\{0,1\}$	4	113–116	4	109–112
$(2,0,2)$	$\{1\}$	$\{1\}$	4	119–122	2	117–118
$(2,1,0)$	$\varnothing$	$\varnothing$	1	124	1	123
$(2,1,1)$	$\{0\}$	$\{1\}$	4	127–130	2	125–126
$(2,1,2)$	$\{0\}$	$\{1\}$	4	133–136	2	131–132
$(2,2,0)$	$\{1\}$	$\{0\}$	4	139–142	2	137–138
$(2,2,1)$	$\{0\}$	$\{0\}$	4	145–148	2	143–144
$(2,2,2)$	$\{0,1\}$	$\varnothing$	4	150–153	1	149
Total			99		54	

S2.27.2. Canonical two-valued colorings: all 153 states

Table S74 explicitly lists all 153 states. The last column is the complete binary vector in vertex order $1, \ldots, 37$, separated after the 16 core entries. Every row has context sum one in all 24 contexts. The support-size distribution is

$$\begin{array}{c|cccccc} m & 12 & 13 & 14 & 15 & 16 & 18 \\ \#\{s : |\operatorname{supp} s| = m\} & 6 & 18 & 27 & 24 & 51 & 27 \end{array} \tag{S14}$$

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$.

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
1	$\{1,2,5,8\}$	$\{21,22,24,25,27,28,30,31,32,33,35,37\}$	16	1100100100000000 0000110110110111110101
2	$\{2,5,8,12,13\}$	$\{17,30,31,32,33,35,37\}$	12	0100100100011000 1000000000000111110101
3	$\{2,5,8,12\}$	$\{17,19,22,25,28,30,31,32,33,35,37\}$	15	0100100100010000 1010010010011110101
4	$\{2,5,8,13\}$	$\{17,18,21,24,27,30,31,32,33,35,37\}$	15	0100100100001000 1100100100011110101
5	$\{2,5,8\}$	$\{17,18,19,21,22,24,25,27,28,30,31,32,33,35,37\}$	18	0100100100000000 1110110110111110101
6	$\{1,2,5,9,15\}$	$\{21,22,24,25,26,28,31,32,35\}$	14	1100100010000010 0000110111010011100100
7	$\{1,2,5,9\}$	$\{21,22,24,25,26,28,30,31,32,33,35,36\}$	16	1100100010000000 0000110111010111110110
8	$\{2,5,9,13,15\}$	$\{17,18,21,24,26,31,32,35\}$	13	0100100010001010 1100100101000011100100
9	$\{2,5,9,13\}$	$\{17,18,21,24,26,30,31,32,33,35,36\}$	15	0100100010001000 1100100101000111110110
10	$\{2,5,9,15\}$	$\{17,18,19,21,22,24,25,26,28,31,32,35\}$	16	0100100010000010 1110110111010011100100
11	$\{2,5,9\}$	$\{17,18,19,21,22,24,25,26,28,30,31,32,33,35,36\}$	18	0100100010000000 1110110111010111110110
12	$\{1,2,5,10,15\}$	$\{21,22,24,25,26,27,31,32,37\}$	14	1100100001000010 0000110111100011100001
13	$\{1,2,5,10\}$	$\{21,22,24,25,26,27,30,31,32,33,36,37\}$	16	1100100001000000 0000110111100111110011
14	$\{2,5,10,12,15\}$	$\{17,19,22,25,26,31,32,37\}$	13	0100100001010010 1010010011000011100001
15	$\{2,5,10,12\}$	$\{17,19,22,25,26,30,31,32,33,36,37\}$	15	0100100001010000 1010010011000111110011
16	$\{2,5,10,15\}$	$\{17,18,19,21,22,24,25,26,27,31,32,37\}$	16	0100100001000010 1110110111100011100001
17	$\{2,5,10\}$	$\{17,18,19,21,22,24,25,26,27,30,31,32,33,36,37\}$	18	0100100001000000 1110110111100111110011
18	$\{1,2,6,8,16\}$	$\{21,22,23,25,27,28,30,33,35\}$	14	1100010100000001 0000111010110100010100
19	$\{1,2,6,8\}$	$\{21,22,23,25,27,28,30,31,33,34,35,37\}$	16	1100010100000000 0000111010111011011101
20	$\{2,6,8,13,16\}$	$\{17,18,21,23,27,30,33,35\}$	13	0100010100001001 1100101000110010010100
21	$\{2,6,8,13\}$	$\{17,18,21,23,27,30,31,33,34,35,37\}$	15	0100010100001000 1100101000110011011101
22	$\{2,6,8,16\}$	$\{17,18,19,21,22,23,25,27,28,30,33,35\}$	16	0100010100000001 1110111010111010010100
23	$\{2,6,8\}$	$\{17,18,19,21,22,23,25,27,28,30,31,33,34,35,37\}$	18	0100010100000000 1110111010111011011101
24	$\{1,2,6,9,15\}$	$\{21,22,23,25,26,28,31,34,35\}$	14	1100010010000010 0000111011010011001100
25	$\{1,2,6,9\}$	$\{21,22,23,25,26,28,30,31,33,34,35,36\}$	16	1100010010000000 0000111011010111011110
26	$\{2,6,9,13,15\}$	$\{17,18,21,23,26,31,34,35\}$	13	0100010010001010 1100101001000011001100
27	$\{2,6,9,13\}$	$\{17,18,21,23,26,30,31,33,34,35,36\}$	15	0100010010001000 1100101001000111011110
28	$\{2,6,9,15\}$	$\{17,18,19,21,22,23,25,26,28,31,34,35\}$	16	0100010010000010 1110111011010011001100
29	$\{2,6,9\}$	$\{17,18,19,21,22,23,25,26,28,30,31,33,34,35,36\}$	18	0100010010000000 1110111010111010111110
30	$\{1,2,6,10,15,16\}$	$\{21,22,23,25,26,27\}$	12	1100010001000011 0000111011100000000000
31	$\{1,2,6,10,15\}$	$\{21,22,23,25,26,27,31,34,37\}$	14	1100010001000010 0000111011100011001001
32	$\{1,2,6,10,16\}$	$\{21,22,23,25,26,27,30,33,36\}$	14	1100010001000001 0000111011100100100010
33	$\{1,2,6,10\}$	$\{21,22,23,25,26,27,30,31,33,34,36,37\}$	16	1100010001000000 0000111011100111011011
34	$\{2,6,10,15,16\}$	$\{17,18,19,21,22,23,25,26,27\}$	14	0100010001000011 1110111011100000000000
35	$\{2,6,10,15\}$	$\{17,18,19,21,22,23,25,26,27,31,34,37\}$	16	0100010001000010 1110111011100011001001
36	$\{2,6,10,16\}$	$\{17,18,19,21,22,23,25,26,27,30,33,36\}$	16	0100010001000001 1110111011100100100010
37	$\{2,6,10\}$	$\{17,18,19,21,22,23,25,26,27,30,31,33,34,36,37\}$	18	0100010001000000 1110111011100111011011
38	$\{1,2,7,8,16\}$	$\{21,22,23,24,27,28,30,32,35\}$	14	1100001100000001 0000111100110101001100
39	$\{1,2,7,8\}$	$\{21,22,23,24,27,28,30,31,32,34,35,37\}$	16	1100001100000000 0000111100111011101101
40	$\{2,7,8,12,16\}$	$\{17,19,22,23,28,30,32,35\}$	13	0100001100010001 1010011000010100100100
41	$\{2,7,8,12\}$	$\{17,19,22,23,28,30,31,32,34,35,37\}$	15	0100001100010000 1010011000011011101101
42	$\{2,7,8,16\}$	$\{17,18,19,21,22,23,24,27,28,30,32,35\}$	16	0100001100000001 1110111100110101001100
43	$\{2,7,8\}$	$\{17,18,19,21,22,23,24,27,28,30,31,32,34,35,37\}$	18	0100001100000000 1110111100111011101101
44	$\{1,2,7,9\}$	$\{21,22,23,24,26,28,30,31,32,34,35,36\}$	16	1100001010000000 0000111101010111011110
45	$\{2,7,9\}$	$\{17,18,19,21,22,23,24,26,28,30,31,32,34,35,36\}$	18	0100001010000000 1110111101010111011110
46	$\{1,2,7,10,16\}$	$\{21,22,23,24,26,27,30,32,36\}$	14	1100001001000001 0000111101100101000010
47	$\{1,2,7,10\}$	$\{21,22,23,24,26,27,30,31,32,34,36,37\}$	16	1100001001000000 0000111101100111011011
48	$\{2,7,10,12,16\}$	$\{17,19,22,23,26,30,32,36\}$	13	0100001001010001 1010011000010100010
49	$\{2,7,10,12\}$	$\{17,19,22,23,26,30,31,32,34,36,37\}$	15	0100001001010000 1010011001000111101011
50	$\{2,7,10,16\}$	$\{17,18,19,21,22,23,24,26,27,30,32,36\}$	16	0100001001000001 1110111101100101000010
51	$\{2,7,10\}$	$\{17,18,19,21,22,23,24,26,27,30,31,32,34,36,37\}$	18	0100001001000000 1110111101100111011011

Continued on the next page.

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
52	$\{1,3,5,8,14\}$	$\{20,22,24,25,27,28,31,33,37\}$	14	1010100100000100 0001010111001010001
53	$\{1,3,5,8\}$	$\{20,22,24,25,27,28,29,31,32,33,35,37\}$	16	1010100100000000 0001010111101110101
54	$\{3,5,8,13,14\}$	$\{17,18,20,24,27,31,33,37\}$	13	0010100100001100 1101000100100010001
55	$\{3,5,8,13\}$	$\{17,18,20,24,27,29,31,32,33,35,37\}$	15	0010100100001000 1101000100101011101
56	$\{3,5,8,14\}$	$\{17,18,19,20,22,24,25,27,28,31,33,37\}$	16	0010100100000100 1111010111001010001
57	$\{3,5,8\}$	$\{17,18,19,20,22,24,25,27,28,29,31,32,33,35,37\}$	18	0010100100000000 1111010111011101101
58	$\{1,3,5,9,14\}$	$\{20,22,24,25,26,28,31,33,36\}$	14	1010100010000100 0001010111010010010
59	$\{1,3,5,9\}$	$\{20,22,24,25,26,28,29,31,32,33,35,36\}$	16	1010100010000000 0001010111011101110
60	$\{3,5,9,13,14\}$	$\{17,18,20,24,26,31,33,36\}$	13	0010100010001100 1101000101000010010
61	$\{3,5,9,13\}$	$\{17,18,20,24,26,29,31,32,33,35,36\}$	15	0010100010001000 1101000101001011110
62	$\{3,5,9,14\}$	$\{17,18,19,20,22,24,25,26,28,31,33,36\}$	16	0010100010000100 1111010111010010010
63	$\{3,5,9\}$	$\{17,18,19,20,22,24,25,26,28,29,31,32,33,35,36\}$	18	0010100010000000 1111010111011101110
64	$\{1,3,5,10\}$	$\{20,22,24,25,26,27,29,31,32,33,36,37\}$	16	1010100001000000 0001010111101101110
65	$\{3,5,10\}$	$\{17,18,19,20,22,24,25,26,27,29,31,32,33,36,37\}$	18	0010100001000000 1111010111101101110
66	$\{1,3,6,8,16\}$	$\{20,22,23,25,27,28,29,33,35\}$	14	1010010100000001 0001011010111000010100
67	$\{1,3,6,8\}$	$\{20,22,23,25,27,28,29,31,33,34,35,37\}$	16	1010010100000000 0001011010111010111101
68	$\{3,6,8,13,16\}$	$\{17,18,20,23,27,29,33,35\}$	13	0010010100001001 1101001000101000010100
69	$\{3,6,8,13\}$	$\{17,18,20,23,27,29,31,33,34,35,37\}$	15	0010010100001000 1101001000101011101
70	$\{3,6,8,16\}$	$\{17,18,19,20,22,23,25,27,28,29,33,35\}$	16	0010010100000001 1111011010111000010100
71	$\{3,6,8\}$	$\{17,18,19,20,22,23,25,27,28,29,31,33,34,35,37\}$	18	0010010100000000 1111011010111010111101
72	$\{1,3,6,9\}$	$\{20,22,23,25,26,28,29,31,33,34,35,36\}$	16	1010010010000000 0001011011011010111110
73	$\{3,6,9,11,13\}$	$\{18,29,31,33,34,35,36\}$	12	0010010010101000 0100000000001011110
74	$\{3,6,9,11\}$	$\{18,19,22,25,28,29,31,33,34,35,36\}$	15	0010010010100000 0110010010011011110
75	$\{3,6,9,13\}$	$\{17,18,20,23,26,29,31,33,34,35,36\}$	15	0010010010001000 1101001001001011110
76	$\{3,6,9\}$	$\{17,18,19,20,22,23,25,26,28,29,31,33,34,35,36\}$	18	0010010010000000 1111011011011011110
77	$\{1,3,6,10,16\}$	$\{20,22,23,25,26,27,29,33,36\}$	14	1010010001000001 0001011011101000010
78	$\{1,3,6,10\}$	$\{20,22,23,25,26,27,29,31,33,34,36,37\}$	16	1010010001000000 0001011011101011011
79	$\{3,6,10,11,16\}$	$\{18,19,22,25,27,29,33,36\}$	13	0010010001100001 0110010010100010010
80	$\{3,6,10,11\}$	$\{18,19,22,25,27,29,31,33,34,36,37\}$	15	0010010001100000 0110010010101011011
81	$\{3,6,10,16\}$	$\{17,18,19,20,22,23,25,26,27,29,33,36\}$	16	0010010001000001 1111011011010010010
82	$\{3,6,10\}$	$\{17,18,19,20,22,23,25,26,27,29,31,33,34,36,37\}$	18	0010010001000000 1111011011011011011
83	$\{1,3,7,8,14,16\}$	$\{20,22,23,24,27,28\}$	12	1010001100000101 0001011100110000000000
84	$\{1,3,7,8,14\}$	$\{20,22,23,24,27,28,31,34,37\}$	14	1010001100000100 0001011100110010001001
85	$\{1,3,7,8,16\}$	$\{20,22,23,24,27,28,29,32,35\}$	14	1010001100000001 0001011100111001001100
86	$\{1,3,7,8\}$	$\{20,22,23,24,27,28,29,31,32,34,35,37\}$	16	1010001100000000 0001011100111011011101
87	$\{3,7,8,14,16\}$	$\{17,18,19,20,22,23,24,27,28\}$	14	0010001100000101 1111011100110000000000
88	$\{3,7,8,14\}$	$\{17,18,19,20,22,23,24,27,28,31,34,37\}$	16	0010001100000100 1111011100111001001001
89	$\{3,7,8,16\}$	$\{17,18,19,20,22,23,24,27,28,29,32,35\}$	16	0010001100000001 1111011100111001001100
90	$\{3,7,8\}$	$\{17,18,19,20,22,23,24,27,28,29,31,32,34,35,37\}$	18	0010001100000000 1111011100111011011101
91	$\{1,3,7,9,14\}$	$\{20,22,23,24,26,28,31,34,36\}$	14	1010001010000100 0001011101001001001010
92	$\{1,3,7,9\}$	$\{20,22,23,24,26,28,29,31,32,34,35,36\}$	16	1010001010000000 0001011101011011011110
93	$\{3,7,9,11,14\}$	$\{18,19,22,24,28,31,34,36\}$	13	0010001010100100 0110010100010010010
94	$\{3,7,9,11\}$	$\{18,19,22,24,28,29,31,32,34,35,36\}$	15	0010001010100000 0110010100011011110
95	$\{3,7,9,14\}$	$\{17,18,19,20,22,23,24,26,28,31,34,36\}$	16	0010001010000100 1111011101001001010
96	$\{3,7,9\}$	$\{17,18,19,20,22,23,24,26,28,29,31,32,34,35,36\}$	18	0010001010000000 1111011101011011110
97	$\{1,3,7,10,16\}$	$\{20,22,23,24,26,27,29,32,36\}$	14	1010001001000001 0001011101101000010
98	$\{1,3,7,10\}$	$\{20,22,23,24,26,27,29,31,32,34,36,37\}$	16	1010001001000000 0001011101101011011
99	$\{3,7,10,11,16\}$	$\{18,19,22,24,27,29,32,36\}$	13	0010001001100001 0110010100101000010
100	$\{3,7,10,11\}$	$\{18,19,22,24,27,29,31,32,34,36,37\}$	15	0010001001100000 0110010100101011011
101	$\{3,7,10,16\}$	$\{17,18,19,20,22,23,24,26,27,29,32,36\}$	16	0010001001000001 1111011101010000010
102	$\{3,7,10\}$	$\{17,18,19,20,22,23,24,26,27,29,31,32,34,36,37\}$	18	0010001001000000 1111011101011011011
103	$\{1,4,5,8,14\}$	$\{20,21,24,25,27,28,30,33,37\}$	14	1001100100000100 0001100111011010010001
104	$\{1,4,5,8\}$	$\{20,21,24,25,27,28,29,30,32,33,35,37\}$	16	1001100100000000 0001100111011110110101
105	$\{4,5,8,12,14\}$	$\{17,19,20,25,28,30,33,37\}$	13	0001100100010100 1011000010010100001
106	$\{4,5,8,12\}$	$\{17,19,20,25,28,29,30,32,33,35,37\}$	15	0001100100010000 1011000010011110110101
107	$\{4,5,8,14\}$	$\{17,18,19,20,21,24,25,27,28,30,33,37\}$	16	0001100100000100 1111100111011010010001
108	$\{4,5,8\}$	$\{17,18,19,20,21,24,25,27,28,29,30,32,33,35,37\}$	18	0001100100000000 1111100111011110110101
109	$\{1,4,5,9,14,15\}$	$\{20,21,24,25,26,28\}$	12	1001100010000110 0001100111010000000000
110	$\{1,4,5,9,14\}$	$\{20,21,24,25,26,28,30,33,36\}$	14	1001100010000100 0001100111010010010010
111	$\{1,4,5,9,15\}$	$\{20,21,24,25,26,28,29,32,35\}$	14	1001100010000010 0001100111011100100100
112	$\{1,4,5,9\}$	$\{20,21,24,25,26,28,29,30,32,33,35,36\}$	16	1001100010000000 0001100111011110110110
113	$\{4,5,9,14,15\}$	$\{17,18,19,20,21,24,25,26,28\}$	14	0001100010000110 1111100111010000000000
114	$\{4,5,9,14\}$	$\{17,18,19,20,21,24,25,26,28,30,33,36\}$	16	0001100010000100 1111100111010010010010

Continued on the next page.

TABLE S74: Canonical enumeration of all 153 two-valued states, in the canonical order defined above. The support is split into core and completion vertices. The last column is $(s_r(1), \ldots, s_r(16)\,;\, s_r(17), \ldots, s_r(37))$. (continued).

r	Core support	Completion support	$\lvert\operatorname{supp} s_r\rvert$	37-bit state vector
115	$\{4,5,9,15\}$	$\{17,18,19,20,21,24,25,26,28,29,32,35\}$	16	0001100010000010 1111100111011001001 00
116	$\{4,5,9\}$	$\{17,18,19,20,21,24,25,26,28,29,30,32,33,35,36\}$	18	0001100010000000 1111100111011101101 10
117	$\{1,4,5,10,15\}$	$\{20,21,24,25,26,27,29,32,37\}$	14	1001100001000010 0001100111101001000 01
118	$\{1,4,5,10\}$	$\{20,21,24,25,26,27,29,30,32,33,36,37\}$	16	1001100001000000 0001100111101101100 11
119	$\{4,5,10,12,15\}$	$\{17,19,20,25,26,29,32,37\}$	13	0001100001010010 1011000011001001000 01
120	$\{4,5,10,12\}$	$\{17,19,20,25,26,29,30,32,33,36,37\}$	15	0001100001010000 1011000011001101100 11
121	$\{4,5,10,15\}$	$\{17,18,19,20,21,24,25,26,27,29,32,37\}$	16	0001100001000010 1111100111101001000 01
122	$\{4,5,10\}$	$\{17,18,19,20,21,24,25,26,27,29,30,32,33,36,37\}$	18	0001100001000000 1111100111101101100 11
123	$\{1,4,6,8\}$	$\{20,21,23,25,27,28,29,30,33,34,35,37\}$	16	1001010100000000 0001101010111001110 01
124	$\{4,6,8\}$	$\{17,18,19,20,21,23,25,27,28,29,30,33,34,35,37\}$	18	0001010100000000 1111101010111001110 01
125	$\{1,4,6,9,15\}$	$\{20,21,23,25,26,28,29,34,35\}$	14	1001010010000010 0001101011010000110 00
126	$\{1,4,6,9\}$	$\{20,21,23,25,26,28,29,30,33,34,35,36\}$	16	1001010010000000 0001101011011001111 10
127	$\{4,6,9,11,15\}$	$\{18,19,21,25,28,29,34,35\}$	13	0001010010100010 0110100010011000011 00
128	$\{4,6,9,11\}$	$\{18,19,21,25,28,29,30,33,34,35,36\}$	15	0001010010100000 0110100010011001111 10
129	$\{4,6,9,15\}$	$\{17,18,19,20,21,23,25,26,28,29,34,35\}$	16	0001010010000010 1111101011011000011 00
130	$\{4,6,9\}$	$\{17,18,19,20,21,23,25,26,28,29,30,33,34,35,36\}$	18	0001010010000000 1111101011011001111 10
131	$\{1,4,6,10,15\}$	$\{20,21,23,25,26,27,29,34,37\}$	14	1001010001000010 0001101011010000100 01
132	$\{1,4,6,10\}$	$\{20,21,23,25,26,27,29,30,33,34,36,37\}$	16	1001010001000000 0001101011010011011 00
133	$\{4,6,10,11,15\}$	$\{18,19,21,25,27,29,34,37\}$	13	0001010001100010 0110100010101000001 00
134	$\{4,6,10,11\}$	$\{18,19,21,25,27,29,30,33,34,36,37\}$	15	0001010001100000 0110100010101100111 01
135	$\{4,6,10,15\}$	$\{17,18,19,20,21,23,25,26,27,29,34,37\}$	16	0001010001000010 1111101011010000100 01
136	$\{4,6,10\}$	$\{17,18,19,20,21,23,25,26,27,29,30,33,34,36,37\}$	18	0001010001000000 1111101011011001111 01
137	$\{1,4,7,8,14\}$	$\{20,21,23,24,27,28,30,34,37\}$	14	1001001100000100 0001101100110100010 01
138	$\{1,4,7,8\}$	$\{20,21,23,24,27,28,29,30,32,34,35,37\}$	16	1001001100000000 0001101100111101011 01
139	$\{4,7,8,12,14\}$	$\{17,19,20,23,28,30,34,37\}$	13	0001001100010100 1011001000010100010 01
140	$\{4,7,8,12\}$	$\{17,19,20,23,28,29,30,32,34,35,37\}$	15	0001001100010000 1011001000011101011 01
141	$\{4,7,8,14\}$	$\{17,18,19,20,21,23,24,27,28,30,34,37\}$	16	0001001100000100 1111101100110100010 01
142	$\{4,7,8\}$	$\{17,18,19,20,21,23,24,27,28,29,30,32,34,35,37\}$	18	0001001100000000 1111101100111101011 01
143	$\{1,4,7,9,14\}$	$\{20,21,23,24,26,28,30,34,36\}$	14	1001001010000100 0001101101010100101 00
144	$\{1,4,7,9\}$	$\{20,21,23,24,26,28,29,30,32,34,35,36\}$	16	1001001010000000 0001101101011101011 10
145	$\{4,7,9,11,14\}$	$\{18,19,21,24,28,30,34,36\}$	13	0001001010100100 0110100100010100101 00
146	$\{4,7,9,11\}$	$\{18,19,21,24,28,29,30,32,34,35,36\}$	15	0001001010100000 0110100100011101111 10
147	$\{4,7,9,14\}$	$\{17,18,19,20,21,23,24,26,28,30,34,36\}$	16	0001001010000100 1111101101010100101 00
148	$\{4,7,9\}$	$\{17,18,19,20,21,23,24,26,28,29,30,32,34,35,36\}$	18	0001001010000000 1111101101011101011 10
149	$\{1,4,7,10\}$	$\{20,21,23,24,26,27,29,30,32,34,36,37\}$	16	1001001001000000 0001101101011010101 11
150	$\{4,7,10,11,12\}$	$\{19,29,30,32,34,36,37\}$	12	0001001001110000 0010000000001101011 11
151	$\{4,7,10,11\}$	$\{18,19,21,24,27,29,30,32,34,36,37\}$	15	0001001001100000 0110100100101101011 11
152	$\{4,7,10,12\}$	$\{17,19,20,23,26,29,30,32,34,36,37\}$	15	0001001001010000 1011001001001101011 11
153	$\{4,7,10\}$	$\{17,18,19,20,21,23,24,26,27,29,30,32,34,36,37\}$	18	0001001001000000 1111101101011010101 11

S2.27.3. A minimum six-state unital, separating family

Define $\Omega_6 = \{\sigma_1, \ldots, \sigma_6\}$ in the following order:

$$(\sigma_1, \ldots, \sigma_6) = (s_3, s_{31}, s_{75}, s_{89}, s_{113}, s_{151}). \tag{S15}$$

Table S75 prints their complete supports. Table S76 gives all 37 profiles $(\sigma_1(i), \ldots, \sigma_6(i))$ and their active state sets. Every profile is nonzero and no two coincide, so Ω_6 is unital and separating. Since five states supply at most $2^5 - 1 = 31 < 37$ nonzero binary profiles, no smaller unital, separating family can exist.

TABLE S75: Complete supports and state vectors of the minimum unital, separating family Ω_6. The full state indices are those of Table S74.

State	Full index	Complete support	Weight	37-bit vector
σ_1	s_3	$\{2,5,8,12,17,19,22,25,28,30,31,32,33,35,37\}$	15	0100100100010000 1010010010011110101
σ_2	s_{31}	$\{1,2,6,10,15,21,22,23,25,26,27,31,34,37\}$	14	1100010001000010 0000111011100010010001
σ_3	s_{75}	$\{3,6,9,13,17,18,20,23,26,29,31,33,34,35,36\}$	15	0010010010001000 1101001001001011110
σ_4	s_{89}	$\{3,7,8,16,17,18,19,20,22,23,24,27,28,29,32,35\}$	16	0010001100000001 1111011100111001001 00
σ_5	s_{113}	$\{4,5,9,14,15,17,18,19,20,21,24,25,26,28\}$	14	0001100010000110 1111100111010000000 0
σ_6	s_{151}	$\{4,7,10,11,18,19,21,24,27,29,30,32,34,36,37\}$	15	0001001001100000 0110100100101101010 11

TABLE S76: All 37 vertex profiles on Ω_6, in two panels. Here $X_i^{(6)} = \{j : \sigma_j(i) = 1\}$, $W_6 = |X_i^{(6)}|$, and W_{153} is the weight across the full state space. The six-bit rows are nonzero and pairwise distinct.

i	Ray	$X_i^{(6)}$	Profile	W_6	W_{153}	i	Ray	$X_i^{(6)}$	Profile	W_6	W_{153}
1	z	$\{2\}$	010000	1	54	20	B_{00}	$\{3,4,5\}$	001110	3	86
2	M_{00}	$\{1,2\}$	110000	2	51	21	B_{01}	$\{2,5,6\}$	010011	3	86
3	M_{01}	$\{3,4\}$	001100	2	51	22	B_{02}	$\{1,2,4\}$	110100	3	86
4	M_{02}	$\{5,6\}$	000011	2	51	23	B_{10}	$\{2,3,4\}$	011100	3	86
5	M_{10}	$\{1,5\}$	100010	2	51	24	B_{11}	$\{4,5,6\}$	000111	3	86
6	M_{11}	$\{2,3\}$	011000	2	51	25	B_{12}	$\{1,2,5\}$	110010	3	86
7	M_{12}	$\{4,6\}$	000101	2	51	26	B_{20}	$\{2,3,5\}$	011010	3	86
8	M_{20}	$\{1,4\}$	100100	2	51	27	B_{21}	$\{2,4,6\}$	010101	3	86
9	M_{21}	$\{3,5\}$	001010	2	51	28	B_{22}	$\{1,4,5\}$	100110	3	86
10	M_{22}	$\{2,6\}$	010001	2	51	29	C_{00}	$\{3,4,6\}$	001101	3	76
11	P_0	$\{6\}$	000001	1	16	30	C_{01}	$\{1,6\}$	100001	2	76
12	P_1	$\{1\}$	100000	1	16	31	C_{02}	$\{1,2,3\}$	111000	3	76
13	P_2	$\{3\}$	001000	1	16	32	C_{10}	$\{1,4,6\}$	100101	3	76
14	Q_0	$\{5\}$	000010	1	26	33	C_{11}	$\{1,3\}$	101000	2	76
15	Q_1	$\{2,5\}$	010010	2	26	34	C_{12}	$\{2,3,6\}$	011001	3	76
16	Q_2	$\{4\}$	000100	1	26	35	C_{20}	$\{1,3,4\}$	101100	3	76
17	A_0	$\{1,3,4,5\}$	101110	4	83	36	C_{21}	$\{3,6\}$	001001	2	76
18	A_1	$\{3,4,5,6\}$	001111	4	83	37	C_{22}	$\{1,2,6\}$	110001	3	76
19	A_2	$\{1,4,5,6\}$	100111	4	83						

S2.27.4. All full 153-state vertex profiles

Table S77 prints the transpose of the complete state table: the full 153-bit profile of every vertex, in three consecutive blocks of 51 states. Thus it also enumerates the complete active state set $X_i = \{s_r : s_r(i) = 1\}$ for every vertex: a one in position r means exactly $s_r \in X_i$. The row weights are 54 for z, 51 for M, 16 for P, 26 for Q, 83 for A, 86 for B, and 76 for C.

TABLE S77: Complete 153-bit vertex profiles, with consecutive 51-bit blocks. Every row is nonzero and all 37 rows are distinct.

i	Ray	W_{153}	$s_1 \ldots s_{51}$	$s_{52} \ldots s_{102}$	$s_{103} \ldots s_{153}$
1	z	54	100001100001100001100001100001111000011000010110000	110000110000101100001000011000011110000110000110000	110000111100001100001011000011000011000011000010000
2	M_{00}	51	111111111111111111111111111111111111111111111111111	000000000000000000000000000000000000000000000000000	000000000000000000000000000000000000000000000000000
3	M_{01}	51	000000000000000000000000000000000000000000000000000	111111111111111111111111111111111111111111111111111	000000000000000000000000000000000000000000000000000
4	M_{02}	51	000000000000000000000000000000000000000000000000000	000000000000000000000000000000000000000000000000000	111111111111111111111111111111111111111111111111111
5	M_{10}	51	111111111111111110000000000000000000000000000000000	111111111111110000000000000000000000000000000000000	111111111111111111000000000000000000000000000000000
6	M_{11}	51	000000000000000011111111111111111111000000000000000	000000000000011111111111111111000000000000000000000	000000000000000000111111111111111000000000000000000
7	M_{12}	51	000000000000000000000000000000000000111111111111111	000000000000000000000000000000001111111111111111111	000000000000000000000000000000001111111111111111111
8	M_{20}	51	111110000000000011111100000000000000011111100000000	111111000000001111110000000000111111100000000000000	111111000000000000011000000000001111110000000000000
9	M_{21}	51	000001111110000000000111111000000000000011000000000	000001111110000000111100000000000011111100000000000	000001111111100000000111111000000000000111100000000
10	M_{22}	51	000000000011111100000000000111111110000000111111000	000000000011000000000011111100000000000111111000000	000000000000111111000000011111100000000000111111000
11	P_0	16	000000000000000000000000000000000000000000000000000	000000000000000000001100001100000000001100001100000	000000000000000000000011000011000000000011000011000
12	P_1	16	011000000000011000000000000000000000011000001100000	000000000000000000000000000000000000000000000000000	001100000000000011000000000000000000001100000000001010
13	P_2	16	010100011000000000011000011000000000000000000000000	001100001100000011000101000000000000000000000000000	000000000000000000000000000000000000000000000000000000
14	Q_0	26	000000000000000000000000000000000000000000000000000000	101010101010000000000000000000011001100101010000000	101010110011000000000000000000000010101010101000000
15	Q_1	26	000001010101010100000001010101100110000000000000000	000000000000000000000000000000000000000000000000000000	000000101010101010100010101010101000000000000000000
16	Q_2	26	000000000000000010101000000101010101010010010100000	000000000000010101000000101010101010000000101010000	000000000000000000000000000000000000000000000000000000
17	A_0	83	011110011110011100111001111000011110011110100111100	111110011101001111000110000110000111100001100001100	111110000111001111010000110000110011110000110001100
18	A_1	83	000110011110000110011100111100001111000011010000110	111110011101001111011110011100000111001111001111101	000011000011110000110100111100111100001100111100101
19	A_2	83	001010000110011110000110000110000111100111101001111	000011000011010000011001010011110000111100111100111	001111000011110011110100111100111100001100111101111
20	B_{00}	86	000000000000000000000000000000000000000000000000000000	111111111111111111111001110011111111111110011110011	111111111111111111111111001110011111111111001110011
21	B_{01}	86	100111111111100111111111111111111111111111001111110011	000000000000000000000000000000000000000000000000000000	110011111111111001111111111111111111001111111110101
22	B_{02}	86	101011100111111111100111100111111111111111111111111111	110011110011111100111010111111111111111111111111111111	000000000000000000000000000000000000000000000000000000
23	B_{10}	86	000000000000000011111111111111111111111111111111111111	000000000000011111100111001111111111100111100111111111	000000000000000000011100111001111111111001110011111111
24	B_{11}	86	100111111111100110000000000000000000011001111110011	111111111111110000000000000000011111111111111111111	110011111111111001100000000000000110011111111110101
25	B_{12}	86	101011100111111111100111100111111111100000000000000	110011110011111100111010111111100000000000000000000	111111111111111111111111111111111111111100000000000000
26	B_{20}	86	000001111111111110000001111111111111000000111111111	000001111111100000010011110011000000011001111100011	000000111111111111110011001111001100000011001110011
27	B_{21}	86	100110000001100111111100000011111111111001100110011	111111000000111111110000011111111111111000000111111	110011000000001100111100000011111111100110000010101
28	B_{22}	86	101011100110000001100111100110000000011111111100000	110011110011001100111010100000011111111111111000000	111111111111110000001111111100000011111111111100000
29	C_{00}	76	000000000000000000000000000000000000000000000000000000	010101010101111111111111111111100110011010101111111	010101001100111111111111111111111110101010101011111
30	C_{01}	76	111110101010101011111110101010011001111111111111111	000000000000000000000000000000000000000000000000000000	111111010101010101011101010101010111111111111111111
31	C_{02}	76	111111111111111101010111111101010101010101011010101	111111111111110101011111101010101010101111111010101	000000000000000000000000000000000000000000000000000000
32	C_{10}	76	111111111111111100000000000000000000111111111111111	010101010101100000000000000000011001101010111111	010101001100111111100000000000000101010101011111
33	C_{11}	76	111110101010101011111101010100110011000000000000000	111111111111111111111111111111110000000000000000000	111111010101010101110101010101010000000000000000000
34	C_{12}	76	000000000000000001010111111101010101010101011010101	000000000000000101011111101010101010101111111010101	000000000000000000001111111111111111111111111111111
35	C_{20}	76	111111111110000001111111111100000000111111110000000	010101010101001111111111100000000110011010101000000	010101001100110000011111111100000001010101010100000
36	C_{21}	76	000000101010101010000000101010011001100000011111111	000000111111110000001111111111100000000111111111111	000000010101010101010001010101010100000011111111111
37	C_{22}	76	111110000001111101010100000001010101010100010101	111111000000110101010000001010101010100000010101	111111000000011111111100000011111111111100000011111

S2.27.5. The six-state double cover

The fractional-coloring certificate uses a second six-state family,

$$\mathcal{R}_6 = \{s_2, s_{30}, s_{73}, s_{99}, s_{109}, s_{139}\}. \tag{S16}$$

Table S78 specifies these states in the branch parametrization. Each of the 37 vertices occurs in exactly two of their supports. Thus weights 1/2 give a fractional coloring of total weight three, and uniform weights 1/6 reproduce the maximally mixed probabilities 1/3 for every ray. The family Ω_6 in Eq. (S15) supplies the separating certificate; $\mathcal{R}_6$ supplies the double-cover certificate.

TABLE S78: Explicit six-state double cover $\mathcal{R}_6$. Omitted P, Q values are zero; completion values follow from the state equations.

j	State	t	ε	Active P subscripts	Active Q subscripts
1	s_2	$(0,0,0)$	0	$\{1,2\}$	$\varnothing$
2	s_{30}	$(0,1,2)$	1	$\varnothing$	$\{1,2\}$
3	s_{73}	$(1,1,1)$	0	$\{0,2\}$	$\varnothing$
4	s_{99}	$(1,2,2)$	0	$\{0\}$	$\{2\}$
5	s_{109}	$(2,0,1)$	1	$\varnothing$	$\{0,1\}$
6	s_{139}	$(2,2,0)$	0	$\{1\}$	$\{0\}$

S2.27.6. Order determination and exact verification

The companion file `sources/hesse/hesse37-canonical-evaluations.txt` records the 24 numbered contexts and all 153 binary state vectors. The parametrization in this appendix defines their canonical order independently of any external program.

Extend the canonical states to the 76 events of $L(H_{37})$ as in Sec. V A. If $X_x = \{r : s_r(x) = 1\}$, then the full profiles satisfy

$$X_x \subseteq X_y \quad \Longleftrightarrow \quad x \leq y. \tag{S17}$$

The verifier checks all $76^2 = 5{,}776$ ordered event pairs against the incidence order in Eq. (S27). Whenever $x \not\leq y$, the companion file `sources/hesse/hesse37-order-witnesses.csv` gives the least canonical index r with $s_r(x) = 1$ and $s_r(y) = 0$. Its event labels `a1`,...,`a37` denote atoms and `c1`,...,`c37` their complements; `0` and `1` denote the bounds. Every one of the 594 distinct nonorthogonal atom pairs also has a joint-one state. Thus the full family is order determining; the six-state family Ω_6 supplies the minimum unital, separating vertex representation in Eq. (S26). The distinct minima for event separation and order determination are certified in Appendix S3.3.

The verifier `hesse37-verify.py` uses exact rational arithmetic in $\mathbb{Q}(\zeta)$, with no numerical tolerance or external solver. It checks distinct rays, all 666 Hermitian inner products, the 24 bases, all graph triangles, linearity, the chromatic number and core criticality, all 153 states, the minimum separating family, both profile tables, the maximum squared ray overlap, the six-state double cover, and all event-order comparisons. It also compares the reconstructed state vectors with the archived enumeration. The accompanying machine-readable data give the same numbering and context ordering as the figure and these appendices. From the extracted joint project directory, run

```
python verification/verify-integrated.py
```

S2.28. Four-dimensional $H_{\mathrm{YO}}^{(4)}$ and H_{65}: states and profiles

This appendix provides the complete enumeration of all two-valued states on the four-dimensional examples $H_{\mathrm{YO}}^{(4)}$ and H_{65}: the 25 two-valued states of the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)}$ (Appendix S2.28.1), and the complete branch-by-branch parametrization of all $24 \times 2^{14} = 393{,}216$ two-valued states—together with the 48 canonical unital and separating states $s_{\alpha,\varepsilon}$ and the 65 vertex state-profiles—of the canonical linear hypergraph H_{65} (Appendix S2.28.2).

S2.28.1. Complete enumeration of the 25 two-valued states on $H_{\mathrm{YO}}^{(4)}$

Table S79 lists all 25 two-valued states $s_1, \ldots, s_{25} : V_{26} \to \{0,1\}$ on $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$. States $s_1, \ldots, s_{24}$ have $s_r(26) = 0$ and correspond to the 24 two-valued states of the completed

Yu–Oh hypergraph H_{YO}^{+} in $\mathbb{R}^3$ (grouped into 12 pairs of states sharing the same support on the three coordinate contexts $\{1,2,3\}$, $\{4,5,6\}$, $\{7,8,9\}$), while s_{25} is the unique apex state with support $\{26\}$. Table S80 displays the transposed 26×25 vertex-by-state binary matrix, listing for each vertex $i \in \{1,\dots,26\}$ its state weight $\sum_{r=1}^{25} s_r(i) \in \{1,3,8,13\}$, the set of states $A_i = \{s_r : s_r(i) = 1\}$, and its 25-bit profile $(s_1(i),\dots,s_{25}(i))$.

Inspection of Table S80 highlights three structural properties of the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$:

1. *Unitality and atom separation:* Every row of Table S80 has positive weight (1, 3, 8, or 13) and all 26 binary rows are distinct ($A_i \neq A_j$ for all $1 \le i < j \le 26$), so $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is nonempty, unital, and separating on atoms.

2. *Exclusion of simultaneous* 1*s on the non-orthogonal triad atoms* $\{10,11,12,13\}$*:* Although the four Yu–Oh triad rays $v_{10}, v_{11}, v_{12}, v_{13}$ are pairwise non-orthogonal ($v_i \cdot v_j = \pm 1 \neq 0$ for $10 \le i < j \le 13$), their active state sets in Table S80 are pairwise disjoint:

$$\begin{aligned} A_{10} &= \{s_{13}, s_{19}, s_{23}\}, \quad A_{11} = \{s_5, s_7, s_{15}\}, \\ A_{12} &= \{s_1, s_3, s_9\}, \qquad A_{13} = \{s_{11}, s_{17}, s_{21}\}. \end{aligned} \tag{S18}$$

   Consequently, $\sum_{i=10}^{13} s(i) \le 1$ holds for every $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$, and no two-valued state assigns $(1,1)$ to any of the $\binom{4}{2} = 6$ non-orthogonal pairs $\{10,11\}$, $\{10,12\}$, $\{10,13\}$, $\{11,12\}$, $\{11,13\}$, $\{12,13\}$.

3. *Failure of order-determinacy:* Even though $A_i \neq A_j$ for all $i \neq j$, the state space $\mathcal{S}(H_{\mathrm{YO}}^{(4)})$ is not order-determining on atoms or compound propositions[4]. In particular, each triad atom $t \in \{10,11,12,13\}$ (*weight* 3) is strictly dominated across all 25 states ($A_t \subsetneq A_w$, i.e., $s(t) \le s(w)$ for all $s \in \mathcal{S}(H_{\mathrm{YO}}^{(4)})$) by three non-orthogonal completion atoms $w \in \{14,\dots,25\}$ (*weight* 13):

$$\begin{aligned} A_{10} &\subsetneq A_{16},\, A_{20},\, A_{24}, \quad A_{11} \subsetneq A_{18},\, A_{21},\, A_{25}, \\ A_{12} &\subsetneq A_{14},\, A_{22},\, A_{23}, \quad A_{13} \subsetneq A_{15},\, A_{17},\, A_{19}, \end{aligned} \tag{S19}$$

   together with compound-event inclusions such as $A_{12} \subsetneq A_3 \,\dot\cup\, A_{14}$, $A_{11} \subsetneq A_3 \,\dot\cup\, A_{25}$, and their unions with the apex set $A_{26} = \{s_{25}\}$.

TABLE S79: Complete enumeration of the 25 two-valued states $s_1,\dots,s_{25} : V_{26} \to \{0,1\}$ on the non-linear warm-up hypergraph $H_{\mathrm{YO}}^{(4)} = (V_{26}, E_{16}^{(4)})$, showing for each state its support on the 13-ray Yu–Oh core $\{1,\dots,13\}$, its forced support on $\{14,\dots,26\}$, its total support size $|S_r|$, and its 26-bit binary vector $(s_r(1\dots13)\, s_r(14\dots25)\, s_r(26))$.

State	Core support	Completion/apex support	$\lvert S_r\rvert$	26-bit state vector
s_1	$\{1,4,8,12\}$	$\{14,15,21,22,23\}$	9	1001000100010 110000011100 0
s_2	$\{1,4,8\}$	$\{14,15,16,17,21,22,23,25\}$	11	1001000100000 111100011101 0
s_3	$\{1,5,7,12\}$	$\{14,18,22,23,24\}$	9	1000101000010 100010001110 0
s_4	$\{1,5,7\}$	$\{14,16,17,18,22,23,24,25\}$	11	1000101000000 101110001111 0
s_5	$\{1,5,9,11\}$	$\{17,18,21,22,25\}$	9	1000100010100 000110011001 0
s_6	$\{1,5,9\}$	$\{14,15,17,18,21,22,24,25\}$	11	1000100010000 110110011011 0
s_7	$\{1,6,8,11\}$	$\{16,18,21,23,25\}$	9	1000010100100 001010010101 0
s_8	$\{1,6,8\}$	$\{14,15,16,18,21,23,24,25\}$	11	1000010100000 111010010111 0
s_9	$\{2,4,7,12\}$	$\{14,19,20,22,23\}$	9	0101001000010 100001101100 0
s_{10}	$\{2,4,7\}$	$\{14,16,17,19,20,22,23,25\}$	11	0101001000000 101101101101 0
s_{11}	$\{2,4,9,13\}$	$\{14,15,17,19,25\}$	9	0101000010001 110101000001 0
s_{12}	$\{2,4,9\}$	$\{14,15,17,19,20,21,22,25\}$	11	0101000010000 110101111001 0
s_{13}	$\{2,6,7,10\}$	$\{14,16,20,24,25\}$	9	0100011001000 101000100011 0
s_{14}	$\{2,6,7\}$	$\{14,16,18,19,20,23,24,25\}$	11	0100011000000 101011100111 0
s_{15}	$\{2,6,9,11\}$	$\{18,19,20,21,25\}$	9	0100010010100 000011110001 0

Continued on the next page.

TABLE S79: All 25 states of $H_{\mathrm{YO}}^{(4)}$ (continued).

State	Core support	Completion/apex support	$\lvert S_r\rvert$	26-bit state vector
s_{16}	$\{2,6,9\}$	$\{14,15,18,19,20,21,24,25\}$	11	0100010010000 110011110011 0
s_{17}	$\{3,4,8,13\}$	$\{15,16,17,19,23\}$	9	0011000100001 011101000100 0
s_{18}	$\{3,4,8\}$	$\{15,16,17,19,20,21,22,23\}$	11	0011000100000 011101111100 0
s_{19}	$\{3,5,7,10\}$	$\{16,17,20,22,24\}$	9	0010101001000 001100101010 0
s_{20}	$\{3,5,7\}$	$\{16,17,18,19,20,22,23,24\}$	11	0010101000000 001111101110 0
s_{21}	$\{3,5,9,13\}$	$\{15,17,18,19,24\}$	9	0010100010001 010111000010 0
s_{22}	$\{3,5,9\}$	$\{15,17,18,19,20,21,22,24\}$	11	0010100010000 010111111010 0
s_{23}	$\{3,6,8,10\}$	$\{15,16,20,21,24\}$	9	0010010101000 011000110010 0
s_{24}	$\{3,6,8\}$	$\{15,16,18,19,20,21,23,24\}$	11	0010010100000 011011110110 0
s_{25}	$\emptyset$	$\{26\}$	1	0000000000000 000000000000 1

TABLE S80: Vertex state profiles across the 25 two-valued states $s_1,\dots,s_{25}$ of $H_{\mathrm{YO}}^{(4)}$, showing for each vertex $i \in \{1,\dots,26\}$ its state weight $\sum_{r=1}^{25} s_r(i)$, the active state indices $\{r : s_r(i)=1\}$, and the 25-bit profile $(s_{1\dots12}(i)\, s_{13\dots24}(i)\, s_{25}(i))$. All 26 rows are nonzero and pairwise distinct.

Vertex i	Weight $\sum_r s_r(i)$	Active states $\{r \in \{1,\dots,25\} : s_r(i)=1\}$	25-bit profile $(s_{1\dots12}(i)\, s_{13\dots24}(i)\, s_{25}(i))$
1	8	$\{1,2,3,4,5,6,7,8\}$	111111110000 000000000000 0
2	8	$\{9,10,11,12,13,14,15,16\}$	000000001111 111100000000 0
3	8	$\{17,18,19,20,21,22,23,24\}$	000000000000 0000111111111 0
4	8	$\{1,2,9,10,11,12,17,18\}$	110000001111 000011000000 0
5	8	$\{3,4,5,6,19,20,21,22\}$	001111000000 000000111100 0
6	8	$\{7,8,13,14,15,16,23,24\}$	000000110000 111100000011 0
7	8	$\{3,4,9,10,13,14,19,20\}$	001100001100 110000110000 0
8	8	$\{1,2,7,8,17,18,23,24\}$	110000110000 000011000011 0
9	8	$\{5,6,11,12,15,16,21,22\}$	000011000011 001100001100 0
10	3	$\{13,19,23\}$	000000000000 100000100010 0
11	3	$\{5,7,15\}$	000010100000 001000000000 0
12	3	$\{1,3,9\}$	101000001000 000000000000 0
13	3	$\{11,17,21\}$	000000000010 000010001000 0
14	13	$\{1,2,3,4,6,8,9,10,11,12,13,14,16\}$	111101011111 110100000000 0
15	13	$\{1,2,6,8,11,12,16,17,18,21,22,23,24\}$	110001010011 000111001111 0
16	13	$\{2,4,7,8,10,13,14,17,18,19,20,23,24\}$	010100110100 110011110011 0
17	13	$\{2,4,5,6,10,11,12,17,18,19,20,21,22\}$	010111000111 000011111100 0
18	13	$\{3,4,5,6,7,8,14,15,16,20,21,22,24\}$	001111110000 011100011101 0
19	13	$\{9,10,11,12,14,15,16,17,18,20,21,22,24\}$	000000001111 011111011101 0
20	13	$\{9,10,12,13,14,15,16,18,19,20,22,23,24\}$	000000001101 111101110111 0
21	13	$\{1,2,5,6,7,8,12,15,16,18,22,23,24\}$	110011110001 001101000111 0
22	13	$\{1,2,3,4,5,6,9,10,12,18,19,20,22\}$	111111001101 000001110100 0
23	13	$\{1,2,3,4,7,8,9,10,14,17,18,20,24\}$	111100111100 010011010001 0
24	13	$\{3,4,6,8,13,14,16,19,20,21,22,23,24\}$	001101010000 110100111111 0
25	13	$\{2,4,5,6,7,8,10,11,12,13,14,15,16\}$	010111110111 111100000000 0
26	1	$\{25\}$	000000000000 000000000000 1

S2.28.2. Complete parametrization of all 393,216 *two-valued states and the* 48 *canonical separating states on* H_{65}

Every two-valued state $s \in \mathcal{S}(H_{65})$ on the canonical linear hypergraph $H_{65} = (V_{65}, E_{29})$ is uniquely indexed by a pair $(\alpha, \boldsymbol{\varepsilon})$ consisting of:

1. a branch index $\alpha \in \{1,\dots,24\}$ selecting one of the 24 four-element core state supports $S_\alpha \subset V_{\mathrm{core}} = \{1,\dots,17\}$; and
2. a 14-bit binary vector $\boldsymbol{\varepsilon} = (\varepsilon_m)_{m\in\mathcal{F}_\alpha} \in \{0,1\}^{14}$ indexed by the 14 *free completion contexts*

$$\mathcal{F}_\alpha = \big\{m \in \{1,\dots,24\} : S_\alpha \cap \{u_m, v_m\} = \emptyset\big\},$$

whose complement is the set of 10 *hit completion contexts* $\mathcal{H}_\alpha = \{1,\dots,24\} \setminus \mathcal{F}_\alpha$.

Explicitly, the state $s_{\alpha,\boldsymbol{\varepsilon}} : V_{65} \to \{0,1\}$ takes values

$$\begin{aligned} s_{\alpha,\boldsymbol{\varepsilon}}(i) &= \mathbf{1}_{i\in S_\alpha} \qquad (1 \le i \le 17), \\ \big(s_{\alpha,\boldsymbol{\varepsilon}}(2m+16),\, s_{\alpha,\boldsymbol{\varepsilon}}(2m+17)\big) &= \begin{cases} (0,0), & m \in \mathcal{H}_\alpha, \\ (1-\varepsilon_m,\, \varepsilon_m), & m \in \mathcal{F}_\alpha. \end{cases} \end{aligned} \tag{S20}$$

Because $|\mathcal{F}_\alpha| = 14$ for every $\alpha \in \{1,\dots,24\}$, each branch α yields $2^{14} = 16{,}384$ states $s_{\alpha,\boldsymbol{\varepsilon}}$, giving $|\mathcal{S}(H_{65})| = 24 \times 2^{14} = 393{,}216$ two-valued states in total. Setting all 14 free bits equal to $\varepsilon \in \{0,1\}$ ($\varepsilon_m = \varepsilon$ for all $m \in \mathcal{F}_\alpha$) yields the 48 *canonical states* $s_{\alpha,0}, s_{\alpha,1}$ of Eq. (S29), which have support size $|S_\alpha| + 14 = 18$:

$$\begin{aligned} \operatorname{supp}(s_{\alpha,0}) &= S_\alpha \cup \big\{2m+16 : m \in \mathcal{F}_\alpha\big\}, \\ \operatorname{supp}(s_{\alpha,1}) &= S_\alpha \cup \big\{2m+17 : m \in \mathcal{F}_\alpha\big\}. \end{aligned} \tag{S21}$$

Table S81 enumerates all 24 branches $\alpha = 1,\dots,24$, specifying S_α, the 10 hit context indices $\mathcal{H}_\alpha \subset \{1,\dots,24\}$, the 14 free context indices $\mathcal{F}_\alpha \subset \{1,\dots,24\}$, and the 24-bit free-context indicator string. Table S82 lists the resulting state profiles and weights for all 65 vertices of H_{65} across both the 48 canonical states $\{s_{\alpha,\varepsilon}\}$ and the full space of 393,216 states: across the 48 canonical states $(s_{1,0}, s_{1,1}, \dots, s_{24,0}, s_{24,1})$, each '1' in row i of Table S82 expands to '(1,1)' for core vertices $i \in \{1,\dots,17\}$, to '(1,0)' for first completion vertices $w_{m,1} = 2m+16$, and to '(0,1)' for second completion vertices $w_{m,2} = 2m+17$ (and each '0' expands to '(0,0)'), making all 65 canonical 48-bit vertex profiles nonzero and pairwise distinct by inspection.

TABLE S81: Complete branch-by-branch parametrization of the $24 \times 2^{14} = 393{,}216$ two-valued states $s_{\alpha,\boldsymbol{\varepsilon}}$ and the 48 canonical states $s_{\alpha,0}, s_{\alpha,1}$ on H_{65} via Eqs. (S20)–(S21). For each $\alpha \in \{1,\dots,24\}$, the table lists the 4-element core support S_α, the 10 hit completion context indices $m \in \mathcal{H}_\alpha$ where $(s(2m+16), s(2m+17)) = (0,0)$ is forced on E_m, the 14 free completion context indices $m \in \mathcal{F}_\alpha$ where $(s(2m+16), s(2m+17)) = (1-\varepsilon_m, \varepsilon_m)$ with $\varepsilon_m \in \{0,1\}$ on E_m, and the 24-bit indicator string $(\mathbf{1}_{1\in\mathcal{F}_\alpha} \dots \mathbf{1}_{12\in\mathcal{F}_\alpha}\, \mathbf{1}_{13\in\mathcal{F}_\alpha} \dots \mathbf{1}_{24\in\mathcal{F}_\alpha})$.

Branch	Core support S_α	Hit contexts $\mathcal{H}_\alpha$	Free contexts $\mathcal{F}_\alpha$	Free indicator $m = 1\dots24$
S_1	{1,5,9,16}	{3,7,9,10,11,13,19,21,22,23}	{1,2,4,5,6,8,12,14,15,16,17,18,20,24}	110111010001 011111010001
S_2	{1,5,10,14}	{3,5,6,7,11,13,17,18,19,23}	{1,2,4,8,9,10,12,14,15,16,20,21,22,24}	110100011101 011100011101
S_3	{1,6,9,16}	{3,7,9,10,11,15,17,21,22,24}	{1,2,4,5,6,8,12,13,14,16,18,19,20,23}	110111010001 110101110010
S_4	{1,6,11,12}	{1,2,3,7,11,13,14,15,17,24}	{4,5,6,8,9,10,12,16,18,19,20,21,22,23}	000111011101 000101111110
S_5	{1,7,10,14}	{3,5,6,7,11,16,17,18,20,21}	{1,2,4,8,9,10,12,13,14,15,19,22,23,24}	110100011101 111000100111
S_6	{1,7,11,12}	{1,2,3,7,11,13,14,16,20,21}	{4,5,6,8,9,10,12,15,17,18,19,22,23,24}	000111011101 001011100111
S_7	{2,5,9,16}	{1,5,9,10,12,13,19,21,22,23}	{2,3,4,6,7,8,11,14,15,16,17,18,20,24}	011101110010 011111010001
S_8	{2,5,11,13}	{1,3,4,5,12,13,15,16,19,23}	{2,6,7,8,9,10,11,14,17,18,20,21,22,24}	010001111110 010011011101
S_9	{2,6,9,16}	{1,5,9,10,12,15,17,21,22,24}	{2,3,4,6,7,8,11,13,14,16,18,19,20,23}	011101110010 110101110010
S_{10}	{2,6,10,15}	{1,5,7,8,12,15,17,19,20,24}	{2,3,4,6,9,10,11,13,14,16,18,21,22,23}	011101001110 110101001110
S_{11}	{2,8,10,15}	{1,5,7,8,12,14,18,19,20,22}	{2,3,4,6,9,10,11,13,15,16,17,21,23,24}	011101001110 101110001011
S_{12}	{2,8,11,13}	{1,3,4,5,12,14,15,16,18,22}	{2,6,7,8,9,10,11,13,17,19,20,21,23,24}	010001111110 100010111011
S_{13}	{3,5,10,14}	{2,5,6,8,9,13,17,18,19,23}	{1,3,4,7,10,11,12,14,15,16,20,21,22,24}	101100100111 011100011101
S_{14}	{3,5,11,13}	{2,3,4,8,9,13,15,16,19,23}	{1,5,6,7,10,11,12,14,17,18,20,21,22,24}	100011100111 010011011101
S_{15}	{3,7,9,17}	{2,8,9,11,12,16,20,21,23,24}	{1,3,4,5,6,7,10,13,14,15,17,18,19,22}	101111100100 111011100100
S_{16}	{3,7,10,14}	{2,5,6,8,9,16,17,18,20,21}	{1,3,4,7,10,11,12,13,14,15,19,22,23,24}	101100100111 111000100111
S_{17}	{3,8,9,17}	{2,8,9,11,12,14,18,22,23,24}	{1,3,4,5,6,7,10,13,15,16,17,19,20,21}	101111100100 101110111000
S_{18}	{3,8,11,13}	{2,3,4,8,9,14,15,16,18,22}	{1,5,6,7,10,11,12,13,17,19,20,21,23,24}	100011100111 100010111011
S_{19}	{4,6,10,15}	{4,6,7,8,10,15,17,19,20,24}	{1,2,3,5,9,11,12,13,14,16,18,21,22,23}	111010001011 110101001110
S_{20}	{4,6,11,12}	{1,2,4,6,10,13,14,15,17,24}	{3,5,7,8,9,11,12,16,18,19,20,21,22,23}	001010111011 000101111110
S_{21}	{4,7,9,17}	{4,6,10,11,12,16,20,21,23,24}	{1,2,3,5,7,8,9,13,14,15,17,18,19,22}	111010111000 111011100100
S_{22}	{4,7,11,12}	{1,2,4,6,10,13,14,16,20,21}	{3,5,7,8,9,11,12,15,17,18,19,22,23,24}	001010111011 001011100111
S_{23}	{4,8,9,17}	{4,6,10,11,12,14,18,22,23,24}	{1,2,3,5,7,8,9,13,15,16,17,19,20,21}	111010111000 101110111000
S_{24}	{4,8,10,15}	{4,6,7,8,10,14,18,19,20,22}	{1,2,3,5,9,11,12,13,15,16,17,21,23,24}	111010001011 101110001011

TABLE S82: Vertex state profiles and weights for all 65 vertices of H_{65}, showing for each core vertex $i \in \{1,\dots,17\}$ and each completion pair $(w_{m,1}, w_{m,2}) = (2m+16, 2m+17)$ $(m = 1,\dots,24)$ the number of active branches W_{24}, the weight $W_{48} = \sum_{\alpha,\varepsilon} s_{\alpha,\varepsilon}(i)$ across the 48 canonical states, the weight $W_{393,216} = \sum_{s\in\mathcal{S}(H_{65})} s(i)$ across all 393,216 states, the active branch indices $\alpha \in \{1,\dots,24\}$ (equal to $Z(e_m)$ for $(w_{m,1}, w_{m,2})$), and the 24-bit branch profile across $S_1,\dots,S_{24}$ (in which each `1` expands to `11` for core vertices i, to `10` for $w_{m,1}$, and to `01` for $w_{m,2}$ across the 48 canonical states).

Vertex / pair	W_{24}	W_{48}	$W_{393,216}$	Active branches α	24-bit branch profile
Core 1	6	12	98,304	$\{1,2,3,4,5,6\}$	11111100 00000000 00000000
Core 2	6	12	98,304	$\{7,8,9,10,11,12\}$	00000011 11110000 00000000
Core 3	6	12	98,304	$\{13,14,15,16,17,18\}$	00000000 00001111 11000000
Core 4	6	12	98,304	$\{19,20,21,22,23,24\}$	00000000 00000000 00111111
Core 5	6	12	98,304	$\{1,2,7,8,13,14\}$	11000011 00001100 00000000
Core 6	6	12	98,304	$\{3,4,9,10,19,20\}$	00110000 11000000 00110000
Core 7	6	12	98,304	$\{5,6,15,16,21,22\}$	00001100 00000011 00001100
Core 8	6	12	98,304	$\{11,12,17,18,23,24\}$	00000000 00110000 11000011
Core 9	8	16	131,072	$\{1,3,7,9,15,17,21,23\}$	10100010 10000010 10001010
Core 10	8	16	131,072	$\{2,5,10,11,13,16,19,24\}$	01001000 01101001 00100001
Core 11	8	16	131,072	$\{4,6,8,12,14,18,20,22\}$	00010101 00010100 01010100
Core 12	4	8	65,536	$\{4,6,20,22\}$	00010100 00000000 00010100
Core 13	4	8	65,536	$\{8,12,14,18\}$	00000001 00010100 01000000
Core 14	4	8	65,536	$\{2,5,13,16\}$	01001000 00001001 00000000
Core 15	4	8	65,536	$\{10,11,19,24\}$	00000000 01100000 00100001
Core 16	4	8	65,536	$\{1,3,7,9\}$	10100010 10000000 00000000
Core 17	4	8	65,536	$\{15,17,21,23\}$	00000000 00000010 10001010
E_1 : (18, 19)	14	14	114,688	$\{1,2,3,5,13,14,15,16,17,18,19,21,23,24\}$	11101000 00001111 11101011
E_2 : (20, 21)	14	14	114,688	$\{1,2,3,5,7,8,9,10,11,12,19,21,23,24\}$	11101011 11110000 00101011
E_3 : (22, 23)	14	14	114,688	$\{7,9,10,11,13,15,16,17,19,20,21,22,23,24\}$	00000010 11101011 10111111
E_4 : (24, 25)	14	14	114,688	$\{1,2,3,4,5,6,7,9,10,11,13,15,16,17\}$	11111110 11101011 10000000
E_5 : (26, 27)	14	14	114,688	$\{1,3,4,6,14,15,17,18,19,20,21,22,23,24\}$	10110100 00000110 11111111
E_6 : (28, 29)	14	14	114,688	$\{1,3,4,6,7,8,9,10,11,12,14,15,17,18\}$	10110111 11110110 11000000
E_7 : (30, 31)	14	14	114,688	$\{7,8,9,12,13,14,15,16,17,18,20,21,22,23\}$	00000011 10011111 11011110
E_8 : (32, 33)	14	14	114,688	$\{1,2,3,4,5,6,7,8,9,12,20,21,22,23\}$	11111111 10010000 00011110
E_9 : (34, 35)	14	14	114,688	$\{2,4,5,6,8,10,11,12,19,20,21,22,23,24\}$	01011101 01110000 00111111
E_{10} : (36, 37)	14	14	114,688	$\{2,4,5,6,8,10,11,12,13,14,15,16,17,18\}$	01011101 01111111 11000000
E_{11} : (38, 39)	14	14	114,688	$\{7,8,9,10,11,12,13,14,16,18,19,20,22,24\}$	00000011 11111101 01110101
E_{12} : (40, 41)	14	14	114,688	$\{1,2,3,4,5,6,13,14,16,18,19,20,22,24\}$	11111100 00001101 01110101
E_{13} : (42, 43)	14	14	114,688	$\{3,5,9,10,11,12,15,16,17,18,19,21,23,24\}$	00101000 11110011 11101011
E_{14} : (44, 45)	14	14	114,688	$\{1,2,3,5,7,8,9,10,13,14,15,16,19,21\}$	11101011 11001111 00101000
E_{15} : (46, 47)	14	14	114,688	$\{1,2,5,6,7,11,13,15,16,17,21,22,23,24\}$	11001110 00101011 10001111
E_{16} : (48, 49)	14	14	114,688	$\{1,2,3,4,7,9,10,11,13,17,19,20,23,24\}$	11110010 11101000 10110011
E_{17} : (50, 51)	14	14	114,688	$\{1,6,7,8,11,12,14,15,17,18,21,22,23,24\}$	10000111 00110110 11001111
E_{18} : (52, 53)	14	14	114,688	$\{1,3,4,6,7,8,9,10,14,15,19,20,21,22\}$	10110111 11000110 00111100
E_{19} : (54, 55)	14	14	114,688	$\{3,4,5,6,9,12,15,16,17,18,20,21,22,23\}$	00111100 10010011 11011110
E_{20} : (56, 57)	14	14	114,688	$\{1,2,3,4,7,8,9,12,13,14,17,18,20,23\}$	11110011 10011100 11010010
E_{21} : (58, 59)	14	14	114,688	$\{2,4,8,10,11,12,13,14,17,18,19,20,23,24\}$	01010001 01111100 11110011
E_{22} : (60, 61)	14	14	114,688	$\{2,4,5,6,8,10,13,14,15,16,19,20,21,22\}$	01011101 01001111 00111100
E_{23} : (62, 63)	14	14	114,688	$\{3,4,5,6,9,10,11,12,16,18,19,20,22,24\}$	00111100 11110001 01110101
E_{24} : (64, 65)	14	14	114,688	$\{1,2,5,6,7,8,11,12,13,14,16,18,22,24\}$	11001111 00111101 01000101

S2.29. Parent-state and deletion certificates

The 49-ray, 87-ray, 40-ray, and 72-ray parents have no context states. The two smaller 68-ray parents are treated in Appendix S2.30.1. The first and last two have the independently verified complete bases printed in Appendix S1.16; the 87-ray parent contains the known 69-ray KS lineage. The 70-ray single-apex lift of the completed Cabello parent instead has the unique state supported at its apex.

For the real Yu–Oh parent, the intermediate 33 rays admit 192 assignments to the twenty bases alone and none after its sixteen additional pair exclusions. The sixteen missing pairs are uniquely completed by rays 34–49. The x-axis-plane extension by rays 26–29 alone has 72

basis assignments; all 24 original states extend. Including its eight additional pair exclusions leaves eight extensions, all with $s(2) = 1$. The z-axis-plane case is identical, with $s(5) = 1$. The two added bases on either axis each exclude twelve of the sixteen original axis-zero states when their further pair exclusions are included; eight original states are excluded by both, leaving none. Thus the axis forcing uses the complete orthogonality constraints, not the added bases alone.

TABLE S83. Context-state counts after deleting each single context of the real 49-ray parent. Every count is positive. Context numbering is Table S21. Deleting a completion context 21–36 leaves its fresh ray in no context; that ray is counted as an unconstrained free vertex, which doubles the count. If the orthogonalities of the deleted context are kept as geometric exclusions, deletions 21–36 leave no state, while deletions 1–20 leave 1, 2, or 4 states.

Deleted context	States	Deleted context	States	Deleted context	States
1	12	13	2	25	2
2	4	14	4	26	2
3	4	15	4	27	2
4	32	16	16	28	2
5	2	17	4	29	2
6	2	18	4	30	2
7	4	19	4	31	2
8	4	20	4	32	2
9	12	21	2	33	2
10	4	22	2	34	2
11	4	23	2	35	2
12	2	24	2	36	2

All complete states of H_{65} are additionally supplied as `data/h65-all-states.csv.gz`. Each row gives its branch index, the fourteen-bit choice index within that branch, and its 65-bit state vector in vertex order. There are exactly $24 \times 2^{14} = 393{,}216$ data rows. This supplements the complete parametrization and the printed 48-state family; it does not assert that 48 is minimal. The 153 Hesse states, their event-order witnesses, and all original printed profiles remain in the accompanying Hesse data.

S2.30. States of the Cabello restrictions and completions

S2.30.1. States of the vertex-deleted 45-ray completion

The nine-context $C_{18,9}$, its eighteen-context faithful completion, and both 68-ray parents have no context states by their retained parity proofs. For the 45-ray construction, use exactly the local labels in Tables S31 and S32. Table S84 lists all twelve admissible core supports, and Table S85 lists the fourteen bridge pairs with their fresh labels and zero-profiles. For a support S_α, let F_α be the bridge indices missed by S_α. On every hit bridge both fresh values are zero; on each missed bridge independently choose exactly one of its two fresh vertices. This enumerates every context state once.

TABLE S84: All twelve core supports; $m = |F_\alpha|$. The final column specifies extendibility under all geometric pair exclusions.

α	S_α	m	Geometric extension
1	$\{1, 6, 11, 12\}$	8	yes
2	$\{1, 10, 14, 18\}$	8	yes
3	$\{3, 7, 10, 14\}$	8	yes
4	$\{3, 8, 11, 13\}$	8	yes
5	$\{3, 13, 14, 17\}$	8	no
6	$\{2, 6, 10, 15\}$	8	yes
7	$\{6, 12, 15, 16\}$	8	no
8	$\{8, 10, 15, 18\}$	8	yes
9	$\{9, 16, 18, 19\}$	9	no
10	$\{9, 17, 18, 19\}$	9	no
11	$\{2, 11, 13, 19\}$	8	yes
12	$\{7, 11, 12, 19\}$	8	yes

TABLE S85: The fourteen bridge pairs, fresh labels, and zero-profiles; each Z_m is the set of missed-bridge branch indices.

m	Bridge	Fresh pair	Z_m
1	$\{1, 2\}$	$\{21, 22\}$	$\{3, 4, 5, 7, 8, 9, 10, 12\}$
2	$\{1, 13\}$	$\{23, 24\}$	$\{3, 6, 7, 8, 9, 10, 12\}$
3	$\{1, 17\}$	$\{25, 26\}$	$\{3, 4, 6, 7, 8, 9, 11, 12\}$
4	$\{2, 14\}$	$\{27, 28\}$	$\{1, 4, 7, 8, 9, 10, 12\}$
5	$\{2, 17\}$	$\{29, 30\}$	$\{1, 2, 3, 4, 7, 8, 9, 12\}$
6	$\{3, 16\}$	$\{31, 32\}$	$\{1, 2, 6, 8, 10, 11, 12\}$
7	$\{6, 17\}$	$\{33, 34\}$	$\{2, 3, 4, 8, 9, 11, 12\}$
8	$\{7, 8\}$	$\{35, 36\}$	$\{1, 2, 5, 6, 7, 9, 10, 11\}$
9	$\{7, 15\}$	$\{37, 38\}$	$\{1, 2, 4, 5, 9, 10, 11\}$
10	$\{7, 16\}$	$\{39, 40\}$	$\{1, 2, 4, 5, 6, 8, 10, 11\}$
11	$\{8, 12\}$	$\{41, 42\}$	$\{2, 3, 5, 6, 9, 10, 11\}$
12	$\{8, 16\}$	$\{43, 44\}$	$\{1, 2, 3, 5, 6, 10, 11, 12\}$
13	$\{10, 19\}$	$\{45, 46\}$	$\{1, 4, 5, 7\}$
14	$\{11, 18\}$	$\{47, 48\}$	$\{3, 5, 6, 7\}$

Taking the first fresh vertex on every missed bridge gives $s_{\alpha,0}$, and taking the second gives $s_{\alpha,1}$. Table S86 explicitly enumerates all 24 canonical context states. Their profiles are nonzero and pairwise distinct on all 45 vertices. The twelve branch supports and fourteen distinct nonempty zero-profiles verify this by Lemma 21(3).

TABLE S86: All 24 canonical context-state supports of the 45-ray example. Some fail geometric pair exclusions.

α	ε	Support
1	0	$\{1,6,11,12,27,29,31,35,37,39,43,45\}$
1	1	$\{1,6,11,12,28,30,32,36,38,40,44,46\}$
2	0	$\{1,10,14,18,29,31,33,35,37,39,41,43\}$
2	1	$\{1,10,14,18,30,32,34,36,38,40,42,44\}$
3	0	$\{3,7,10,14,21,23,25,29,33,41,43,47\}$
3	1	$\{3,7,10,14,22,24,26,30,34,42,44,48\}$
4	0	$\{3,8,11,13,21,25,27,29,33,37,39,45\}$
4	1	$\{3,8,11,13,22,26,28,30,34,38,40,46\}$
5	0	$\{3,13,14,17,21,35,37,39,41,43,45,47\}$
5	1	$\{3,13,14,17,22,36,38,40,42,44,46,48\}$
6	0	$\{2,6,10,15,23,25,31,35,39,41,43,47\}$
6	1	$\{2,6,10,15,24,26,32,36,40,42,44,48\}$
7	0	$\{6,12,15,16,21,23,25,27,29,35,45,47\}$
7	1	$\{6,12,15,16,22,24,26,28,30,36,46,48\}$
8	0	$\{8,10,15,18,21,23,25,27,29,31,33,39\}$
8	1	$\{8,10,15,18,22,24,26,28,30,32,34,40\}$
9	0	$\{9,16,18,19,21,23,25,27,29,33,35,37,41\}$
9	1	$\{9,16,18,19,22,24,26,28,30,34,36,38,42\}$
10	0	$\{9,17,18,19,21,23,27,31,35,37,39,41,43\}$
10	1	$\{9,17,18,19,22,24,28,32,36,38,40,42,44\}$
11	0	$\{2,11,13,19,25,31,33,35,37,39,41,43\}$
11	1	$\{2,11,13,19,26,32,34,36,38,40,42,44\}$
12	0	$\{7,11,12,19,21,23,25,27,29,31,33,43\}$
12	1	$\{7,11,12,19,22,24,26,28,30,32,34,44\}$

Ten branches miss eight pairs and two miss nine. Therefore

$$|\mathcal{S}_{\mathrm{ctx}}| = 10\,2^8 + 2\,2^9 = 3{,}584, \qquad |\mathcal{S}_{\mathrm{geom}}| = 8\,2^8 = 2048. \tag{S22}$$

The geometric branches are $1,2,3,4,6,8,11,12$; arbitrary fresh choices within each of these branches remain geometrically compatible. The other branches are excluded by the four forced pairs. Both full enumerations are supplied as compressed CSV data, in addition to this complete finite parametrization. Every geometric state vanishes on 9, 16, and 17, so this family is neither unital nor separating.

For an explicit counterexample, the context state supported at

$$\{6,12,15,16,21,23,25,27,29,35,45,47\}$$

selects both 16 and 21, although $v_{16} \cdot v_{21} = 0$.

S2.30.2. Deleting one of the nine Cabello contexts

Number the nine contexts as the first nine rows of Table S29. For every deletion there are 26 context states, six geometric states, and a proper four-coloring of the remaining context graph. The color strings in Table S87 use the ordered ray list $(1,2,3,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20)$. The geometric states vanish precisely on the omitted context.

TABLE S87: All single-context deletion counts and four-coloring witnesses for $C_{18,9}$.

e	Omitted context	$\lvert\mathcal{S}_{\mathrm{ctx}}\rvert$	$\lvert\mathcal{S}_{\mathrm{geom}}\rvert$	Four-color string
1	$\{9,10,12,13\}$	26	6	012102203020112333
2	$\{9,11,14,15\}$	26	6	012021231012102333
3	$\{10,11,16,17\}$	26	6	012021320012102333
4	$\{2,3,12,18\}$	26	6	013103302120113322
5	$\{1,3,15,19\}$	26	6	103130320102113232
6	$\{1,2,17,20\}$	26	6	330130230103112222
7	$\{6,7,13,18\}$	26	6	012221213030102333
8	$\{6,8,14,19\}$	26	6	012202230013112333
9	$\{7,8,16,20\}$	26	6	012021320012113332

For deletion of context 9, $\{7,8,16,20\}$, Table S88 explicitly enumerates all 26 supports; exactly the six marked rows satisfy every geometric pair exclusion. All nine full deletion-state lists accompany the data.

TABLE S88: All 26 states after deleting context 9; geometric compatibility is marked explicitly.

j	Support	Geometric
1	$\{1,6,11,12\}$	yes
2	$\{1,7,8,11,12\}$	no
3	$\{1,7,12,14,16\}$	no
4	$\{1,8,9,16,18\}$	no
5	$\{1,10,14,18\}$	yes
6	$\{2,6,10,15\}$	yes
7	$\{2,7,8,10,15\}$	no
8	$\{2,8,13,15,16\}$	no
9	$\{2,7,9,16,19\}$	no
10	$\{2,11,13,19\}$	yes
11	$\{3,6,9,17\}$	no
12	$\{3,7,8,9,17\}$	no
13	$\{3,13,14,17\}$	yes
14	$\{6,12,15,17\}$	no
15	$\{7,8,12,15,17\}$	no
16	$\{9,17,18,19\}$	yes
17	$\{3,6,9,16,20\}$	no
18	$\{3,7,8,9,16,20\}$	no
19	$\{3,7,10,14,20\}$	no
20	$\{3,8,11,13,20\}$	no
21	$\{3,13,14,16,20\}$	no
22	$\{6,12,15,16,20\}$	no
23	$\{7,11,12,19,20\}$	no
24	$\{7,8,12,15,16,20\}$	no
25	$\{9,16,18,19,20\}$	no
26	$\{8,10,15,18,20\}$	no

S2.30.3. All single-ray deletions and the alternative parity subsystem

The next table records the two span failures and all forced-zero core rays after geometric pair exclusion for every possible single-ray deletion. An entry $\{u,v\} : z$ means $v_z \in \mathrm{span}(v_u, v_v)$. Every row has twelve compatible core states, 3,584 abstract completion states, and 2,048 geometric completion states, with exactly four forced extra pairs and no other extra orthogonality in the supplied realization. The complete core supports and 24 canonical states for every row are also supplied in the data.

TABLE S89: All eighteen vertex-deletion checks; the completed geometric families fail unitality and separation in every case.

Deleted ray	First span failure	Second span failure	Forced-zero core rays
1	$\{15,19\}:2$	$\{17,20\}:3$	$\{2,3,13\}$
2	$\{12,18\}:1$	$\{17,20\}:3$	$\{1,3,14\}$
3	$\{12,18\}:1$	$\{15,19\}:2$	$\{1,2,16\}$
6	$\{13,18\}:8$	$\{14,19\}:7$	$\{7,8,17\}$
7	$\{13,18\}:8$	$\{16,20\}:6$	$\{6,8,15\}$
8	$\{14,19\}:7$	$\{16,20\}:6$	$\{6,7,12\}$
9	$\{12,13\}:11$	$\{14,15\}:10$	$\{10,11,20\}$
10	$\{12,13\}:11$	$\{16,17\}:9$	$\{9,11,19\}$
11	$\{14,15\}:10$	$\{16,17\}:9$	$\{9,10,18\}$
12	$\{2,3\}:13$	$\{9,10\}:18$	$\{8,13,18\}$
13	$\{6,7\}:12$	$\{9,10\}:18$	$\{1,12,18\}$
14	$\{6,8\}:15$	$\{9,11\}:19$	$\{2,15,19\}$
15	$\{1,3\}:14$	$\{9,11\}:19$	$\{7,14,19\}$
16	$\{7,8\}:17$	$\{10,11\}:20$	$\{3,17,20\}$
17	$\{1,2\}:16$	$\{10,11\}:20$	$\{6,16,20\}$
18	$\{2,3\}:13$	$\{6,7\}:12$	$\{11,12,13\}$
19	$\{1,3\}:14$	$\{6,8\}:15$	$\{10,14,15\}$
20	$\{1,2\}:16$	$\{7,8\}:17$	$\{9,16,17\}$

For the alternative subsystem, take parent variant 2 of Table S35 and retain vertices 1–17 and 66–68. It is stateless. Each ray deletion instead leaves the positive geometric state count below; restriction of any such state shows that no eighteen-ray subset can be KS.

TABLE S90: Geometric state counts after every single-ray deletion of the twenty-ray parity subsystem.

Deleted ray	States	Deleted ray	States
1	8	2	8
3	8	4	28
5	28	6	8
7	8	8	8
9	12	10	12
11	12	12	10
13	10	14	10
15	10	16	10
17	10	66	8
67	8	68	8

S3. SUPPLEMENTARY PROOFS, REDUCTIONS, AND EXACT VERIFICATION CERTIFICATES

S3.1. Formulas used by the finite certificates

The following formulas restore the explicit conventions used by the later certificates. State indices are local to the corresponding complete table. For Hesse, the six-state vertex-separating family, its seven-state event-separating extension, and the eighteen-state order-determining family are respectively

$$\Omega_6 = \{s_3, s_{31}, s_{75}, s_{89}, s_{113}, s_{151}\}. \tag{S23}$$

$$\Omega_7 = \Omega_6 \cup \{s_{84}\} \tag{S24}$$

$$\begin{aligned}\Omega_{18} = \{&s_2, s_8, s_{20}, s_{30}, s_{40}, s_{45},\\ &s_{54}, s_{64}, s_{73}, s_{79}, s_{83}, s_{93},\\ &s_{105}, s_{109}, s_{119}, s_{123}, s_{127}, s_{150}\}.\end{aligned} \tag{S25}$$

The members of Ω_6 are denoted $\sigma_1, \dots, \sigma_6$ in the displayed order. Their atom profiles and the nontrivial atom–coatom order are

$$\kappa(i) = (\sigma_1(i), \dots, \sigma_6(i)). \tag{S26}$$

$$a_i \le a_j^{\perp} \quad \Longleftrightarrow \quad \{i, j\} \in E(\Gamma(H_{37})). \tag{S27}$$

For H_{65}, the following core coloring extends to the completed hypergraph. A base-compatible independent support S_α is extended canonically using $\varepsilon \in \{0, 1\}$. In the plane formula, η_A, η_B are an integer orthogonal basis and $n_A = \|\eta_A\|^2$, $n_B = \|\eta_B\|^2$.

$$\begin{aligned}(c(1), \dots, c(17)) = (&0, 1, 2, 3, 0, 1, 2, 4, 0, 1, 3,\\ &3, 4, 2, 1, 0, 2).\end{aligned} \tag{S28}$$

$$\begin{aligned}&\big(s_{\alpha,\varepsilon}(w_{m,1}),\ s_{\alpha,\varepsilon}(w_{m,2})\big)\\ &\quad = \begin{cases}(0, 0), & |S_\alpha \cap \{u_m, v_m\}| = 1,\\ (1-\varepsilon,\ \varepsilon), & S_\alpha \cap \{u_m, v_m\} = \emptyset.\end{cases}\end{aligned} \tag{S29}$$

$$\begin{aligned}w_{m,1}(a, b) &= a\,\eta_A + b\,\eta_B,\\ w_{m,2}(a, b) &= -n_B b\,\eta_A + n_A a\,\eta_B\end{aligned} \tag{S30}$$

The eight rays adjoined to completed Yu–Oh and the nine uncovered pairs of the Cabello subsystem are fixed by the following local labels. The last display records the completed historical Γ_1 TITS incidence.

$$\begin{aligned}&(0, 1, 2), (0, 2, -1), (0, 1, -2), (0, 2, 1),\\ &(1, 2, 0), (2, -1, 0), (1, -2, 0), (2, 1, 0).\end{aligned} \tag{S31}$$

$$\begin{aligned}&\{1, 13\}, \{2, 14\}, \{3, 16\}, \{6, 17\}, \{7, 15\},\\ &\{8, 12\}, \{9, c\}, \{10, b\}, \{11, a\}.\end{aligned} \tag{S32}$$

$$E(I_{16,9}) = E(B_{13,7}) \cup \{\{4, 11, 24\}, \{4, 10, 18\}\}. \tag{S33}$$

a. Fractional value of the faithful Cabello completion. The geometric orthogonality graph of $C_{18,9}$ has independence number four and exactly 54 independent four-sets, with each of its 18 rays belonging to twelve of them. Assigning weight $1/12$ to each set gives a fractional cover of total weight $9/2$, matching $18/4$. Fresh pair completion preserves this value by the main completion proposition, proving $\chi_f(C_{36,18}) = 9/2$ as recorded in the main comparison. The complete 54-set certificate is in the electronic audit data.

S3.2. Snark obstructions and quantitative robustness bounds

S3.2.1. Cubic graphs, Blanuša snarks, and Isaacs dot products

For a finite simple cubic graph X, let $H(X)$ be the context hypergraph whose vertices are the edges of X and whose contexts are the three edges incident with each vertex of X. Then $\Gamma(H(X)) = L(X)$ and $\chi(H(X)) = \chi'(X)$. An orthogonal ray assignment for $H(X)$ in $\mathbb{F}^3$ is an assignment of nonzero vectors to the edges of X such that incident edges are orthogonal.

Silva and Ibiapina[5] (Question 6) asked whether an orthogonal vector edge coloring in $\mathbb{R}^3$ can exist when $\chi'(X) = 4$. Theorem 11 excludes the Petersen graph P even over $\mathbb{C}$. Using the multipole forcing law of Theorem 11(1), we can extend this exclusion to the two 18-vertex Blanuša snarks[6] and the infinite family of Isaacs dot-product snarks[7].

Recall that for a cubic graph G with two nonadjacent edges $e_1 = \{x_1, y_1\}$ and $e_2 = \{x_2, y_2\}$, the Isaacs dot product $G \cdot P$ is formed by removing e_1, e_2 from G, removing two adjacent vertices u, v from the Petersen graph P (leaving the 4-pole $P \setminus \{u, v\}$ whose dual hypergraph is precisely H_8), and attaching the two dangling edges $\{2, 8\}$ of u to $\{x_1, y_1\}$ and the two dangling edges $\{13, 14\}$ of v to $\{x_2, y_2\}$; the two distinct choices of nonadjacent edges in $G = P$ produce the two Blanuša snarks B_1 and B_2[6–8].

Corollary 1 (Obstruction for Blanuša and dot-product snarks)**.** *If a cubic graph G admits no orthogonal vector edge coloring in $\mathbb{C}^3$, then neither does any Isaacs dot product $G \cdot P$. In particular, both 18-vertex Blanuša snarks $B_1, B_2 = P \cdot P$ and all iterated dot products $X = P \cdot P \cdots P$ satisfy $\chi(H(X)) = \chi'(X) = 4 > 3 = \omega(L(X))$ and admit no orthogonal vector edge coloring in $\mathbb{C}^3$.*

Proof. In any orthogonal vector edge coloring of $G \cdot P$ in $\mathbb{C}^3$, the restriction to the embedded copy of $H_8 = H(P \setminus \{u, v\})$ satisfies $[v_2] = [v_8]$ and $[v_{13}] = [v_{14}]$ by Theorem 11(1). Assigning the common ray $[v_2] = [v_8]$ to $e_1 = \{x_1, y_1\}$ and the common ray $[v_{13}] = [v_{14}]$ to $e_2 = \{x_2, y_2\}$ preserves orthogonality at x_1, y_1, x_2, y_2 and therefore yields a valid orthogonal vector edge coloring of G in $\mathbb{C}^3$. Since P admits no such coloring by Theorem 11, induction excludes B_1, B_2 and all iterated dot products with P. □

Generalized Petersen graphs[9] and flower snarks provide further families for comparing real, complex, injective, and edge-faithful realizability.

S3.2.2. Explicit analytic lower bound and tetrahedral tight-frame upper bounds for Δ_3

Let A be the 30 edges of $\Gamma(G_{32})$ and define the minimum total squared orthogonality error in $\mathbb{C}^3$:

$$\Delta_3 = \min_{\substack{v_i \in \mathbb{C}^3 \\ \|v_i\|=1 \ (1 \le i \le 15)}} \sum_{\{i,j\} \in A} |\langle v_i, v_j \rangle|^2. \tag{S34}$$

By compactness of $(S^5)^{15}$ and Theorem 11, $\Delta_3 > 0$. We now establish both a certified rational analytic lower bound and explicit tetrahedral tight-frame upper bounds.

Proposition 2 (Quantitative bounds on Δ_3). *Every assignment of unit vectors $v_1, \dots, v_{15} \in \mathbb{C}^3$ satisfies*

$$\varepsilon \coloneqq \max_{\{i,j\}\in A} |\langle v_i, v_j\rangle| > \frac{1}{120}, \tag{S35}$$

and the minimum total squared error in Eq. (S34) satisfies

$$\begin{aligned}\frac{1}{14400} < \Delta_3 &\le \min_{a^2+b^2+c^2=1} F(a,b,c) \\ &\le \frac{9275914990340646}{10001032226635921} < 0.927496 < \frac{18}{19},\end{aligned} \tag{S36}$$

where

$$F(a,b,c) = 12a^2 + 12(ab+bc-ca)^2 + 6(b^2-a^2-c^2)^2. \tag{S37}$$

Proof. To prove the lower bound (S35), let $\varepsilon = \max_{\{i,j\}\in A} |\langle v_i, v_j\rangle|$. For any context $e = \{x, y, z\}$, the Gram matrix of $\{v_x, v_y, v_z\}$ has diagonal entries 1 and off-diagonal entries of modulus at most ε, so Gershgorin's circle theorem implies $\|I_3 - (v_x v_x^\dagger + v_y v_y^\dagger + v_z v_z^\dagger)\|_{\mathrm{op}} \le 2\varepsilon$. Hence for any unit vectors $u, w \in \mathbb{C}^3$,

$$\Big|1 - \big(|\langle u, x\rangle|^2 + |\langle u, y\rangle|^2 + |\langle u, z\rangle|^2\big)\Big| \le 2\varepsilon, \tag{S38}$$

and if $\{u, x\}, \{w, z\} \in A$, then

$$\big|\langle u, w\rangle - \langle u, y\rangle\langle y, w\rangle\big| \le 4\varepsilon. \tag{S39}$$

In $\{5, 6, 7\}$ and $\{9, 10, 11\}$, the vectors $w_7 = v_{12} - v_5\langle v_5, v_{12}\rangle$ and $w_9 = v_4 - v_{11}\langle v_{11}, v_4\rangle$ satisfy $\|w_7 - v_7\langle v_7, v_{12}\rangle\| \le 3\varepsilon$ and $\|w_9 - v_9\langle v_9, v_4\rangle\| \le 3\varepsilon$, so $|\langle w_7, w_9\rangle| \le 7\varepsilon$. Expanding,

$$\begin{aligned}\langle w_7, w_9\rangle = {} & \langle v_{12}, v_4\rangle - \langle v_{12}, v_{11}\rangle\langle v_{11}, v_4\rangle - \langle v_{12}, v_5\rangle\langle v_5, v_4\rangle \\ & + \langle v_{12}, v_5\rangle\langle v_5, v_{11}\rangle\langle v_{11}, v_4\rangle.\end{aligned}$$

The two middle terms are at most ε in modulus, since $|\langle v_{12}, v_{11}\rangle| \le \varepsilon$ and $|\langle v_5, v_4\rangle| \le \varepsilon$. Eq. (S39) on $\{1, 2, 3\}$ replaces each of $\langle v_{12}, v_4\rangle$, $\langle v_{12}, v_5\rangle$, $\langle v_5, v_{11}\rangle$, and $\langle v_{11}, v_4\rangle$ by its product through v_2 with error at most 4ε, and hence the triple product with error at most 12ε, since all factors have modulus at most one. The main terms combine to

$$\big(1 + |\langle v_2, v_5\rangle|^2 |\langle v_2, v_{11}\rangle|^2\big)\langle v_{12}, v_2\rangle\langle v_2, v_4\rangle,$$

whose real prefactor is at least one, so

$$\big|\langle v_{12}, v_2\rangle\langle v_2, v_4\rangle\big| \le (7 + 2 + 4 + 12)\varepsilon = 25\varepsilon.$$

By reflection symmetry, we may assume $|\langle v_2, v_4\rangle| \le 5\sqrt{\varepsilon}$. Applying Eq. (S38) to v_4 in $\{1, 2, 3\}$ and to v_2 in $\{3, 4, 5\}$ gives $|\langle v_1, v_4\rangle|^2 \ge 1 - 27\varepsilon - \varepsilon^2$ and $|\langle v_2, v_5\rangle|^2 \ge 1 - 27\varepsilon - \varepsilon^2$, so $\min_\theta \|v_1 - e^{i\theta} v_4\| \le \sqrt{54\varepsilon + 2\varepsilon^2}$ and $\min_\theta \|v_2 - e^{i\theta} v_5\| \le \sqrt{54\varepsilon + 2\varepsilon^2}$. Since $|\langle v_4, v_{10}\rangle| \le \varepsilon$ and $|\langle v_5, v_7\rangle| \le \varepsilon$, Eq. (S38) applied to v_1 in $\{9, 10, 11\}$ and to v_2 in $\{7, 8, 9\}$ gives

$$|\langle v_i, v_9\rangle|^2 \ge 1 - 2\varepsilon - \varepsilon^2 - \big(\varepsilon + \sqrt{54\varepsilon + 2\varepsilon^2}\big)^2 \qquad (i = 1, 2).$$

Comparing the sum of these two inequalities with $|\langle v_1, v_9\rangle|^2 + |\langle v_2, v_9\rangle|^2 \le 1 + 2\varepsilon$ from $\{1, 2, 3\}$ gives

$$1 \le 6\varepsilon + 2\varepsilon^2 + 2\big(\varepsilon + \sqrt{54\varepsilon + 2\varepsilon^2}\big)^2.$$

The right-hand side is strictly increasing for $\varepsilon > 0$, and at $\varepsilon = 1/120$ we have $54\varepsilon + 2\varepsilon^2 = 3241/7200 < (27/40)^2$, so the right-hand side is strictly bounded above by

$$\frac{6}{120} + \frac{2}{120^2} + 2\left(\frac{1}{120} + \frac{27}{40}\right)^2 = \frac{1417}{1440} < 1,$$

certifying $\varepsilon > 1/120$ and $\Delta_3 \geq \varepsilon^2 > 1/14400$.

For the upper bounds, set $v_4 = e_1$, $v_8 = e_2$, $v_{12} = e_3$, and for any unit vector $(a, b, c) \in S^2$, define the remaining 12 unit vectors as representatives of the tetrahedral A_4-orbit of rays:

$$\begin{aligned} v_1 &= (-b, -c, -a)^T, & v_2 &= (c, -a, b)^T, \\ v_3 &= (-a, -b, c)^T, & v_5 &= (-a, b, c)^T, \\ v_6 &= (-b, c, -a)^T, & v_7 &= (c, a, b)^T, \\ v_9 &= (-c, -a, b)^T, & v_{10} &= (-a, b, -c)^T, \\ v_{11} &= (b, -c, -a)^T, & v_{13} &= (a, b, c)^T, \\ v_{14} &= (b, c, -a)^T, & v_{15} &= (-c, a, b)^T. \end{aligned} \tag{S40}$$

These 15 vectors form a tight frame $\sum_{i=1}^{15} v_i v_i^T = 5I_3$, and the sum of squared overlaps on the 30 edges of $\Gamma(G_{32})$ equals $F(a, b, c)$ in Eq. (S37). Evaluating Eq. (S40) at the normalized integer direction $(a, b, c) = (1, 3, -3)/\sqrt{19}$ yields rational squared overlaps with $\Delta_3 \leq 342/361 = 18/19 \approx 0.947368$ and $\max_{\{i,j\} \in A} |\langle v_i, v_j \rangle|^2 = 1/19$; evaluating at the refined normalized integer direction $(a, b, c) = (2332, 7101, -6644)/\sqrt{100005161}$ yields the certified rational bound in Eq. (S36). □

Numerical minimization of F on S^2 (as well as unrestricted local searches over $(S^5)^{15}$) finds an attained candidate minimum near 0.927495724691 at $(a, b, c) \approx (0.23319713, 0.71005263, -0.66441280)$; certifying whether the unrestricted global minimum Δ_3 coincides with $\min_{S^2} F$ remains an open optimization problem. Choosing instead the rational unit vector $(a, b, c) = \frac{1}{3}(1, 2, -2)$ in Eq. (S40) makes $ab + bc - ca = 0$, so four of the ten contexts of G_{32} ($\{1, 2, 3\}, \{5, 6, 7\}, \{9, 10, 11\}, \{13, 14, 15\}$) become exact orthonormal bases of $\mathbb{R}^3$, all 15 rays are distinct, and $\Delta_3 \leq 38/27$ with $\max_{\{i,j\} \in A} |\langle v_i, v_j \rangle| = 1/3$.

S3.3. Hesse H_{37}: minimum state families

All state indices in this appendix use the canonical enumeration of Appendix S2.27. The minima for unital vertex separation and event separation have the counting lower bounds $2^5 - 1 < 37$ and $2^6 < 76$. The families in Eqs. (S23) and (S24) attain those bounds: their profiles give respectively 37 distinct nonzero atom profiles and 76 distinct event profiles. The seven-state family still has 462 ordered event pairs for which inclusion of the profiles does not imply the lattice order. The family in Eq. (S25) has no such pair and hence gives the upper bound eighteen for order determination. This appendix supplies a computer-assisted proof of the matching lower bound.

S3.3.1. Reduction to families of seventeen states

Let $D = \{P_0, P_1, P_2, Q_0, Q_1, Q_2\}$. For each unordered pair $\{u, v\} \subset D$, order determination requires a state with $s(u) = s(v) = 1$: the pair is nonorthogonal, so $u \not\leq v^\perp$. Every state takes value one on at most two members of D. Consequently the fifteen pairs require fifteen different states. Table S91 gives their complete witness classes. These classes are disjoint, their union T has 27 states, and a choice of one witness from each class has $1^3 2^{12} = 4{,}096$ possibilities.

TABLE S91: Complete joint-one witness classes for the fifteen pairs in D. Each entry lists all canonical indices r for which $s_r(u) = s_r(v) = 1$.

Pair $\{u, v\}$	Witness indices r
$\{P_0, P_1\}$	150
$\{P_0, P_2\}$	73
$\{P_0, Q_0\}$	93, 145
$\{P_0, Q_1\}$	127, 133
$\{P_0, Q_2\}$	79, 99
$\{P_1, P_2\}$	2
$\{P_1, Q_0\}$	105, 139
$\{P_1, Q_1\}$	14, 119
$\{P_1, Q_2\}$	40, 48
$\{P_2, Q_0\}$	54, 60
$\{P_2, Q_1\}$	8, 26
$\{P_2, Q_2\}$	20, 68
$\{Q_0, Q_1\}$	109, 113
$\{Q_0, Q_2\}$	83, 87
$\{Q_1, Q_2\}$	30, 34

The nine nonorthogonal pairs $\{z, C_{ak}\}$ require further joint-one witnesses. No state in T witnesses any of them. Indeed, if a state activates a P vertex then $s(z) = 0$. If it instead activates two Q vertices, the state equations force the selected M vertices into the remaining Q column, and all nine C values are zero. Any state outside T witnesses at most six of the nine pairs: for $s(z) = 1$, the three selected M vertices force at least three C values to vanish. Thus at least two states outside T are necessary.

It follows that an order-determining family of at most seventeen states would have exactly seventeen members: one from each of the fifteen classes in Table S91, and two outside T that together witness all nine $\{z, C_{ak}\}$ pairs. Enumerating the $\binom{126}{2}$ unordered pairs outside T leaves exactly 132 with this property. The reduction therefore leaves precisely

$$4{,}096 \cdot 132 = 540{,}672 \tag{S41}$$

candidate families. The nine pairs alone do not require three extra states; for example, s_1 and s_{72} jointly witness all nine. The remaining order requirements must also be checked.

S3.3.2. Exhaustive order certificate

For each of the 5,407 ordered event pairs $x \not\leq y$, define its nonempty witness set

$$W_{xy} = \{r \in \{1, \ldots, 153\} : s_r(x) = 1,\ s_r(y) = 0\}. \tag{S42}$$

A family is order determining exactly when its index set meets every W_{xy}. Repeated sets and proper supersets can be discarded, because meeting an inclusion-minimal set also meets every set containing it. There are exactly 81 inclusion-minimal witness sets. The exhaustive check forms each of the 540,672 families in Eq. (S41) and counts the minimal witness sets that it meets. The largest count is 75, strictly below 81. Hence no family of seventeen states is order determining. Together with Ω_{18}, this proves the minimum eighteen.

The accompanying Hesse extension checker, invoked by the joint entry point, reconstructs the contexts and all 153 states from their symbolic parametrization, checks their agreement with `hesse37-data.json`, and performs this finite search using integer bit sets and the Python standard library. No numerical tolerance, optimization solver, or unreported search cutoff is used. It also verifies the printed seven- and eighteen-state families and every entry of Table S91. From the extracted joint project directory, run the joint entry point, without Python's optimization option:

```
python verification/verify-integrated.py
```

The generated file `verification/_work/hesse/hesse37-extensions-certificate.json` records all 81 minimal witness sets, a representative event pair for each, all 132 possible extra pairs, the coverage-count distribution over the 540,672 candidates, and all

event profiles for Ω_6, Ω_7, and Ω_{18}. The generated file `verification/_work/hesse/hesse37-order18-witnesses.csv` records a witness in Ω_{18} for every $x \not\leq y$. These two files use the same numbered event labels as the full-family certificate in Appendix S2.27.6.

The program additionally checks Berge girth five from the incidence graph, computes every meet and join in the 76-event order, verifies the orthomodular law, and verifies the projector sum in Eq. (30) exactly in $\mathbb{Q}(\zeta)$. The nonexistence of an injective real representation is proved analytically in Sec. V A.

S3.4. Four-dimensional $H_{\mathrm{YO}}^{(4)}$ and H_{65}: completion and verification

The coordinates in Table S17 can be reproduced by the following finite selection rule. Normalize an integer vector by dividing by the greatest common divisor of its entries and choosing the sign so that its first nonzero entry is positive. Form the set of normalized nonzero vectors obtained from $\{-3,-2,\dots,3\}^4$, ordered by squared norm, then maximum absolute entry, then lexicographically. Process bridge edges in the order $e_{13},\dots,e_{24},e_1,\dots,e_{12}$. For each pair, choose η_A to be the first vector in this ordered pool orthogonal to its endpoints, and η_B the first such vector also orthogonal to η_A.

For all $(a,b) \in \{-7,-6,\dots,7\}^2 \setminus \{(0,0)\}$, form the pair in Eq. (S30) and normalize both vectors. Discard pairs that repeat an already placed ray or have an orthogonality to any already placed vertex other than the two endpoints. Among the remaining ordered pairs choose the lexicographically smallest score

$$\left(\max_{j,k} |(w_{m,j})_k|,\ \|w_{m,1}\|^2 + \|w_{m,2}\|^2,\ w_{m,1}, w_{m,2} \right). \tag{S43}$$

In Eq. (S43), the order on the last two entries is the usual lexicographic order on their integer coordinates. This rule yields exactly the 48 completion vectors in Table S17. It is a recipe for this specific example; the general existence assertion is established independently by Lemma 21.

The companion verifier parses the integrated coordinate and state tables, reproduces this completion rule and all 393,216 state extensions, and checks the maximal cliques, fractional-coloring certificate, atom-order failure, forced hub-completion zero, and warm-up data. It is included with the complete state file and invoked by the joint entry point in Sec. S3.7.

S3.5. Exact Cabello completion and forced-zero certificates

For each of the nine uncovered pairs in Eq. (S32), rational row reduction verifies that its two endpoint vectors span no third core ray. If two endpoint planes were orthogonal, their four endpoints would form another full basis. Exact four-clique enumeration gives only the original nine bases, all of whose pairs are already covered. This excludes such an orthogonal-plane pair. The integer vectors in Table S28 directly verify the resulting faithful completion: 36 distinct projective rays, eighteen full bases, and 108 orthogonal pairs, with every pair covered by exactly one listed basis. The nine-context graph and the full eighteen-ray orthogonality graph both fail four-colorability and admit five-colorings, as checked by exact finite search in the companion certificate. Lemma 21(2) transfers the chromatic value to the completion.

For the vertex-deleted core with c removed, rational row reduction finds exactly the span failures

$$v_{16} \in \mathrm{span}(v_1, v_2), \qquad v_{17} \in \mathrm{span}(v_7, v_8).$$

There is no orthogonal-plane-pair obstruction. The finite-avoidance choice in Lemma 21 can therefore avoid all extra incidences except those forced by these two spans; the explicit 45-ray inventory has exactly the four pairs printed in Sec. S1.17.2. The two fresh vectors

completing $\{1,2\}$ must both be orthogonal to 16, and those completing $\{7,8\}$ must both be orthogonal to 17.

The twelve rows of Table S84 give a short complete proof of the resulting obstruction. Whenever 16 is selected, neither 1 nor 2 is selected. The context completing $\{1,2\}$ then needs one fresh value equal to one, while geometric exclusion with 16 requires both to be zero. Hence every geometric extension has $s(16) = 0$. The same argument with $\{7,8\}$ gives $s(17) = 0$. These implications use the displayed core states in addition to the span incidences; the span incidences alone would not imply them. The retained basis $\{10, 11, 16, 17\}$ gives $s(10)+s(11) = 1$. If $s(9) = 1$, the other two hub bases give $s(10) = s(11) = 0$, a contradiction. Thus 9, 16, and 17 have identical zero profiles.

Conversely, each of the eight indicated core rows avoids 16 and 17, so all its fresh-pair choices satisfy the four extra exclusions. Each misses eight bridges. This proves the exact count $8\,2^8 = 2{,}048$, as well as failure of geometric unitality and separation. The two excluded core rays remain impossible for every fresh-pair completion of these fixed planes; additional accidental orthogonalities cannot restore them. Table S89 supplies the individual span checks for all eighteen deletions. Their full core tables, exact integer completions, and extension counts are included in `data/cabello18-review-checks.json`.

To audit the ancestry result, enumerate every four-clique after adjoining either three-ray choice to all 65 original rays. In both cases there are exactly 35 bases, 195 geometric pairs, and 192 context-covered pairs; the maximum basis intersection is two. The first union contains the displayed eighteen-ray parity proof. The second union contains the twenty-ray proof with multiplicities $4, 4, 2, \ldots, 2$ and eleven contexts. Each of its twenty ray deletions has a compatible assignment by Table S90. All twenty rays lie in Peres's 24-ray set. Enumerating the $\binom{24}{4} = 10{,}626$ twenty-ray subsets of that set, with all bases they contain, gives exactly 96 ray-critical KS subsets, each with eleven bases and all mutually isomorphic; the twenty-ray proof is one of them. Since Peres's set contains Kernaghan's critical twenty-ray set and 95 others[3], the twenty-ray proof is isomorphic to Kernaghan's[10]. All coordinates, full basis lists, and the preserved H_{65} restriction are checked independently of the unrestricted ancestry existence proposition.

S3.6. Transverse extension and criticality certificates

For $d = 3$, number the two copies of an n-vertex input as i and $n + i$, and the fresh transverse vertex as $2n + i$. The transverse contexts are $\{i, n + i, 2n + i\}$ and the horizontal contexts are the two numbered copies of the original list. This explicitly specifies all 45 vertices and 33 contexts for $T(H_9)$, and all 75 vertices and 57 contexts for $T(H^+_{\mathrm{YO}})$.

For each deleted original context e, a strong three-coloring of $H \setminus \{e\}$ having two equal colors and a singleton color on e gives a state after deletion of one horizontal copy of e. Use the repeated-color indicator on the deleted copy and the singleton-color indicator on the intact copy. Both have exactly one active vertex in each of their required contexts and their supports are disjoint; the remaining transverse values are forced. For deletion in the other copy, interchange the two indicators. Table S92 supplies an explicit coloring for every required context of both inputs. The color strings use original vertex order, and their labels 0,1,2 may be permuted.

TABLE S92: Explicit deletion colorings with pattern (1,1,2) and the number of states after deleting the corresponding first-copy horizontal context.

Input	Deleted original context	Color string	Extension states
H_9	$\{1,2,3\}$	212010201201122	20
H_9	$\{1,11,12\}$	201020120121221	20
H_9	$\{2,8,15\}$	021020120121220	12
H_9	$\{3,4,5\}$	012120102012212	20
H_9	$\{4,10,13\}$	012102120121000	12
H_9	$\{5,6,7\}$	012010120121220	20
H_9	$\{6,12,14\}$	012102102012202	12
H_9	$\{7,8,9\}$	012010200121222	20
H_9	$\{9,10,11\}$	012010201121222	20
H^+_{YO}	$\{1,2,3\}$	200021210112020102011222 1	16
H^+_{YO}	$\{1,10,19\}$	210021201212020001011202 1	8
H^+_{YO}	$\{1,13,20\}$	120210201012120011200220 1	8
H^+_{YO}	$\{2,5,8\}$	021021012120201122211001 2	24
H^+_{YO}	$\{3,11,14\}$	012021102120200122210001 1	8
H^+_{YO}	$\{3,12,25\}$	021210102101222020210101 0	8
H^+_{YO}	$\{4,5,6\}$	021200210102121110220221 0	16
H^+_{YO}	$\{4,10,18\}$	021210201202121010120201 0	8
H^+_{YO}	$\{4,11,24\}$	021201012120201120211000 2	8
H^+_{YO}	$\{6,12,17\}$	021210102100222110210101 2	8
H^+_{YO}	$\{6,13,22\}$	021012102120200112210001 2	8
H^+_{YO}	$\{7,8,9\}$	021012002120101112222001 2	16
H^+_{YO}	$\{7,11,15\}$	021201012100221120211001 2	8
H^+_{YO}	$\{7,13,21\}$	021012102120100112220001 2	8
H^+_{YO}	$\{9,10,23\}$	021210201102121010220201 0	8
H^+_{YO}	$\{9,12,16\}$	021210102102222010210101 0	8

Deleting one transverse context T_v instead permits an ordered pair of original state supports S_1, S_2 exactly when $S_1 \cap S_2 \subseteq \{v\}$. In these non-three-colorable inputs disjoint pairs do not exist, so the intersection must equal $\{v\}$; the isolated fresh vertex has two choices. For H_9 this gives four states for each $v = 1, \ldots, 12$ and twelve states for each $v = 13, 14, 15$. Together with Table S92, this proves criticality of all 33 contexts. For completed Yu–Oh it gives zero states at $v = 1, \ldots, 13$ and eight states at $v = 14, \ldots, 25$. Thus horizontal criticality does not imply criticality of this full transverse construction. The corresponding G_{32} extension has 45 vertices and 35 contexts and is stateless. Its distinct six state supports have pairwise intersection size one; for H_9 the distinct eight supports have intersection size one in 21 pairs and size three in seven pairs.

S3.7. Data and verification

The companion electronic package contains coordinate and context inventories, complete state and coloring enumerations, verification programs, exact parent inventories, and finite certificates. Vertex labels and state orderings agree with the tables in this supplement. The verification programs check the common Yu–Oh numbering, the published 49-ray parent map, the parent vectors and contexts, the full H_{65} state inventory, and cross-reference targets. The finite lower-bound certificate for the eighteen-state Hesse minimum is the exhaustive 540,672-candidate argument of Sec. S3.3.

Run the companion verification entry point from the project directory:

```
python verification/verify-integrated.py
```

The exact checks use rational or $\mathbb{Q}(\zeta)$ arithmetic, with $\zeta^2 + \zeta + 1 = 0$. Reports and hashes accompany the source.

The finite checks certify the instances and certificates they explicitly test. The analytic realization and existence results also rely on their proofs in the text.

S3.8. Historical Kochen–Specker gadget certificates (1965 Fig. 1 and 1967 Γ_1)

S3.8.1. The 1965 triangles and the completed historical TITS

Identifying the repeated occurrences of vertices 1 and 2 in Kochen–Specker's 1965 Fig. 1 gives

$$\{134, 325, 2\,9\,11, 11\,1\,10, 479, 5\,8\,10, 678\}. \tag{S44}$$

These are also the triples used in the displayed function definitions on p. 182 of Ref.[11]. Their four state supports are

$$\{1,5,6,9\}, \quad \{2,4,6,10\}, \quad \{3,6,9,10\}, \quad \{4,5,6,11\}.$$

The TIFS between 1 and 2 is nonvacuous. Directly, if both were true, contexts 134, 325, 2 9 11, and 11 1 10 would make 3,4,5,9,10,11 false; contexts 479 and 5 8 10 would then make both 7 and 8 true, contradicting 678.

Every state has 6 true and 7,8 false, so no three-coloring can exist: the three color-class states of any coloring would need to be disjoint. The graph is abstract rather than an injective ray example. In a purported injective realization in $\mathbb{C}^3$, vertices 1 and 9 are both orthogonal to 4 and 11. Distinct rays 4 and 11 span a plane, whose orthogonal complement is one-dimensional; thus rays 1 and 9 coincide. This proves the obstruction even over the complex field. No claim that the 1965 authors presented this graph as a Hilbert-space realization is intended.

For the 1967 Γ_1[12] (pp. 68,70), the correspondence between its original vertices and $I_{16,9}$ is

Γ_1	a_0	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9
$I_{16,9}$	10	9	1	7	3	8	2	11	4	24

The already complete triangles are $\{a_1, a_3, a_5\}$, $\{a_2, a_4, a_6\}$, and $\{a_7, a_8, a_9\}$. The six remaining edges

$$a_0a_1, \ a_0a_2, \ a_0a_8, \ a_3a_7, \ a_4a_7, \ a_5a_6$$

are completed by fresh vertices $23, 19, 18, 15, 14, 5$, respectively. The resulting nine contexts are exactly Eq. (S33). This also identifies its directed TITS $a_0 \to a_9$ with $10 \to 24$.

[1] M. Pavičić, J.-P. Merlet, B. McKay, and N. D. Megill, Journal of Physics A: Mathematical and General **38**, 1577 (2005), arXiv:quant-ph/0409014.
[2] J. Bub, Foundations of Physics **26**, 787 (1996).
[3] A. Cabello, J. M. Estebaranz, and G. García-Alcaine, Physics Letters A **212**, 183 (1996), arXiv:quant-ph/9706009.
[4] K. Svozil and J. Tkadlec, Journal of Mathematical Physics **37**, 5380 (1996).
[5] A. Silva and A. Ibiapina, "On orthogonal vector edge coloring," (2019), arXiv:1909.01918 [cs.DM].
[6] D. Blanuša, Glasnik Matematičko-Fizički i Astronomski, Serija II **1**, 31 (1946).
[7] R. Isaacs, The American Mathematical Monthly **82**, 221 (1975).
[8] G. Brinkmann, M. Preissmann, and D. Sasaki, Discrete Mathematics & Theoretical Computer Science **17**, 369 (2015).
[9] G. Schrag, Discrete Mathematics **70**, 185 (1988).
[10] M. Kernaghan, Journal of Physics A: Mathematical and General **27**, L829 (1994).
[11] S. Kochen and E. P. Specker, in *The Theory of Models, Proceedings of the 1963 International Symposium at Berkeley*, edited by J. W. Addison, L. Henkin, and A. Tarski (North Holland, Amsterdam, New York, Oxford, 1965) pp. 177–189.
[12] S. Kochen and E. P. Specker, Journal of Mathematics and Mechanics **17**, 59 (1967).