

Abstraction of Graph Theory

Wael Ali

Section (2)

- A **simple graph** G consists of a non-empty finite set $V(G)$ of elements called vertices (or nodes), and a finite set $E(G)$ of distinct unordered pairs of distinct elements of $V(G)$ called edges.
- A **graph** G consists of a non-empty finite set $V(G)$ of elements called vertices, and a finite family $E(G)$ of unordered pairs of (not necessarily distinct) elements of $V(G)$ called edges.
- Two graphs G_1 and G_2 are **isomorphic** if there is a one-one correspondence between the vertices of G_1 and those of G_2 such that the number of edges joining any two vertices of G_1 is equal to the number of edges joining the corresponding vertices of G_2 .




Figure 1: Two simple graphs that are isomorphic to each other

- A graph is **connected** if it cannot be expressed as the union of two graphs, and **disconnected**.
- Two vertices v and w of a graph G are **adjacent** if there is an edge vw joining them, and the vertices v and w are then **incident** with such an edge. Similarly, two distinct edges e and f are adjacent if they have a vertex in common.






Figure 2: Each of G_1 and G_2 is a component of $G_1 \cup G_2$

- The **degree** of a vertex v of G is the number of edges incident with v , and is written $deg(v)$. *Loop-edge* increase node-degree by 2. vertex of degree 0 is an **isolated vertex** and a vertex of degree 1 is an **end-vertex**.
- **Handshaking lemma**; in any graph the sum of all the vertex-degrees is an even number - in fact, twice the number of edges, since each edge contributes exactly 2 to the sum.
- A **subgraph** of a graph G is a graph, each of whose vertices belongs to $V(G)$ and each of whose edges belongs to $E(G)$.
- **Matrix representation** one way to represent graph is by its **adjacency matrix** A , and its **incidence matrix** M as follows;



Exercise 2

2a Write down the vertex-set and edge-set of each graph in Fig 2.5



The first graph G_1 is $(V(G_1), E(G_1))$

$$\begin{aligned} V(G_1) &= \{x, y, z, u, v, w\} \\ E(G_1) &= \{\{x, u\}, \{x, v\}, \{x, w\}, \\ &\quad \{y, u\}, \{y, v\}, \{y, w\}, \\ &\quad \{z, u\}, \{z, v\}, \{z, w\}\} \end{aligned}$$

The second graph G_2 is $(V(G_2), E(G_2))$

$$\begin{aligned} V(G_2) &= \{n, m, q, r, l, p\} \\ E(G_2) &= \{\{n, r\}, \{n, p\}, \{n, q\}, \\ &\quad \{m, r\}, \{m, p\}, \{m, q\}, \\ &\quad \{l, r\}, \{l, p\}, \{l, q\}\} \end{aligned}$$

2b Draw;

- (i) a simple graph.
 - (ii) a non-simple graph with no loops.
 - (iii) a non-simple graph with no multiple edges, each having 5 vertices each having 5 vertices and 8 edges.
-

2c Draw;

- (i) Draw a graph on six vertices whose degrees are 5,5,5,5,3,3; does there exist a simple graph with these degrees?




Figure 3: (i) a simple graph




Figure 4: (ii) a non-simple graph with no loops




Figure 5: (iii) a 5-vertices and 8-degrees each

- (ii) How does the answer to part (i) changed if the degrees are 5, 5, 4, 3, 3,2?
-




Figure 6: (i) a non-simple graph with 6-vertices with degrees $[5,5,5,5,3,3]$

There isn't a simple graph with last mentioned degrees for a 6-vertices graph. But if we just remove the red-arc and one of the blue-arcs, we then get a simple graph as shown in the next figure.





Figure 7: (i) a simple graph with 6-vertices with degrees $[5,5,4,3,3,2]$

(2d) Verify that handshaking lemma is hold for figure 2.1

As we can see in the last figure, the sum



of all vertex-degrees is $2 + 2 + 3 + 1 = 8$ which is an even number.

- (2f) (i) By suitably lettering the vertices, show that the two graphs in Fig. 2.20 are isomorphic.
- (ii) Explain why two graphs in Fig. 2.21 are not isomorphic.

-
- (i) As shown in the following figure 8 the two graphs are labeled with the same letters in a way to emphasize that they are both isomorphic to each other.
 - (ii) As shown in the following figure 9, we cannot find the red part of the first graph as a subset in the second graph.




Figure 8: (i)





Figure 9: (ii) not isomorphic graphs

Section (3)

- **Null graphs**, a graph whose edge-set is empty. **Complete graph**, a simple graph in which each pair of distinct vertices are adjacent, in this case k -vertex must have $n(n-1)/2$ degree as n the total number of vertices.
- **Regular graph**, a graph in which each vertex has the same degree. **Platonic graphs**, formed from the vertices and edges of the five regular (Platonic) solids - the tetrahedron, octahedron, cube.
- **Bipartite graph**, If the vertex set of a graph G can be split into two disjoint sets A and B so that each edge of G joins a vertex of A and a vertex of B . A **complete bipartite graph** is a bipartite graph in which each vertex in A is joined to each vertex in B by just one edge.
- **Cubes**, the k -cube Q_k is the graph whose vertices correspond to the sequences (a_1, a_2, \dots, a_k) , where each $a_i = 0$ or 1 , and whose edges join those sequences that differ in just one place. You should check that Q_k has 2^k vertices and $k2^{k-1}$ edges, and is regular of degree k .



- **Complement of a simple graph**, if G is a simple graph with vertex set $V(G)$, its complement \bar{G} is the simple graph with vertex set $V(G)$ in which two vertices are adjacent if and only if they are not adjacent in G .

Exercise 3

- (3a) Draw the following graphs:

- the null graph N_5 .
- the complete graph K_6 .
- the complete bipartite graph $K_{2,4}$.
- the union of $K_{1,3}$ and W_4 .
- the complement of the cycle graph C_4 .




Figure 10: (i) Null graph N_5




Figure 11: (ii) Complete graph K_6




Figure 12: (iii) Complete bipartite graph $K_{2,4}$




Figure 13: (iv) Union of $K_{1,3}$ and W_4




Figure 14: (iv) Complement of cycle graph C_4

- (3c) Draw the graphs $K_{2,2,2}$, and $K_{3,3,2}$, and write down the number of edges of $K_{3,4,5}$.

The graphs $K_{2,2,2}$ and $K_{3,3,2}$ are shown in the following figures, respectively.
For the graph $K_{3,4,5}$, there is 47 edges.




Figure 15: $K_{2,2,2}$




Figure 16: $K_{3,3,2}$

- (3g) A simple graph that is isomorphic to its complement is self-complementary.

- (i) Prove that, if G is self-complementary, then G has $4k$ or $4k + 1$ vertices, where k is an integer,
- (ii) Find all self-complementary graphs with 4 and 5 vertices,
- (iii) Find a self-complementary graph with 8 vertices.

- (i) Proof

If G is a self-complementary with n vertices, and

$$G \cup \overline{G} = K_n.$$

But we know that, the total number of edge in the complete graph K_n i.e. $|E(K_n)|$ is $n(n-1)/2$, that is,

$$|E(G)| = |E(\overline{G})| = \frac{n(n-1)}{4}.$$

In other words, n or $n-1$ must be divisible by 4, that is, when n is $4k$ or $4k+1$.




Figure 17: (ii) 4 and 5 vertices self-complementary graphs




Figure 18: (iii) a self-complementary graph with 8 vertices