



Inverse Problems 1: convolution and deconvolution

Lesson 7: Tikhonov regularization, part I

Luca Ratti

September 24, 2019

University of Helsinki

Table of contents

1. Review
2. Regularization Theory
3. Tikhonov regularization
4. Tikhonov regularization - implementation

Review

Singular Value Decomposition

Linear inverse problems

Deconvolution can be seen as a linear inverse problem:

given $m \in \mathbb{R}^n$, find $f \in \mathbb{R}^n$ such that $Af = m$, being $A \in \mathbb{R}^{n \times n}$

Singular Value Decomposition

Linear inverse problems

Deconvolution can be seen as a linear inverse problem:

given $m \in \mathbb{R}^n$, find $f \in \mathbb{R}^n$ such that $Af = m$, being $A \in \mathbb{R}^{n \times n}$

Singular Value Decomposition is a tool to analyze the well-posedness of a linear inverse problem:

Singular Value Decomposition

$$A = UDV^T$$

U, V orthogonal matrices ($U^T = UU^T = V^T V = VV^T = I$),

$$D = \text{diag}\{d_1, \dots, d_n\}, \quad d_1 \geq d_2 \geq \dots \geq d_n$$

Singular Value Decomposition

Linear inverse problems

Deconvolution can be seen as a linear inverse problem:

given $m \in \mathbb{R}^n$, find $f \in \mathbb{R}^n$ such that $Af = m$, being $A \in \mathbb{R}^{n \times n}$

Singular Value Decomposition is a tool to analyze the well-posedness of a linear inverse problem:

Singular Value Decomposition

$$A = UDV^T$$

U, V orthogonal matrices ($U^T = UU^T = V^T V = VV^T = I$),

$$D = \text{diag}\{d_1, \dots, d_n\}, \quad d_1 \geq d_2 \geq \dots \geq d_n$$

- if $d_n > 0$, the problem is well posed: $\forall m \in \mathbb{R}^n \quad \exists! f \in \mathbb{R}^n$ s.t. $Af = m$. Moreover, $f = A^{-1}m$ and $\|f - f_\delta\| \leq \|A^{-1}\| \|m - m_\delta\|$.
- if $d_r > 0$ and $d_{r+1} = \dots = d_n = 0$, then existence, uniqueness and stability fail.

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

The operator \mathcal{A} is associated to an ill-posed problem

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

The operator \mathcal{A} is associated to an ill-posed problem

- The approximation by discretization causes "*stabilty*". In theory, for the discretized problem in \mathbb{R}^n , $d_n > 0$.

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

The operator \mathcal{A} is associated to an ill-posed problem

- The approximation by discretization causes "*stabilty*". In theory, for the discretized problem in \mathbb{R}^n , $d_n > 0$.
- The approximation of \mathcal{A} implies that $d_n \rightarrow 0$ as $n \rightarrow \infty$.

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

The operator \mathcal{A} is associated to an ill-posed problem

- The approximation by discretization causes "stabilty". In theory, for the discretized problem in \mathbb{R}^n , $d_n > 0$.
- The approximation of \mathcal{A} implies that $d_n \rightarrow 0$ as $n \rightarrow \infty$.
- The arithmetic of the computer is finite: if d_n is too small, it is indistinguishable from 0 (*machine epsilon*)

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

The operator \mathcal{A} is associated to an ill-posed problem

- The approximation by discretization causes "stabilty". In theory, for the discretized problem in \mathbb{R}^n , $d_n > 0$.
- The approximation of \mathcal{A} implies that $d_n \rightarrow 0$ as $n \rightarrow \infty$.
- The arithmetic of the computer is finite: if d_n is too small, it is indistinguishable from 0 (*machine epsilon*)

Conclusions

The matrix A is, in theory, always invertible. Nevertheless, $\|A^{-1}\| = \frac{1}{d_n} \rightarrow \infty$ as $n \rightarrow \infty \Rightarrow$ **ill-conditioning**.

What about deconvolution?

Main issue

The convolution matrix is an approximation in $\mathbb{R}^{n \times n}$ of a continuous operator \mathcal{A} operating on real-valued function. As $n \rightarrow \infty$, A converges to \mathcal{A} (in a suitable sense).

The operator \mathcal{A} is associated to an ill-posed problem

- The approximation by discretization causes "stability". In theory, for the discretized problem in \mathbb{R}^n , $d_n > 0$.
- The approximation of \mathcal{A} implies that $d_n \rightarrow 0$ as $n \rightarrow \infty$.
- The arithmetic of the computer is finite: if d_n is too small, it is indistinguishable from 0 (*machine epsilon*)

Conclusions

The matrix A is, in theory, always invertible. Nevertheless,

$\|A^{-1}\| = \frac{1}{d_n} \rightarrow \infty$ as $n \rightarrow \infty \Rightarrow$ **ill-conditioning**.

For large n , due to finite precision, $d_n = 0$. \Rightarrow **ill-posedness**.

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Solution

$$f = A^+m$$

Problem

$$\text{find } f \text{ s.t. } Af = m$$

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Solution

$$f = A^{-1}m$$

$$f^+ = A^+m$$

Problem

$$\text{find } f \text{ s.t. } Af = m$$

find

$$f^+ = \arg \min_{g \in \mathbb{R}^n} \{\|g\|, g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}\}$$

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Solution

$$f = A^{-1}m$$

Problem

$$\text{find } f \text{ s.t. } Af = m$$

$$f^+ = A^+m$$

find

$$f^+ = \arg \min_{g \in \mathbb{R}^n} \{\|g\|, g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}\}$$

- the first problem can be ill-posed, the second one is always well-posed

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Solution

$$f = A^{-1}m$$

Problem

$$\text{find } f \text{ s.t. } Af = m$$

$$f^+ = A^+m$$

find

$$f^+ = \arg \min_{g \in \mathbb{R}^n} \{\|g\|, g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}\}$$

- the first problem can be ill-posed, the second one is always well-posed
- if A is invertible, $A^{-1} = A^+$ (the second problem is a generalization of the first one)

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Solution

$$f = A^{-1}m$$

Problem

$$\text{find } f \text{ s.t. } Af = m$$

$$f^+ = A^+m$$

find

$$f^+ = \arg \min_{g \in \mathbb{R}^n} \{\|g\|, g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}\}$$

- the first problem can be ill-posed, the second one is always well-posed
- if A is invertible, $A^{-1} = A^+$ (the second problem is a generalization of the first one)
- even the second problem can be ill-conditioned: $\|A^+\| = \frac{1}{d_r}$ can be very large

Moore-Penrose pseudoinverse

The cure of ill-posedness is represented by the pseudoinverse:

$$A^+ = VD^+U^T, \quad D^+ = \text{diag}\left(\frac{1}{d_1}, \dots, \frac{1}{d_r}, 0, \dots, 0\right)$$

Solution

$$f = A^{-1}m$$

Problem

$$\text{find } f \text{ s.t. } Af = m$$

$$f^+ = A^+m$$

find

$$f^+ = \arg \min_{g \in \mathbb{R}^n} \{\|g\|, g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}\}$$

- the first problem can be ill-posed, the second one is always well-posed
- if A is invertible, $A^{-1} = A^+$ (the second problem is a generalization of the first one)
- even the second problem can be ill-conditioned: $\|A^+\| = \frac{1}{d_r}$ can be very large

from now on, we will not distinguish between f and f^+

Pseudoinverse - interpretation

If A is not invertible, ill-posedness can be treated by relying on:

Pseudoinverse - interpretation

If A is not invertible, ill-posedness can be treated by relying on:

1. Data fidelity

Existence fails: $\text{range}(A) \neq \mathbb{R}^n \Rightarrow \exists m \in \mathbb{R}^n : \nexists f \in \mathbb{R}^n : Af = m$
(you might never notice it if you do inverse crime). What to do?

Extract any possible information from the data

$$g \in \arg \min_{h \in \mathbb{R}^n} \{ \|Ah - m\| \}$$

Pseudoinverse - interpretation

If A is not invertible, ill-posedness can be treated by relying on:

1. Data fidelity

Existence fails: $\text{range}(A) \neq \mathbb{R}^n \Rightarrow \exists m \in \mathbb{R}^n : \nexists f \in \mathbb{R}^n : Af = m$
(you might never notice it if you do inverse crime). What to do?

Extract any possible information from the data

$$g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}$$

2. A priori knowledge

Uniqueness fails: $\ker(A) \neq \{0\} \Rightarrow \exists k \in \mathbb{R}^n : Af = A(f + k)$.

What to do? Use any a priori information at disposal on the solution E.g. the solution is smooth, has a known average, has small norm.

$$f^+ = \arg \min_{g \in \mathbb{R}^n} \{\|g\|, g \in \arg \min_{h \in \mathbb{R}^n} \{\|Ah - m\|\}\}$$

Regularization Theory

Regularization

The theory of Regularization of (linear, finite-dimensional) inverse problems is based on:

Regularization

The theory of Regularization of (linear, finite-dimensional) inverse problems is based on:

1. a family of regularization functionals $R_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^n$ varying with $\alpha > 0$ s.t.

$$R_\alpha m \rightarrow A^+ m \quad \text{as } \alpha \rightarrow 0, \quad \forall m \in \mathbb{R}^n.$$

We will mainly focus on linear regularization functionals, which can therefore be represented by a matrix in $\mathbb{R}^{n \times n}$.

Regularization

The theory of Regularization of (linear, finite-dimensional) inverse problems is based on:

1. a family of **regularization functionals** $R_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^n$ varying with $\alpha > 0$ s.t.

$$R_\alpha m \rightarrow A^+ m \quad \text{as } \alpha \rightarrow 0, \quad \forall m \in \mathbb{R}^n.$$

We will mainly focus on **linear** regularization functionals, which can therefore be represented by a matrix in $\mathbb{R}^{n \times n}$.

2. a suitable **parameter choice rule** $\alpha = \alpha(\delta)$ ensuring

$$R_{\alpha(\delta)} m_\delta \rightarrow A^+ m \quad \text{as } \delta \rightarrow 0, \quad \forall m_\delta : \|m_\delta - m\| \leq \delta, \quad \forall m \in \mathbb{R}^n$$

Regularization

The theory of Regularization of (linear, finite-dimensional) inverse problems is based on:

1. a family of **regularization functionals** $R_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^n$ varying with $\alpha > 0$ s.t.

$$R_\alpha m \rightarrow A^+ m \quad \text{as } \alpha \rightarrow 0, \quad \forall m \in \mathbb{R}^n.$$

We will mainly focus on **linear** regularization functionals, which can therefore be represented by a matrix in $\mathbb{R}^{n \times n}$.

2. a suitable **parameter choice rule** $\alpha = \alpha(\delta)$ ensuring

$$R_{\alpha(\delta)} m_\delta \rightarrow A^+ m \quad \text{as } \delta \rightarrow 0, \quad \forall m_\delta : \|m_\delta - m\| \leq \delta, \quad \forall m \in \mathbb{R}^n$$

A desirable property is that R_α are **better-conditioned** than A^+ :

$$\|R_\alpha\| \leq \|A^+\| \quad \forall \alpha > 0$$

Regularization

The theory of Regularization of (linear, finite-dimensional) inverse problems is based on:

1. a family of **regularization functionals** $R_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^n$ varying with $\alpha > 0$ s.t.

$$R_\alpha m \rightarrow A^+ m \quad \text{as } \alpha \rightarrow 0, \quad \forall m \in \mathbb{R}^n.$$

We will mainly focus on **linear** regularization functionals, which can therefore be represented by a matrix in $\mathbb{R}^{n \times n}$.

2. a suitable **parameter choice rule** $\alpha = \alpha(\delta)$ ensuring

$$R_{\alpha(\delta)} m_\delta \rightarrow A^+ m \quad \text{as } \delta \rightarrow 0, \quad \forall m_\delta : \|m_\delta - m\| \leq \delta, \quad \forall m \in \mathbb{R}^n$$

A desirable property is that R_α are **better-conditioned** than A^+ :

$$\|R_\alpha\| \leq \|A^+\| \quad \forall \alpha > 0$$

Remark: a parameter choice $\alpha = \alpha(\delta)$ is defined *a priori*, since it holds for any m and any perturbation m_δ . We will focus more on a *posteriori* (heuristic) rules $\alpha = \alpha(\delta, m_\delta)$.

Graphical interpretation

Compute $f_{\alpha,\delta} = R_{\alpha}m_{\delta}$ instead of $f_{\delta} = A^{+}m_{\delta}$. Error bound for the approximation of $f = f^{+} = A^{+}m$ (remember $\|m - m_{\delta}\| \leq \delta$):

Graphical interpretation


Compute $f_{\alpha,\delta} = R_{\alpha}m_{\delta}$ instead of $f_{\delta} = A^{+}m_{\delta}$. Error bound for the approximation of $f = f^{+} = A^{+}m$ (remember $\|m - m_{\delta}\| \leq \delta$):

$$\|f^{+} - A^{+}m_{\delta}\| \leq \|A^{+}\|\delta, \quad \|f^{+} - R_{\alpha}m_{\delta}\| \leq \|f^{+} - R_{\alpha}m\| + \|R_{\alpha}\|\delta.$$

Graphical interpretation

Compute $f_{\alpha,\delta} = R_\alpha m_\delta$ instead of $f_\delta = A^+ m_\delta$. Error bound for the approximation of $f = f^+ = A^+ m$ (remember $\|m - m_\delta\| \leq \delta$):


$$\|f^+ - A^+ m_\delta\| \leq \|A^+\| \delta, \quad \|f^+ - R_\alpha m_\delta\| \leq \|f^+ - R_\alpha m\| + \|R_\alpha\| \delta.$$



Graphical interpretation

Compute $f_{\alpha,\delta} = R_\alpha m_\delta$ instead of $f_\delta = A^+ m_\delta$. Error bound for the approximation of $f = f^+ = A^+ m$ (remember $\|m - m_\delta\| \leq \delta$):


$$\|f^+ - A^+ m_\delta\| \leq \|A^+\| \delta, \quad \|f^+ - R_\alpha m_\delta\| \leq \|f^+ - R_\alpha m\| + \|R_\alpha\| \delta.$$



Graphical interpretation

Compute $f_{\alpha,\delta} = R_\alpha m_\delta$ instead of $f_\delta = A^+ m_\delta$. Error bound for the approximation of $f = f^+ = A^+ m$ (remember $\|m - m_\delta\| \leq \delta$):


$$\|f^+ - A^+ m_\delta\| \leq \|A^+\| \delta, \quad \|f^+ - R_\alpha m_\delta\| \leq \|f^+ - R_\alpha m\| + \|R_\alpha\| \delta.$$



Graphical interpretation

Compute $f_{\alpha,\delta} = R_{\alpha}m_{\delta}$ instead of $f_{\delta} = A^{+}m_{\delta}$. Error bound for the approximation of $f = f^{+} = A^{+}m$ (remember $\|m - m_{\delta}\| \leq \delta$):


$$\|f^{+} - A^{+}m_{\delta}\| \leq \|A^{+}\|\delta, \quad \|f^{+} - R_{\alpha}m_{\delta}\| \leq \|f^{+} - R_{\alpha}m\| + \|R_{\alpha}\|\delta.$$



Graphical interpretation

Compute $f_{\alpha,\delta} = R_\alpha m_\delta$ instead of $f_\delta = A^+ m_\delta$. Error bound for the approximation of $f = f^+ = A^+ m$ (remember $\|m - m_\delta\| \leq \delta$):


$$\|f^+ - A^+ m_\delta\| \leq \|A^+\| \delta, \quad \|f^+ - R_\alpha m_\delta\| \leq \|f^+ - R_\alpha m\| + \|R_\alpha\| \delta.$$



Graphical interpretation

Compute $f_{\alpha,\delta} = R_\alpha m_\delta$ instead of $f_\delta = A^+ m_\delta$. Error bound for the approximation of $f = f^+ = A^+ m$ (remember $\|m - m_\delta\| \leq \delta$):

$$\|f^+ - A^+ m_\delta\| \leq \|A^+\| \delta, \quad \|f^+ - R_\alpha m_\delta\| \leq \|f^+ - R_\alpha m\| + \|R_\alpha\| \delta.$$




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :


Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :



Graphical interpretation gets complicated

The previous pictures showed a single α at a time. Let's visualize the error bound as a function of δ and α :




Graphical interpretation - parameter choice rule

The parameter choice rule is a relationship $\alpha = \alpha(\delta)$ ensuring that, along the curve $\|f^+ - R_{\alpha(\delta)}m_\delta\|$, we are converging to 0


Graphical interpretation - parameter choice rule

The parameter choice rule is a relationship $\alpha = \alpha(\delta)$ ensuring that, along the curve $\|f^+ - R_{\alpha(\delta)}m_\delta\|$, we are converging to 0




Graphical interpretation - parameter choice rule

The parameter choice rule is a relationship $\alpha = \alpha(\delta)$ ensuring that, along the curve $\|f^+ - R_{\alpha(\delta)}m_\delta\|$, we are converging to 0



Graphical interpretation - parameter choice rule

The parameter choice rule is a relationship $\alpha = \alpha(\delta)$ ensuring that, along the curve $\|f^+ - R_{\alpha(\delta)}m_\delta\|$, we are converging to 0



Tikhonov regularization

Tikhonov regularization

Tikhonov regularization consists in a specific choice of the regularization functionals, denoted by T_α .

Tikhonov regularization

Tikhonov regularization consists in a specific choice of the regularization functionals, denoted by T_α .

Motivation: the pseudoinverse relies on a combination of **data fidelity** (minimize $\|Af - m\|$) and **a priori knowledge** (in particular, minimize $\|f\|$).

Tikhonov regularization

Tikhonov regularization consists in a specific choice of the regularization functionals, denoted by T_α .

Motivation: the pseudoinverse relies on a combination of **data fidelity** (minimize $\|Af - m\|$) and **a priori knowledge** (in particular, minimize $\|f\|$).

Tikhonov regularization

$$T_\alpha m = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m\|^2 + \alpha \|f\|^2 \}$$

Tikhonov regularization

Tikhonov regularization consists in a specific choice of the regularization functionals, denoted by T_α .

Motivation: the pseudoinverse relies on a combination of **data fidelity** (minimize $\|Af - m\|$) and **a priori knowledge** (in particular, minimize $\|f\|$).

Tikhonov regularization

$$T_\alpha m = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m\|^2 + \alpha \|f\|^2 \}$$

Our goals:

- find an explicit expression for the matrix representing T_α ;
- check that this choice is a regularization strategy;
- implement and see the benefits of regularization.

A first explicit expression

Theorem

Let $A = UDV^T$, being U, V orthogonal matrices and $D = \text{diag}(d_1, \dots, d_n)$, such that $d_1 \geq \dots \geq d_n$. Suppose that $\exists r \leq n$: $d_r > 0$ and $d_{r+1} = \dots = d_n = 0$. Let

$$\mathcal{D}_\alpha^+ = \text{diag} \left(\frac{d_1}{d_1^2 + \alpha}, \dots, \frac{d_n}{d_n^2 + \alpha} \right).$$

Then, $T_\alpha = V\mathcal{D}_\alpha^+U^T$

A first explicit expression

Theorem

Let $A = UDV^T$, being U, V orthogonal matrices and $D = \text{diag}(d_1, \dots, d_n)$, such that $d_1 \geq \dots \geq d_n$. Suppose that $\exists r \leq n$: $d_r > 0$ and $d_{r+1} = \dots = d_n = 0$. Let

$$\mathcal{D}_\alpha^+ = \text{diag} \left(\frac{d_1}{d_1^2 + \alpha}, \dots, \frac{d_n}{d_n^2 + \alpha} \right).$$

Then, $T_\alpha = V\mathcal{D}_\alpha^+U^T$

Preliminary remarks:

- in case $d_k = 0$, then $\frac{d_k}{d_k^2 + \alpha} = 0$;
- if $\alpha \rightarrow 0$ and $d_k \neq 0$, $\frac{d_k}{d_k^2 + \alpha} \rightarrow \frac{1}{d_k}$;

A first explicit expression

Theorem

Let $A = UDV^T$, being U, V orthogonal matrices and $D = \text{diag}(d_1, \dots, d_n)$, such that $d_1 \geq \dots \geq d_n$. Suppose that $\exists r \leq n$: $d_r > 0$ and $d_{r+1} = \dots = d_n = 0$. Let

$$\mathcal{D}_\alpha^+ = \text{diag} \left(\frac{d_1}{d_1^2 + \alpha}, \dots, \frac{d_n}{d_n^2 + \alpha} \right).$$

Then, $T_\alpha = V\mathcal{D}_\alpha^+U^T$


Comparison with TSVD:

we use a different manipulation on the singular values:

$$d_\alpha^{\text{Tikh}}(d) = \frac{d}{d^2 + \alpha},$$

whereas in TSVD we have:

$$d_\alpha^{\text{TSVD}}(d) = \frac{1}{d} \chi_{(\alpha, \infty)}(d)$$



Explicit expression - proof (part I)

- Since the matrix V is orthogonal, its columns consist in an orthonormal basis of \mathbb{R}^n . Hence, every vector $f \in \mathbb{R}^n$ can be expressed as $f = Va$, being $a \in \mathbb{R}^n$.

Explicit expression - proof (part I)

- Since the matrix V is orthogonal, its columns consist in an orthonormal basis of \mathbb{R}^n . Hence, every vector $f \in \mathbb{R}^n$ can be expressed as $f = Va$, being $a \in \mathbb{R}^n$.
- Let $T_\alpha m = Va^*$. Then, we can reformulate Tikhonov regularization as follows: find

$$a^* = \arg \min_{a \in \mathbb{R}^n} \{ \|AVa - m\|^2 + \alpha \|Va\|^2 \}$$

Explicit expression - proof (part I)

- Since the matrix V is orthogonal, its columns consist in an orthonormal basis of \mathbb{R}^n . Hence, every vector $f \in \mathbb{R}^n$ can be expressed as $f = Va$, being $a \in \mathbb{R}^n$.
- Let $T_\alpha m = Va^*$. Then, we can reformulate Tikhonov regularization as follows: find

$$a^* = \arg \min_{a \in \mathbb{R}^n} \{ \|AVa - m\|^2 + \alpha \|Va\|^2 \}$$

- By the orthogonality of U and V , we notice that

$$\begin{aligned} \|AVa - m\|^2 + \alpha \|Va\|^2 &= \|UDV^T Va - m\|^2 + \alpha a^T V^T Va \\ &= \|UDa - UU^T m\|^2 + \alpha \|a\|^2 = \|Da - U^T m\|^2 + \alpha \|a\|^2 \\ &= \|Da - m'\|^2 + \alpha \|a\|^2, \end{aligned}$$

where we have defined $m' = U^T m$.

Explicit expression - proof (part II)

- D is a diagonal matrix; suppose $d_{r+i} = \dots = d_n = 0$. By the definition of the norm of a vector:

$$\|Da - m'\|^2 + \alpha\|a\|^2 = \sum_{j=1}^n (d_j a_j - m'_j)^2 + \alpha \sum_{j=i}^n a_j^2$$

Explicit expression - proof (part II)

- D is a diagonal matrix; suppose $d_{r+i} = \dots = d_n = 0$. By the definition of the norm of a vector:

$$\begin{aligned}\|Da - m'\|^2 + \alpha\|a\|^2 &= \sum_{j=1}^n (d_j a_j - m'_j)^2 + \alpha \sum_{j=1}^n a_j^2 \\ &= \sum_{j=1}^r (d_j^2 a_j^2 - 2d_j a_j m'_j + m_j'^2) + \sum_{j=r+1}^n m_j'^2 + \alpha \sum_{j=1}^n a_j^2\end{aligned}$$

Explicit expression - proof (part II)

- D is a diagonal matrix; suppose $d_{r+i} = \dots = d_n = 0$. By the definition of the norm of a vector:

$$\begin{aligned}\|Da - m'\|^2 + \alpha\|a\|^2 &= \sum_{j=1}^n (d_j a_j - m'_j)^2 + \alpha \sum_{j=1}^n a_j^2 \\&= \sum_{j=1}^r (d_j^2 a_j^2 - 2d_j a_j m'_j + m_j'^2) + \sum_{j=r+1}^n m_j'^2 + \alpha \sum_{j=1}^n a_j^2 \\&= \sum_{j=1}^r \left((d_j^2 + \alpha) a_j^2 - 2d_j a_j m'_j \right) + \alpha \sum_{j=r+1}^n a_j^2 + \sum_{j=1}^n m_j'^2\end{aligned}$$

Explicit expression - proof (part II)

- D is a diagonal matrix; suppose $d_{r+i} = \dots = d_n = 0$. By the definition of the norm of a vector:

$$\begin{aligned}\|Da - m'\|^2 + \alpha\|a\|^2 &= \sum_{j=1}^n (d_j a_j - m'_j)^2 + \alpha \sum_{j=1}^n a_j^2 \\&= \sum_{j=1}^r (d_j^2 a_j^2 - 2d_j a_j m'_j + m_j'^2) + \sum_{j=r+1}^n m_j'^2 + \alpha \sum_{j=1}^n a_j^2 \\&= \sum_{j=1}^r \left((d_j^2 + \alpha) a_j^2 - 2d_j a_j m'_j \right) + \alpha \sum_{j=r+1}^n a_j^2 + \sum_{j=1}^n m_j'^2 \\&= \sum_{j=1}^r (d_j^2 + \alpha) \left(a_j^2 - 2 \frac{d_j a_j m'_j}{d_j^2 + \alpha} \right) + \alpha \sum_{j=r+1}^n a_j^2 + \sum_{j=1}^n m_j'^2\end{aligned}$$

Explicit expression - proof (part II)

- D is a diagonal matrix; suppose $d_{r+i} = \dots = d_n = 0$. By the definition of the norm of a vector:

$$\begin{aligned}\|Da - m'\|^2 + \alpha\|a\|^2 &= \sum_{j=1}^n (d_j a_j - m'_j)^2 + \alpha \sum_{j=1}^n a_j^2 \\&= \sum_{j=1}^r (d_j^2 a_j^2 - 2d_j a_j m'_j + m_j'^2) + \sum_{j=r+1}^n m_j'^2 + \alpha \sum_{j=1}^n a_j^2 \\&= \sum_{j=1}^r \left((d_j^2 + \alpha) a_j^2 - 2d_j a_j m'_j \right) + \alpha \sum_{j=r+1}^n a_j^2 + \sum_{j=1}^n m_j'^2 \\&= \sum_{j=1}^r (d_j^2 + \alpha) \left(a_j^2 - 2 \frac{d_j a_j m'_j}{d_j^2 + \alpha} \right) + \alpha \sum_{j=r+1}^n a_j^2 + \sum_{j=1}^n m_j'^2 \\&= \sum_{j=1}^r (d_j^2 + \alpha) \left(a_j - \frac{d_j m'_j}{d_j^2 + \alpha} \right)^2 - \sum_{j=1}^r \frac{d_j^2 m_j'^2}{(d_j^2 + \alpha)^2} + \alpha \sum_{j=r+1}^n a_j^2 + \sum_{j=1}^n m_j'^2\end{aligned}$$

Explicit expression - proof (part III)

- We are looking for a vector $a^* \in \mathbb{R}^n$ minimizing

$$\sum_{j=1}^r (d_j^2 + \alpha) \left(a_j - \frac{d_j m'_j}{d_j^2 + \alpha} \right)^2 + \alpha \sum_{j=r+1}^n a_j^2 + R(D, m', \alpha),$$

where we collected in $R(D, m', \alpha)$ all the terms independent of a .

Explicit expression - proof (part III)

- We are looking for a vector $a^* \in \mathbb{R}^n$ minimizing

$$\sum_{j=1}^r (d_j^2 + \alpha) \left(a_j - \frac{d_j m'_j}{d_j^2 + \alpha} \right)^2 + \alpha \sum_{j=r+1}^n a_j^2 + R(D, m', \alpha),$$

where we collected in $R(D, m', \alpha)$ all the terms independent of a .

- The solution can be computed by hand selecting

$$a_j = \frac{d_j m'_j}{d_j^2 + \alpha} \text{ for } j = 1, \dots, r; \quad a_j = 0 \text{ for } j = r+1, \dots, n;$$

the first formula can be used in both cases ($d_{r+1} = \dots = d_n = 0$).

Explicit expression - proof (part III)

- We are looking for a vector $a^* \in \mathbb{R}^n$ minimizing

$$\sum_{j=1}^r (d_j^2 + \alpha) \left(a_j - \frac{d_j m'_j}{d_j^2 + \alpha} \right)^2 + \alpha \sum_{j=r+1}^n a_j^2 + R(D, m', \alpha),$$

where we collected in $R(D, m', \alpha)$ all the terms independent of a .

- The solution can be computed by hand selecting

$$a_j = \frac{d_j m'_j}{d_j^2 + \alpha} \text{ for } j = 1, \dots, r; \quad a_j = 0 \text{ for } j = r+1, \dots, n;$$

the first formula can be used in both cases ($d_{r+1} = \dots = d_n = 0$).

- We conclude $a^* = \mathcal{D}_\alpha^+ m' = \mathcal{D}_\alpha^+ U^T m$, hence $T_\alpha m = V \mathcal{D}_\alpha^+ U^T m$.

Theoretical results

Employing the explicit representation $T_\alpha = V\mathcal{D}_\alpha^+U^T$, we are able to prove the following results:

Theoretical results

Employing the explicit representation $T_\alpha = V\mathcal{D}_\alpha^+U^T$, we are able to prove the following results:

- the matrix T_α is better-conditioned than A^+ . Indeed,

$$\|T_\alpha\| = \frac{d_r}{d_r^2 + \alpha} \leq \frac{1}{d_r};$$

Theoretical results

Employing the explicit representation $T_\alpha = V\mathcal{D}_\alpha^+U^T$, we are able to prove the following results:

- the matrix T_α is **better-conditioned** than A^+ . Indeed,

$$\|T_\alpha\| = \frac{d_r}{d_r^2 + \alpha} \leq \frac{1}{d_r};$$

- the operators T_α are **regularization operators**. Indeed,

$$\|T_\alpha m - f^+\| \leq \frac{\alpha}{d_r(d_r^2 + \alpha)} \|m\|,$$

hence, for any $m \in \mathbb{R}^n$, $T_\alpha \rightarrow A^+$ as $\alpha \rightarrow 0$.

Theoretical results

Employing the explicit representation $T_\alpha = V\mathcal{D}_\alpha^+U^T$, we are able to prove the following results:

- the matrix T_α is **better-conditioned** than A^+ . Indeed,

$$\|T_\alpha\| = \frac{d_r}{d_r^2 + \alpha} \leq \frac{1}{d_r};$$

- the operators T_α are **regularization operators**. Indeed,

$$\|T_\alpha m - f^+\| \leq \frac{\alpha}{d_r(d_r^2 + \alpha)} \|m\|,$$

hence, for any $m \in \mathbb{R}^n$, $T_\alpha \rightarrow A^+$ as $\alpha \rightarrow 0$.

- an a priori **parameter choice** is available for α , but we are not interested on it (theoretically demanding, sub-optimal with respect to the a posteriori rules we will learn)

Tikhonov regularization - implementation

Implementation

Example

1. Set $n = 200$. Define in Matlab the signal $f \in \mathbb{R}^n$ such that:

$$f = \begin{cases} 1 & \text{if } n_1 \leq n \leq n_2, \\ 3 \frac{n - n_3}{n_4 - n_3} & \text{if } n_3 \leq n \leq n_4, \\ -1 - \cos\left(\frac{2\pi n}{n_6 - n_5}\right) & \text{if } n_5 \leq n \leq n_6, \end{cases}$$

being

$$\begin{aligned} n_1 &= 2 \left\lfloor \frac{n}{15} \right\rfloor, \quad n_2 = 3 \left\lfloor \frac{n}{15} \right\rfloor, \\ n_3 &= 4 \left\lfloor \frac{n}{15} \right\rfloor, \quad n_4 = 8 \left\lfloor \frac{n}{15} \right\rfloor, \\ n_5 &= 10 \left\lfloor \frac{n}{15} \right\rfloor, \quad n_6 = 14 \left\lfloor \frac{n}{15} \right\rfloor. \end{aligned}$$

Implementation


Example

1. Set $n = 200$. Define in Matlab the signal $f \in \mathbb{R}^n$ such that:

$$f = \begin{cases} 1 & \text{if } n_1 \leq n \leq n_2, \\ 3 \frac{n - n_3}{n_4 - n_3} & \text{if } n_3 \leq n \leq n_4, \\ -1 - \cos\left(\frac{2\pi n}{n_6 - n_5}\right) & \text{if } n_5 \leq n \leq n_6, \end{cases}$$

being

$$\begin{aligned} n_1 &= 2 \left\lfloor \frac{n}{15} \right\rfloor, & n_2 &= 3 \left\lfloor \frac{n}{15} \right\rfloor, \\ n_3 &= 4 \left\lfloor \frac{n}{15} \right\rfloor, & n_4 &= 8 \left\lfloor \frac{n}{15} \right\rfloor, \\ n_5 &= 10 \left\lfloor \frac{n}{15} \right\rfloor, & n_6 &= 14 \left\lfloor \frac{n}{15} \right\rfloor. \end{aligned}$$



Implementation

Example

2. Define the point spread function by normalizing the vector $[1, 4, 8, 16, 19, 15, 10, 7, 1]$. Create the convolution matrix associated to the zero-padding case.
3. Generate the noisy measurement $m_\delta = Af + \delta r$, being r a Gaussian random vector of n dimensions, $\|r\| = 1$. Consider $\delta = 10^{-5}$.
4. Compute the pseudoinverse A^+ of A and find $f_\delta = A^+ m_\delta$.
5. Select $\alpha = 0.1$. Compute T_α and find $f_{\alpha,\delta} = T_\alpha m_\delta$.
6. Compare $f_{\alpha,\delta}$ and f_δ by graphical inspection and by computing their normalized error with respect to $f^+ = f$.
7. Repeat the previous experiment with the following couples (δ, α) :
 $(10^{-5}, 10)$, $(10^{-5}, 10^{-6})$, $(10^{-5}, 10^{-14})$; $(10^{-1}, 0.1)$, $(10^{-3}, 0.1)$, $(10^{-8}, 0.1)$.

Results interpretation

