



Inverse Problems 1: convolution and deconvolution

Lesson 8: Tikhonov regularization, part II

Luca Ratti

September 25, 2019

University of Helsinki

1. Normal equations for Tikhonov regularization
2. Generalized Tikhonov regularization
3. Choice of the regularization parameter

Normal equations for Tikhonov regularization

Tikhonov regularization

$$T_{\alpha}m = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m\|^2 + \alpha \|f\|^2 \}$$

Tikhonov regularization

$$T_{\alpha}m = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m\|^2 + \alpha \|f\|^2 \}$$

How to compute $T_{\alpha}m$?

Theorem

Let $A = UDV^T$, being U , V orthogonal matrices and $D = \text{diag}(d_1, \dots, d_n)$, such that $d_1 \geq \dots \geq d_n$. Suppose that $\exists r \leq n$: $d_r > 0$ and $d_{r+1} = \dots = d_n = 0$. Let

$$\mathcal{D}_{\alpha}^{+} = \text{diag} \left(\frac{d_1}{d_1^2 + \alpha}, \dots, \frac{d_n}{d_n^2 + \alpha} \right).$$

Then, $T_{\alpha} = V\mathcal{D}_{\alpha}^{+}U^T$

Tikhonov regularization

$$T_{\alpha}m = \arg \min_{f \in \mathbb{R}^n} \{\|Af - m\|^2 + \alpha\|f\|^2\}$$

How to compute $T_{\alpha}m$?

Theorem

Let $A = UDV^T$, being U, V orthogonal matrices and $D = \text{diag}(d_1, \dots, d_n)$, such that $d_1 \geq \dots \geq d_n$. Suppose that $\exists r \leq n$: $d_r > 0$ and $d_{r+1} = \dots = d_n = 0$. Let

$$\mathcal{D}_{\alpha}^{+} = \text{diag} \left(\frac{d_1}{d_1^2 + \alpha}, \dots, \frac{d_n}{d_n^2 + \alpha} \right).$$

Then, $T_{\alpha} = V\mathcal{D}_{\alpha}^{+}U^T$

Issue: the computation of U, D, V is numerically expensive.

Normal equations

Theorem

The solution $T_\alpha m$ can be computed as

$$T_\alpha m = (A^T A + \alpha I)^{-1} A^T m.$$

Normal equations

Theorem

The solution $T_\alpha m$ can be computed as

$$T_\alpha m = (A^T A + \alpha I)^{-1} A^T m.$$

Proof

- Define the quadratic functional to be minimized:

$$Q_\alpha(f) = \|Af - m\|^2 + \alpha\|f\|^2.$$

Normal equations

Theorem

The solution $T_\alpha m$ can be computed as

$$T_\alpha m = (A^T A + \alpha I)^{-1} A^T m.$$

Proof

- Define the quadratic functional to be minimized:

$$Q_\alpha(f) = \|Af - m\|^2 + \alpha\|f\|^2.$$

- Then, if $f_\alpha = T_\alpha m$ is the minimum of Q_α , the following optimality condition must be satisfied:

$$\left. \frac{d}{dt} Q_\alpha(f_\alpha + tw) \right|_{t=0} = 0 \quad \forall w \in \mathbb{R}^n$$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$Q_{\alpha}(f_{\alpha} + tw) = \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2$$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned} Q_\alpha(f_\alpha + tw) &= \|A(f_\alpha + tw) - m\|^2 + \alpha\|f_\alpha + tw\|^2 \\ &= \langle (Af_\alpha - m) + tAw, (Af_\alpha - m) + tAw \rangle + \alpha\langle f_\alpha + tw, f_\alpha + tw \rangle \end{aligned}$$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned}Q_{\alpha}(f_{\alpha} + tw) &= \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2 \\&= \langle (Af_{\alpha} - m) + tAw, (Af_{\alpha} - m) + tAw \rangle + \alpha\langle f_{\alpha} + tw, f_{\alpha} + tw \rangle \\&= \|Af_{\alpha} - m\|^2 + 2t\langle Af_{\alpha} - m, Aw \rangle + t^2\|Aw\|^2\end{aligned}$$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned}Q_{\alpha}(f_{\alpha} + tw) &= \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2 \\&= \langle (Af_{\alpha} - m) + tAw, (Af_{\alpha} - m) + tAw \rangle + \alpha\langle f_{\alpha} + tw, f_{\alpha} + tw \rangle \\&= \|Af_{\alpha} - m\|^2 + 2t\langle Af_{\alpha} - m, Aw \rangle + t^2\|Aw\|^2 \\&\quad + \alpha\|f_{\alpha}\|^2 + 2t\alpha\langle f_{\alpha}, w \rangle + t^2\alpha\|w\|^2\end{aligned}$$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned}Q_{\alpha}(f_{\alpha} + tw) &= \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2 \\&= \langle (Af_{\alpha} - m) + tAw, (Af_{\alpha} - m) + tAw \rangle + \alpha\langle f_{\alpha} + tw, f_{\alpha} + tw \rangle \\&= \|Af_{\alpha} - m\|^2 + 2t\langle Af_{\alpha} - m, Aw \rangle + t^2\|Aw\|^2 \\&\quad + \alpha\|f_{\alpha}\|^2 + 2t\alpha\langle f_{\alpha}, w \rangle + t^2\alpha\|w\|^2\end{aligned}$$

- the computation of the derivative implies

$$\frac{d}{dt}Q_{\alpha}(f_{\alpha} + tw) = 2\langle Af_{\alpha} - m, Aw \rangle + 2t\|Aw\|^2 + 2\alpha\langle f_{\alpha}, w \rangle + 2\alpha t\|w\|^2$$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned}Q_{\alpha}(f_{\alpha} + tw) &= \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2 \\&= \langle (Af_{\alpha} - m) + tAw, (Af_{\alpha} - m) + tAw \rangle + \alpha\langle f_{\alpha} + tw, f_{\alpha} + tw \rangle \\&= \|Af_{\alpha} - m\|^2 + 2t\langle Af_{\alpha} - m, Aw \rangle + t^2\|Aw\|^2 \\&\quad + \alpha\|f_{\alpha}\|^2 + 2t\alpha\langle f_{\alpha}, w \rangle + t^2\alpha\|w\|^2\end{aligned}$$

- the computation of the derivative implies

$$\frac{d}{dt}Q_{\alpha}(f_{\alpha} + tw) = 2\langle Af_{\alpha} - m, Aw \rangle + 2t\|Aw\|^2 + 2\alpha\langle f_{\alpha}, w \rangle + 2\alpha t\|w\|^2$$

- evaluating in $t = 0$, $\langle Af_{\alpha} - m, Aw \rangle + \alpha\langle f_{\alpha}, w \rangle = 0 \quad \forall w \in \mathbb{R}^n$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned}Q_{\alpha}(f_{\alpha} + tw) &= \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2 \\&= \langle (Af_{\alpha} - m) + tAw, (Af_{\alpha} - m) + tAw \rangle + \alpha\langle f_{\alpha} + tw, f_{\alpha} + tw \rangle \\&= \|Af_{\alpha} - m\|^2 + 2t\langle Af_{\alpha} - m, Aw \rangle + t^2\|Aw\|^2 \\&\quad + \alpha\|f_{\alpha}\|^2 + 2t\alpha\langle f_{\alpha}, w \rangle + t^2\alpha\|w\|^2\end{aligned}$$

- the computation of the derivative implies

$$\frac{d}{dt}Q_{\alpha}(f_{\alpha} + tw) = 2\langle Af_{\alpha} - m, Aw \rangle + 2t\|Aw\|^2 + 2\alpha\langle f_{\alpha}, w \rangle + 2\alpha t\|w\|^2$$

- evaluating in $t = 0$, $\langle Af_{\alpha} - m, Aw \rangle + \alpha\langle f_{\alpha}, w \rangle = 0 \quad \forall w \in \mathbb{R}^n$
- using the transpose, $\langle A^T Af_{\alpha} - A^T m + \alpha f_{\alpha}, w \rangle = 0 \quad \forall w \in \mathbb{R}^n$

Normal equations - proof

- We rewrite the quadratic functional as follows:

$$\begin{aligned}Q_{\alpha}(f_{\alpha} + tw) &= \|A(f_{\alpha} + tw) - m\|^2 + \alpha\|f_{\alpha} + tw\|^2 \\&= \langle (Af_{\alpha} - m) + tAw, (Af_{\alpha} - m) + tAw \rangle + \alpha\langle f_{\alpha} + tw, f_{\alpha} + tw \rangle \\&= \|Af_{\alpha} - m\|^2 + 2t\langle Af_{\alpha} - m, Aw \rangle + t^2\|Aw\|^2 \\&\quad + \alpha\|f_{\alpha}\|^2 + 2t\alpha\langle f_{\alpha}, w \rangle + t^2\alpha\|w\|^2\end{aligned}$$

- the computation of the derivative implies

$$\frac{d}{dt}Q_{\alpha}(f_{\alpha} + tw) = 2\langle Af_{\alpha} - m, Aw \rangle + 2t\|Aw\|^2 + 2\alpha\langle f_{\alpha}, w \rangle + 2\alpha t\|w\|^2$$

- evaluating in $t = 0$, $\langle Af_{\alpha} - m, Aw \rangle + \alpha\langle f_{\alpha}, w \rangle = 0 \quad \forall w \in \mathbb{R}^n$
- using the transpose, $\langle A^T Af_{\alpha} - A^T m + \alpha f_{\alpha}, w \rangle = 0 \quad \forall w \in \mathbb{R}^n$
- in conclusion, $(A^T A + \alpha I)f_{\alpha} = A^T m$,
and the matrix $A^T A + \alpha I$ is surely invertible.

Stacked version

The following formulation is equivalent to the normal equations:

Stacked form

$T_\alpha m$ is the unique **least squares** solution of $\begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} f = \begin{bmatrix} m \\ 0 \end{bmatrix}$.

Stacked version

The following formulation is equivalent to the normal equations:

Stacked form

$T_\alpha m$ is the unique **least squares** solution of $\begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} f = \begin{bmatrix} m \\ 0 \end{bmatrix}$.

Proof

- Let $\tilde{A} = \begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} \in \mathbb{R}^{2n \times n}$ and $\tilde{m} = \begin{bmatrix} m \\ 0 \end{bmatrix} \in \mathbb{R}^{2n \times 1}$.

Stacked version

The following formulation is equivalent to the normal equations:

Stacked form

$T_\alpha m$ is the unique **least squares** solution of $\begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} f = \begin{bmatrix} m \\ 0 \end{bmatrix}$.

Proof

- Let $\tilde{A} = \begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} \in \mathbb{R}^{2n \times n}$ and $\tilde{m} = \begin{bmatrix} m \\ 0 \end{bmatrix} \in \mathbb{R}^{2n \times 1}$.
- Analogously to the previous proof, one can show that the least squares solution (i.e., $f = \arg \min \{\|\tilde{A}f - \tilde{m}\|^2\}$) is the solution of the normal equations: $\tilde{A}^T \tilde{A}f = \tilde{A}^T \tilde{m}$

Stacked version

The following formulation is equivalent to the normal equations:

Stacked form

$T_\alpha m$ is the unique **least squares** solution of $\begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} f = \begin{bmatrix} m \\ 0 \end{bmatrix}$.

Proof

- Let $\tilde{A} = \begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} \in \mathbb{R}^{2n \times n}$ and $\tilde{m} = \begin{bmatrix} m \\ 0 \end{bmatrix} \in \mathbb{R}^{2n \times 1}$.
- Analogously to the previous proof, one can show that the least squares solution (i.e., $f = \arg \min \{\|\tilde{A}f - \tilde{m}\|^2\}$) is the solution of the normal equations: $\tilde{A}^T \tilde{A}f = \tilde{A}^T \tilde{m}$
- we are not going to prove why such solution is unique

Stacked version

The following formulation is equivalent to the normal equations:

Stacked form

$T_\alpha m$ is the unique **least squares** solution of $\begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} f = \begin{bmatrix} m \\ 0 \end{bmatrix}$.

Proof

- Let $\tilde{A} = \begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} \in \mathbb{R}^{2n \times n}$ and $\tilde{m} = \begin{bmatrix} m \\ 0 \end{bmatrix} \in \mathbb{R}^{2n \times 1}$.
- Analogously to the previous proof, one can show that the least squares solution (i.e., $f = \arg \min \{\|\tilde{A}f - \tilde{m}\|^2\}$) is the solution of the normal equations: $\tilde{A}^T \tilde{A} f = \tilde{A}^T \tilde{m}$
- we are not going to prove why such solution is unique
- One easily notices that:

$$\tilde{A}^T \tilde{A} = \begin{bmatrix} A^T & \sqrt{\alpha}I \end{bmatrix} \begin{bmatrix} A \\ \sqrt{\alpha}I \end{bmatrix} = A^T A + \alpha I, \quad \tilde{A}^T \tilde{m} = \begin{bmatrix} A^T & \sqrt{\alpha}I \end{bmatrix} \begin{bmatrix} m \\ 0 \end{bmatrix} = A^T m.$$

Example 1

1. Set $n = 100$ and define the signal f and the point spread function PSF as in the previous lecture. Create the convolution matrix A .
2. Generate the noisy measurement $m_\delta = Af + \delta r$, being r a Gaussian random vector of n dimensions, $\|r\| = 1$. Consider $\delta = 10^{-2}$.
3. Compute the Tikhonov regularized solution associated to $\alpha = 0.1$ by employing the singular values, by the normal equations and by their stacked version.
4. Compare the results obtained by comparing the norm of the difference between the three version and by estimating the computational time required (use the commands `tic` and `toc`).
5. Replicate the same analysis for $n = 250, 500, 1000, 2000$.

Generalized Tikhonov regularization

Generalized Tikhonov

Remind: the formulation of Tikhonov involves a **data fidelity** term $\|Af - m\|$ and some **a priori knowledge** encoded by $\|f\|$. We assume to know that, among the possible solutions of $Af = m$, we are promoting the one of smallest norm.

Generalized Tikhonov

Remind: the formulation of Tikhonov involves a **data fidelity** term $\|Af - m\|$ and some **a priori knowledge** encoded by $\|f\|$. We assume to know that, among the possible solutions of $Af = m$, we are promoting the one of smallest norm.

We can generalize the argument to incorporate different a priori information:

- we are interested in the solution f **close to a known signal f_*** (it minimizes $\|f - f_*\|$);

Generalized Tikhonov

Remind: the formulation of Tikhonov involves a **data fidelity** term $\|Af - m\|$ and some **a priori knowledge** encoded by $\|f\|$. We assume to know that, among the possible solutions of $Af = m$, we are promoting the one of smallest norm.

We can generalize the argument to incorporate different a priori information:

- we are interested in the solution f **close to a known signal f_*** (it minimizes $\|f - f_*\|$);
- we want the **norm $\|Lf\|^2$ to be small**, being L an invertible matrix (it can represent, e.g., a differential operator)

Generalized Tikhonov

Remind: the formulation of Tikhonov involves a **data fidelity** term $\|Af - m\|$ and some **a priori knowledge** encoded by $\|f\|$. We assume to know that, among the possible solutions of $Af = m$, we are promoting the one of smallest norm.

We can generalize the argument to incorporate different a priori information:

- we are interested in the solution f **close to a known signal f_*** (it minimizes $\|f - f_*\|$);
- we want the **norm $\|Lf\|^2$ to be small**, being L an invertible matrix (it can represent, e.g., a differential operator)

Generalized Tikhonov regularization

$$T_\alpha m = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m\|^2 + \alpha \|L(f - f_*)\|^2 \}$$

Explicit formulas

It is easy to replicate the computation leading to the normal equations, obtaining

Generalized normal equations

The solution $T_\alpha m$ can be computed as

$$T_\alpha = (A^T A + \alpha L^T L)^{-1} (A^T m + \alpha L^T L f_*),$$

Explicit formulas

It is easy to replicate the computation leading to the normal equations, obtaining

Generalized normal equations

The solution $T_\alpha m$ can be computed as

$$T_\alpha = (A^T A + \alpha L^T L)^{-1} (A^T m + \alpha L^T L f_*),$$

or alternatively

Stacked form

$$T_\alpha m \text{ is the least squares solution of } \begin{bmatrix} A \\ \sqrt{\alpha} L \end{bmatrix} f = \begin{bmatrix} m \\ \sqrt{\alpha} L f_* \end{bmatrix}.$$

Exercise: try to prove both formulas

Implementation

Example 2

1. Set $n = 200$, let $n_1 = 2 \lfloor \frac{n}{15} \rfloor$, $n_4 = 8 \lfloor \frac{n}{15} \rfloor$, $n_5 = 10 \lfloor \frac{n}{15} \rfloor$, $n_6 = 14 \lfloor \frac{n}{15} \rfloor$ and define the signal f as

$$f = \begin{cases} 6 & \text{if } n_1 \leq n \leq n_4, \\ 4 - \cos\left(\frac{2\pi n}{n_6 - n_5}\right) & \text{if } n_5 \leq n \leq n_6, \\ 5 & \text{elsewhere.} \end{cases}$$

2. create PSF by triple auto-convolution and normalization of $[1, 1, 1]$.
3. Generate the noisy measurement $m_\delta = Af + \delta r$, being r a Gaussian random vector, $\|r\| = 1$. Consider $\delta = 10^{-2}$.
4. Compute the Tikhonov regularized solution with $\alpha = 0.1$.
5. Compute the generalized Tikhonov solution by choosing $L = I$ and $f_* = 5$.

Example 3

1. Create f as in Example 2, subtracting 5. Take PSF and m_δ as in Example 2.
2. Compute the Tikhonov regularized solution associated to $\alpha = 0.1$.
3. Compute the generalized Tikhonov solution by choosing $f_* = 0$ and by considering a matrix L obtained by discretization of the derivative operator by means of a finite difference scheme:

$$\frac{df}{dx}(x_i) \approx (Lf)_i = \frac{f(x_i) - f(x_{i-1})}{\Delta x} = \frac{f_i - f_{i-1}}{\Delta x}.$$

Hint: in our examples, $\Delta x = 1$. Then

$$(Lf)_i = f_i - f_{i-1} = \begin{bmatrix} -1 & 1 \end{bmatrix} \begin{bmatrix} f_{i-1} \\ f_i \end{bmatrix} = [0 \dots -1 \quad 1 \dots 0] f$$

Choice of the regularization parameter

If we knew the solution...

In applications, the noise level δ is fixed (and, approximatively, known).

Issue: how to choose the parameter α ?

If we knew the solution...

In applications, the noise level δ is fixed (and, approximatively, known).

Issue: how to choose the parameter α ?

- if α is too large, we lose fidelity with respect to the data
- if α is too small, we lose the benefits of regularization

If we knew the solution...

In applications, the noise level δ is fixed (and, approximatively, known).

Issue: how to choose the parameter α ?

- if α is too large, we lose fidelity with respect to the data
- if α is too small, we lose the benefits of regularization

For Tikhonov, there exists an **a priori** parameter choice rule $\alpha = \alpha(\delta)$ which holds $\forall m_\delta : \|m - m_\delta\| \leq \delta$. Such rule is sub-optimal in most of the cases. **A posteriori** rule: decide α according to m_δ .


If we knew the solution...

In applications, the noise level δ is fixed (and, approximatively, known).

Issue: how to choose the parameter α ?

- if α is too large, we lose fidelity with respect to the data
- if α is too small, we lose the benefits of regularization

For Tikhonov, there exists an **a priori** parameter choice rule $\alpha = \alpha(\delta)$ which holds $\forall m_\delta : \|m - m_\delta\| \leq \delta$. Such rule is sub-optimal in most of the cases. **A posteriori** rule: decide α according to m_δ .



Idea

Take α such that $\|T_\alpha m_\delta - f\|$ is minimized.


If we knew the solution...

In applications, the noise level δ is fixed (and, approximatively, known).

Issue: how to choose the parameter α ?

- if α is too large, we lose fidelity with respect to the data
- if α is too small, we lose the benefits of regularization

For Tikhonov, there exists an **a priori** parameter choice rule $\alpha = \alpha(\delta)$ which holds $\forall m_\delta : \|m - m_\delta\| \leq \delta$. Such rule is sub-optimal in most of the cases. **A posteriori** rule: decide α according to m_δ .



Idea

Take α such that $\|T_\alpha m_\delta - f\|$ is minimized.

Big problem: we need to know the exact solution!

Morozov's discrepancy principle

Consider the **residual**

$$r(\alpha) = \|Af_{\alpha,\delta} - m_\delta\|, \quad f_{\alpha,\delta} = T_\alpha m_\delta.$$

Morozov's discrepancy principle

Consider the **residual**

$$r(\alpha) = \|Af_{\alpha,\delta} - m_\delta\|, \quad f_{\alpha,\delta} = T_\alpha m_\delta.$$

Idea: we know that the measurement m_δ is reliable up to a certain precision δ (because $\|m_\delta - m\| \leq \delta$), so we are not likely to gain any meaningful information if $r(\alpha) < \delta \dots$

Morozov's discrepancy principle

Consider the **residual**

$$r(\alpha) = \|Af_{\alpha,\delta} - m_\delta\|, \quad f_{\alpha,\delta} = T_\alpha m_\delta.$$

Idea: we know that the measurement m_δ is reliable up to a certain precision δ (because $\|m_\delta - m\| \leq \delta$), so we are not likely to gain any meaningful information if $r(\alpha) < \delta$...

Morozov's discrepancy principle

Select α s.t. $r(\alpha) = \delta$.

Morozov's discrepancy principle

Consider the **residual**

$$r(\alpha) = \|Af_{\alpha,\delta} - m_\delta\|, \quad f_{\alpha,\delta} = T_\alpha m_\delta.$$

Idea: we know that the measurement m_δ is reliable up to a certain precision δ (because $\|m_\delta - m\| \leq \delta$), so we are not likely to gain any meaningful information if $r(\alpha) < \delta$...

Morozov's discrepancy principle

Select α s.t. $r(\alpha) = \delta$.

Is it always possible to solve
 $r(\alpha) = \delta$?

Morozov's discrepancy principle

Consider the **residual**

$$r(\alpha) = \|Af_{\alpha,\delta} - m_\delta\|, \quad f_{\alpha,\delta} = T_\alpha m_\delta.$$

Idea: we know that the measurement m_δ is reliable up to a certain precision δ (because $\|m_\delta - m\| \leq \delta$), so we are not likely to gain any meaningful information if $r(\alpha) < \delta$...

Morozov's discrepancy principle

Select α s.t. $r(\alpha) = \delta$.

Is it always possible to solve
 $r(\alpha) = \delta$?

Theory: $r(\alpha)$ is an increasing function of $\alpha \in (0, \infty)$.

Its codomain is $(\|Pm_\delta\|, \|m_\delta\|)$, where P is the orthogonal projection on $\text{range}(A)^\perp$.

Morozov's discrepancy principle

Consider the **residual**

$$r(\alpha) = \|Af_{\alpha,\delta} - m_\delta\|, \quad f_{\alpha,\delta} = T_\alpha m_\delta.$$

Idea: we know that the measurement m_δ is reliable up to a certain precision δ (because $\|m_\delta - m\| \leq \delta$), so we are not likely to gain any meaningful information if $r(\alpha) < \delta \dots$


Morozov's discrepancy principle

Select α s.t. $r(\alpha) = \delta$.

Is it always possible to solve $r(\alpha) = \delta$?

Theory: $r(\alpha)$ is an increasing function of $\alpha \in (0, \infty)$.

Its codomain is $(\|Pm_\delta\|, \|m_\delta\|)$, where P is the orthogonal projection on $\text{range}(A)^\perp$.



Morozov's discrepancy principle - in practice

Practical remarks:

- in application, δ is not always well known: we don't need high accuracy when solving $r(\alpha) = \delta$
- $f_{\alpha,\delta}$ is not extremely sensitive with respect to α
- alternative formulations of Morozov: $\|Af_{\alpha,\delta} - m\|^2 = \delta^2$ (which is equivalent), $r(\alpha) = \tau\delta$ (*rule of thumb*: $\tau = 1.2$).

Morozov's discrepancy principle - in practice

Practical remarks:

- in application, δ is not always well known: we don't need high accuracy when solving $r(\alpha) = \delta$
- $f_{\alpha,\delta}$ is not extremely sensitive with respect to α
- alternative formulations of Morozov: $\|Af_{\alpha,\delta} - m\|^2 = \delta^2$ (which is equivalent), $r(\alpha) = \tau\delta$ (*rule of thumb*: $\tau = 1.2$).

How to solve $r(\alpha) = \delta$?

1. generate several possible α (the command `logspace` is very useful), evaluate r in those points, minimize $|r(\alpha) - \delta|$;
2. iterative algorithms, e.g. (geometric) bisection, allow to compute very few values of $r(\alpha)$ to satisfy Morozov's principle up to a fixed tolerance.

A different idea

The parameter choice should detect the **perfect balance** between the two terms appearing in Tikhonov regularization:

$$f_{\alpha,\delta} = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m_\delta\|^2 + \alpha \|f\|^2 \}.$$

L-curve method

A different idea

The parameter choice should detect the **perfect balance** between the two terms appearing in Tikhonov regularization:

$$f_{\alpha,\delta} = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m_\delta\|^2 + \alpha \|f\|^2 \}.$$

What does *perfect balance* mean? We need to understand the tradeoff between data fidelity and a priori knowledge reliability.

L-curve method

A different idea

The parameter choice should detect the **perfect balance** between the two terms appearing in Tikhonov regularization:

$$f_{\alpha,\delta} = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m_\delta\|^2 + \alpha \|f\|^2 \}.$$

What does *perfect balance* mean? We need to understand the tradeoff between data fidelity and a priori knowledge reliability.

This is done by defining a **parametric curve** $(X(\alpha), Y(\alpha))$:

$$X(\alpha) = \log(\|Af_{\alpha,\delta} - m\|).$$

$$Y(\alpha) = \log(\|f_{\alpha,\delta}\|),$$

L-curve method

A different idea

The parameter choice should detect the **perfect balance** between the two terms appearing in Tikhonov regularization:


$$f_{\alpha,\delta} = \arg \min_{f \in \mathbb{R}^n} \{ \|Af - m_\delta\|^2 + \alpha \|f\|^2 \}.$$

What does *perfect balance* mean? We need to understand the tradeoff between data fidelity and a priori knowledge reliability.

This is done by defining a **parametric curve** $(X(\alpha), Y(\alpha))$:

$$X(\alpha) = \log(\|Af_{\alpha,\delta} - m\|).$$

$$Y(\alpha) = \log(\|f_{\alpha,\delta}\|),$$



If the curve $(X(\alpha), Y(\alpha))$ has a shape similar to the letter L , then its "vertex" represent a good balance: a bigger α would cause a dramatic increase of the residual with no benefit for regularity, a smaller one would do the opposite.

Warning: that doesn't happen in any inverse problem!

If the curve $(X(\alpha), Y(\alpha))$ has a shape similar to the letter L , then its "vertex" represent a good balance: a bigger α would cause a dramatic increase of the residual with no benefit for regularity, a smaller one would do the opposite.

Warning: that doesn't happen in any inverse problem!

How to select the vertex?

- select some values of α (e.g. using logspace), compute X, Y and select the vertex by inspection;
- select some values of α , compute X, Y and select the point which is the closest to the origin of the graphic;
- more involved rules based on the slope of the curve.

Example 4

1. Create f as in Example 2, subtracting 5. Take PSF and m_δ as in Example 2, but use $\delta = 10^{-3}$.
2. Create a sampling of values of 60 α s from 10^{-9} to 10^{-2} , equally spaced in a logarithmic scale.
3. By employing the knowledge of the true solution, find which α gives the smallest relative error.
4. Show the residual as a function of α and automatically pick α satisfying Morozov's discrepancy principle. Is it close to the real value giving the smallest error?
5. Compute $X(\alpha)$, $Y(\alpha)$ and plot the L curve. Automatically select the vertex of the curve and the associated α . Is it close to the real value giving the smallest error?