



# Inverse Problems 1: convolution and deconvolution

## Lesson 1: introduction to convolution

---

Luca Ratti

September 3, 2019

University of Helsinki

# Table of contents

---



1. Motivation
2. Convolution in 1D: mathematical description
3. Convolution in 1D: examples
4. Convolution in 2D: a brief excursion
5. About the course

# Motivation

---

# Examples of convolution

Convolution is a mathematical model describing physical phenomena like image blurring and averaging of signals



Defocus aberration

Credits: <https://en.wikipedia.org/wiki/Deconvolution>

# Examples of convolution



Convolution is a mathematical model describing physical phenomena like image blurring and averaging of signals



Motion blur

# Examples of convolution

Convolution is a mathematical model describing physical phenomena like image blurring and averaging of signals



Blurred audio signal

# Deconvolution

The task of deconvolution is to restore the original signal from the blurred one.



Original

# Deconvolution

The task of deconvolution is to restore the original signal from the blurred one.



Original



Blurred



# Deconvolution

The task of deconvolution is to restore the original signal from the blurred one.



Original



Blurred



Reconstructed

# What is convolution?


## Main idea

Convolution occurs when the value of a signal in one point is influenced by the values of the points close by

# What is convolution?

## Main idea

Convolution occurs when the value of a signal in one point is influenced by the values of the points close by




An example from optics

# What is convolution?

## Main idea

Convolution occurs when the value of a signal in one point is influenced by the values of the points close by




An example from optics

# What is convolution?

## Main idea

Convolution occurs when the value of a signal in one point is influenced by the values of the points close by



An example from optics

## Convolution in 1D: mathematical description

---


## A first example

Replace each element of a vector with the average of the one before, itself and the one after.

## A first example

Replace each element of a vector with the average of the one before, itself and the one after.

The idea is rather simple...






## A first example

Replace each element of a vector with the average of the one before, itself and the one after.

The idea is rather simple...




... how to express it  
in mathematical terms?

$$c_4 = ???$$

## A first example

Replace each element of a vector with the average of the one before, itself and the one after.

The idea is rather simple...




... how to express it  
in mathematical terms?

$$c_4 = \frac{1}{3}(f_3 + f_4 + f_5)$$

## A first example

Replace each element of a vector with the average of the one before, itself and the one after.

The idea is rather simple...




... how to express it  
in mathematical terms?

$$c_i = \frac{1}{3}(f_{i-1} + f_i + f_{i+1})$$

## A first example

Replace each element of a vector with the average of the one before, itself and the one after.

The idea is rather simple...



... how to express it  
in mathematical terms?

$$c_i = \frac{1}{3}f_{i-1} + \frac{1}{3}f_i + \frac{1}{3}f_{i+1}$$

# Point Spread Function

The effect of the proximal points is described by a vector defined as  
Point Spread Function:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

# Point Spread Function

The effect of the proximal points is described by a vector defined as  
Point Spread Function:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

# Point Spread Function

The effect of the proximal points is described by a vector defined as **Point Spread Function**:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

We assume  $p$  satisfy a normalization property:  $\sum_{\ell=-\nu}^{\nu} p_{\ell} = 1$ .

# Point Spread Function

The effect of the proximal points is described by a vector defined as **Point Spread Function**:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

We assume  $p$  satisfy a normalization property:  $\sum_{\ell=-\nu}^{\nu} p_{\ell} = 1$ .

Examples:

- no convolution effect:  $p = [1]$ ;



# Point Spread Function

The effect of the proximal points is described by a vector defined as **Point Spread Function**:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

We assume  $p$  satisfy a normalization property:  $\sum_{\ell=-\nu}^{\nu} p_{\ell} = 1$ .

Examples:

- no convolution effect:  $p = [1]$ ;
- average over 3 elements:  $p = [\frac{1}{3}, \frac{1}{3}, \frac{1}{3}]$ ;

# Point Spread Function

The effect of the proximal points is described by a vector defined as **Point Spread Function**:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

We assume  $p$  satisfy a normalization property:  $\sum_{\ell=-\nu}^{\nu} p_{\ell} = 1$ .

Examples:

- no convolution effect:  $p = [1]$ ;
- average over 3 elements:  $p = [\frac{1}{3}, \frac{1}{3}, \frac{1}{3}]$ ;
- average over 5 elements:  $p = [\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}]$ ;

# Point Spread Function

The effect of the proximal points is described by a vector defined as **Point Spread Function**:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

We assume  $p$  satisfy a normalization property:  $\sum_{\ell=-\nu}^{\nu} p_{\ell} = 1$ .

Examples:

- no convolution effect:  $p = [1]$ ;
- average over 3 elements:  $p = [\frac{1}{3}, \frac{1}{3}, \frac{1}{3}]$ ;
- average over 5 elements:  $p = [\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}]$ ;
- average over 5 elements, different weights:  $p = [\frac{1}{10}, \frac{2}{10}, \frac{4}{10}, \frac{2}{10}, \frac{1}{10}]$

# Point Spread Function

The effect of the proximal points is described by a vector defined as **Point Spread Function**:

$$p \in \mathbb{R}^m, \text{ with } m = 2\nu + 1$$

The vector is typically indexed as follows:

$$p = [p_{-\nu}, p_{-\nu+1}, \dots, p_{-1}, p_0, p_1, \dots, p_{\nu-1}, p_{\nu}]$$

We assume  $p$  satisfy a normalization property:  $\sum_{\ell=-\nu}^{\nu} p_{\ell} = 1$ .

Examples:

- no convolution effect:  $p = [1]$ ;
- average over 3 elements:  $p = [\frac{1}{3}, \frac{1}{3}, \frac{1}{3}]$ ;
- average over 5 elements:  $p = [\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}]$ ;
- average over 5 elements, different weights:  $p = [\frac{1}{10}, \frac{2}{10}, \frac{4}{10}, \frac{2}{10}, \frac{1}{10}]$
- average over 5 elements, asymmetric:  $p = [\frac{1}{10}, \frac{1}{10}, \frac{3}{10}, \frac{3}{10}, \frac{2}{10}]$

# Mathematical formulation

**1D-signal:** consider a vector  $f \in \mathbb{R}^n$ .

**Point Spread Function:** take  $p \in \mathbb{R}^m$ ,  $m = 2\nu + 1$ .

# Mathematical formulation

**1D-signal:** consider a vector  $f \in \mathbb{R}^n$ .

**Point Spread Function:** take  $p \in \mathbb{R}^m$ ,  $m = 2\nu + 1$ .

The **convolution** between  $f$  and  $p$  is a vector  $p * f \in \mathbb{R}^n$  such that:

## Convolution formula

$$\begin{aligned}(p * f)_j &= \sum_{\ell=-\nu}^{\nu} p_{\ell} f_{j-\ell} \\ &= p_{-\nu} f_{j+\nu} + \dots + p_0 f_j + \dots + p_{\nu} f_{j-\nu}\end{aligned}$$

# Mathematical formulation

**1D-signal:** consider a vector  $f \in \mathbb{R}^n$ .

**Point Spread Function:** take  $p \in \mathbb{R}^m$ ,  $m = 2\nu + 1$ .

The **convolution** between  $f$  and  $p$  is a vector  $p * f \in \mathbb{R}^n$  such that:

## Convolution formula

$$\begin{aligned}(p * f)_j &= \sum_{\ell=-\nu}^{\nu} p_{\ell} f_{j-\ell} \\ &= p_{-\nu} f_{j+\nu} + \dots + p_0 f_j + \dots + p_{\nu} f_{j-\nu}\end{aligned}$$


**Problem:** the formula requires the knowledge of the values  $f_{-\nu+1}, f_{-\nu+2}, \dots, f_0$  and also  $f_{n+1}, \dots, f_{n+\nu}$ . How to define them? There is not an unique convention (next lesson).

## Convolution in 1D: examples

---




## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1



Convolution with a symmetric filter (zero extension of  $f$ )


## Example 1



Convolution with a symmetric filter (zero extension of  $f$ )




## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 1




Convolution with a symmetric filter (zero extension of  $f$ )

## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2



Convolution with an asymmetric filter (periodic extension of  $f$ )


## Example 2



Convolution with an asymmetric filter (periodic extension of  $f$ )




## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2



Convolution with an asymmetric filter (periodic extension of  $f$ )

## Example 2




Convolution with an asymmetric filter (periodic extension of  $f$ )

## Convolution in 2D: a brief excursion

---


## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.




## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.






## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.





## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.





## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



## Example 2



The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2



The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.





2D convolution



## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2


The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.




2D convolution



## Example 2

The same idea holds for 2D convolution, as explained by the illustration: in this case the signal is a matrix and the **point spread function** is a smaller square matrix.



2D convolution

## About the course

---

# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)

# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)
- Algebraic detour: the cause of ill-posedness; Singular Value Decomposition and Truncated SVD (2-3)

# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)
- Algebraic detour: the cause of ill-posedness; Singular Value Decomposition and Truncated SVD (2-3)
- Regularization: Tikhonov (4)

# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)
- Algebraic detour: the cause of ill-posedness; Singular Value Decomposition and Truncated SVD (2-3)
- Regularization: Tikhonov (4)
- Regularization: sparsity and Total Variation (5)

# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)
- Algebraic detour: the cause of ill-posedness; Singular Value Decomposition and Truncated SVD (2-3)
- Regularization: Tikhonov (4)
- Regularization: sparsity and Total Variation (5)
- Compressed Sensing, a brief overview (6)

# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)
- Algebraic detour: the cause of ill-posedness; Singular Value Decomposition and Truncated SVD (2-3)
- Regularization: Tikhonov (4)
- Regularization: sparsity and Total Variation (5)
- Compressed Sensing, a brief overview (6)
- 2D convolution and deconvolution (6)



# Outline of the course

- Introduction to convolution 1D: matrix representation of the inverse problem, naïve inversion (1)
- Algebraic detour: the cause of ill-posedness; Singular Value Decomposition and Truncated SVD (2-3)
- Regularization: Tikhonov (4)
- Regularization: sparsity and Total Variation (5)
- Compressed Sensing, a brief overview (6)
- 2D convolution and deconvolution (6)
- Convolutional Neural Networks (7)

# Main aims of the course

In the end of the course, the teaching staff would like you to:

# Main aims of the course

In the end of the course, the teaching staff would like you to:

1. have a solid grasp of the mathematical concepts of **convolution** and **deconvolution**;

# Main aims of the course

In the end of the course, the teaching staff would like you to:

1. have a solid grasp of the mathematical concepts of **convolution** and **deconvolution**;
2. get a flavour of what **inverse problems** are, and of one of the most important mathematical theory to tackle them: **regularization**;

# Main aims of the course

In the end of the course, the teaching staff would like you to:

1. have a solid grasp of the mathematical concepts of **convolution** and **deconvolution**;
2. get a flavour of what **inverse problems** are, and of one of the most important mathematical theory to tackle them: **regularization**;
3. be able to **implement** what you have learnt (hands-on approach, homework program tailored to the course);

# Main aims of the course

In the end of the course, the teaching staff would like you to:

1. have a solid grasp of the mathematical concepts of **convolution** and **deconvolution**;
2. get a flavour of what **inverse problems** are, and of one of the most important mathematical theory to tackle them: **regularization**;
3. be able to **implement** what you have learnt (hands-on approach, homework program tailored to the course);
4. (optional, but recommended) actively explore the field of deconvolution, by a small project, and learn how to report scientific findings.

# Practical information

- Lectures: Tue 12:15 - 14, Wed 14:15 - 16;
- Matlab exercises: online (MOOC) + exercise session: Thu 12:15-14;
- completion: home exam
- project: (optional) related to the topics of Inverse Problems 1 or Inverse Problems 2;
- material and additional information: **https://courses.helsinki.fi/en/mast31401/130419888;**
- contact: heli.virtanen@helsinki.fi