

# Introduction to graph theory

**Sylwia Cichacz**

Akademia Górniczo-Hutnicza w Krakowie

September 1, 2021, Kraków

## Definition:

A **graph**  $G = (V, E)$  consists of two finite sets  $V$  and  $E$ . The elements of  $V$  are called the **vertices** and the elements of  $E$  the **edges** of  $G$ .

## Definition:

A **graph**  $G = (V, E)$  consists of two finite sets  $V$  and  $E$ . The elements of  $V$  are called the **vertices** and the elements of  $E$  the **edges** of  $G$ .

Each edge is a two-element subset of  $V$ . For notation convenience, instead of representing the edge  $\{x, y\}$  we will denote it by  $xy$ .

## Definition:

A **graph**  $G = (V, E)$  consists of two finite sets  $V$  and  $E$ . The elements of  $V$  are called the **vertices** and the elements of  $E$  the **edges** of  $G$ .

$|V|$  – **order of graph** ( $|G|$ ),

## Definition:

A **graph**  $G = (V, E)$  consists of two finite sets  $V$  and  $E$ . The elements of  $V$  are called the **vertices** and the elements of  $E$  the **edges** of  $G$ .

$|V|$  – **order of graph** ( $|G|$ ),


$|E|$  – **size of graph** ( $\|G\|$ ).

## Definition:

A **graph**  $G = (V, E)$  consists of two finite sets  $V$  and  $E$ . The elements of  $V$  are called the **vertices** and the elements of  $E$  the **edges** of  $G$ .

$|V|$  – **order of graph** ( $|G|$ ),

$|E|$  – **size of graph** ( $\|G\|$ ).



- A **directed graph** or a **digraph** – by replacing the set  $E$  with a set of ordered pairs of vertices  $((x, y) \neq (y, x))$

- A **directed graph** or a **digraph** – by replacing the set  $E$  with a set of ordered pairs of vertices  $((x, y) \neq (y, x))$
- A **mixed graph** – if  $E$  contains both directed and undirected edges.



- A **directed graph** or a **digraph** – by replacing the set  $E$  with a set of ordered pairs of vertices  $((x, y) \neq (y, x))$
- A **mixed graph** – if  $E$  contains both directed and undirected edges.
- A **multigraph** – if  $E$  is a multiset.

- A **directed graph** or a **digraph** – by replacing the set  $E$  with a set of ordered pairs of vertices  $((x, y) \neq (y, x))$
- A **mixed graph** – if  $E$  contains both directed and undirected edges.
- A **multigraph** – if  $E$  is a multiset.
- A **pseudograph** – if  $E$  has edges that connect a vertex to itself (a **loop**).

- A **directed graph** or a **digraph** – by replacing the set  $E$  with a set of ordered pairs of vertices  $((x, y) \neq (y, x))$
- A **mixed graph** – if  $E$  contains both directed and undirected edges.
- A **multigraph** – if  $E$  is a multiset.
- A **pseudograph** – if  $E$  has edges that connect a vertex to itself (a **loop**).
- A **simple graph** a finite graph without loops or multiple edges.

**Definition:**

Let  $G = (V, E)$  be a graph

## Definition:

Let  $G = (V, E)$  be a graph

$v, w \in V(G)$  – **adjacent**  $\Leftrightarrow \exists e = vw \in E(G)$ , in this case we also say that  $x$  is a **neighbor** of  $y$ .

## Definition:

Let  $G = (V, E)$  be a graph

$v, w \in V(G)$  – **adjacent**  $\Leftrightarrow \exists e = vw \in E(G)$ , in this case we also say that  $x$  is a **neighbor** of  $y$ .

$v, w \in V(G)$  – **incident** with  $e = vw \in E(G)$ . We also say that  $u$  and  $v$  are the **endpoints** of  $e$ .

## Definition:

Let  $G = (V, E)$  be a graph

$v, w \in V(G)$  – **adjacent**  $\Leftrightarrow \exists e = vw \in E(G)$ , in this case we also say that  $x$  is a **neighbor** of  $y$ .

$v, w \in V(G)$  – **incident** with  $e = vw \in E(G)$ . We also say that  $u$  and  $v$  are the **endpoints** of  $e$ .

$e_1, e_2 \in E(G)$  – **incident**  $\Leftrightarrow e_1 \cap e_2 \neq \emptyset$

## Definition:

Let  $G = (V, E)$  be a graph

$v, w \in V(G)$  – **adjacent**  $\Leftrightarrow \exists e = vw \in E(G)$ , in this case we also say that  $x$  is a **neighbor** of  $y$ .

$v, w \in V(G)$  – **incident** with  $e = vw \in E(G)$ . We also say that  $u$  and  $v$  are the **endpoints** of  $e$ .

$e_1, e_2 \in E(G)$  – **incident**  $\Leftrightarrow e_1 \cap e_2 \neq \emptyset$  ( $\exists v \in V(G) : v \in e_1$  and  $v \in e_2$ ).



**Definition:**

Let  $G = (V, E)$  be a graph

## Definition:

Let  $G = (V, E)$  be a graph

The **neighborhood** of a vertex  $v \in V(G)$ , denoted by  $N(v)$  is a set of all vertices adjacent to  $v$ , i.e:

## Definition:

Let  $G = (V, E)$  be a graph

The **neighborhood** of a vertex  $v \in V(G)$ , denoted by  $N(v)$  is a set of all vertices adjacent to  $v$ , i.e:

$$N(v) = \{w \in V(G) : vw \in E(G)\}.$$

## Definition:

Let  $G = (V, E)$  be a graph

The **neighborhood** of a vertex  $v \in V(G)$ , denoted by  $N(v)$  is a set of all vertices adjacent to  $v$ , i.e:

$$N_G(v) = N(v) = \{w \in V(G) : vw \in E(G)\}.$$

**Definition:**

Let  $G = (V, E)$  be a graph

**Definition:**

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ .

**Definition:**

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)  
 $\deg(v) = |N(v)|$ .

## Definition:

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)

$$\deg(v) = |N(v)|.$$

If  $d(v) = 0$ , then  $v$  – **isolated vertex**.



## Definition:

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)

$$\deg(v) = |N(v)|.$$

If  $d(v) = 0$ , then  $v$  – **isolated vertex**.

The **maximum degree** of a graph  $G$ :

## Definition:

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)

$$\deg(v) = |N(v)|.$$

If  $d(v) = 0$ , then  $v$  – **isolated vertex**.

The **maximum degree** of a graph  $G$ :

$$\Delta(G) = \max\{d(v) : v \in V\},$$

## Definition:

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)

$$\deg(v) = |N(v)|.$$

If  $d(v) = 0$ , then  $v$  – **isolated vertex**.

The **maximum degree** of a graph  $G$ :

$$\Delta(G) = \max\{d(v) : v \in V\},$$

The **minimum degree** of a graph  $G$ :

## Definition:

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)

$$\deg(v) = |N(v)|.$$

If  $d(v) = 0$ , then  $v$  – **isolated vertex**.

The **maximum degree** of a graph  $G$ :

$$\Delta(G) = \max\{d(v) : v \in V\},$$

The **minimum degree** of a graph  $G$ :

$$\delta(G) = \min\{d(v) : v \in V\},$$

## Definition:

Let  $G = (V, E)$  be a graph

The **degree** of a vertex  $v \in V(G)$  ( $\deg(v)$ ,  $d_G(v)$ ) is the number of edges incident with  $v$ . Alternatively (in a simple graph)

$$\deg(v) = |N(v)|.$$

If  $d(v) = 0$ , then  $v$  – **isolated vertex**.

The **maximum degree** of a graph  $G$ :


$$\Delta(G) = \max\{d(v) : v \in V\},$$

The **minimum degree** of a graph  $G$ :


$$\delta(G) = \min\{d(v) : v \in V\},$$

$G$  –  **$r$ -regular**, if  $\forall v \in V(G) \ d(v) = r$ .

# 3-regular graph



# 3-regular graph



Petersen graph

## Theorem: handshake lemma

The degree sum formula states that for a given graph  $G = (V, E)$


$$\sum_{v \in V(G)} \deg(v) = 2|E(G)|.$$



## Theorem: handshake lemma

The degree sum formula states that for a given graph  $G = (V, E)$

$$\sum_{v \in V(G)} \deg(v) = 2|E(G)|.$$



*From Newsweek.*

## Corollary:

In any graph the number of vertices having the odd degree is even.

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:**

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

$V = V_1 \cup V_2$ , where

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

$V = V_1 \cup V_2$ , where

$V_1$  – the set of vertices of odd degree

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

$V = V_1 \cup V_2$ , where

$V_1$  – the set of vertices of odd degree

$V_2$  – the set of vertices of even degree

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

$V = V_1 \cup V_2$ , where

$V_1$  – the set of vertices of odd degree

$V_2$  – the set of vertices of even degree

Thus



## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

$V = V_1 \cup V_2$ , where

$V_1$  – the set of vertices of odd degree

$V_2$  – the set of vertices of even degree

Thus

$$2|E(G)| = \sum_{v \in V_1} \deg(v) + \sum_{u \in V_2} \deg(u)$$

## Corollary:

In any graph the number of vertices having the odd degree is even.

**Proof:** Let  $G = (V, E)$

$V = V_1 \cup V_2$ , where

$V_1$  – the set of vertices of odd degree

$V_2$  – the set of vertices of even degree

Thus

$$2|E(G)| = \sum_{v \in V_1} \deg(v) + \sum_{u \in V_2} \deg(u) \Rightarrow \sum_{v \in V_1} \deg(v) =$$

$$2|E(G)| - \underbrace{\sum_{u \in V_2} \deg(u)}_{\text{even number}}.$$

$$\underbrace{\hspace{10em}}_{\text{even number}}$$

**Definition:**

A **degree sequence** of a simple graph  $G$  is the set of degrees of all vertices in  $V$  written in non-increasing order.

**Definition:**

A **degree sequence** of a simple graph  $G$  is the set of degrees of all vertices in  $V$  written in non-increasing order.

It is clear that each simple graph has exactly one degree sequence, but that the converse need not hold.

# A degree sequence

## Definition:

A **degree sequence** of a simple graph  $G$  is the set of degrees of all vertices in  $V$  written in non-increasing order.

## Definition:

A **graphic sequence** is a sequence of numbers which can be the degree sequence of some graph.

# A degree sequence

## Definition:

A **degree sequence** of a simple graph  $G$  is the set of degrees of all vertices in  $V$  written in non-increasing order.

## Definition:

A **graphic sequence** is a sequence of numbers which can be the degree sequence of some graph.

## Example:

Is  $(4, 1, 1)$  graphic?

# A degree sequence

## Definition:

A **degree sequence** of a simple graph  $G$  is the set of degrees of all vertices in  $V$  written in non-increasing order.

## Definition:

A **graphic sequence** is a sequence of numbers which can be the degree sequence of some graph.

## Example:

Is  $(4, 3, 2, 2, 1)$  graphic?

## Theorem: Erdős, Gallai, 1960

A sequence  $\{d_1, \dots, d_n\}$  is graphic iff the sum of the elements is even and the sequence obeys the property

$$\sum_{i=1}^r d_i \leq r(r-1) + \sum_{i=r+1}^n \min\{r, d_i\}$$

for each integer  $r \leq n-1$ .



## Theorem: Erdős, Gallai, 1960

A sequence  $\{d_1, \dots, d_n\}$  is graphic iff the sum of the elements is even and the sequence obeys the property

$$\sum_{i=1}^r d_i \leq r(r-1) + \sum_{i=r+1}^n \min\{r, d_i\}$$

for each integer  $r \leq n-1$ .

## Theorem: Erdős, Gallai, 1960

A sequence  $\{d_1, \dots, d_n\}$  is graphic iff the sum of the elements is even and the sequence obeys the property

$$\sum_{i=1}^r d_i \leq r(r-1) + \sum_{i=r+1}^n \min\{r, d_i\}$$

for each integer  $r \leq n-1$ .

## Theorem: Havel (1955) and Hakimi (1962)

A degree sequence with  $n \geq 3$  and  $d_1 \geq 1$  is graphical iff the sequence  $\{d_{2-1}, d_{3-1}, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_p\}$  is graphical.

## Theorem: Havel (1955) and Hakimi (1962)

A degree sequence with  $n \geq 3$  and  $d_1 \geq 1$  is graphical iff the sequence  $\{d_{2-1}, d_{3-1}, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_p\}$  is graphical.

## Theorem: Havel (1955) and Hakimi (1962)

A degree sequence with  $n \geq 3$  and  $d_1 \geq 1$  is graphical iff the sequence  $\{d_{2-1}, d_{3-1}, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_p\}$  is graphical.

In addition, they showed that if a degree sequence is graphic, then there exists a graph  $G$  such that the node of highest degree is adjacent to the  $\Delta(G)$  next highest degree vertices of  $G$ .

## Theorem: Havel (1955) and Hakimi (1962)

A degree sequence with  $n \geq 3$  and  $d_1 \geq 1$  is graphical iff the sequence  $\{d_{2-1}, d_{3-1}, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_p\}$  is graphical.

In addition, they showed that if a degree sequence is graphic, then there exists a graph  $G$  such that the node of highest degree is adjacent to the  $\Delta(G)$  next highest degree vertices of  $G$ .

## Example:

Check if the sequence  $(4, 3, 2, 2, 1)$  is graphical.

# A degree sequence

## Theorem: Havel (1955) and Hakimi (1962)

A degree sequence with  $n \geq 3$  and  $d_1 \geq 1$  is graphical iff the sequence  $\{d_{2-1}, d_{3-1}, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_p\}$  is graphical.


### Example:

$(4, 3, 2, 2, 1)$    ①   ②   ③   ④   ⑤   ⑥

## Definition:

A **walk** consists of an alternating sequence of vertices and edges consecutive elements of which are incident


$(v_0, e_1, v_1, e_2, v_2, \dots, v_{k-1}, e_k, v_k)$ , that begins and ends with a vertex.



## Definition:

A **walk** consists of an alternating sequence of vertices and edges consecutive elements of which are incident

$(v_0, e_1, v_1, e_2, v_2, \dots, v_{k-1}, e_k, v_k)$ , that begins and ends with a vertex. A **trail** is a walk without repeated edges.






## Definition:

A **walk** consists of an alternating sequence of vertices and edges consecutive elements of which are incident

$(v_0, e_1, v_1, e_2, v_2, \dots, v_{k-1}, e_k, v_k)$ , that begins and ends with a vertex. A **trail** is a walk without repeated edges. A **path** is a walk without repeated vertices.




## Definition:

A **walk** consists of an alternating sequence of vertices and edges consecutive elements of which are incident

$(v_0, e_1, v_1, e_2, v_2, \dots, v_{k-1}, e_k, v_k)$ , that begins and ends with a vertex. A **trail** is a walk without repeated edges. A **path** is a walk without repeated vertices.

If a walk (resp. trail, path) begins at  $x$  and ends at  $y$  then it is an  $x - y$  walk (resp.  $x - y$  trail, resp.  $x - y$  path).




## Definition:

A **walk** consists of an alternating sequence of vertices and edges consecutive elements of which are incident

$(v_0, e_1, v_1, e_2, v_2, \dots, v_{k-1}, e_k, v_k)$ , that begins and ends with a vertex. A **trail** is a walk without repeated edges. A **path** is a walk without repeated vertices.

If a walk (resp. trail, path) begins at  $x$  and ends at  $y$  then it is an  $x - y$  walk (resp.  $x - y$  trail, resp.  $x - y$  path). The **length of a walk** – the number of edges in the walk.




## Definition:

A **walk** consists of an alternating sequence of vertices and edges consecutive elements of which are incident

$(v_0, e_1, v_1, e_2, v_2, \dots, v_{k-1}, e_k, v_k)$ , that begins and ends with a vertex. A **trail** is a walk without repeated edges. A **path** is a walk without repeated vertices.


If a walk (resp. trail, path) begins at  $x$  and ends at  $y$  then it is an  $x - y$  walk (resp.  $x - y$  trail, resp.  $x - y$  path). The **length of a walk** – the number of edges in the walk.

$P_n$  – a path of length  $n$ .



## Definition:


A walk is **closed**, if  $v_0 = v_k$ , **open** for  $v_0 \neq v_k$ .



**Definition:**

A walk is **closed**, if  $v_0 = v_k$ , **open** for  $v_0 \neq v_k$ .

By  $C_n$  we denote a cycle with  $n$  edges ( $n$  vertices).





## Definition:

A graph  $G$  is **bipartite** if the vertices of  $G$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that no two vertices in the same subset are adjacent. It is denoted by  $G = (V_1, V_2; E)$ .

## Definition:

A graph  $G$  is **bipartite** if the vertices of  $G$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that no two vertices in the same subset are adjacent. It is denoted by  $G = (V_1, V_2; E)$ .





## Definition:

A graph  $G$  is **bipartite** if the vertices of  $G$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that no two vertices in the same subset are adjacent. It is denoted by  $G = (V_1, V_2; E)$ .

## Definition:

A **complete bipartite graph**  $G = (V_1, V_2; E)$  is a bipartite graph such that for any two vertices  $v_1 \in V_1$  and  $v_2 \in V_2$ ,  $v_1 v_2 \in E$ . The complete bipartite graph with **partitions** of size  $|V_1| = m$  and  $|V_2| = n$  is denoted by  $K_{m,n}$ .

## Definition:

A graph  $G$  is **bipartite** if the vertices of  $G$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that no two vertices in the same subset are adjacent. It is denoted by  $G = (V_1, V_2; E)$ .

## Definition:

A **complete bipartite graph**  $G = (V_1, V_2; E)$  is a bipartite graph such that for any two vertices  $v_1 \in V_1$  and  $v_2 \in V_2$ ,  $v_1 v_2 \in E$ . The complete bipartite graph with **partitions** of size  $|V_1| = m$  and  $|V_2| = n$  is denoted by  $K_{m,n}$ .

$K_{1,n}$  is a **star**.

## Definition:

A graph  $G$  is **bipartite** if the vertices of  $G$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that no two vertices in the same subset are adjacent. It is denoted by  $G = (V_1, V_2; E)$ .

## Definition:

A **complete bipartite graph**  $G = (V_1, V_2; E)$  is a bipartite graph such that for any two vertices  $v_1 \in V_1$  and  $v_2 \in V_2$ ,  $v_1 v_2 \in E$ . The complete bipartite graph with **partitions** of size  $|V_1| = m$  and  $|V_2| = n$  is denoted by  $K_{m,n}$ .

## Theorem:



A graph  $G$  is bipartite iff  $G$  has no cycle of odd size.

# More examples of graphs:

- 1 The **complete graph** is a graph of order  $n$  that is an  $(n - 1)$ -regular graph. It is denoted by  $K_n$ .


# More examples of graphs:

- 1 The **complete graph** is a graph of order  $n$  that is an  $(n - 1)$ -regular graph. It is denoted by  $K_n$ .
- 2 The **empty graph**  $E_n = (V, \emptyset)$ .  $E_n$  is 0-regular graph.




# More examples of graphs:

- 1 The **complete graph** is a graph of order  $n$  that is an  $(n - 1)$ -regular graph. It is denoted by  $K_n$ .
- 2 The **empty graph**  $E_n = (V, \emptyset)$ .  $E_n$  is 0-regular graph.



$E_6$




$K_4$


- 3 The **complement of a graph**  $G$  is the graph  $\bar{G}$  with vertex set  $V(\bar{G}) = V(G)$  and edge set  $E(\bar{G}) = \{xy : x, y \in V(G), xy \notin E(G)\}$ .

# More examples of graphs:

- 1 The **complete graph** is a graph of order  $n$  that is an  $(n - 1)$ -regular graph. It is denoted by  $K_n$ .
- 2 The **empty graph**  $E_n = (V, \emptyset)$ .  $E_n$  is 0-regular graph.



$E_6$



$K_4$

- 3 The **complement of a graph**  $G$  is the graph  $\bar{G}$  with vertex set  $V(\bar{G}) = V(G)$  and edge set  $E(\bar{G}) = \{xy : x, y \in V(G), xy \notin E(G)\}$ .  
 $\bar{K}_n = E_n$

# More examples of graphs:

- 1 The **complete graph** is a graph of order  $n$  that is an  $(n - 1)$ -regular graph. It is denoted by  $K_n$ .
- 2 The **empty graph**  $E_n = (V, \emptyset)$ .  $E_n$  is 0-regular graph.
- 3 The **complement of a graph**  $G$  is the graph  $\bar{G}$  with vertex set  $V(\bar{G}) = V(G)$  and edge set  $E(\bar{G}) = \{xy : x, y \in V(G), xy \notin E(G)\}$ .  
 $\bar{K}_n = E_n$



$\bar{K}_{2,3}$



# More examples of graphs:

- 1 The **complete graph** is a graph of order  $n$  that is an  $(n - 1)$ -regular graph. It is denoted by  $K_n$ .
- 2 The **empty graph**  $E_n = (V, \emptyset)$ .  $E_n$  is 0-regular graph.
- 3 The **complement of a graph**  $G$  is the graph  $\bar{G}$  with vertex set  $V(\bar{G}) = V(G)$  and edge set  $E(\bar{G}) = \{xy : x, y \in V(G), xy \notin E(G)\}$ .  
 $\bar{K}_n = E_n$



$\bar{K}_{2,3}$

**Note:**  $\bar{\bar{G}} = G$ .

## Definition:

The **adjacency matrix** of a graph  $G$  with the vertex set  $V = \{v_1, v_2, \dots, v_n\}$  is a binary square  $n \times n$  matrix  $A(G) = (a_{ij})_{n \times n}$  such that


$$a_{i,j} = \begin{cases} 1 & v_i v_j \in E(G) \\ 0 & v_i v_j \notin E(G). \end{cases}$$

# Matrices associated with graphs

## Definition:

The **adjacency matrix** of a graph  $G$  with the vertex set  $V = \{v_1, v_2, \dots, v_n\}$  is a binary square  $n \times n$  matrix  $A(G) = (a_{ij})_{n \times n}$  such that

$$a_{i,j} = \begin{cases} 1 & v_i v_j \in E(G) \\ 0 & v_i v_j \notin E(G). \end{cases}$$



## Definition:

The **adjacency matrix** of a graph  $G$  with the vertex set  $V = \{v_1, v_2, \dots, v_n\}$  is a binary square  $n \times n$  matrix  $A(G) = (a_{ij})_{n \times n}$  such that

$$a_{i,j} = \begin{cases} 1 & v_i v_j \in E(G) \\ 0 & v_i v_j \notin E(G). \end{cases}$$

**Observation:**

## Definition:

The **adjacency matrix** of a graph  $G$  with the vertex set  $V = \{v_1, v_2, \dots, v_n\}$  is a binary square  $n \times n$  matrix  $A(G) = (a_{ij})_{n \times n}$  such that

$$a_{i,j} = \begin{cases} 1 & v_i v_j \in E(G) \\ 0 & v_i v_j \notin E(G). \end{cases}$$

## Observation:

- For a simple graph the adjacency matrix is symmetric.

## Definition:

The **adjacency matrix** of a graph  $G$  with the vertex set  $V = \{v_1, v_2, \dots, v_n\}$  is a binary square  $n \times n$  matrix  $A(G) = (a_{ij})_{n \times n}$  such that

$$a_{i,j} = \begin{cases} 1 & v_i v_j \in E(G) \\ 0 & v_i v_j \notin E(G). \end{cases}$$

## Observation:

- For a simple graph the adjacency matrix is symmetric.
- $\sum_{j=1}^m a_{i,j} = \deg(v_i)$ .

## Definition:

For a simple graph  $G = (V, E)$  the **incidence matrix** a  $n \times m$  matrix  $B(G) = (b_{ij})_{n \times m}$ , where  $n$  and  $m$  are the numbers of vertices and edges respectively ( $|V| = n$  and  $|E| = m$ ), such that


$$b_{i,j} = \begin{cases} 1 & v_i \in e_j \\ 0 & v_i \notin e_j. \end{cases}$$

# Matrices associated with graphs

## Definition:

For a simple graph  $G = (V, E)$  the **incidence matrix** is a  $n \times m$  matrix  $B(G) = (b_{ij})_{n \times m}$ , where  $n$  and  $m$  are the numbers of vertices and edges respectively ( $|V| = n$  and  $|E| = m$ ), such that

$$b_{i,j} = \begin{cases} 1 & v_i \in e_j \\ 0 & v_i \notin e_j. \end{cases}$$





## Definition:

For a simple graph  $G = (V, E)$  the **incidence matrix** a  $n \times m$  matrix  $B(G) = (b_{ij})_{n \times m}$ , where  $n$  and  $m$  are the numbers of vertices and edges respectively ( $|V| = n$  and  $|E| = m$ ), such that

$$b_{i,j} = \begin{cases} 1 & v_i \in e_j \\ 0 & v_i \notin e_j. \end{cases}$$

## Observation:

## Definition:

For a simple graph  $G = (V, E)$  the **incidence matrix** a  $n \times m$  matrix  $B(G) = (b_{ij})_{n \times m}$ , where  $n$  and  $m$  are the numbers of vertices and edges respectively ( $|V| = n$  and  $|E| = m$ ), such that

$$b_{i,j} = \begin{cases} 1 & v_i \in e_j \\ 0 & v_i \notin e_j. \end{cases}$$

## Observation:

- In every column of the matrix  $B(G)$  there are exactly two 1.

## Definition:

For a simple graph  $G = (V, E)$  the **incidence matrix** a  $n \times m$  matrix  $B(G) = (b_{ij})_{n \times m}$ , where  $n$  and  $m$  are the numbers of vertices and edges respectively ( $|V| = n$  and  $|E| = m$ ), such that

$$b_{i,j} = \begin{cases} 1 & v_i \in e_j \\ 0 & v_i \notin e_j. \end{cases}$$

## Observation:


- In every column of the matrix  $B(G)$  there are exactly two 1.
- $\sum_{j=1}^m b_{i,j} = \deg(v_i)$ .

**Definition:**

A graph  $G' = (V', E')$  is a **subgraph** of another graph  $G = (V, E)$  iff  $V' \subset V$  and  $E' \subset E$ .

## Definition:

A graph  $G' = (V', E')$  is a **subgraph** of another graph  $G = (V, E)$  iff  $V' \subset V$  and  $E' \subset E$ .




Graph G

# Subgraphs

## Definition:

A graph  $G' = (V', E')$  is a **subgraph** of another graph  $G = (V, E)$  iff  $V' \subset V$  and  $E' \subset E$ .




Graph  $G$

Is  $G' = (V', E')$  such that  $V' = \{a, b, c\}$  and  $E' = \{e_5\}$  a subgraph of  $G$ ?

No

## Definition:

A graph  $G' = (V', E')$  is a **subgraph** of another graph  $G = (V, E)$  iff  $V' \subset V$  and  $E' \subset E$ .



Graph  $G$

Is  $G' = (V', E')$  such that  $V' = \{a, b, c\}$  and  $E' = \{e_1\}$  a subgraph of  $G$ ?

Yes


**Definition:**

Let  $G = (V, E)$  be any graph, and let  $S \subset V$  be any subset of vertices of  $G$ . Then the **induced subgraph**  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .



## Definition:


Let  $G = (V, E)$  be any graph, and let  $S \subset V$  be any subset of vertices of  $G$ . Then the **induced subgraph**  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .



Graph  $G$

## Definition:

Let  $G = (V, E)$  be any graph, and let  $S \subset V$  be any subset of vertices of  $G$ . Then the **induced subgraph**  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .




Graph  $G$

Is  $G' = (V', E')$  such that  $V' = \{a, b, c\}$  and  $E' = \{e_1\}$  an induced subgraph  $G$  (i.e.  $G' = G[V']$ )?

## Definition:

Let  $G = (V, E)$  be any graph, and let  $S \subset V$  be any subset of vertices of  $G$ . Then the **induced subgraph**  $G[S]$  is the graph whose vertex set is  $S$  and whose edge set consists of all of the edges in  $E$  that have both endpoints in  $S$ .



Graph  $G$


Is  $G' = (V', E')$  such that  $V' = \{a, b, c\}$  and  $E' = \{e_1, e_2, e_3\}$  an induced subgraph  $G$  (i.e.  $G' = G[V']$ )?

## Definition:

A graph is **connected** if any two vertices are joint by a path. If a graph not connected then is **disconnected**.

## Definition:


A graph is **connected** if any two vertices are joint by a path. If a graph not connected then is **disconnected**.



Graph  $G$

## Definition:

A graph is **connected** if any two vertices are joint by a path. If a graph not connected then is **disconnected**.




Graph  $G$

Is  $G$  connected?

YES

## Definition:

A graph is **connected** if any two vertices are joint by a path. If a graph not connected then is **disconnected**.



Graph G NO

Is  $G$  connected?


## Definition:

For a disconnected graph  $G$  every maximal (with respect to inclusion) connected subgraph is called a **connected component** of  $G$ .




## Definition:

For a disconnected graph  $G$  every maximal (with respect to inclusion) connected subgraph is called a **connected component** of  $G$ .



## Definition:


For a disconnected graph  $G$  every maximal (with respect to inclusion) connected subgraph is called a **connected component** of  $G$ .



Is  $G[V']$  such that  $V' = \{a, b, c\}$  a connected component?

## Definition:


For a disconnected graph  $G$  every maximal (with respect to inclusion) connected subgraph is called a **connected component** of  $G$ .



Is  $G' = (V', E')$  such that  $V' = \{a, b, c, d\}$  and  $E' = \{e_1, e_2, e_3\}$  a connected component?

## Definition:

For a disconnected graph  $G$  every maximal (with respect to inclusion) connected subgraph is called a **connected component** of  $G$ .




Is  $G[V']$  such that  $V' = \{a, b, c, d\}$  a connected component?

## Definition:

The **distance between two vertices  $x$  and  $y$** , denoted by  **$\text{dist}(x, y)$**  is the length of a shortest path joining them.


## Definition:

The **distance between two vertices  $x$  and  $y$** , denoted by  $\text{dist}(x, y)$  is the length of a shortest path joining them.



## Definition:


The **distance between two vertices  $x$  and  $y$** , denoted by  $\text{dist}(x, y)$  is the length of a shortest path joining them.




$$\text{dist}(b, d) = 3?$$

## Definition:

The **distance between two vertices  $x$  and  $y$** , denoted by  $\text{dist}(x, y)$  is the length of a shortest path joining them.



$\text{dist}(b, f) = ?$  



# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$

# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$



$$G_1 \cong G_2.$$

# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$

$$G_1 \cong G_2.$$





# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$

$$G_1 \cong G_2.$$





$$f(u) = a,$$

# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$

$$G_1 \cong G_2.$$





$$f(u) = a, f(v) = b,$$

# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$

$$G_1 \cong G_2.$$





$$f(u) = a, f(v) = b, f(w) = d,$$

# Isomorphism of graphs

Two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a bijection  $f: V_1 \rightarrow V_2$  that preserves the adjacency

$$uv \in E_1 \iff f(u)f(v) \in E_2.$$

$$G_1 \cong G_2.$$



$$f(u) = a, f(v) = b, f(w) = d, f(t) = c.$$