

## Lecture 10: 23 Aug, 2017

Lecturer: Swaprava Nath

Scribe(s): Asim Unmesh

**Disclaimer:** These notes aggregate content from several texts and have not been subjected to the usual scrutiny deserved by formal publications. If you find errors, please bring to the notice of the Instructor.

## 10.1 Outcome Equivalence of Behavioral and Mixed Strategy

In the context of *Extensive Form Games* (EFG) *Behavioral Strategy* for an EFG is determined by a probability distribution over moves for each node the player is going to play at. Thus the player is randomizing at each node rather than playing a deterministic choice at each node.

A Behavioral Strategy  $b_i$  and a Mixed Strategy  $\sigma_i$  are **Outcome Equivalent** if for all  $\sigma_{-i}$ , the probability distribution induced over the terminal vertices are the same for  $(b_i, \sigma_{-i})$  and  $(\sigma_i, \sigma_{-i})$ .

**Theorem 10.1 (Kuhn 1953)** *In Games with Perfect Recall every mixed strategy is Outcome Equivalent to behavioral strategies.*

## 10.2 Belief

We now define the set of all *Information Set* of player  $i$ .

$$I_i = I_i^1, I_i^2, \dots, I_i^{k(i)}$$

In an *Imperfect Information Extensive Form Game* (IIIEFG), the belief of player  $i$  is a map

$$\mu_i^j : I_i^j \rightarrow [0, 1]$$

such that,

$$\sum_{x \in I_i^j} \mu_i^j(x) = 1$$

## 10.3 Bayesian Belief of Player i

$\mu_i$  is *Bayesian* with respect to  $\sigma$ , if it is derived from mixed strategy profile  $\sigma$  using Bayes rule. That is:

$$\mu_i(x) = \frac{P_\sigma(x)}{\sum_{y \in I_i^j} P_\sigma(y)} \quad x \in I_i^j \quad \forall j = 1, \dots, k(i)$$

## 10.4 Sequential Rationality

A strategy  $\sigma_i$  of player  $i$  at an Information Set  $I_i^j$  is sequentially rational given  $\sigma_{-i}$  and beliefs  $\mu_i$  if  $\forall \sigma'_i$

$$\sum_{x \in I_i^j} \mu_i(x) U_i(\sigma_i, \sigma_{-i} | x) \geq \sum_{x \in I_i^j} \mu_i(x) U_i(\sigma'_i, \sigma_{-i} | x)$$

## 10.5 Football Cricket Game example for Sequential Rationality




Figure 10.1: Football-Cricket Game for 2 players

In the above figure you can see Football Cricket game played between two players 1 and 2.

$$\mu_2(a) = p = 0.5$$

$$\mu_2(b) = 1 - p = 0.5$$

$$\begin{aligned}
& \sum_{x \in I_2^1} \mu_2(x) U_2(\sigma_1, \sigma_2 | x) \\
&= 0.5[q \cdot 1 + (1-q) \cdot 0] + 0.5[2 \times (1-q)] \\
&= 0.5[2 - q]
\end{aligned}$$

Thus, to maximise his utility given his belief about the moves of player 1, which is  $(0.5, 0.5)$ , it will be sequentially rational for player 2 to keep  $q = 0$ , i.e., play football with zero probability.

So,  $\sigma = ((0.5, 0.5), (0, 1))$

## 10.6 Perfect Bayesian Equilibrium(PBE)

An assessment  $(\sigma, \mu)$  is a PBE if for every player  $i$

1.  $\mu_i$  is Bayesian with respect to  $\sigma$ .
2.  $\sigma_i$  is Sequentially Rational given  $\sigma_{-i}$  and  $\mu_i$  at every information set of  $i$ .

**Theorem 10.2** Every Perfect Bayesian Equilibrium(PBE) is a Mixed Strategy Nash Equilibrium(MSNE).