Roll No.	160554	Dept.	CS	Section	A-4	

Instructions:

1. This question paper contains a total of **1** page (**1** side of paper).
2. Write your name, roll number, department, and section on every side of every sheet of this booklet
3. Write final answers neatly with a blue/black pen in the given boxes.
4. **Answers written outside the box will NOT be graded.**

Total **10 Marks**

Q. 1: Write the output of the following program in the appropriate box and answer the question. (2+2+2 = 6 Marks)

```
#include<stdio.h>
void func1(int *arr){
  for(int i=1; i<=4;i++)
    (i-1)[arr] = (i%4);
}
void func2 (int a[1] , int arr){
  a[0] = arr + 2;
}
int main(){
  int arr[5] = {1,1,1,1,1};
  func1(arr+1);
  for(int i=0; i<5; i++)
        printf("%d ", i[arr] );
  printf("\n");
  func2(arr, *arr);
  for(int i=0; i<5; i++)
        printf("%d ", i[arr] );
  return 0;
}
```

Output
1 2 3 0 1
3 2 3 0 1

Will the answer change if we write func2 as:

```
void func2 (int a[1] , int arr){
 int temp = arr + 2;
 a = &temp;
}
```

Explain your answer

No, answer will not change Because Both Statement's meaning is same.

Q. 2: Write the output of the following sequence of instructions **long arr2[60]; int arr1[50], a1=(&arr2[40]), a2=(&arr2[30]), b1=(&arr1[40]), b2=(&arr1[30]); printf("%d %d", a1-a2 , b1-b2);** Explain your answer. (2+2 Marks)

10 10

Because $a_1 = 40$, & $a_2 = 30$, Because & shows the position of that integer variable in an array

Same as, $b_1 = 40$, $b_2 = 30$