

# Projective, Injective, and Flat Modules

Swayam Chube

December 11, 2024

## §1 PROJECTIVE MODULES

**DEFINITION 1.1.** An  $A$ -module  $M$  is said to be *projective* if the functor  $\text{Hom}_A(M, -) : \mathcal{M}\text{od}_A \rightarrow \mathcal{M}\text{od}_A$  is exact.

### §§ Kaplansky's Theorem

**THEOREM 1.2.** Let  $(A, \mathfrak{m}, k)$  be a local ring. If  $M$  is a projective  $A$ -module, then  $M$  is free.

## §2 FLAT MODULES

**DEFINITION 2.1.** An  $A$ -module  $M$  is said to be *flat* if the functor  $- \otimes_A M : \mathcal{M}\text{od}_A \rightarrow \mathcal{M}\text{od}_A$  is exact.

**DEFINITION 2.2.** Let  $M$  be an  $A$ -module and  $\sum_{i=1}^n f_i x_i = 0$  be a relation in  $M$  for  $f_i \in A$  and  $x_i \in M$ . We say that the relation is *trivial* if there exists an integer  $m \geq 0$ , elements  $y_j \in M$  for  $1 \leq j \leq m$  and  $a_{ij} \in A$  for  $1 \leq i \leq n$  and  $1 \leq j \leq m$  such that

$$x_i = \sum_{j=1}^m a_{ij} y_j \quad \forall 1 \leq i \leq n \quad \text{and} \quad 0 = \sum_{i=1}^n a_{ij} f_i \quad \forall 1 \leq j \leq m.$$

**LEMMA 2.3 (EQUATIONAL CRITERION OF FLATNESS).** An  $A$ -module  $M$  is flat if and only if every relation in  $M$  is trivial.

*Proof.* Suppose  $M$  is flat and  $\sum_{i=1}^n f_i x_i = 0$  is a relation in  $M$ . Let  $\mathfrak{a} = (f_1, \dots, f_n) \subseteq A$  and consider the  $A$ -linear surjection  $A^n = \bigoplus_{i=1}^n A e_i \rightarrow I$  given by  $e_i \mapsto f_i$  whose kernel is  $K \subseteq A^n$ . That is,  $0 \rightarrow K \rightarrow A^n \rightarrow \mathfrak{a} \rightarrow 0$ . Since  $M$  is flat, tensoring with  $M$  preserves exactness and we have an exact sequence

$$0 \longrightarrow K \otimes_A M \longrightarrow A^n \otimes_A M \longrightarrow \mathfrak{a} \otimes_A M \longrightarrow 0.$$

Note that the natural map  $\mathfrak{a} \otimes_A M \rightarrow R \otimes_A M$  is injective due to the flatness of  $M$ . Consequently,  $\sum_{i=1}^n f_i \otimes x_i$  maps to 0 in  $R \otimes_A M$  and hence, must be zero in  $\mathfrak{a} \otimes_A M$ . The

exactness of the above sequence furnishes an element  $\sum_{j=1}^m k_j \otimes y_j \in K \otimes_A M$  that maps to 0 in  $A^n \otimes_A M$ .

Each  $k_j$  can be written in the form

$$\sum_{i=1}^n a_{ij} e_i \quad \forall 1 \leq j \leq m,$$

and hence, the image of  $\sum_{j=1}^m k_j \otimes y_j$  in  $A^n \otimes_A M$  is

$$\sum_{j=1}^m \sum_{i=1}^n a_{ij} e_i \otimes y_j = \sum_{i=1}^n e_i \otimes \left( \sum_{j=1}^m a_{ij} y_j \right) = 0,$$

and the conclusion follows.

Conversely, suppose every relation in  $M$  is trivial and let  $\mathfrak{a}$  be a finitely generated ideal of  $A$ . It suffices to show that  $\text{Tor}_1^A(A/\mathfrak{a}, M) = 0$ , which is equivalent (from the Tor long exact sequence) to showing that the map  $\mathfrak{a} \otimes_A M \rightarrow A \otimes_A M$  is injective.

Suppose  $\sum_{i=1}^n f_i \otimes x_i \in \mathfrak{a} \otimes_A M$  maps to 0 in  $A \otimes_A M$ . Then,  $\sum_{i=1}^n f_i x_i = 0$  in  $M$ , consequently, there is an  $m \geq 0$ ,  $y_j \in M$ ,  $a_{ij} \in M$  for  $1 \leq i \leq n$  and  $1 \leq j \leq m$  such that

$$x_i = \sum_{j=1}^m a_{ij} y_j \quad \forall 1 \leq i \leq n \quad \text{and} \quad 0 = \sum_{i=1}^n a_{ij} f_i \quad \forall 1 \leq j \leq m.$$

Consequently, in  $\mathfrak{a} \otimes_A M$ ,

$$\sum_{i=1}^n f_i \otimes x_i = \sum_{i=1}^n f_i \otimes \left( \sum_{j=1}^m a_{ij} y_j \right) = \left( \sum_{i=1}^n a_{ij} f_i \right) \otimes y_j = 0.$$

This proves injectivity, thereby completing the proof. ■

**LEMMA 2.4.** Let  $(A, \mathfrak{m}, k)$  be a local ring and  $M$  a flat  $A$ -module. If  $x_1, \dots, x_n \in M$  are such that their images  $\bar{x}_1, \dots, \bar{x}_n \in M/\mathfrak{m}M$  are linearly independent over  $k$ , then  $x_1, \dots, x_n$  are linearly independent over  $A$ .

*Proof.* We prove this statement by induction on  $n$ . If  $n = 1$ , then  $a \in A$  is such that  $ax_1 = 0$  and  $\bar{x}_1 \neq 0$ . From Lemma 2.3, there are  $b_1, \dots, b_m \in A$  and  $y_1, \dots, y_m \in M$  such that

$$x_1 = \sum_{j=1}^m b_j y_j \quad \text{and} \quad ab_j = 0 \quad \forall 1 \leq j \leq m.$$

Since  $x_1 \notin \mathfrak{m}M$ , it follows that at least one of the  $b_j$ 's must be a unit, whence  $a = 0$ .

Now, suppose  $n > 1$  and there is a relation  $\sum_{i=1}^n a_i x_i = 0$  in  $M$ . From Lemma 2.3, there is an  $m \geq 0$ ,  $y_j \in M$ , and  $b_{ij} \in A$  for  $1 \leq i \leq n$  and  $1 \leq j \leq m$  such that

$$x_i = \sum_{j=1}^m b_{ij} y_j \quad \forall 1 \leq i \leq n \quad \text{and} \quad 0 = \sum_{i=1}^n b_{ij} a_i \quad \forall 1 \leq j \leq m.$$

Since  $x_n \notin \mathfrak{m}M$ , at least one of the  $b_{nj}$ 's must be a unit, whence we can write

$$a_n = \sum_{i=1}^{n-1} c_i a_i,$$

for some  $c_i \in A$  for  $1 \leq i \leq n-1$ . Therefore, we have

$$0 = \sum_{i=1}^n a_i x_i = \sum_{i=1}^{n-1} a_i (x_i + c_i x_n).$$

Since  $\bar{x}_1, \dots, \bar{x}_{n-1}$  are  $k$ -linearly independent in  $M/\mathfrak{m}M$ , we see that  $\bar{x}_1 + \bar{c}_1 \bar{x}_n, \dots, \bar{x}_{n-1} + \bar{c}_{n-1} \bar{x}_n$  must also be  $k$ -linearly independent. Due to the induction hypothesis,  $a_1 = \dots = a_{n-1} = 0$  and hence,  $a_n = 0$ . This completes the proof. ■

**THEOREM 2.5.** Let  $(A, \mathfrak{m}, k)$  be a local ring. If  $M$  is a finitely generated flat  $A$ -module, then  $M$  is free.

*Proof.* Let  $x_1, \dots, x_n \in M$  be a minimal generating set, that is,  $\bar{x}_1, \dots, \bar{x}_n$  are  $k$ -linearly independent in  $M/\mathfrak{m}M$ . Due to the preceding lemma,  $x_1, \dots, x_n$  are linearly independent over  $A$ , and hence,  $M$  is a free  $A$ -module. ■

## §§ Cartier's Theorem

**THEOREM 2.6 (CARTIER).** Let  $M$  be a finitely generated module over an integral domain  $A$ . If for every  $\mathfrak{m} \in \text{MaxSpec}(A)$ ,  $M_{\mathfrak{m}}$  is free as an  $A_{\mathfrak{m}}$ -module, then  $M$  is a projective  $A$ -module.

*Proof.* First show that  $M$  is a torsion-free  $A$ -module. Suppose  $am = 0$  for some  $0 \neq a \in A$  and  $m \in M$ . Let  $\mathfrak{a}$  be the annihilator of  $m$  in  $A$  and  $\mathfrak{m}$  a maximal ideal containing  $\mathfrak{a}$ . Note that  $\frac{a}{1} \frac{m}{1} = 0$  in  $M_{\mathfrak{m}}$ , which is free over  $A_{\mathfrak{m}}$ , an integral domain, whence, is torsion free. That is,  $\frac{m}{1} = 0$ , whence, there is some  $s \in A \setminus \mathfrak{m}$  such that  $sm = 0$ , which is absurd, since  $\mathfrak{a} \subseteq \mathfrak{m}$ . This shows that  $M$  is torsion-free.

Now, choose a set of generators  $\{m_i : 1 \leq i \leq n\}$  for  $M$  over  $A$ . Let  $\mathcal{P}$  be the collection of  $A$ -endomorphisms of  $M$  which are of the form

$$m \mapsto \sum_{i=1}^n f_i(m) m_i,$$

where  $f_1, \dots, f_n : M \rightarrow A$  are  $A$ -module homomorphisms. Note that  $\mathcal{P}$  is an  $A$ -submodule of  $\text{End}_A(M)$ . We shall show that  $\text{id}_M \in \mathcal{P}$ .

Let  $\mathfrak{m}$  be a maximal ideal of  $A$ . We know that  $M_{\mathfrak{m}}$  is free as an  $A_{\mathfrak{m}}$ -module and hence, there are  $A_{\mathfrak{m}}$ -module homomorphisms  $f_i : M_{\mathfrak{m}} \rightarrow A_{\mathfrak{m}}$  such that

$$m' = \sum_{i=1}^n f_i'(m') \frac{m_i}{1} \quad \forall m' \in M_{\mathfrak{m}}.$$

To see that this is possible, first consider an  $A_{\mathfrak{m}}$ -basis  $\{e_i : 1 \leq i \leq N\}$  for  $M_{\mathfrak{m}}$ . We can write

$$e_i = \sum_{j=1}^n a_{ij} \frac{m_j}{1} \quad \forall 1 \leq i \leq N.$$

Further, there are  $A_{\mathfrak{m}}$ -linear maps  $f_i : M_{\mathfrak{m}} \rightarrow A_{\mathfrak{m}}$  such that

$$m' = \sum_{j=1}^N f_j(m') e_j.$$

Set

$$f'_j(m') = \sum_{i=1}^N a_{ij} f_i(m') \quad \forall m' \in M_{\mathfrak{m}}.$$

Then,

$$\sum_{j=1}^n f'_j(m') \frac{m_j}{1} = \sum_{i=1}^N \sum_{j=1}^n a_{ij} f_i(m') \frac{m_j}{1} = \sum_{i=1}^N f_i(m') e_i = m'.$$

Coming back, since  $M$  is torsion-free, the canonical map  $M \rightarrow M_{\mathfrak{m}}$  is an injective map of  $A$ -modules. Further, we can find an  $s \in A \setminus \mathfrak{m}$  such that  $sf'_i \left( \frac{m_j}{1} \right) \in A$  for  $1 \leq i, j \leq n$ .

Note that  $m' \mapsto sf'_i(m')$  is  $A_{\mathfrak{m}}$ -linear as a map  $M_{\mathfrak{m}} \rightarrow A_{\mathfrak{m}}$ , and hence, is  $A$ -linear. The restriction of this map to  $M \subseteq M_{\mathfrak{m}}$  takes values in  $A$ . Thus, we can identify  $sf'_i$  with an  $A$ -linear map  $M \rightarrow A$ . Further, for every  $m \in M$ , we have

$$sm = \sum_{i=1}^n sf'_i(m) m_i.$$

That is,  $s \cdot \mathbf{id}_M \in \mathcal{P}$ . Now, let  $\mathfrak{a}$  be the collection of all  $a \in A$  such that  $a \cdot \mathbf{id}_M \in \mathcal{P}$ . Then  $\mathfrak{a}$  is an ideal of  $A$ . If  $\mathfrak{a}$  were a proper ideal, it would be contained in a maximal ideal  $\mathfrak{m}$ . But from our preceding conclusion, there is some  $s \in A \setminus \mathfrak{m}$  such that  $s \cdot \mathbf{id}_M \in \mathcal{P}$ , a contradiction. Thus,  $\mathfrak{a} = A$ , in particular,  $\mathbf{id}_M \in \mathcal{P}$ .

Finally, we show that  $M$  is projective. We have shown that there are  $A$ -linear maps  $f_i : M \rightarrow A$  such that

$$m = \sum_{i=1}^n f_i(m) m_i \quad \forall m \in M.$$

Let  $F$  be the free module  $\bigoplus_{i=1}^n Ae_i$  and let  $g : F \rightarrow M$  be given by  $e_i \mapsto m_i$  and  $f : M \rightarrow F$  given by

$$f(m) = \sum_{i=1}^n f_i(m) e_i.$$

By our construction,  $g \circ f = \mathbf{id}_M$ , and hence  $M$  is a direct summand of  $F$ , i.e.  $M$  is projective. ■

**COROLLARY.** A finitely generated flat module over an integral domain is projective.

*Proof.* Follows from Theorem 2.6 and Theorem 2.5. ■

## §§ Finitely Presented Modules and Flatness

**THEOREM 2.7.** Let  $M$  be a finitely presented  $A$ -module and  $N$  be any  $A$ -module. If  $B$  is a flat  $A$ -algebra, then there is a natural isomorphism

$$\mathrm{Hom}_A(M, N) \otimes_A B \cong \mathrm{Hom}_B(M \otimes_A B, N \otimes_A B).$$

*Proof.* Fixing  $N$  and  $B$ , there are contravariant functors  $\mathcal{F}, \mathcal{G} : \mathfrak{Mod}_A^{op} \rightarrow \mathfrak{Mod}_B$  given by

$$\mathcal{F}(M) = \mathrm{Hom}_A(M, N) \otimes_A B \quad \mathcal{G}(M) = \mathrm{Hom}_B(M \otimes_A B, N \otimes_A B).$$

Define the natural transformation  $\lambda : \mathcal{F} \Rightarrow \mathcal{G}$  given by

$$\lambda_M(f \otimes b) = b \cdot (f \otimes \mathbf{id}_B).$$

We first show that this is natural in  $M$ . Indeed, suppose  $\varphi : M' \rightarrow M$  is  $A$ -linear, we wish to show that

$$\begin{array}{ccc} \mathcal{F}(M) & \longrightarrow & \mathcal{F}(M') \\ \lambda_M \downarrow & & \downarrow \lambda_{M'} \\ \mathcal{G}(M) & \longrightarrow & \mathcal{G}(M') \end{array}$$

commutes. Consider  $f \otimes b \in \mathcal{F}(M)$ , which maps to  $f \circ \varphi \otimes b \in \mathcal{F}(M')$ , which maps to  $b \cdot (f \circ \varphi \otimes \mathbf{id}_B) \in \mathcal{G}(M')$ . On the other hand, under  $\lambda_M$ ,  $f \otimes b$  maps to  $b \cdot (f \otimes \mathbf{id}_B) \in \mathcal{G}(M)$ , which maps to  $b \cdot (f \circ \varphi \otimes \mathbf{id}_B)$ , which shows commutativity.

Next, suppose  $M = A^n$  were free of finite rank. In this case, there is a sequence of isomorphisms

$$\mathrm{Hom}_A(A^n, N) \otimes_A B \cong N^n \otimes_A B \cong (N \otimes_A B)^n \cong \mathrm{Hom}_B(B^n, N \otimes_A B) \cong \mathrm{Hom}_B(A^n \otimes_A B, N \otimes_A B).$$

Under the above isomorphism,  $f \otimes b$  first maps to  $(f(e_1), \dots, f(e_n))^T \otimes b$  in  $N^n \otimes_A B$ . Under the second map, it goes to  $(f(e_1) \otimes b, \dots, f(e_n) \otimes b)^T$  in  $(N \otimes_A B)^n$ . Under the third map it goes to the unique morphism  $g : B^n \rightarrow N \otimes_A B$  that sends  $e_i \mapsto f(e_i) \otimes b$ .

Consider the map  $b \cdot (f \otimes \mathbf{id}_B) \in \mathrm{Hom}_B(A^n \otimes_A B, N \otimes_A B)$ . Under this map,  $e_i \in B^n$  is the same as  $e_i \otimes 1 \in A^n \otimes B$ , which maps to  $b \cdot (f(e_i) \otimes 1) = f(e_i) \otimes b \in N \otimes_A B$ . It follows that this is the same as the aforementioned  $g$ . Thus,  $\lambda_M$  is an isomorphism in this case.

Finally, there is an exact sequence  $A^m \rightarrow A^n \rightarrow M \rightarrow 0$  since  $M$  is finitely presented. This fits into a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}(M) & \longrightarrow & \mathcal{F}(A^n) & \longrightarrow & \mathcal{F}(A^m) \\ & & \downarrow \lambda & & \downarrow \lambda & & \downarrow \lambda \\ 0 & \longrightarrow & \mathcal{G}(M) & \longrightarrow & \mathcal{G}(A^n) & \longrightarrow & \mathcal{G}(A^m) \end{array}$$

where the last two  $\lambda$ 's are isomorphisms. Due to the Five Lemma (after adding another column of zeros to the left), we see that  $\lambda_M : \mathcal{F}(M) \rightarrow \mathcal{G}(M)$  must be an isomorphism, thereby completing the proof. ■

**COROLLARY.** Let  $M$  be a finitely presented  $A$ -module and  $N$  be any  $A$ -module. Then for every  $\mathfrak{p} \in \text{Spec}(A)$ ,

$$\text{Hom}_A(M, N)_{\mathfrak{p}} \cong \text{Hom}_{A_{\mathfrak{p}}}(M_{\mathfrak{p}}, N_{\mathfrak{p}}).$$

*Proof.* Note that the localization functor at  $\mathfrak{p} \in \text{Spec}(A)$  is naturally isomorphic to  $-\otimes_A A_{\mathfrak{p}}$ .  $\blacksquare$

**THEOREM 2.8.** Let  $M$  be a finitely presented  $A$ -module. Then the following are equivalent

- (a)  $M$  is projective.
- (b)  $M_{\mathfrak{p}}$  is a free  $A_{\mathfrak{p}}$ -module for all  $\mathfrak{p} \in \text{Spec}(A)$ .
- (c)  $M_{\mathfrak{m}}$  is a free  $A_{\mathfrak{m}}$ -module for all  $\mathfrak{m} \in \text{MaxSpec}(A)$ .

*Proof.* That  $(a) \implies (b) \implies (c)$  is obvious. It suffices to show that  $(c) \implies (a)$ . To this end, we shall show that  $\text{Hom}_A(M, -)$  is an exact functor. We know that  $\text{Hom}_A(M, -)$  is left exact so let  $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$  be a short exact sequence. Upon application of the above functor, note that we have an exact sequence

$$0 \longrightarrow \text{Hom}_A(M, N') \longrightarrow \text{Hom}_A(M, N) \longrightarrow \text{Hom}_A(M, N'') \rightarrow K \rightarrow 0,$$

where  $K$  is the cokernel. Localizing the above sequence at a maximal ideal  $\mathfrak{m}$  and using the exactness of localization and the preceding result, we have an exact sequence

$$0 \longrightarrow \text{Hom}_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}, N'_{\mathfrak{m}}) \longrightarrow \text{Hom}_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}, N_{\mathfrak{m}}) \longrightarrow \text{Hom}_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}, N''_{\mathfrak{m}}) \rightarrow K_{\mathfrak{m}} \rightarrow 0.$$

But since  $M_{\mathfrak{m}}$  is a free  $A_{\mathfrak{m}}$ -module, the functor  $\text{Hom}_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}, -)$  is exact, whence  $K_{\mathfrak{m}} = 0$  for every  $\mathfrak{m} \in \text{MaxSpec}(A)$ . This shows that  $K = 0$ , that is,  $M$  is projective.  $\blacksquare$

### §3 INJECTIVE MODULES

**DEFINITION 3.1.** An  $A$ -module  $M$  is said to be *injective* if the (contravariant) functor  $\text{Hom}_A(-, M) : \mathfrak{Mod}_A^{op} \rightarrow \mathfrak{Mod}_A$  is exact.

**THEOREM 3.2 (BAER'S CRITERION).** An  $A$ -module  $E$  is injective if and only if for every ideal  $\mathfrak{a} \triangleleft A$ , every  $A$ -linear map  $\mathfrak{a} \rightarrow E$  can be extended to an  $A$ -linear map  $A \rightarrow E$ .

*Proof.* The forward direction is tautological. We prove the converse. Suppose  $N \leq M$  are  $A$ -modules and  $\alpha : N \rightarrow E$  is an  $A$ -linear map. We shall extend  $\alpha$  to a map  $M \rightarrow E$ .

Let  $\Sigma$  be the collection of all pairs  $(N', \alpha')$  where  $N \leq N' \leq M$  and  $\alpha' : N' \rightarrow E$  is  $A$ -linear such that  $\alpha'|_N = \alpha$ . Using a standard Zorn argument,  $\Sigma$  admits a maximal element  $\alpha' : N' \rightarrow E$  extending  $\alpha$ . We contend that  $N' = M$ .

Suppose not. Then choose some  $x \in M \setminus N'$  and let  $\mathfrak{a} = (N' :_A x) \triangleleft A$ . Consider the composite map  $\mathfrak{a} \xrightarrow{x} N' \xrightarrow{\alpha'} E$ , which extends to a map  $f : A \rightarrow E$  and set  $N'' = N' + Ax \leq M$ . Define  $\alpha'' : N'' \rightarrow E$  by

$$\alpha''(n' + ax) = \alpha'(n') + f(a).$$

This is well defined, for if  $n'_1 + a_1x = n'_2 + a_2x$ , then  $(a_1 - a_2)x = n'_2 - n'_1$ , i.e.  $(a_1 - a_2) \in \mathfrak{a}$  and hence,

$$f(a_1 - a_2) = \alpha'((a_1 - a_2)x) = \alpha'(n'_2 - n'_1).$$

But note that  $(N', \alpha') < (N'', \alpha'')$  in  $\Sigma$ , a contradiction. Thus  $N' = M$  and we are done. ■

**COROLLARY.** Let  $A$  be a noetherian ring. If  $\{E_i : i \in I\}$  is a collection of injective  $A$ -modules, then  $E = \bigoplus_{i \in I} E_i$  is an injective  $A$ -module.

*Proof.* Let  $\mathfrak{a} \trianglelefteq A$  and  $f : \mathfrak{a} \rightarrow E$  be  $A$ -linear. Note that  $\mathfrak{a} = (a_1, \dots, a_n)$  is finitely generated, and each  $f(a_i)$  has support contained in a finite subset of  $I$ . Thus,  $f(\mathfrak{a})$  is contained in a direct sum of a finite subset of  $\{E_i : i \in I\}$ . But note that a finite direct sum of injectives is injective over any ring, and hence,  $f$  can be extended to all of  $A$ , thereby completing the proof. ■

**COROLLARY.** Let  $A$  be a PID. An  $A$ -module  $E$  is injective if and only if it is divisible.

*Proof.* Immediate from Theorem 3.2. ■

## §§ Injective Hulls

**DEFINITION 3.3.** Let  $M \leq E$  be  $A$ -modules. Then  $E$  is said to be an *essential extension* of  $M$  if every non-zero submodule of  $E$  intersects  $M$  non-trivially. We denote this by  $M \leq_e E$ .

**REMARK 3.4.** The above is equivalent to requiring that for every  $x \in E \setminus \{0\}$ , there is an  $a \in A \setminus \{0\}$  such that  $ax \in M \setminus \{0\}$ .

We note some trivial properties of essential extensions before proceeding.

**PROPOSITION 3.5.** Let  $L \leq M \leq N$  be  $A$ -modules. Then

$$L \leq_e M \text{ and } M \leq_e N \iff L \leq_e N.$$

*Proof.* Straightforward. ■

**PROPOSITION 3.6.** Let  $M \leq E$  be  $A$ -modules. Consider the set

$$\mathcal{E} = \{N \leq E : M \leq_e N\}.$$

Then  $\mathcal{E}$  has a maximal element.

*Proof.* Standard application of Zorn's lemma. ■

**PROPOSITION 3.7.** If  $N_1 \leq_e M_1$  and  $N_2 \leq_e M_2$ , then  $N_1 \oplus N_2 \leq_e M_1 \oplus M_2$ .

*Proof.* Trivial. ■

**REMARK 3.8.** Before we proceed, we make an important observation. Suppose  $M \leq_e N$  and suppose there is a commutative diagram:

$$\begin{array}{ccc} & N & \\ \uparrow & \searrow f & \\ M & \longrightarrow & E. \end{array}$$

We claim that  $f$  is injective. Indeed, due to the commutativity of the diagram,  $\ker f \cap M = 0$ , but since  $M \leq_e N$ , we have that  $\ker f = 0$ .

**DEFINITION 3.9.** Let  $M \leq E$  be  $A$ -modules. Then  $E$  is said to be an *injective hull* of  $M$  if  $E$  is an injective  $A$ -module and  $M \leq_e E$ . It is customary to denote  $E$  by  $E_A(M)$ .

**PROPOSITION 3.10.** Suppose  $M \leq E$  and  $N \leq F$  are  $A$ -modules such that  $E$  and  $F$  are injective hulls of  $M$  and  $N$  respectively. Then  $E \oplus F$  is an injective hull of  $M \oplus N$ .

*Proof.* Obviously  $E \oplus F$  is injective and due to the preceding result, an essential extension of  $M \oplus N$ . The conclusion follows. ■

**PROPOSITION 3.11.** An  $A$ -module  $E$  is injective if and only if  $E$  has no proper essential extensions.

*Proof.* Suppose  $E$  were injective and  $E \leq_e M$ . Then, there is a submodule  $N$  of  $M$  such that  $M = E \oplus N$ . If  $N$  were non-trivial, then it would intersect  $E$  trivially, thus  $N$  must be trivial and  $E = M$ .

Conversely, suppose  $E$  has no proper essential extensions. There is an injective module  $I$  such that  $E \hookrightarrow I$ . We shall show that  $E$  is a direct summand of  $I$ . Indeed, consider the collection

$$\Sigma = \{N \leq I : E \cap N = 0\}.$$

A standard application of Zorn's lemma furnishes a maximal element  $N$  of  $\Sigma$ . Note that if  $M$  is a submodule of  $I$  properly containing  $N$ , then  $E \cap M \neq 0$ . The canonical projection  $I \twoheadrightarrow I/N$  restricts to an injective map on  $E$  and any submodule of  $I/N$  is of the form  $M/N$  for some  $M$  containing  $N$ . Thus, it follows that  $E \hookrightarrow I/N$  is an essential extension. But since  $E$  does not admit any proper essential extensions, we must have that the aforementioned map is surjective, that is,  $E + N = I$ , whence  $E \oplus N = I$  and hence,  $E$  is injective. ■

**THEOREM 3.12.** Let  $M \leq E$  be  $A$ -modules. The following are equivalent:

- (a)  $E$  is an injective hull of  $M$ .
- (b)  $E$  is a minimal injective  $A$ -module containing  $M$ .
- (c)  $E$  is a maximal essential extension of  $M$ .



*Proof.* (a)  $\implies$  (b) Suppose  $I$  is an injective module such that  $M \leq I \leq E$ . Since  $M \leq_e E$ , we have that  $I \leq_e E$ . But due to Proposition 3.11, we see that  $I = E$ .

(b)  $\implies$  (c) Let  $N \leq E$  be a maximal element of  $\{N \leq E : M \leq_e N\}$ . We contend that  $N$  has no proper essential extensions. Suppose  $f : N \hookrightarrow L$  is an essential extension. Then, there is a map  $L \rightarrow E$  making

$$\begin{array}{ccccc} & & & & E \\ & & & \nearrow & \uparrow \\ 0 & \longrightarrow & N & \xrightarrow{f} & L \end{array}$$

commute. We claim that the map  $L \rightarrow E$  is injective. Indeed, if  $0 \neq x \in L$  maps to 0, then there is an  $0 \neq a \in A$  such that  $0 \neq ax \in f(N)$ . But since  $N \hookrightarrow E$ , we have that  $ax = 0$ , a contradiction. Thus, in  $E$ ,  $L = N$ , since  $N$  has no proper essential extensions in  $E$ . Consequently,  $N$  has no proper essential extensions, that is,  $N$  is injective, whence  $N = E$ .

(c)  $\implies$  (a) Injectivity follows from the fact that  $E$  has no proper essential extensions due to maximality.  $\blacksquare$

**THEOREM 3.13.** Let  $M$  be an  $A$ -module. Then there exists an injective hull  $M \hookrightarrow E$ , which is unique up to isomorphism.

*Proof.* Let  $I$  be an injective module such that  $M \hookrightarrow I$ . Using (b)  $\implies$  (c) of the proof of Theorem 3.12, we see that a maximal essential extension  $E$  of  $M$  contained in  $I$  is an injective hull.

It remains to establish uniqueness. Suppose  $M \hookrightarrow E'$  is another injective hull. Then, there is a commutative diagram

$$\begin{array}{ccc} & & E' \\ & \nearrow & \uparrow \\ M & \hookrightarrow & E \end{array}$$

with the induced map  $E \rightarrow E'$  injective as argued in the preceding proof. The maximality of essentialness and transitivity of essentialness both imply that  $E \rightarrow E'$  must be an isomorphism.  $\blacksquare$

**THEOREM 3.14 (CANTOR-SCHRÖDER-BERNSTEIN).** If  $M$  and  $N$  are injective  $A$ -modules with injective  $A$ -linear maps  $M \hookrightarrow N$  and  $N \hookrightarrow M$ , then  $M \cong N$ .

*Proof.* We may suppose that  $N \leq M$ , whence there is a submodule  $P$  of  $M$  such that  $M = N \oplus P$  where  $P$  is injective too. Let  $f : M \rightarrow N$  be an injective  $A$ -linear map.

Note first that if  $x_0 + f(x_1) + \cdots + f^{(n)}(x_n) = 0$  where  $x_i \in P$ , then all  $x_i = 0$ . Indeed,  $f(x_1) + \cdots + f^{(n)}(x_n) \in \text{im}(f) \subseteq N$  and  $x_0 \in P$ , whence  $x_0 = 0$ . Since  $f$  is injective, we have  $x_1 + \cdots + f^{(n-1)}(x_n) = 0$ . Working downwards, we have our conclusion.

Now, set  $X = P \oplus f(P) \oplus f^{(2)}(P) \oplus \cdots \subseteq M$  and let  $E = E_A(f(X)) \subseteq N$  an injective hull. Write  $N = E \oplus Q$ . Since  $X = P \oplus f(X)$ , we have

$$E(X) \cong E(P \oplus f(X)) \cong E(P) \oplus E(f(X)) \cong P \oplus E.$$

On the other hand, since  $f$  is injective,

$$E(X) \cong E(f(X)) = E \implies P \oplus E \cong E.$$

Consequently,

$$M = N \oplus P = Q \oplus E \oplus P \cong Q \oplus E \cong N,$$

thereby completing the proof. ■

**PROPOSITION 3.15.** Let  $A$  be a noetherian ring and  $M$  an  $A$ -module. Then  $\text{Ass}_A(E(M)) = \text{Ass}_A(M)$ . In particular,  $E(A/\mathfrak{p}) = \{\mathfrak{p}\}$  for every  $\mathfrak{p} \in \text{Spec}(A)$ .

*Proof.* Since  $M \hookrightarrow E(M)$ , we have that  $\text{Ass}_A(M) \subseteq \text{Ass}_A(E(M))$ . Conversely, suppose  $\mathfrak{p} \in \text{Ass}_A(E(M))$ , that is,  $R/\mathfrak{p} \hookrightarrow E(M)$  and identify  $R/\mathfrak{p}$  with a submodule of  $E(M)$ . Since  $M \leq_e E(M)$ ,  $(R/\mathfrak{p}) \cap M \neq 0$ . Choosing a non-zero  $x$  in the intersection, we have that  $\text{Ann}_A(x) = \mathfrak{p}$ , that is,  $\mathfrak{p} \in \text{Ass}_A(M)$ . This completes the proof. ■

**DEFINITION 3.16.** A nonzero  $A$ -module  $M$  is said to be *decomposable* if there are nonzero submodules  $N_1, N_2 \leq M$  such that  $M = N_1 \oplus N_2$ . An  $A$ -module that is not decomposable is said to be *indecomposable*.

**THEOREM 3.17 (MATLIS).** Let  $A$  be a noetherian ring and  $M$  an  $A$ -module. Then,

- (a)  $E$  is an indecomposable injective  $A$ -module if and only if  $E \cong E(A/\mathfrak{p})$  for some  $\mathfrak{p} \in \text{Spec}(A)$ .
- (b)  $E_A(A/\mathfrak{p}) \not\cong E_A(A/\mathfrak{q})$  if  $\mathfrak{p} \neq \mathfrak{q} \in \text{Spec}(A)$ .
- (c) every injective  $A$ -module can be written as a direct sum of indecomposable  $A$ -modules.

*Proof.* (a) Suppose  $E$  is an indecomposable injective  $A$ -module and choose some  $\mathfrak{p} \in \text{Ass}_A(E)$ . There is an injection  $A/\mathfrak{p} \hookrightarrow E$ , which extends to an injection (due to Remark 3.8)  $E(A/\mathfrak{p}) \hookrightarrow E$ . Since  $E$  is indecomposable,  $E \cong E(A/\mathfrak{p})$ .

Conversely, we must show that  $E = E(A/\mathfrak{p})$  is indecomposable. Suppose  $E = E_1 \oplus E_2$ . The map  $A/\mathfrak{p} \hookrightarrow E_1 \oplus E_2$  sends  $\bar{1} \in A/\mathfrak{p}$  to some  $(x_1, x_2) \in E_1 \oplus E_2$ . Then,

$$\mathfrak{p} = \text{Ann}_A((x_1, x_2)) = \text{Ann}_A(x_1) \cap \text{Ann}_A(x_2),$$

whence, we may suppose without loss of generality that  $\mathfrak{p} = \text{Ann}_A(x_1)$ . Consequently, the composition  $A/\mathfrak{p} \hookrightarrow E \twoheadrightarrow E_1$  is injective. This means that  $E \twoheadrightarrow E_1$  is a lift of an injection  $A/\mathfrak{p} \hookrightarrow E_1$ , whence  $E \twoheadrightarrow E_1$  must be injective (due to Remark 3.8), that means  $E_2 = 0$ , as desired.

(b) Follows from the fact that  $\text{Ass}_A(E(A/\mathfrak{p})) = \{\mathfrak{p}\}$ .

(c) This is another standard Zorn argument. Begin with the collection

$$\Sigma = \{ \{E_i\}_{i \in I} : \text{each } E_i \text{ is indecomposable injective, and their sum is direct} \}.$$

Choose a maximal element  $\{E_i\}_{i \in J}$  in  $\Sigma$  and let  $I = \bigoplus_{i \in J} E_i$ . Suppose  $I \neq E$ . Since  $I$  is injective (owing to  $A$  being noetherian), we can write  $E = I \oplus E'$ . Since  $E' \neq 0$ , it has an associated prime,  $\mathfrak{p}$ . We can then write  $E' = E(A/\mathfrak{p}) \oplus E''$ , contradicting the maximality of  $\{E_i\}_{i \in J}$ . This completes the proof. ■

## §4 UNCATEGORIZED

### §§ Eakin-Nagata Theorem

**THEOREM 4.1 (FORMANEK).** Let  $A$  be a ring, and  $B$  a finitely generated faithful  $A$ -module. Suppose the set of  $A$ -submodules  $\Sigma = \{aB : a \trianglelefteq A\}$  has the ascending chain condition, then  $A$  is noetherian.

*Proof.* It suffices to show that  $B$  is a noetherian  $A$ -module since it is finitely generated and faithful. Suppose not. Then consider the collection

$$\Gamma = \{aB : a \trianglelefteq A, B/aB \text{ is a non-noetherian } A\text{-module}\},$$

which contains  $(0)$  and hence is non-empty. Since  $\Sigma$  has the ascending chain condition, so does  $\Gamma$ , whence, it contains a maximal element  $aB$ .

Replacing  $B$  by  $B/aB$ , we see that  $B$  is a non-noetherian  $A$ -module. This may not be faithful and hence, replace  $A$  by  $A/\text{Ann}_A(B)$ . Then,  $B$  is a finite, non-noetherian, faithful  $A$ -module such that for every ideal  $0 \neq a \triangleleft A$ ,  $B/aB$  is a noetherian  $A$ -module.

Next, set

$$\mathfrak{M} = \{N \leq B : B/N \text{ is a faithful } A\text{-module}\},$$

which is non-empty, since  $\{0\} \in \mathfrak{M}$ . Suppose  $B$  is generated as an  $A$ -module by  $b_1, \dots, b_n$ . It is not hard to argue that

$$N \in \mathfrak{M} \iff \forall a \in A \setminus \{0\}, \{ab_1, \dots, ab_n\} \not\subseteq N.$$

It follows that every chain in  $\mathfrak{M}$  has a maximal element and hence Zorn's Lemma applies to furnish a maximal element  $N_0 \in \mathfrak{M}$ .

If  $B/N_0$  is a noetherian  $A$ -module, then  $A$  is noetherian since  $B/N_0$  is faithful and finite. If not, replace  $B$  with  $B/N_0$ , which is still a finite faithful  $A$ -module and satisfies:

- (1)  $B$  is a non-noetherian  $A$ -module.
- (2) for any ideal  $0 \neq a \trianglelefteq A$ ,  $B/aB$  is a noetherian  $A$ -module.
- (3) for any submodule  $0 \neq N$  of  $B$ ,  $B/N$  is not faithful as an  $A$ -module.

Now, let  $N$  be a non-zero submodule of  $B$ . Due to (3), there is a  $0 \neq a \in A$  such that  $aB \subseteq N$ . Due to (2),  $B/aB$  is a noetherian  $A$ -module with  $N/aB$  as a submodule. Thus,  $N/aB$  is a noetherian, in particular, a finite  $A$ -module. Since  $aB$  is also finite as an  $A$ -module, we have that  $N$  is a finite  $A$ -module. Hence,  $B$  is a noetherian  $A$ -module, which is absurd. This completes the proof. ■

**THEOREM 4.2 (EAKIN-NAGATA).** Let  $A \subseteq B$  be an extension of rings such that  $B$  is a finite  $A$ -module. If  $B$  is a noetherian ring, then so is  $A$ .

*Proof.* Note that  $B$  is a finite, faithful  $A$ -module, since  $1 \in B$ . The conclusion follows from Theorem 4.1. ■