

Homological methods in Commutative Algebra

Swayam Chube

January 6, 2025

CONTENTS

1	Regular Sequences	1
1.1	Regular sequences and the Koszul complex	1
1.2	Gorenstein Rings	8
2	Regular Rings	10
2.1	Regular Rings	10

§1 REGULAR SEQUENCES

§§ Regular sequences and the Koszul complex

DEFINITION 1.1. Let A be a ring and M an A -module. An element $a \in A$ is said to be *M -regular* if a is a non zero-divisor on M . A sequence a_1, \dots, a_n of elements of A is an *M -sequence* if

- (1) Each a_i is $M/(a_1, \dots, a_{i-1})M$ -regular.
- (2) $M \neq (a_1, \dots, a_n)M$.

DEFINITION 1.2. Let A be a ring and $x_1, \dots, x_n \in A$. We define a complex K_\bullet by setting $K_0 = A$, $K_p = 0$ for $p > n$ or $p < 0$, and

$$K_p = \bigoplus_{1 \leq i_1 < \dots < i_p \leq n} Ae_{i_1} \wedge \dots \wedge e_{i_p}.$$

For $1 \leq p \leq n$, define $K_p \rightarrow K_{p-1}$ by

$$d(e_{i_1} \wedge \dots \wedge e_{i_p}) = \sum_{i=1}^p (-1)^{r-1} x_{i_r} e_{i_1} \wedge \dots \wedge \widehat{e}_{i_r} \wedge \dots \wedge e_{i_p},$$

and extend linearly to K_p . This is known as the *Koszul complex*.

PROPOSITION 1.3. The Koszul complex is indeed a complex.

Proof. $d \circ d : K_1 \rightarrow K_{-1}$ is obviously the zero map. Now, let $p \geq 2$, we shall show that $(d \circ d)(e_{i_1} \wedge \cdots \wedge e_{i_p}) = 0$. Note that the above can be written as a linear combination of the basis elements of K_{p-2} . Consider the basis element $e_{i_1} \wedge \cdots \wedge \widehat{e}_{i_a} \wedge \cdots \wedge \widehat{e}_{i_b} \wedge \cdots \wedge e_{i_p}$. We shall show that its coefficient is 0.

Indeed, its coefficient is contributed by

$$e_{i_1} \wedge \cdots \wedge \widehat{e}_{i_a} \wedge \cdots \wedge e_{i_p} \quad \text{and} \quad e_{i_1} \wedge \cdots \wedge \widehat{e}_{i_b} \wedge \cdots \wedge e_{i_p},$$

each of which has coefficient $(-1)^{a-1}x_{i_a}$ and $(-1)^{b-1}x_{i_b}$ respectively. The coefficient of our desired basis element in the differential of the first is $(-1)^{b-2}x_{i_b}$ and in the second is $(-1)^{a-1}x_{i_a}$. Thus, the coefficient of our desired basis element in the differential of $e_{i_1} \wedge \cdots \wedge e_{i_p}$ is

$$(-1)^{a-1}x_{i_a}(-1)^{b-2}x_{i_b} + (-1)^{b-1}x_{i_b}(-1)^{a-1}x_{i_a} = 0,$$

thereby completing the proof. ■

DEFINITION 1.4. Let C_\bullet and D_\bullet be complexes of A -modules. Define their *tensor product* $(C \otimes D)_\bullet$ by

$$(C \otimes D)_n = \bigoplus_{i+j=n} C_i \otimes_A D_j.$$

The boundary maps are given by $d : (C \otimes D)_n \rightarrow (C \otimes D)_{n-1}$

$$d(x \otimes y) = dx \otimes y + (-1)^i x \otimes dy \quad x \in C_i, y \in C_j.$$

PROPOSITION 1.5. There is an isomorphism of complexes $(C \otimes D)_\bullet \cong (D \otimes C)_\bullet$.

Proof. If $x \otimes y \in (C \otimes D)_n$ with $x \in C_i$ and $y \in D_j$, then send this element to $(-1)^{ij}y \otimes x \in (D \otimes C)_n$. It is not hard to check that this is indeed a chain map. That this is an isomorphism of chain complexes follows from the fact that for every n , $(C \otimes D)_n \rightarrow (D \otimes C)_n$ is an isomorphism. ■

PROPOSITION 1.6. Let $x_1, \dots, x_n \in A$. Then $K_\bullet(x_1, \dots, x_n) \cong K_\bullet(x_1) \otimes \cdots \otimes K_\bullet(x_n)$ as complexes.

Proof. We prove this by induction on n . The base case with $n = 1$ is tautological. Suppose now that $n \geq 1$. We shall show that $K_\bullet(x_1, \dots, x_n) \otimes K_\bullet(x_{n+1}) \cong K_\bullet(x_1, \dots, x_{n+1})$. Write the complex $K_\bullet(x_{n+1})$ as

$$0 \longrightarrow Ae_{n+1} \xrightarrow{e_{n+1} \mapsto x_{n+1}} A \longrightarrow 0.$$

Then, $(K(x_1, \dots, x_n) \otimes K(x_{n+1}))_p = (K_p(x_1, \dots, x_n) \otimes A) \oplus (K_{p-1}(x_1, \dots, x_n) \otimes Ae_{n+1})$. There is a natural isomorphism

$$(K(x_1, \dots, x_n) \otimes K(x_{n+1}))_p \longrightarrow K_p(x_1, \dots, x_{n+1}),$$

which sends $e_{i_1} \wedge \cdots \wedge e_{i_p} \otimes 1$ to $e_{i_1} \wedge \cdots \wedge e_{i_p}$ in $K_p(x_1, \dots, x_n)$, and sends $e_{i_1} \wedge \cdots \wedge e_{i_{p-1}} \otimes e_{n+1}$ to $e_{i_1} \wedge \cdots \wedge e_{i_{p-1}} \wedge e_{n+1}$ in $K_p(x_1, \dots, x_{n+1})$.

It remains to check that the map defined above is indeed a chain map. Indeed, under the differential in the tensor complex, $e_{i_1} \wedge \cdots \wedge e_{i_p} \otimes 1$ maps to $d(e_{i_1} \wedge \cdots \wedge e_{i_p}) \otimes 1$, which maps to $e(e_{i_1} \wedge \cdots \wedge e_{i_p})$ under the aforementioned isomorphism. On the other hand, the starting element maps to $e_{i_1} \wedge \cdots \wedge e_{i_p}$ under the isomorphism first and then maps to $d(e_{i_1} \wedge \cdots \wedge e_{i_p})$ under the differential.

Next, if we begin with $e_{i_1} \wedge \cdots \wedge e_{i_{p-1}} \otimes e_{n+1}$, then under the differential, it maps to

$$d(e_{i_1} \wedge \cdots \wedge e_{i_{p-1}}) \otimes e_{n+1} + (-1)^{p-1} x_{n+1} e_{i_1} \wedge \cdots \wedge e_{i_{p-1}} \otimes 1,$$

which maps to

$$d(e_{i_1} \wedge \cdots \wedge e_{i_{p-1}}) \wedge e_{n+1} + (-1)^{p-1} x_{n+1} e_{i_1} \wedge \cdots \wedge e_{i_{p-1}}$$

under the isomorphism. On the other hand, the starting element maps to $e_{i_1} \wedge \cdots \wedge e_{i_{p-1}} \wedge e_{n+1}$ under the isomorphism, which maps to the above element under the differential. This completes the proof. \blacksquare

DEFINITION 1.7. Let $\underline{x} = x_1, \dots, x_n$ be a sequence in A . For an A -module M , set

$$K_\bullet(\underline{x}, M) = K(\underline{x}) \otimes M.$$

The homology groups of this complex are denoted by $H_p(\underline{x}, M)$. Similarly, for a complex C_\bullet of A -modules, set $C_\bullet(\underline{x}) = C_\bullet \otimes K_\bullet(\underline{x})$.

PROPOSITION 1.8. Let $\underline{x} = x_1, \dots, x_n$ be a sequence in A . Then

$$H_0(\underline{x}, M) = M/(\underline{x})M \quad H_n(\underline{x}, M) \cong \{\xi \in M : x_1\xi = \cdots = x_n\xi = 0\}.$$

Proof. The assertion about $H_0(\underline{x}, M)$ is trivial. $H_n(\underline{x}, M)$ is precisely the kernel of the map $K_n(\underline{x}, M) \rightarrow K_{n-1}(\underline{x}, M)$, which is given by

$$\xi e_1 \wedge \cdots \wedge e_n \longmapsto \sum_{i=1}^n (-1)^{i-1} x_i \xi e_1 \wedge \cdots \wedge \widehat{e_i} \wedge \cdots \wedge e_n,$$

where $\xi e_{i_1} \wedge \cdots \wedge e_{i_p} \in K_p(\underline{x}, M)$ is shorthand for $e_{i_1} \wedge \cdots \wedge e_{i_p} \otimes \xi \in K_p(\underline{x}, M)$.

The right hand side of the above equation is zero if and only if each $x_i \xi$ is zero, whence the conclusion follows. \blacksquare

THEOREM 1.9. Let C_\bullet be a complex of A -modules and $x \in A$. Then, there is a short exact sequence of complexes

$$0 \rightarrow C_\bullet \rightarrow C_\bullet(x) \rightarrow C'_\bullet \rightarrow 0,$$

where $C'_{p+1} = C_p$ is the (upward) shift of the complex C_\bullet . The homology long exact sequence so obtained looks like

$$\cdots \rightarrow H_p(C_\bullet) \rightarrow H_p(C_\bullet(x)) \rightarrow H_{p-1}(C_\bullet) \xrightarrow{(-1)^{p-1}x} H_{p-1}(C_\bullet) \rightarrow \cdots$$

Further, we have $x \cdot H_p(C_\bullet(x)) = 0$ for all $p \in \mathbb{Z}$.

Proof. Denote the Koszul complex $K_\bullet(x)$ by

$$\cdots \rightarrow 0 \rightarrow Ae_1 \xrightarrow{e_1 \mapsto x} A \rightarrow 0.$$

Thus, we can identify $C_\bullet(x)$ with $C_p \oplus C_{p-1}$ with the boundary map as

$$d(\xi, \eta) = (d\xi + (-1)^{p-1}x\eta, d\eta) \in C_{p-1} \oplus C_{p-2}.$$

Hence, we have a short exact sequence

$$\begin{array}{ccccccc} \cdots & \longrightarrow & C_p & \longrightarrow & C_{p-1} & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & C_p \oplus C_{p-1} & \longrightarrow & C_{p-1} \oplus C_{p-2} & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & C_{p-1} & \longrightarrow & C_{p-2} & \longrightarrow & \cdots \end{array}$$

That the above commutes is straightforward. It remains to compute the boundary map from $H_{p-1}(C_\bullet) = H_p(C'_\bullet)$ to $H_{p-1}(C_\bullet)$.

Choose a cycle $\eta \in C'_p = C_{p-1}$, that is, $d\eta = 0$. This lifts to $(0, \eta) \in C_p \oplus C_{p-1}$, which maps to $((-1)^{p-1}x\eta, 0) \in C_{p-1} \oplus C_{p-2}$, which again lifts to $(-1)^{p-1}x\eta$ in C_{p-1} , which is a cycle in C_{p-1} . Hence, the induced map on homologies is multiplication by $(-1)^{p-1}x$.

Finally, we must show that x annihilates $H_p(C_\bullet(x))$ for all p . Choose a cycle $(\xi, \eta) \in C_p \oplus C_{p-1}$, that is, $d\eta = 0$, and $d\xi = (-1)^p x\eta$. Hence,

$$C_{p+1} \ni d(0, (-1)^p \xi) = ((-1)^p x\xi, (-1)^p d\xi) = x \cdot (\xi, \eta).$$

Thus, x annihilates $[(\xi, \eta)] \in H_p(C_\bullet(x))$, whence annihilates all of $H_p(C_\bullet(x))$. ■

COROLLARY. Let $\underline{x} = x_1, \dots, x_n$ be a sequence in A . Then (\underline{x}) annihilates $H_p(\underline{x}, M)$ for every $p \in \mathbb{Z}$.

Proof. It suffices to show that x_n annihilates $H_p(\underline{x}, M)$ since the Koszul complex is invariant under permutation of the sequence \underline{x} . But this is obvious, since $K_\bullet(\underline{x}, M)$ is isomorphic to $K_\bullet(x_1, \dots, x_{n-1}, M) \otimes K_\bullet(x_n)$ due to the commutativity of tensor products of complexes. We are done by invoking the preceding theorem with $C_\bullet = K_\bullet(x_1, \dots, x_{n-1}, M)$ and $x = x_n$. ■

THEOREM 1.10. Let A be a ring, M an A -module, and x_1, \dots, x_n an M -sequence. Then

$$H_p(\underline{x}, M) = 0 \quad \forall p > 0, \quad H_0(\underline{x}, M) = M/(\underline{x})M.$$

Proof. Induct on n . The base case with $n = 1$ follows from the fact that $H_1(x_1, M) = (0 :_M x_1) = 0$, since x_1 is M -regular. Now, suppose $n > 1$. If $p > 1$, then there is an exact sequence furnished by Theorem 1.9 by taking $C_\bullet = K_\bullet(x_1, \dots, x_{n-1}, M)$ and $x = x_n$:

$$0 = H_p(x_1, \dots, x_{n-1}, M) \longrightarrow H_p(x_1, \dots, x_n, M) \longrightarrow H_{p-1}(x_1, \dots, x_{n-1}, M) = 0,$$

whence $H_p(\underline{x}, M) = 0$. It remains to establish that $H_1(\underline{x}, M) = 0$. Set $M_i = M/(x_1, \dots, x_i)M$ with the convention that $M_0 = M$. The above long exact sequence again furnishes

$$0 = H_1(x_1, \dots, x_{n-1}, M) \rightarrow H_1(\underline{x}, M) \rightarrow H_0(x_1, \dots, x_{n-1}, M) = M_{n-1} \xrightarrow{x_n} M_{n-1}.$$

But since x_n is a non zero-divisor on M_{n-1} , we see that $H_1(\underline{x}, M) = 0$ as desired. \blacksquare

THEOREM 1.11. Suppose one of the following two conditions holds:

- (α) (A, \mathfrak{m}) is a Noetherian local ring, $x_1, \dots, x_n \in \mathfrak{m}$, and M is a finite A -module.
- (β) A is an \mathbb{N} -graded ring, M is an \mathbb{N} -graded A -module, and x_1, \dots, x_n are homogeneous elements of positive degree.

Then, if $H_1(\underline{x}, M) = 0$ and $M \neq 0$, then x_1, \dots, x_n is an M -sequence.

Proof. Induction on n . If $n = 1$, then $0 = H_1(x_1, M) = (0 :_M x_1)$, whence x_1 is a non zero-divisor on M . Now suppose $n > 1$. Again, we make use of the exact sequence associated with $K_\bullet(x_1, \dots, x_{n-1}, M) \otimes K_\bullet(x_n)$ to get

$$H_1(x_1, \dots, x_{n-1}, M) \xrightarrow{-x_n} H_1(x_1, \dots, x_{n-1}, M) \rightarrow H_1(\underline{x}, M) = 0.$$

But since $H_i(x_1, \dots, x_{n-1}, M)$ is a finite A -module in case (α) or a \mathbb{N} -graded module in case (β), the above surjection implies, due to Nakayama, that $H_1(x_1, \dots, x_{n-1}, M) = 0$. The induction hypothesis then implies x_1, \dots, x_{n-1} is an M -sequence.

Now, continuing the above long exact sequence, we get

$$0 = H_1(\underline{x}, M) \rightarrow H_0(x_1, \dots, x_{n-1}, M) = M_{n-1} \xrightarrow{x_n} M_{n-1},$$

where $M_{n-1} = M/(x_1, \dots, x_{n-1})M$. The above sequence implies x_n is M_{n-1} -regular, whence x_1, \dots, x_n is an M -sequence, as desired. \blacksquare

THEOREM 1.12. Let A be a Noetherian ring, M a finite A -module, and I an ideal of A such that $M \neq IM$. For a given integer $n > 0$, the following conditions are equivalent:

- (1) $\text{Ext}_A^i(N, M) = 0$ for all $i < n$ and for any finite A -module N with $\text{Supp}(N) \subseteq V(I)$.
- (2) $\text{Ext}_A^i(A/I, M) = 0$ for all $i < n$.
- (3) $\text{Ext}_A^i(N, M) = 0$ for all $i < n$ and for some finite A -module N with $\text{Supp}(N) = V(I)$.
- (4) There exists an M -sequence of length n contained in I .

Proof. (1) \Rightarrow (2) \Rightarrow (3) is clear. (3) \Rightarrow (4) First, we show that I contains an M -regular element. Suppose not, then due to prime avoidance, I must be contained in some associated prime $\mathfrak{p} \in \text{Ass}_A(M)$. Thus, there is an injective map $A/\mathfrak{p} \hookrightarrow M$, which upon localizing at \mathfrak{p} , we see that $\text{Hom}_{A_{\mathfrak{p}}}(\kappa(\mathfrak{p}), M_{\mathfrak{p}}) \neq 0$. Now, $\mathfrak{p} \in V(I) = \text{Supp}(N)$, whence $N_{\mathfrak{p}} \neq 0$, and hence, due to Nakayama's lemma, $N_{\mathfrak{p}}/\mathfrak{p}N_{\mathfrak{p}} \neq 0$ (since $N_{\mathfrak{p}}$ is a finite $A_{\mathfrak{p}}$ -module). Then,

$N_{\mathfrak{p}}/\mathfrak{p}N_{\mathfrak{p}}$ is a non-zero $\kappa(\mathfrak{p})$ -vector space, and consequently, $\text{Hom}_{A_{\mathfrak{p}}}(N_{\mathfrak{p}}/\mathfrak{p}N_{\mathfrak{p}}, \kappa(\mathfrak{p})) \neq 0$ (choose a basis and project onto a coordinate). Now, we can form the composition

$$N_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}/\mathfrak{p}N_{\mathfrak{p}} \rightarrow \kappa(\mathfrak{p}) \hookrightarrow M_{\mathfrak{p}}.$$

The first two maps are surjections and hence, the composition is non-zero. It follows that $\text{Hom}_{A_{\mathfrak{p}}}(N_{\mathfrak{p}}, M_{\mathfrak{p}}) \neq 0$. Since N is finite over a Noetherian ring, we have

$$(\text{Hom}_A(N, M))_{\mathfrak{p}} = \text{Hom}_{A_{\mathfrak{p}}}(N_{\mathfrak{p}}, M_{\mathfrak{p}}) \neq 0,$$

whence $\text{Ext}_A^0(N, M) = \text{Hom}_A(N, M) \neq 0$, a contradiction to (3). Hence, I contains an M -regular element, say f . If $n = 1$, then we are already done. If $n > 1$, then set $M_1 = M/fM$ and consider the short exact sequence

$$0 \rightarrow M \xrightarrow{f} M \rightarrow M_1 \rightarrow 0.$$

The long exact sequence using $\text{Ext}_A(N, -)$ gives

$$\cdots \rightarrow \text{Ext}_A^{i-1}(N, M) \xrightarrow{f} \text{Ext}_A^{i-1}(N, M) \rightarrow \text{Ext}_A^{i-1}(N, M_1) \rightarrow \text{Ext}_A^i(N, M) \rightarrow \cdots.$$

For $1 \leq i < n$, this implies $\text{Ext}_A^{i-1}(N, M_1) = 0$, and due to the induction hypothesis, there is an M_1 -sequence f_2, \dots, f_n in I . Thus, f_1, \dots, f_n is an M -sequence in I .

(4) \Rightarrow (1). Induction on n . We shall deal with the base case later. Suppose $n > 1$. Let $\underline{x} = x_1, \dots, x_n$ be an M -sequence in I . Set $M_1 = M/x_1M$ which fits into a short exact sequence $0 \rightarrow M \xrightarrow{x_1} M \rightarrow M_1 \rightarrow 0$. The sequence x_2, \dots, x_n is an M_1 -sequence in I , whence due to the inductive hypothesis, $\text{Ext}_A^i(N, M_1) = 0$ for all $i < n - 1$. The long exact sequence corresponding to $\text{Ext}_A(N, -)$ gives us

$$0 = \text{Ext}_A^{i-1}(N, M_1) \rightarrow \text{Ext}_A^i(N, M) \xrightarrow{x_1} \text{Ext}_A^i(N, M)$$

for all $0 \leq i < n$, with the convention that $\text{Ext}_A^{-1}(N, M_1) = 0$. But note that $\text{Ext}_A^i(N, -)$ is annihilated by $\text{Ann}_A(N)$. But since $\text{Supp}(N) = V(\text{Ann}_A(N)) \subseteq V(I)$, we conclude that $I \subseteq \sqrt{I} \subseteq \sqrt{\text{Ann}_A(N)}$. In particular, a sufficiently large power of x_1 annihilates N , whence, annihilates $\text{Ext}_A^i(N, M)$. But since multiplication by x_1 is injective, we must have that $\text{Ext}_A^i(N, M) = 0$ for $i < n$, thereby completing the proof. \blacksquare

THEOREM 1.13. Let A be a Noetherian ring, I an ideal of A , and M a finite A -module such that $M \neq IM$. Then the length of any maximal M -sequence contained in I is the same, say n , and n is determined by

$$\text{Ext}_A^i(A/I, M) = 0 \quad \forall i < n \quad \text{and} \quad \text{Ext}_A^n(A/I, M) \neq 0.$$

We write $n = \text{depth}(I, M)$ and call n the *I-depth* of M .

Proof. Let $\underline{a} = a_1, \dots, a_n$ be a maximal M -sequence in I . Suppose $\text{Ext}_A^n(A/I, M) = 0$. Define $M_i = M/(a_1, \dots, a_i)M$. Using the short exact sequence $0 \rightarrow M \xrightarrow{a_1} M \rightarrow M_1 \rightarrow 0$, we have an exact sequence

$$0 = \text{Ext}_A^{n-1}(A/I, M) \rightarrow \text{Ext}_A^{n-1}(A/I, M_1) \rightarrow \text{Ext}_A^n(A/I, M) = 0,$$

whence $\text{Ext}_A^{n-1}(A/I, M_1) = 0$; and since a_2, \dots, a_n is an M_1 -sequence, $\text{Ext}_A^i(A/I, M_1) = 0$ for $i < n - 1$. Arguing similarly, we get that $\text{Ext}_A^0(A/I, M_n) = 0$. Due to the preceding theorem, I must contain an M_n -regular element, contradicting the maximality of \underline{a} . Thus, $\text{Ext}_A^n(A/I, M) \neq 0$ and $\text{Ext}_A^i(A/I, M) = 0$ for $i < n$.

On the other hand, if $\underline{b} = b_1, \dots, b_m$ is a maximal M -sequence, then due to the above paragraph, $\text{Ext}_A^m(A/I, M) \neq 0$ and $\text{Ext}_A^i(A/I, M) = 0$ for $i < m$. In particular, this means that $m = n$.

Finally, suppose n satisfies the conditions given in the theorem. Then, due to the preceding theorem, there is an M -sequence $\underline{a} = a_1, \dots, a_n$ in I . Further, since $\text{Ext}_A^n(A/I, M) \neq 0$, this sequence must be maximal, else it could be extended and again, due to the preceding theorem $\text{Ext}_A^n(A/I, M) = 0$. This completes the proof. \blacksquare

REMARK 1.14. The above theorem can be phrased more succinctly as

$$\text{depth}(I, M) = \inf \left\{ i : \text{Ext}_A^i(A/I, M) \neq 0 \right\}.$$

In particular, if (A, \mathfrak{m}, k) is a Noetherian local ring, then we write $\text{depth}(\mathfrak{m}, M)$ as $\text{depth } M$ and

$$\text{depth } M = \inf \left\{ i : \text{Ext}_A^i(k, M) \neq 0 \right\}.$$

THEOREM 1.15 (DEPTH SENSITIVITY OF KOSZUL COMPLEX). Let A be a Noetherian ring, $I = (y_1, \dots, y_n)$ an ideal of A , and M a finite A -module such that $M \neq IM$. If

$$q = \sup \{ i : H_i(\underline{y}, M) \neq 0 \},$$

then $\text{depth}(I, M) = n - q$.

Proof. We shall argue by induction on $s = \text{depth}(I, M)$. If $s = 0$, then every element of I is a zero-divisor on M , whence by prime avoidance, there is an associated prime $\mathfrak{p} \in \text{Ass}_A(M)$ such that $I \subseteq \mathfrak{p}$. By definition, there is some $0 \neq \zeta \in M$ such that $\mathfrak{p} = \text{Ann}_A(\zeta)$, and hence, $I\zeta = 0$. Recall that $H_n(\underline{y}, M) = (0 :_M (\underline{y})) = (0 :_M I) \neq 0$, since it contains ζ . Thus, $q = n$.

Now, suppose $s > 0$, then $H_n(\underline{y}, M) = 0$, since some element of I is a non zero-divisor on M . In particular, this means $q < n$. Let $\underline{x} = x_1, \dots, x_s$ be a maximal M -sequence in I . There is a short exact sequence $0 \rightarrow M \xrightarrow{x_1} M \rightarrow M_1 \rightarrow 0$, where $M_1 = M/x_1M$. Since every element in the Koszul complex $K_\bullet(\underline{y})$ is a free module, tensoring with the above short exact sequence will give a short exact sequence of complexes

$$0 \rightarrow K_\bullet(\underline{y}, M) \xrightarrow{x_1} K_\bullet(\underline{y}, M) \rightarrow K_\bullet(\underline{y}, M_1) \rightarrow 0.$$

The associated long exact sequence looks like

$$H_i(\underline{y}, M) \xrightarrow{x_1} H_i(\underline{y}, M) \rightarrow H_i(\underline{y}, M_1) \rightarrow H_{i-1}(\underline{y}, M) \xrightarrow{x_1} H_{i-1}(\underline{y}, M)$$

for all i . Recall that $I = (\underline{y})$ annihilates $H_i(\underline{y}, M)$ for all i , and hence the image of the first map and the kernel of the last map in the above sequence is 0, thereby giving us a short exact sequence

$$0 \rightarrow H_i(\underline{y}, M) \rightarrow H_i(\underline{y}, M_1) \rightarrow H_{i-1}(\underline{y}, M) \rightarrow 0, \quad \forall i \in \mathbb{Z}.$$

Now, note that if $H_i(\underline{y}, M_1) = 0$, then $H_i(\underline{y}, M) = H_{i-1}(\underline{y}, M) = 0$. Hence, $H_{q+1}(\underline{y}, M_1) \neq 0$, but for $i > q + 1$, $H_i(\underline{y}, M_1) = 0$. Now, $\text{depth}(I, M_1) = s - 1$, since x_2, \dots, x_n is a maximal M_1 -sequence in I , for if not, then the original sequence \underline{x} could be extended to a larger M -sequence in I . By the induction hypothesis, we have $q + 1 = n - (s - 1)$, and thus, $s = n - q$. ■

REMARK 1.16. In other words, $\text{depth}(I, M)$ is the number of successive zero terms from the left in the sequence

$$H_n(\underline{y}, M), H_{n-1}(\underline{y}, M), \dots, H_0(\underline{y}, M) = M/IM \neq 0.$$

§§ Gorenstein Rings

LEMMA 1.17. Let A be a ring, M an A -module, and $n \geq 0$ an integer. Then

$$\text{inj dim } M \leq n \iff \text{Ext}_A^{n+1}(A/I, M) = 0 \quad \text{for all ideals } I.$$

If A is Noetherian, then we can replace “for all ideals” by “for all prime ideals” in the above equivalence.

Proof. The forward direction is trivial by considering an injective resolution of length $\leq n$ and constructing the left derived functors of $\text{Hom}_A(A/I, -)$.

We prove the converse. If $n = 0$, then $\text{Ext}_A^1(A/I, M) = 0$, which is equivalent to Baer’s criterion for injectivity. Thus M is injective, that is, $\text{inj dim } M = 0 \leq n$. Now, suppose $n > 0$. Consider an injective resolution of length $n - 1$ and let K be the cokernel of the last map. That is,

$$0 \rightarrow M \rightarrow E^0 \rightarrow E^1 \rightarrow \dots \rightarrow E^{n-1} \rightarrow K_n \rightarrow 0,$$

where every E^i is injective. We claim that K is injective. To see this, break down the above exact sequence into short exact sequences of the form

$$0 \rightarrow K_0 \rightarrow E^0 \rightarrow K_1 \rightarrow 0 \quad 0 \rightarrow K_1 \rightarrow E^1 \rightarrow K_2 \rightarrow 0,$$

and so on, with the convention that $K_0 = M$. The long exact sequence for $\text{Ext}_A(A/I, -)$ on the first short exact sequence gives

$$0 = \text{Ext}_A^n(A/I, E^0) \rightarrow \text{Ext}_A^n(A/I, K_1) \rightarrow \text{Ext}_A^{n+1}(A/I, K_0) = 0,$$

whence $\text{Ext}_A^n(A/I, K_1) = 0$. Proceeding similarly with the other exact sequences, one can show that $\text{Ext}_A^n(A/I, K_n) = 0$, for every ideal I of A . Hence, K_n is injective, i.e., $\text{inj dim } M \leq n$. ■

LEMMA 1.18. Let A be a ring, M and N two A -modules, and $x \in A$. Suppose that x is both A -regular and M -regular, and that $xN = 0$. Set $B = A/xA$ and $\overline{M} = M/xM$. Then

- (1) $\text{Hom}_A(N, M) = 0$ and $\text{Ext}_A^{n+1}(N, M) \cong \text{Ext}_B^n(N, \overline{M})$ for all $n \geq 0$.
- (2) $\text{Ext}_A^n(M, N) \cong \text{Ext}_B^n(\overline{M}, N)$ for all $n \geq 0$.
- (3) $\text{Tor}_n^A(M, N) \cong \text{Tor}_n^B(\overline{M}, N)$ for all $n \geq 0$.

Proof. (1) If $f : N \rightarrow M$ is A -linear, then for any $n \in N$, $xf(n) = f(xn) = 0$, and since x is M -regular, $f(n) = 0$. Thus $f = 0$, as desired. Now, set $T^n(N) = \text{Ext}_A^{n+1}(N, M)$. Then, the collection $(T^n)_{n \geq 0}$ is a contravariant δ -functor from the category \mathfrak{Mod}_B to the category \mathfrak{Mod}_A . Further, the short exact sequence

$$0 \rightarrow M \xrightarrow{x} M \rightarrow \overline{M} \rightarrow 0$$

furnishes a long exact sequence

$$0 = \text{Hom}_A(N, M) \rightarrow \text{Hom}_A(N, \overline{M}) \xrightarrow{\delta} \text{Ext}_A^1(N, M) \xrightarrow{x} \text{Ext}_A^1(N, M) \rightarrow \cdots$$

Since x annihilates N , it must annihilate $\text{Ext}_A^1(N, M)$, and so the above exact sequence reduces to

$$0 \rightarrow \text{Hom}_A(N, \overline{M}) \xrightarrow{\delta} \text{Ext}_A^1(N, M) \rightarrow 0.$$

Thus δ is a natural isomorphism between the functors T^0 and $\text{Ext}_A^1(-, M)$. Now, it suffices to show that the collection $(T^n)_{n \geq 0}$ constitutes a universal δ -functor, whence it suffices to show that $T^n(P) = 0$ for every projective B -module P and $n \geq 1$; since then it would be cofaceable by projectives and due to a theorem of Grothendieck, it would be universal.

This is equivalent to showing that $\text{Ext}_A^n(P, M) = 0$ where P is a direct sum of copies of A/xA and $n \geq 2$. But note that $\text{proj dim}_A A/xA \leq 1$, and hence $\text{Ext}_A^n(A/xA, M) = 0$ for all A -modules M and $n \geq 2$, as desired. This proves (1).

- (2) We contend that $\text{Tor}_n^A(M, B) = 0$ for all $n > 0$. Since $\text{proj dim}_A B \leq 1$, it immediately follows that $\text{Tor}_n^A(M, B) = 0$ for $n > 1$. For $n = 1$, the short exact sequence

$$0 \rightarrow A \xrightarrow{x} A \rightarrow B \rightarrow 0$$

furnishes a long exact sequence

$$0 = \text{Tor}_1^A(M, A) \rightarrow \text{Tor}_1^A(M, B) \rightarrow M \xrightarrow{x} M \rightarrow \overline{M} \rightarrow 0.$$

Since x is M -regular, we have that $\text{Tor}_1^A(M, A) = 0$.

Now, let $P_\bullet \rightarrow M \rightarrow 0$ be a free resolution of M . Because of the preceding paragraph, the sequence $P_\bullet \otimes_A B \rightarrow M \otimes_A B \rightarrow 0$ is exact, so that $P_\bullet \otimes B$ is a free resolution of

the B -module $M \otimes B \cong \overline{M}$. From the Hom-Tensor adjunction, note that there are natural isomorphisms

$$\mathrm{Hom}_A(P_\bullet, N) = \mathrm{Hom}_A(P_\bullet, \mathrm{Hom}_B(B, N)) \cong \mathrm{Hom}_B(P_\bullet \otimes_A B, N).$$

Therefore,

$$\mathrm{Ext}_A^n(M, N) = H^n(\mathrm{Hom}_A(P_\bullet, N)) = H^n(\mathrm{Hom}_B(P_\bullet \otimes_A B, N)) = \mathrm{Ext}_B^n(\overline{M}, N),$$

as desired.

(3) Continuing with the notation of (2), we have

$$\mathrm{Tor}_n^A(M, N) = H_n(P_\bullet \otimes_A N) = H_n((P_\bullet \otimes_A B) \otimes_B N) = \mathrm{Tor}_n^B(\overline{M}, N),$$

thereby completing the proof. ■

§2 REGULAR RINGS

§§ Regular Rings

DEFINITION 2.1. Let (A, \mathfrak{m}, k) be a local ring and let M be a finite A -module. An exact sequence

$$\cdots \rightarrow L_i \xrightarrow{d_i} L_{i-1} \xrightarrow{d_{i-1}} \cdots \rightarrow L_1 \xrightarrow{d_1} L_0 \xrightarrow{\bar{\varepsilon}} M \rightarrow 0$$

is called a *minimal (free) resolution* of M if

- each L_i is a finite free A -module
- $0 = \bar{d}_i : L_i \otimes_A k \rightarrow L_{i-1} \otimes_A k$, or equivalently $d_i L_i \subseteq \mathfrak{m} L_{i-1}$ for all $i \geq 1$, and
- $\bar{\varepsilon} : L_0 \otimes_A k \rightarrow M \otimes_A k$ is an isomorphism.

It is easy to see that a minimal free resolution exists for every finite module over a Noetherian local ring; at each stage simply take a minimal generating set of the kernel and continue.

LEMMA 2.2. Let (A, \mathfrak{m}, k) be a local ring, and M a finite A -module. Suppose L_\bullet is a minimal resolution of M ; then

- (1) $\dim_k \mathrm{Tor}_i^A(M, k) = \mathrm{rank} L_i$ for all i .
- (2) $\mathrm{proj} \dim_A M = \sup \{i : \mathrm{Tor}_i^A(M, k) \neq 0\} \leq \mathrm{proj} \dim_A k$,
- (3) if $M \neq 0$ and $\mathrm{proj} \dim_A M = r < \infty$, then for any finite A -module $N \neq 0$, we have $\mathrm{Ext}_A^r(M, N) \neq 0$.

Proof. (1) This follows immediately from the fact that $\bar{d}_i = 0$ for all $i \geq 1$.

- (2) The second inequality is straightforward. For if $\text{proj dim}_A k = \infty$, then there is nothing to prove. If $\text{proj dim}_A k < \infty$, then take a projective resolution of this length and tensor with A to conclude.

From (1) it immediately follows that $\text{proj dim}_A M \leq \sup \{i: \text{Tor}_i^A(M, k) \neq 0\}$, since this quantity is precisely the length of the minimal free resolution of M . If $\text{proj dim}_A M = \infty$, then there is nothing to prove. If $\text{proj dim}_A M < \infty$, then take a projective resolution of M achieving this length and tensor with k whence it follows that $\sup \{i: \text{Tor}_i^A(M, k) \neq 0\} \leq \text{proj dim}_A M$, as desired.

- (3) Applying $\text{Hom}_A(-, N)$ to the resolution $L_\bullet \rightarrow M$, we obtain a complex which ends with

$$\text{Hom}_A(L_{r-1}, N) \xrightarrow{d_r^*} \text{Hom}_A(L_r, N) \rightarrow 0,$$

where $\text{Ext}_A^r(M, N)$ is the cokernel of the above map. Since each L_i is free, we can write $\text{Hom}_A(L_i, N)$ as a direct sum of some copies of N and we can express every boundary map $d_i : L_i \rightarrow L_{i-1}$ as a matrix with entries in \mathfrak{m} . It follows that d_i^* is given by the same matrix (with entries in \mathfrak{m}). Hence, the image of d_r^* is contained in $\mathfrak{m} \text{Hom}_A(L_r, N)$, which is properly contained in $\text{Hom}_A(L_r, N)$ by Nakayama's lemma. This completes the proof. ■

THEOREM 2.3 (AUSLANDER-BUCHSBAUM). Let A be a Noetherian local ring and $M \neq 0$ a finite A -module. If $\text{proj dim}_A M < \infty$, then

$$\text{proj dim}_A M + \text{depth } M = \text{depth } A.$$

Proof. ■