Results


Descriptives

	Dose	Happiness	Puppy_love
N	Control	9	9
	15 mins	8	8
	30 mins	13	13
Missing	Control	0	0
	15 mins	0	0
	30 mins	0	0
Mean	Control	3.22	3.44
	15 mins	4.88	3.13
	30 mins	4.85	2.00
Median	Control	2.00	4.00
	15 mins	4.50	2.50
	30 mins	4.00	2.00
Standard deviation	Control	1.79	2.07
	15 mins	1.46	1.73
	30 mins	2.12	1.63
Minimum	Control	2.00	1.00
	15 mins	3.00	1.00
	30 mins	2.00	0.00
Maximum	Control	7.00	7.00
	15 mins	7.00	6.00
	30 mins	9.00	5.00
Skewness	Control	1.43	0.319
	15 mins	0.651	0.635
	30 mins	0.550	0.271
Std. error skewness	Control	0.717	0.717
	15 mins	0.752	0.752
	30 mins	0.616	0.616
Kurtosis	Control	1.34	-0.885
	15 mins	-0.732	-0.796
	30 mins	-0.161	-0.980
Std. error kurtosis	Control	1.40	1.40
	15 mins	1.48	1.48
	30 mins	1.19	1.19
Shapiro-Wilk W	Control	0.760	0.921
	15 mins	0.872	0.919
	30 mins	0.935	0.912
Shapiro-Wilk p	Control	0.007	0.400
	15 mins	0.156	0.425
	30 mins	0.395	0.195


Plots


Happiness





Puppy_love






Relationships, Prediction, and Group Comparisons


You have entered a numeric dependent variable, one or more categorical (nominal/ordinal) independent variables, and one or more numeric control variables. Hence, an ANCOVA seems to be a good option for you! In order to run this analysis in jamovi, go to: ANOVA > ANCOVA


- Drop your numeric dependent variable in the box below Dependent Variable
- Drop your nominal/ordinal independent variables in the box below Fixed Factors
- Drop your numeric control variables in the box below Covariates

Scatter Plots of Bivariate Relationships - Dependent/Independent Variables



Scatter Plots of Bivariate Relationships - Dependent/Control Variables





	Sum of Squares	df	Mean Square	F	р	η²p
Dose	12.8	2	6.38	1.98	0.158	0.128
Residuals	87.1	27	3.23			

[3]


Assumption Checks

Normality tests

	statistic	р
Shapiro-Wilk	0.941	0.096
Kolmogorov-Smirnov	0.185	0.258
Anderson-Darling	0.624	0.095

Note. Additional results provided by moretests

Scatterplot



ANCOVA

ANCOVA - Happiness

	Sum of Squares	df	Mean Square	F	р	η²p	ω^2
Dose	36.6	2	18.28	7.48	0.003	0.384	0.234
Puppy_love	17.2	1	17.18	7.03	0.014	0.227	0.109
Dose ∦ Puppy_love	20.4	2	10.21	4.18	0.028	0.258	0.115
Residuals	58.6	24	2.44				

[3]

Assumption Checks

Homogeneity of Variances Tests

	Statistic	df	df2	р
Levene's	8.19	2	27	0.002
Bartlett's	1.11	2		0.574


Note. Additional results provided by moretests

Normality tests

	statistic	р
Shapiro-Wilk	0.948	0.145
Kolmogorov-Smirnov	0.141	0.594
Anderson-Darling	0.691	0.064

Note. Additional results provided by moretests

Q-Q Plot



Contrasts

Contrasts - Dose

	Estimate	SE	t	р
15 mins - Control	1.72	1.60	1.07	0.294
30 mins - Control	4.69	1.27	3.69	0.001

Post Hoc Tests

Post Hoc Comparisons - Dose


Comparison		rison						
Dose		Dose	Mean Difference	SE	df	t	P _{tukey}	p _{bonferroni}
Control	-	15 mins	-1.874	0.794	24.0	-2.360	0.067	0.080
	-	30 mins	-2.006	0.732	24.0	-2.739	0.030	0.034
15 mins	-	30 mins	-0.132	0.743	24.0	-0.178	0.983	1.000

Note. Comparisons are based on estimated marginal means

[4]

Estimated Marginal Means

Dose



Estimated Marginal Means - Dose

			95% Confidence Interval		
Dose	Mean	SE	Lower	Upper	
Control	2.68	0.555	1.54	3.82	
15 mins	4.55	0.569	3.38	5.73	
30 mins	4.69	0.478	3.70	5.67	

[4]

References

[1] The jamovi project (2021). jamovi. (Version 2.2) [Computer Software]. Retrieved from https://www.jamovi.org.

[2] R Core Team (2021). *R: A Language and environment for statistical computing*. (Version 4.0) [Computer software]. Retrieved from https://cran.r-project.org. (R packages retrieved from MRAN snapshot 2021-04-01).

[3] Fox, J., & Weisberg, S. (2020). *car: Companion to Applied Regression*. [R package]. Retrieved from https://cran.r-project.org/package=car.

[4] Lenth, R. (2020). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. [R package]. Retrieved from https://cran.r-project.org/package=emmeans.