

Figure 10.11 The facial nerve. (Reprinted from Netter Anatomy Illustration Collection. ©Elsevier Inc. All Rights Reserved.)

upon that muscle beneath the platysma to the parotid gland, where it divides into an anterior and a posterior branch.

THE FACIAL NERVE

Important anatomic landmarks to the main trunk of the facial nerve include the tympanomastoid suture, the tragal pointer, and the posterior belly of the digastric muscle (Fig. 10.12). The main trunk of the facial nerve exits the stylomastoid foramen, where it can be found 10 mm medial and inferior to the cartilaginous tragal pointer of the external auditory canal. The tympanomastoid suture line is a useful landmark and lies between the mastoid and tympanic segments of the temporal bone. The tympanomastoid suture is approximately 6–8 mm lateral to the stylomastoid foramen. The styloid process is deeper to the nerve trunk and usually lies 5 cm deep to the skin surface in adults. In children, the facial nerve is more superficial and surgical access should take into consideration its superficial location to avoid injury. The facial nerve supplies the postauricular muscles, stylohyoid muscle and posterior belly of the digastric muscle before it bifurcates into its main branches. The facial nerve pierces the parotid gland and bisects the gland along a plane separating the deep and superficial lobes of the gland. In the parotid gland, it divides to two major divisions: the temporal-facial and the cervico-facial branches.

Significant variability in facial nerve branching exists, even between the left and right sides of the face in the same




Figure 10.12 Branches of the facial nerve exposed after superficial parotidectomy on a cadaver. Superficial temporal artery. 1. Frontal branch, 2. Zygomatic branch, 3. Buccal branch, 4. Marginal mandibular branch. The cervical branch is not shown.

individual.²³ After its bifurcation at the pes anserinus, five major branches of the facial nerve exist: the frontal (i.e., temporal), zygomatic, buccal, marginal mandibular, and cervical branches. The frontal branch of the facial nerve is located along a line extending from 0.5 cm below the tragus to a point 1 ± 0.45 cm superior to the supraorbital rim. After exiting the parotid gland, the frontal branch of the facial nerve crosses over the zygomatic arch within the innominate fascia (see Fascia of the head and neck, above), where it remains for a distance 1.5–3.0 cm above the superior border of the arch. After that transition point, the nerve becomes more superficial and travels on the undersurface of the superficial temporal fascia (TPF) until reaching the frontalis muscle.^{1,2} On average, three branches enter the undersurface of the frontalis muscle.²³ However, the marginal mandibular branch of the facial nerve runs below the inferior border of the mandible in 81% of dissected specimens, whereas anterior to the facial artery, it was found above the inferior border in 100% of specimens. Two major branches existed in 67% of individuals; a single marginal mandibular branch was seen in 21%, and three branches were found in 9% of specimens.²⁴

CONCLUSION

The head and neck regional anatomy is a complex area. This chapter has focused on the relevant anatomy related to head and neck reconstruction, and particularly, the approach to vascular and neural structures.

REFERENCES

1. Yang D, Yang JF, Morris S. The fascial planes of the temporal region related to the frontal branch of the facial nerve. *Plast Reconstr Surg* 2011;127(2):991–2.
2. Agarwal CA, Mendenhall SD, Foreman KB, et al. The course of the frontal branch of the facial nerve in relation to fascial planes: an anatomic study. *Plast Reconstr Surg* 2010;125(2):532–7.
3. Yang D, Tang M, Geddes C, et al. Vascular anatomy of the integument of the head and neck. In: Blondeel PN, Morris SF, Hallock GG, et al., editors. *Perforator flaps: anatomy, techniques and clinical applications*, vol. 1. 2nd ed. St Louis, MO: Quality Medical Publishing; 2013. p. 189–221.
4. Magden O, Edizer M, Tayfur V, et al. Anatomic study of the vasculature of the submental artery flap. *Plast Reconstr Surg* 2004;114(7):1719–23.
5. Niranjan NS. An anatomical study of the facial artery. *Ann Plast Surg* 1988;21(1):14–22.
6. Pribaz J, Stephens W, Crespo L, et al. A new intraoral flap: facial artery musculomucosal (FAMM) flap. *Plast Reconstr Surg* 1992;90(3):421–9.
7. Curran AJP, Neligan P, Gullane PJ. Submental artery island flap. *Laryngoscope* 1997;107(11):1545–9.
8. Martin D, Pascal JF, Baudet J, et al. The submental island flap: a new donor site. Anatomy and clinical applications as a free or pedicled flap. *Plast Reconstr Surg* 1993;92(5):867–73.
9. Becker C, Vasile JV, Levine JL, et al. Microlymphatic surgery for the treatment of iatrogenic lymphedema. *Clin Plast Surg* 2012;39(4):385–98.
10. Neligan PC, She-Yue H, Gullane PJ. Reverse flow as an option in microvascular recipient anastomosis. *Plast Reconstr Surg* 1997;100:1780–7.
11. Ozgur Z, Govsa F, Celik S, et al. Clinically relevant variations of the superior thyroid artery: an anatomic guide for surgical neck dissection. *Surg Radiol Anat* 2009;31(3):151–9.
12. Gilbert R, Neligan P. Microsurgical laryngotracheal reconstruction. *Clin Plast Surg* 2005;32(3):293–301.
13. Pinar YA, Govsa F. Anatomy of the superficial temporal artery and its branches: its importance for surgery. *Surg Radiol Anat* 2006;28(3):248–53.
14. Tayfur V, Edizer M, Magden O. Anatomic bases of superficial temporal artery and temporal branch of facial nerve. *J Craniofac Surg* 2010;21(6):1945–7.
15. Hanasono MM, Barnea Y, Skoracki RJ. Microvascular surgery in the previously operated and irradiated neck. *Microsurgery* 2009;29(1):1–7.
16. Sapountzis S, Singhal D, Rashid A, et al. Lymph node flap based on the right transverse cervical artery as a donor site for lymph node transfer. *Ann Plast Surg* 2014;73(4):398–401.
17. Yu P. The transverse cervical vessels as recipient vessels for previously treated head and neck cancer patients. *Plast Reconstr Surg* 2005;115(5):1253–8.
18. Huelke DF. A study of the transverse cervical and dorsal scapular arteries. *Anat Rec* 1958;132(3):233–45.
19. Faltaous AA, Yetman RJ. The submental artery flap: an anatomic study. *Plast Reconstr Surg* 1996;97(1):56–62.
20. Mulholland S, Boyd JB, McCabe S, et al. Recipient vessels in head and neck microsurgery: radiation effect and vessel access. *Plast Reconstr Surg* 1993;92(4):628–32.
21. Boyd B, Mulholland S, Gullane P, et al. Reinnervated lateral antebrachial cutaneous neurosome flaps in oral reconstruction: are we making sense? *Plast Reconstr Surg* 1994;93:1350–9.
22. Brenner E, Schoeller T. Masseteric nerve: a possible donor for facial nerve anastomosis? *Clin Anat* 1998;11(6):396–400.
23. Tzafetta K, Terzis JK. Essays on the facial nerve: Part I. Microanatomy. *Plast Reconstr Surg* 2010;125(3):879–89.
24. Dingman RO, Grabb WC. Surgical anatomy of the mandibular ramus of the facial nerve based on the dissection of 100 facial halves. *Plast Reconstr Surg* 1962;29:266–72.