

Aesthetic Procedure Statistics and Overview

Cosmetic Surgery National Data Bank Statistics. American Society for Aesthetic Plastic Surgery. http://www.surgery.org/media/statistics.

Fedok FG, Advances in minimally invasive facial rejuvenation. Current Opin Otolaryngol Head Neck Surg. 2008;16(4):359–368.

Small R. Aesthetic procedures in office practice. Am Fam Physician. 2009;80(11):1231–1237.

Consultation

Small R. Aesthetic Principles and Consultation. In: Usatine R, Pfenninger J, Stuhlberg D, Small R, eds. Dermatologic and Cosmetic Procedures in Office Practice. Philadelphia, PA. Elsevier. 2011;230–240.

Small R. Aesthetic Procedures Introduction. In: Mayeaux E, ed. The Essential Guide to Primary Care Procedures. Philadelphia, PA. Lippincott Williams & Wilson. 2009;195–199.

Facial Anatomy

Goldberg RA. The three periorbital hollows: a paradigm for periorbital rejuvenation. Plast Reconstr Surg. 2005;116(6):1796–1804.

Netter FH. Atlas of Human Anatomy. 4th Edn. Philadelphia, PA. Saunders. 2006; 24, 25, 35-50.

Dermal Fillers General

Alam M, Gladstone H, Kramer EM, et al. ASDS guidelines of care: injectable fillers. Dermatol Surg. 2008;34(Suppl 1):S115–S148.

Elson M. Dermal Fillers. In: Pfenninger JL, Folwer GC, eds. Procedures for Primary Care, 3rd ed. Mosby/Elsevier, Philadelphia, PA, 2011;373–378.

Fagien S, Klein AW. A brief overview and history of temporary fillers: evolution, advantages, and limitations. Plast Reconstr Surg. 2007;120(6 Suppl):8S–16S.

Goldman MP. Optimizing the use of fillers for facial rejuvenation: the right tool for the right job. Cosmetic Dermatology. 2007;20(7 S3):14–26.

Narins RS, Donofrio LM. Update on Fillers. Dermatol Surg. 2010;36:729.

Niamtu J. Facial aging and regional enhancement with injectable fillers. Cosmetic Dermatology. 2007; 20(5 Suppl):S14–S20.

Sadick NS. Soft tissue augmentation: selection, mode of operation, and proper use of injectable agents. Cosmetic Dermatology. 2007;20(5 S2):8–13.

Small R. Dermal Fillers. In: Usatine R, Pfenninger J, Stuhlberg D, Small R, eds. Dermatologic and Cosmetic Procedures in Office Practice. Philadelphia, PA. Elsevier. 2011;298–308.

Small R. Dermal Fillers for Facial Rejuvenation. In: Mayeaux E, ed. The Essential Guide to Primary Care Procedures. Philadelphia, PA. Lippincott Williams & Wilson. 2009;214–233.

Werschler WP, Kane M. Optimal use of facial filling agents: understanding the products. Cosmetic Dermatology. 2007;20(5 Suppl):S4–S47.

Hyaluronic Acid Dermal Fillers

Andre P. New trends in face rejuvenation by hyaluronic acid injections. J Cosmet Dermatol. 2008;7(4):251–258. Born T. Hyaluronic acids. Clin Plast Surg. 2006;33(4):525–538.

172 Bibliography

Brandt F, Bank D, Cross SL, et al. A lidocaine-containing formulation of large-gel particle hyaluronic acid alleviates pain. Dermatol Surg. 2010;36(Suppl 3):1876–1885.

Carruthers J, Carruthers A, Tezel A, et al. Volumizing with a 20-mg/mL smooth, highly cohesive, viscous hyaluronic acid filler and its role in facial rejuvenation therapy. Dermatol Surg. 2010;36(Suppl 3):1886–1892.

Dover JS, Rubin MG, Bhatia AC. Review of the efficacy, durability, and safety data of two nonanimal stabilized hyaluronic acid fillers from a prospective, randomized, comparative, multicenter study. Dermatol Surg. 2009;35(Suppl 1):322–330.

Green MS. Not all hyaluronic acid dermal fillers are equal. Cosmetic Dermatology. 2007;20(11):724–729.

Kablik J, Monheit GD, Yu L, et al. Comparative physical properties of hyaluronic acid dermal fillers. Derm Surg. 2009;(35 Suppl 1):302–312.

Monheit GD, Coleman KM. Hyaluronic acid fillers. Dermatol Ther. 2006;19(3):141-150.

Weinkle SH, Bank DE, Boyd CM, et al. A multi-center, double-blind, randomized controlled study of the safety and effectiveness of Juvederm injectable gel with and without lidocaine. J Cosmet Dermatol. 2009;8(3):205–210.

Calcium Hydroxylapatite Dermal Fillers

Berlin AL, Hussain M, Goldberg DJ. Calcium hydroxylapatite filler for facial rejuvenation: a histologic and immunohistochemical analysis. Dermatol Surg. 2008;34(Suppl):S64–S67.

Busso M, Voigts R. An investigation of changes in physical properties of injectable calcium hydroxylapatite in a carrier gel when mixed with lidocaine and with lidocaine/epinephrine. Dermatol Surg. 2008;34(Suppl 1):S16–S23.

Carruthers A, Liebeskind M, Carruthers J, et al. Radiographic and computed tomographic studies of calcium hydroxylapatite for treatment of HIV-associated facial lipoatrophy and correction of nasolabial folds. Dermatol Surg. 2008;34(Suppl 1):S78–S84.

Jansen DA, Graivier MH. Evaluation of a calcium hydroxylapatite-based implant (Radiesse) for facial soft-tissue augmentation. Plast Reconstr Surg. 2006;118(3 Suppl):22S-30S.

Marmur E, Green L, Busso M. Controlled, randomized study of pain levels in subjects treated with calcium hydroxylapatite premixed with lidocaine for correction of nasolabial folds. Dermatol Surg. 2010;36(3):309–315.

Anesthesia

Foley K, Pianalto D. Facial Anesthesia. Emerg Med. 2005;37(6):30-34.

Henning JS, Firoz BF. The use of a cooling device as an analgesic before injectable local anesthesia in the pediatric population. Dermatol Surg. 2010;36(4):520–523.

Kawesk S. Topical anesthetic creams [Safety and Efficacy Report]. Plast Reconstr Surg. 2008;121(6):2161–2165. Lee MS. Topical triple-anesthetic gel compared with 3 topical anesthetics. Cosmetic Dermatology. 2003;26(61); 35–38.

Niamtu J, III. Simple technique for lip and nasolabial fold anesthesia for injectable fillers. Dermatol Surg. 2005;31(10):1330–1332.

Salam G, et al. Regional anesthesia for office procedures: Part 1. Head and neck surgeries. Am Fam Phys. 2004;69(3):585–590.

Small R. Anesthesia for Cosmetic Procedures. In: Usatine R, Pfenninger J, Stuhlberg D, Small R, eds. Dermatologic and Cosmetic Procedures in Office Practice. Philadelphia, PA: Elsevier; 2011;241–247.

Smith KC. Ice anesthesia for injection of dermal fillers. Dermatol Surg. 2010;36:812–814.

Weiss RA, Lavin PT. Reduction of pain and anxiety prior to botulinum toxin injections with a new topical anesthetic method. Ophthal Plast Reconstr Surg. 2009;25(3):173–177.

Storing and Reusing Dermal Fillers

Safo PK, Wahlgren C, Obagi S. Safety of Storing and Reusing Hyaluronic Acid Fillers: A Retrospective Chart Review. Cosmet Dermatol. 2011;24:22–27.

Dermal Filler Treatment Areas

Nasolabial Folds

Alam M, Yoo SS. Technique for calcium hydroxylapatite injection for correction of nasolabial fold depressions. J Am Acad Dermatol. 2007;56(2):285–289.

Graivier MH. Correcting nasolabial folds. Cosmetic Surg Times. 2006;1-8.

Heden P, Fagrell D, Jernbeck J, et al. Injection of stabilized hyaluronic acid-based gel of non-animal origin for the correction of nasolabial folds: comparison with and without lidocaine. Dermatol Surg. 2010;36:775–781.

Lupo MP, Smith SR, Thomas JA, et al. Effectiveness of Juvederm Ultra Plus dermal filler in the treatment of severe nasolabial folds. Plast Reconstr Surg. 2008;121(1):289–297.

Marmur ES, Taylor SC, Grimes PE, et al. Six-month safety results of calcium hydroxylapatite for treatment of nasolabial folds in Fitzpatrick skin types IV to VI. Dermatol Surg. 2009;35(Suppl 2):1641–1645.

Moers-Carpi M, Vogt S, Santos BM, et al. A multicenter, randomized trial comparing calcium hydroxylapatite to two hyaluronic acids for treatment of nasolabial folds. Dermatol Surg. 33(Suppl 2):S144–S151.

Bibliography 173

Narins RS, Brandt FS, Leyden J, et al. A randomized, double-blind, multicenter comparison of the efficacy and tolerability of Restylane versus Zyplast for the correction of nasolabial folds. Dermatol Surg. 2003;29(6):588–595.

Smith SR, Busso M, McClaren M, et al. A randomized, bilateral, prospective comparison of calcium hydroxylapatite microspheres versus human-based collagen for the correction of nasolabial folds. Dermatol Surg. 2007;33(Suppl 2):S112–S121.

Weiss RA, Bank D, Brandt FS. Randomized, double-blind, split-face study of small-gel-particle hyaluronic acid with and without lidocaine during correction of nasolabial folds. Dermatol Surg. 2010;36:750–759.

Lip Augmentation

Ali MJ, Ende K, Maas CS. Perioral rejuvenation and lip augmentation. Facial Plast Surg Clin North Am. 2007;15(4):491–500, vii.

Lanigan S. An observational study of a 24 mg/mL hyaluronic acid with pre-incorporated lidocaine for lip definition and enhancement. J Cosmet Dermatol. 2011;10(1):11–14.

Solish N, Swift A. An open-label, pilot study to assess the effectiveness and safety of hyaluronic acid gel in the restoration of soft tissue fullness of the lips. J Drugs Dermatol. 2011;10(2):145–149.

Other Facial Areas

Belmontesi M, Grover R, Verpaele A. Transdermal injection of Restylane SubQ for aesthetic contouring of the cheeks, chin, and mandible. Aesthet Surg J. 2006;26(1S):S28–S34.

Busso M, Karlsburg PL. Cheek augmentation and rejuvenation using injectable calcium hydroxylapatite (radiesse). Cosmetic Dermatology. 2006;19(9):583–588.

Lowe NJ, Grover R. Injectable hyaluronic acid implant for malar and mental enhancement. Dermatol Surg. 2006;32(7):881–885.

Sclafani AP, Kwak E. Alternative management of the aging jawline and neck. Facial Plast Surg. 2005;21(1):47–54. Verpaele A, Strand A. Restylane SubQ, a non-animal stabilized hyaluronic acid gel for soft tissue augmentation of the mid- and lower face. Aesthet Surg J. 2006;26(1S):S10–S17.

Weinkle S. Injection techniques for revolumization of the perioral region with hyaluronic acid. J Drugs Dermatol. 2010;9(4):367–371.

Combining Aesthetic Treatments

Carruthers A, Carruthers J, Monheit GD, et al. Multicenter, randomized, parallel-group study of the safety and effectiveness of onabotulinumtoxinA and hyaluronic acid dermal fillers (24-mg/ml smooth, cohesive gel) alone and in combination for lower facial rejuvenation. Dermatol Surg. 2010;36(Suppl 4):2121–2134.

Carruthers J, Carruthers A, Maberley D. Deep resting glabellar rhytides respond to BTX-A and Hylan B. Derm Surg. 2003;29(5):539–544.

Carruthers JD, Carruthers A. A prospective, randomized, parallel group study analyzing the effect of BTX-A (Botox) and nonanimal sourced hyaluronic acid (NASHA, Restylane) in combination compared with NASHA (Restylane) alone in severe glabellar rhytides in adult female subjects: treatment of severe glabellar rhytides with a hyaluronic acid derivative compared with the derivative and BTX-A. Dermatol Surg. 2003;29(8):802–809.

Carruthers JD, Glogau RG, Blitzer A. Advances in facial rejuvenation: botulinum toxin type a, hyaluronic acid dermal fillers, and combination therapies—consensus recommendations. Plast Reconstr Surg. 2008;121(5 Suppl):5S–30S.

Coleman KR, Carruthers J. Combination therapy with BOTOX and fillers: the new rejuvnation paradigm. Dermatol Ther. 2006;19(3):177–188.

Godin MS, Majmundar MV, Chrzanowski DS, et al. Use of radiesse in combination with restylane for facial augmentation. Arch Facial Plast Surg. 2006;8(2)92–97.

Maas CS. Botulinum neurotoxins and injectable fillers: minimally invasive management of the aging upper face. Facial Plast Surg Clin North Am. 2006;14(3):241–245.

Small R, Hoang D. Combining Cosmetic Treatments. In: Usatine R, Pfenninger J, Stuhlberg D, Small R, eds. Dermatologic and Cosmetic Procedures in Office Practice. Philadelphia, PA. Elsevier. 2011;377–382.

Complications Dermal Fillers

Bachmann F, Erdmann R, Hartmann V, et al. The spectrum of adverse reactions after treatment with injectable fillers in the glabellar region: results from the Injectable Filler Safety Study. Dermatol Surg. 2009;35(Suppl 2):1629–1634. Baumann LS, Romanelli P, Marangoni O, et al. Keratoacanthomas as complication of dermal filler injections. Cosmetic Dermatology. 2007;20(7):450–452.

Brody HJ. Use of hyaluronidase in the treatment of granulomatous hyaluronic acid reactions or unwanted hyaluronic acid misplacement. Dermatol Surg. 2005;31(8 Pt 1):893–897.

Cassuto D, Marangoni O, De SG, et al. Advanced laser techniques for filler-induced complications. Dermatol Surg. 2009;35(Suppl 2):1689–1695.

Chabra I, Obagi S. Severe site reaction after injecting hyaluronic acid-based soft tissue filler. Cosmetic Dermatology. 2011;24(1):14–21.

174 Bibliography

Cohen JL. Understanding, avoiding, and managing dermal filler complications. Dermatol Surg. 2008;34(Suppl 1): S92–S99.

Coleman KR, Carruthers J. Combination therapy with BOTOX and fillers: the new rejuvenation paradigm. Dermatol Ther. 2006;19(3):177–188.

Cox SE. Clinical experience with filler complications. Dermatol Surg. 2009;35:1661–1666.

DeLorenzi C, Weinberg M, Solish N, et al. The long-term efficacy and safety of a subcutaneously injected large-particle stabilized hyaluronic acid-based gel of nonanimal origin in esthetic facial contouring. Dermatol Surg. 2009;35(Suppl 1):313–321.

Geisler D, Shumer S, Elson M. Delayed hypersensitivity reaction to restylane. Cosmetic Dermatology. 2007;20(12):784–786.

Gladstone HB, Cohen JL. Adverse effects when injecting facial fillers. Semin Cutan Med Surg. 2007;26(1):34–39. Glogau RG, Kane MA. Effect of injection techniques on the rate of local adverse events in patients implanted with nonanimal hyaluronic acid gel dermal fillers. Dermatol Surg. 2008;34(Suppl 1):S105–S109.

Goldman MP. Superficial nodularity of hydroxylapatite filler to fill the infraorbital hollow. Dermatol Surg. 2010;36:822–824.

Grunebaum LD, Allemann IB, Dayan S, et al. The risk of alar necrosis associated with dermal filler injection. Dermatol Surg. 2009;35:1635–1640.

Hirsch RJ, Cohen JL, Carruthers JD. Successful management of an unusual presentation of impending necrosis following a hyaluronic acid injection embolus and a proposed algorithm for management with hyaluronidase. Dermatol Surg. 2007;33(3):357–360.

Hirsch RJ, Narurkar V, Carruthers J. Management of injected hyaluronic acid induced Tyndall effects. Lasers Surg Med. 2006;38(3):202–204.

Junkins-Hopkins JM. Filler complications. J Am Acad Dermatol. 2010;63(4):703-705.

Lee A, Grummer SE, Kriegel D, et al. Hyaluronidase. Dermatol Surg. 2010;36(7):1071–1077.

Lowe NJ, Maxwell CA, Patnaik R. Adverse reactions to dermal fillers: review. Dermatol Surg. 2005;31 (11 Pt 2):1616–1625.

Narins RS, Coleman WP, III, Glogau RG. Recommendations and treatment options for nodules and other filler complications. Dermatol Surg. 2009;35(Suppl 2):1667–1671.

Narins RS, Jewell M, Rubin M, et al. Clinical conference: management of rare events following dermal fillers-Focal necrosis and angry red bumps. Dermatol Surg. 2006;32(3):426–434.

Rohrich RJ, Monheit G, Nguyen AT, et al. Soft-tissue filler complications: the important role of biofilms. Plast Reconstr Surg. 2010;125(4):1250–1256.

Requena L, Requena C, Christensen L, et al. Adverse reactions to injectable soft tissue fillers. J Am Acad Dermatol. 2011;64(1):1–34.

Schanz S, Schippert W, Ulmer A, et al. Arterial embolization caused by injection of hyaluronic acid (Restylane). Brit J Dermatol. 2002;146(5):928–929.

Sclafani AP, Fagien S. Treatment of injectable soft tissue filler complications. Dermatol Surg. 2009;35 (Suppl 2):1672–1680.

Sperling B, Bachmann F, Hartmann V, et al. The current state of treatment of adverse reactions to injectable fillers. Dermatol Surg. 2010;36(Suppl 3):1895–1904.

Van DS, Hays GP, Caglia AE, et al. Severe acute local reactions to a hyaluronic acid-derived dermal filler. J Clin Aesthet Dermatol. 2010;3(5):32–35.

 $\label{lem:adaptive_problem} \begin{tabular}{ll} Vartanian AJ, Frankel AS, Rubin MG. Injected hyaluronidase reduces restylane-mediated cutaneous augmentation. Arch Facial Plast Surg. 2005;7(4):231–237. \end{tabular}$

Zielke H, Wolber L, Wiest L, et al. Risk profiles of different injectable fillers: results from the Injectable Filler Safety Study (IFS Study). Dermatol Surg. 2008;34(3):326–335.

Complications Anesthesia

Hahn IH, Hoffman RS, Nelson LS. EMLA-induced methemoglobinemia and systemic topical anesthetic toxicity. J Emerg Med. 2004;26(1):85–88.

Kaweski S. Topical anesthetic creams. Plast Reconstr Surg. 2008;121(6):2161–2165.

Lacy CF, Armstrong LL, Goldman MP, et al. Drug Information Handbook. 20th ed. Hudson: Lexicomp Inc, 2011.
Marra DE, Yip D, Fincher EF, et al. Systemic toxicity from topically applied lidocaine in conjunction with fractional photothermolysis. Arch Dermatol. 142(8):1024–1026, 2006.

Physicians Desk Reference. 65th ed. Montvale, NJ. Thompson PDR. 2011.

Touma S, Jackson JB. Lidocaine and prilocaine toxicity in a patient receiving treatment for molusca contagiosa. J Am Acad Dermatol. 2001;44(2 suppl):399–400.