

# **Applying effective environmental forecasts to Management Strategy Evaluation of yellowtail flounder**

SOPHIE WULFING<sup>1\*</sup> CHENGXUE LI<sup>3</sup> SCOTT I. LARGE<sup>2</sup> VINCENT SABA<sup>2</sup> GAVIN FAY<sup>1</sup>

<sup>1</sup>School for Marine Science and Technology, University of Massachusetts Dartmouth,  
University of New Hampshire, 02747, MA, USA

<sup>2</sup>NOAA, National Marine Fisheries Service, Woods Hole, MA 02543, USA

<sup>3</sup>School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY  
11794, USA

\* Corresponding authors: Sophie Wulfinf (SophieWulfinf@gmail.com)

# 1 ABSTRACT

2 The US Northeast Continental shelf is the fastest warming marine ecosystem in the United  
3 States, so understanding this change is imperative to maintaining the sustainability and  
4 productivity of this region’s fishery. Fished stocks in the region have exhibited different  
5 distributional shifts in response to changing ocean temperatures, chemistry, and currents.  
6 Management Strategy Evaluation (MSE) is a technique in fisheries management to simulate  
7 and compare the likely outcomes of different management actions, assess trade-offs of dif-  
8 ferent management options, and has been used to create more climate-ready management  
9 plans. The Georges Bank Yellowtail Flounder (*Limanda ferruginea*) is an important stock  
10 to the Northeast groundfish fishery and its recruitment has been shown to be impacted by  
11 bottom ocean temperatures. However, predicting future environmental covariates such as  
12 bottom temperature can be challenging, as these conditions are subject to high stochas-  
13 ticity. For this reason, our goal is to analyze the different choices made when projecting  
14 or forecasting future environmental conditions within assessments, what subsequent catch  
15 advice is generated from the MSE, and how this affects long-term fishery dynamics. Results  
16 focusing on recruitment-environment relationships have shown that different projection de-  
17 cisions, including integrating environmental forecasts, have a limited impact on the model  
18 performance and output, and we will expand on these findings by comparing the MSE results  
19 under different climate scenarios, informed by ocean model forecast products.

## 20 2 INTRODUCTION

21 As the world’s oceans are complex physical and chemical systems, climate change has im-  
22 pacted oceans in a variety of ways such as increased water temperature and acidification,  
23 and changes in ocean currents (Fabry et al., 2008; Guinotte & Fabry, 2008; Abraham et al.,  
24 2013; Sydeman et al., 2015; Wilson et al., 2016; Poloczanska et al., 2016; García Molinos et

al., 2017; Friedland et al., 2022; Cheng et al., 2024). As a result, different marine species are projected to have complex and varying responses to these environmental shifts depending on species' life histories, thermal tolerances, mobility, and other aspects of the species' biology (Guinotte & Fabry, 2008; Chown et al., 2010; Hönisch et al., 2012; Punt et al., 2014b; Stewart et al., 2014a; Sydeman et al., 2015; Poloczanska et al., 2016; Wilson et al., 2016; Chaikin et al., 2024). This creates compounding challenges when managing commercially-important fisheries. As some species will shift ranges northward or to deeper waters in search of colder temperatures, this will disrupt historical food chains as predators and prey may shift in different directions, or range shifts will introduce new interactions between predator and prey species that have historically not overlapped (Overholtz et al., 2011; Albouy et al., 2014; Stewart et al., 2014b; Sydeman et al., 2015; Wilson et al., 2016; García Molinos et al., 2017; Chaikin et al., 2024). Further, as fish populations may experience distribution shifts, annual growth patterns, or vary in abundance, this has potential to increase catch effort as historical fishing grounds and times may now not line up with fish stocks' new life history dynamics (Overholtz et al., 2011; Poloczanska et al., 2016; Tommasi et al., 2017; Fulton, 2021). New species may also shift their ranges into a commonly fished area, potentially opening up new commercial opportunities (Stewart et al., 2014a).

The US Northeast continental shelf has warmed faster than any other marine ecosystem in the United States (V. S. Saba et al., 2016; Kleisner et al., 2017; V. Saba et al., 2023). The varying responses to climate change across fished species has presented challenges in accurately assessing stock health of various species in the region, however these environmental covariates have rarely been incorporated into stock assessments in the Northeast U.S. (Hare et al., 2016; Kleisner et al., 2017; Mazur et al., 2023). These covariates have been shown to reduce retrospective patterns and increase projection accuracy in New England populations such as Atlantic Cod, haddock, yellowtail flounder, and winter flounder (Buckley et al., 2004; Klein et al., 2017). However, incorporating environmental covariates into stock assessments is difficult and a very clear and strong relationship between environmental change and population

dynamics must be shown or else improper use of an environmental covariate could lead to poorer rather than better estimation performance (Haltuch & Punt, 2011a; Punt et al., 2014b; Ehrlén & Morris, 2015; Britten et al., n.d.; Miller et al., 2023a).

Management Strategy Evaluations (MSEs) are decision making frameworks that allow assessors to simulate and compare the sustainability and tradeoffs associated with different management decisions (Hollowed et al., 2011; Tommasi et al., 2017; Mazur et al., 2023; V. Saba et al., 2023). Applying climate change covariates and MSE to New England groundfish has shown to improve the robustness of stock assessments (Mazur et al., 2023). However, applications of MSEs can still be challenging, as climate change shifts population baselines and creates dynamics that are harder to predict in biological systems. This demonstrates a need for further research on how to incorporate these environmental conditions into the MSE framework (Hollowed et al., 2011; Punt et al., 2016; Stram & Holsman, 2022a; V. Saba et al., 2023). Populations' responses to climate change can be complex, as fish exhibit different sensitivities to environmental conditions depending on their lifestage, and different environmental changes (i.e. increased sea surface or bottom temperature, changes in salinity, and altered currents) will have different effects on populations (Hollowed et al., 2011).

Further, decisions on how to project environmental conditions are important to consider when predicting stock dynamics. With recent improvements, ocean dynamic models are able to offer improved predictions of future stock dynamics and are being used in stocks worldwide. However, the spatial and temporal scale of these models is relevant (Hobday et al., 2016). Future climate is also difficult to estimate as there is a lot of uncertainty and climate change is a very chaotic process (Punt et al., 2014b). As assessments attempt to capture future population trends, forecasting environmental trends is also imperative, however this can be challenging as optimal decisions can change depending on scale, species, and which environmental covariates are used (Kell et al., 2005; Hollowed et al., 2011; Tommasi et al., 2017; Stram & Holsman, 2022b). In a simulation-estimation experiment, Britten et al. (2024) found that there was no discernable difference if projections continued the AR1

process or if the environmental covariate projection was calculated as the mean of the last 5 years. Further, Kell et al. (2005) showed that climate change has a limited effect in the short term for Atlantic cod, and only long term simulations showed an affect on stock dynamics and recovery. This project aims to build off of this finding by expanding the projections tested, simulating over a greater number of years, and seeing how catch advice and fishery dynamics may respond to changing projections.

The yellowtail flounder (*Limanda ferruginea*) has a long history of harvest in the US Atlantic Coast. However due to stock collapse in the 1980s, it is now primarily caught as bycatch in other surrounding fisheries (CITE RTA). Yellowtail flounder is managed as three distinct stocks in the Northeast US; Southern New England, Gulf of Maine, and Georges Bank. Yellowtail flounder are sensitive to water temperatures at the larval phase (Laurence & Howell, 1981). The Georges Bank stock has been shown to shift its distribution northward in response to warming ocean temperatures (Murawski, 1993; Nye et al., 2009; Hyun et al., 2014). This stock has been in decline starting in the 1970s, leading to a collapse in the early 1990s (Stone et al., 2004). Recently, ocean bottom temperature has been shown to have a strong link to the recruitment of Georges Bank yellowtail flounder, and the incorporation of this environmental covariate has been shown to improve stock assessment accuracy and prediction success [Johnson et al. (1999); Kerr et al. (2022); CITE RESEARCH TRACK]. Because this relationship between environment and stock dynamics has a clear linkage, inclusion of environmental covariates have improved assessment success, and this stock has a long history of data collection, we will use Georges Bank yellowtail flounder as a case study. We aim to incorporate bottom temperature into stock assessments and the Management Strategy Evaluation framework in order to assess different choices when projecting bottom temperature in Georges Bank and how this affects forecasts in assessments and subsequent tactical fishery management advice of Georges Bank yellowtail flounder.

### 3 METHODS

To conduct the MSE process, we used trawl survey data from the Northeast Fisheries Science Center (NEFSC) and outputs from the Woods Hole Assessment Model (WHAM). WHAM was developed by researchers at NEFSC, and is a state-space stock assessment model which incorporates climate variables into fishery processes (Stock & Miller, 2021). The stock data used was collected by the NEFSC fall and spring surveys, the DFO spring survey, as well as commercial data, which spanned from 1973 to 2022. Choices for weights-at-age, natural mortality, fleet selectivity, recruitment and maturity were derived from results from the 2024 research track assessment on yellowtail flounder (CITE ASSESSMENT, see carvalho et al 2017 from class reading for good presentation of wham choices) and these numbers are [Will make supplemental section with research track info and param ests over time for the](#) available in the supplemental material. The output of the WHAM model was then used in the R package whamMSE (CITE CHENG) to simulate the MSE process. This package creates an Operating Model (OM) which simulates fishery dynamics, including fishing and stock dynamics. As a state space model, the OM incorporates process-error in its dynamics, but the values generated by the OM represent the ‘true’ values of the fishery. The Estimation Model (EM) simulates the survey and assessment process in the fishery, including observation errors. Stock assessments were tested with the 0.75 Fmsy NEFMC control rule. We tested a 50 year MSE period where assessments were conducted every three years.

For the environmental covariate of bottom temperature, we used a dataset developed by Du Pontavice et al. (2023), which is also used in the Georges Bank yellowtail flounder stock assessment CITE ASSESSMENT. We projected the ‘real-world- temperature by conducting a linear regression on the existing data points and then projecting future temperatures along this trendline with added variation around a normal distribution. The package whamMSE was used to conduct the Management Strategy Evaluation CITECHENG. In this simulation, the environmental covariate was treated as a latent variable in the operating model, therefore process error of the ecov was turned off in the OM so only the specified projection was

modeled in the “real world” scenario of our simulation. First, the Wood’s Hole Assessment Model was fitted to real Yellowtail flounder data from the NEFSC survey to serve as the basis for the OM. This base-OM was used to estimate parameters for numbers-at-age, selectivity, recruitment, and ecological covariate. These parameters were then treated as ‘truth’ in our simulation and were used in the second OM we developed for the projections. This second OM, with all of the parameters already estimated, fixes the environmental covariate at a true state with no process error, however observation error is still included in this process. This allowed us to run multiple simulations of various temperature projections in our Estimation Model while maintaining the same ‘real-world’ state.

We then tested various EMs that were identical except for this temperature projection. Instead of using a single time series throughout the projection, all EMs were changed to simulate real-life stock assessment processes, where the three-year projection was recalculated at each assessment year step given the previous three years of data. Descriptions of temperature projections are described in figure 1. For the first experiment, we first compared EM temperature projections with varying levels of bias. We developed four scenarios: High Ecov, where the overall trend of the OM temperature projection was doubled to simulate a high bias in projections; Medium Ecov, where the EM uses the same temperature as the OM; Medium-Low Ecov, where the overall trend was multiplied by -1 to simulate low bias in projections; and Low Ecov, where the overall trend was multiplied by -2 to simulate a very low bias in projections. Each simulation was run 100 times to ensure robustness of model results. Any non-converged model resulted in the elimination of that whole simulation.

This simulation was then repeated with the OM changing it’s baseline projection to use the High, Medium, Medium-Low, and Low temperature projections to compare how hypothetical future climate scenarios may change the outcome of having a biased projection. Each of these OMs were tested against each climate projection in the EMs.

After comparing projections through the MSE process, the experiment was repeated except

Will standardize colors of each em across figures




Figure 1: Temperature projections of the different Bias tests. The medium projection follows the overall trend of historical data, and this was used as a baseline for the other bias projections

I realize now I later switch this with saying the 'good' and 'bad' projections later. I will rephrase

we used only the original OM temperature projection (the one that maintains the trend from the overall data set). For the EMs, the two extreme bias projections were kept (High Ecov and Low Ecov), along with various projection choices that, at each assessment year, self-update with the previous years' temperature information. A new temperature projection is then calculated from that in the EM and projected for the next three years until the next assessment. The different projection choices used in the experiment are described in figure 2.

After the MSE process was conducted in both experiments, we compared the outputs of each model and the subsequent catch advice to emerge from the model. We then compared estimated natural and fishing mortality, spawning stock biomass, and estimated recruitment in each model in order to understand how different assumptions in creating environmental projections propagates reflects real-world dynamics and how this ultimately affects management decisions.

## 4 RESULTS

Before assessing the accuracy of each model, each EM simulated was tested for convergence. If any of the EM models did not converge, the whole seed was removed from the results to ensure comparability across results. Convergence checks were used from the `r` packages `tmb` and `nlminb` CITE PACKAGES.

### 4.1 Bias Tests

The estimated parameters for the environmental covariate are shown in figure 3 compared to that of the OM in each bias test (the black vertical line). Across the EMs, the accuracy of the estimated parameter (in relation to the 'real' one calculated by the OM) was generally higher when projections overestimated the temperature timeseries, and lower for low bias

This feels mess and and too many lines on one page. Let me know if you have ideas on how to make neater. Do we just want to look at projection years?




Figure 2: Temperature projections used in each EM. AR1 (Auto-Regressive1) - This projection uses an updated mean at each timestep for the projection. NewWind (New Window) - This process uses the mean of the previous 5 years in its projection. HistAvg (Historical average) - Uses the average of all datapoints collected. NewTrend (New Trend) - Similar to new window, this process takes the last 5 years of data and uses the linear regression of the last 5 years as its projection. TermYear (Terminal Year) - Uses the terminal year of historical data across the projection period. GoodProj (Good projection) - Uses the same time series as the OM. BadProj (Bad projection) - Same as the MedLow bias in the previous experiment, this projection tests projection with a low bias. NoTemp - (No temperature, not pictured) This projection does not consider environmental covariates in its projection.

179 projections. Observing how this parameter is estimated over time, projections that were  
 180 similar to the OM or had slight bias remained around the calculated OM parameter, while  
 181 the projections with high bias tended to diverge away from this ‘true’ mean (figure 4).  
 182 However, the subsequent catch advice from the EMs, spawning stock biomass (SSB), fishing  
 183 mortality ( $\bar{F}$ ), and predicted catch from the OM stays relatively consistent across EMs over  
 184 the time series. These figures are provided in the supplemental material.




Figure 3: Estimated environmental parameter from each EM compared to the ‘true’ value in the OM on the y axis. Colors correspond to if the EM had a high temperature bias (red), a low temperature bias (blue), or if the EM intended to accurately reflect climate dynamics (i.e. AR1 process or temperature projection that matched that of the OM) (green). The dashed black line indicates where these two estimations would be equal

Is this the appropriate amount to talk about the bias tests because we decided there's not a huge real-world application




Figure 4: Calculated environmental covariates from the EM updated at each assessment period. The “True” OM value is shown in the horizontal dashed line

Table 1: Ecov parameter estimates, standard deviation, and rho for each EM for projection

| EM_name          | OM_ecov | mean_ecov | PercentChange | sd_ecov |
|------------------|---------|-----------|---------------|---------|
| Mod_AR1Ecov      | -0.397  | -0.455    | 14.610        | 0.186   |
| Mod_BadProj      | -0.397  | -0.349    | -12.091       | 0.156   |
| Mod_GoodProj     | -0.397  | -0.278    | -29.975       | 0.138   |
| Mod_HistAvgEcov  | -0.397  | -0.535    | 34.761        | 0.188   |
| Mod_RecTrendEcov | -0.397  | -0.278    | -29.975       | 0.124   |
| Mod_RecWindEcov  | -0.397  | -0.402    | 1.259         | 0.147   |
| Mod_TermYear     | -0.397  | -0.326    | -17.884       | 0.134   |

Will fix labels

## 4.2 Projection Tests

When comparing the parameter for the environmental covariate across different projections (table 1), the most accurate estimation of the ecological covariate was the projection that used recent window approach, with only a 1.3% change from the ‘true’ value. Both the recent trend projection and the ‘good’ projection estimated this parameter poorly, with an almost 30% difference between the true covariate and the estimation in the EM. The calculated parameter estimates at each assessment year show that the Historical average and AR1 process tended to underestimate the ecological covariate while the other projections generally over-estimated this trend (figure 5). However, figure 6 shows the catch advice, SSB,  $\bar{F}$ , and predicted catch from the OM of each projection test. Similar to the bias tests, these model outputs were very similar and all converged to similar estimations.

Differences in estimated recruitment between the EM and OM is shown in figure 7, where there is noticeable variation between the EM estimations and that of the OM but for most years, the EMs tended to under estimate recruitment compared to that of the OM. Comparing the calculated variation in recruitment (figure 8) between models that used an ecological covariate and the No Ecov projection of the same seed, we can estimate the percentage of this variation that is captured in the ecological covariate. All models had a mean percentage of variation between 5% and 10%. The AR1 process and the ‘Bad’ projection both had the highest range of recruitment variations, with the AR1 process reaching a maximum of 20.6




Figure 5: Estimated ecological covariates in for the different temperature projection choices in the EM compared to the “true” value in the OM.

I removed the ribbons on these graphs with error because it got messy and some catches had SEs that made it below 0. Let me know if you want me to put ribbons back in because you specifically said you wanted them




Figure 6: Catch advice (top), SSB (second), FBar (third), and predicted catch (bottom) calculated from each EM over the projection period

204 % of variation in recruitment being explained by the ecological covariate. The Recent Trend  
 205 and ‘Good’ projection displayed the smallest range of variation that was captured by the  
 206 ecological covariate.




Figure 7: Percent difference of the calculated recruitment per assessment year in each EM compared to the OM.

207 Looking at the stock status, all of the EMs under-estimated the yellowtail’s overfished status,  
 208 but all models projected that this stock would remain overfished for most of the projection  
 209 period (figure 9). Similarly, the EMs also under- estimated the overfishing status, but this  
 210 varied between status being over and under the overfishing threshold (the red horizontal line)  
 211 over the course of the time series. In this case, each EM overestimated the FXSPR reference  
 212 point when compared to the reference point used in the operating model (figure 10).

213 In the model performance (figure 11), however, these differences have a minimal impact on  
 214 the overall accuracy of these models. With the exception of the model without an ecological




Figure 8: Percent difference of the recruitment deviation per assessment year in each EM compared to the OM.

Want thoughts on this figure (see meeting notes I sent you)




Figure 9: Overfished (left) and overfishing (right) status over the projected time series for each EM. Overfished and overfishing thresholds are shown in the horizontal red line




Figure 10: Percent difference of the calculated FXSPR reference point in each EM compared to the OM.

fix caption. This is not % diff but % variation explained by ecov

covariate and the model with a biased projection, all models performed similarly when comparing SSB, Catch, and  $\bar{F}$  of the first and terminal years of the projection. The calculated SSB,  $(\bar{F})$ , and predicted catch from the terminal three years of the projection period show similar patterns of similarity across EMs (figure 12).

## 5 DISCUSSION

The results from the bias test show that, while incorporating the ecological covariate helps model performance overall, the trajectory of the projection of temperature used in predicting future simulations will predictably under or over estimate the ecological parameter depending on the type of bias. However, the overall result had a limited impact on the SSB,  $\bar{F}$ , and subsequent catch. This is because regardless of the bias in the projection, indicating that bottom temperature's overall affect on recruitment may have a limited impact on the status of the fishery over time. Kell et al. (2005) found that the specific climate scenario modeled in the OM had a limited affect on the recoverability of Atlantic cod stocks.

However, this test was more of a hypothetical exploration of how diverging temperature projections changes the outputs of the model. In reality, stock assessments use hindcast environmental information, so it is not realistic that an assessment would continue to use a temperature trajectory that diverges away from the real environmental conditions. Instead, stock assessments incorporate the most recent data and use this information to make predictions until the next assessment. This is why our projections test updated its temperature predictions with the previous three years of information. Similar to the bias tests, we found that incorporating the environmental covariate resulted in slightly different dynamics, but overall followed similar trajectories. Some models appeared to do better with some metrics (e.g. recent window projections resulted in the most accurate ecological covariate, whereas the 'good' projection performed the most poorly here, it estimated the smallest range of recruitment deviations being explained by the ecological covariate), but this had a limited




Figure 11: Relative performance of each EM

Will change this to terminal year like we talked




Figure 12: Fbar, SSB, and catch of terminal 3 years of MSE in projections test

impact in the overall fishery dynamics, even in the long term. This is in line with previous literature for yellowtail flounder, which has shown that there is little discernible difference between projections that used a continued estimated environmental process, ones that fixed the environmental process at recent 5 year average, and a third EM which used the observed environmental index (Britten et al., 2024). This research also noted that the specific stock-recruitment relationship defined in the model will change the environmental affect size due to the placement of the environmental covariate in the stock-recruit equation. McClatchie et al. (2010) found that for Pacific sardine, *Sardinops sagax*, temperature actually did not have a relationship with recruitment, and the correlation between temperature parameters and recruitment depended on the amount of autocorrelation in the temperature time series, although the specific mechanism that links Pacific sardine recruitment is less clear than with yellowtail flounder. This may also be due to the amount of recruitment variation that was explained by the ecological covariate. In these simulations, the percentage of variation explained by the ecological covariate had a maximum of 20.6% in one simulation, yet De Oliveira & Butterworth (2005) found that management strategies do not show any benefits to risk and average catch if the ecological covariate explains less than 50% of recruitment variation. None of the simulations in this experiment exceeded 50%.

This could be the result of the type of link that the environmental covariate makes with the population. For Georges Bank Yellowtail, the environmental covariate has a lagged controlling effect on recruitment (CITE RT). Over time, the negative effects of high temperature could be masked by density-dependent mortality, or other factors that occur during the lag.

Kell et al. (2005) found that the long-term performance and calculated reference points of Atlantic cod stock depended on if ocean temperatures were linked to juvenile survival versus the carrying capacity of the stock. Further, there may be a greater difference in model outputs if the environmental covariate had a direct effect on distribution, carrying capacity, catchability, mortality, behavioral, migratory, spawning, or metabolic processes; life history traits that can all be affected by environmental change (Punt et al., 2014a; Stock & Miller, 2021), indicating a need for further research on projection choices for environmental covariates that affect other parts of life history. Catch advice is not very dependent on recruitment, and because this environmental link is also lagged, these combined factors could be masking the effect. Further, Haltuch & Punt (2011b) has shown that incorrectly including environmental covariates in the stock does not have a large effect on biomass reference points or relative errors of the model. A review by Punt et al. (2014a) has also shown that short and medium-term fisheries management simulations are generally not greatly improved by incorporating environmental covariates. This research served to test a longer-term simulation and found a similar result. However, if there is a high temporal contrast in the ‘real world’ or OM, incorporating the environmental covariate in the estimation model can improve model accuracy (Miller et al., 2023b). This could indicate that these results may change depending on the amount of future variation in ocean bottom temperatures.

Future ocean conditions are expected to be highly variable in coming years due to climate change, (Wang et al., 2010), so understanding potential impacts on fishery dynamics is vital. Overall, we found that projection decisions had a limited effect on catch advice and fishery dynamics in an MSE framework, and this could be due to the nature of the environmental link to Georges Bank Yellowtail flounder, indicating a need for further research on how this type of link will affect model dynamics over different projection decisions.

WOULD THE DIFFERENCE BETWEEN REC TREND AND REC WINDOW BE  
[whoop ignore this las bit. Notes for me, I thought they were commented out](#)  
 MORE ABOUT THE GENERAL TREND OF THE PROJECTION (I.E. RECENTLY  
 INCREASING THEREFORE REC TREND WITH GENERALLY INCREASE?) WHAT

288 IS THE RECRUITMENT DEVIATIONS GRAPH TELLING UUUUUU gET ADVICE  
289 ON TEH REF POINTS FIGURE GENERALLY UNDER EST RECRUITMENT AND  
290 STOCK STATUS

## References

- Abraham, J. P., Baringer, M., Bindoff, N. L., Boyer, T., Cheng, L. J., Church, J. A., Conroy, J. L., Domingues, C. M., Fasullo, J. T., Gilson, J., Goni, G., Good, S. A., Gorman, J. M., Gouretski, V., Ishii, M., Johnson, G. C., Kizu, S., Lyman, J. M., Macdonald, A. M., . . . Willis, J. K. (2013). A review of global ocean temperature observations: Implications for ocean heat content estimates and climate change. *Reviews of Geophysics*, 51(3), 450–483. <https://doi.org/10.1002/rog.20022>
- Albouy, C., Velez, L., Coll, M., Colloca, F., Le Loc’h, F., Mouillot, D., & Gravel, D. (2014). From projected species distribution to food-web structure under climate change. *Global Change Biology*, 20(3), 730–741. <https://doi.org/10.1111/gcb.12467>
- Britten, G. L., Brooks, E., & Miller, T. (n.d.). *Identification and performance of stock-recruitment functions in state space assessment models*.
- Britten, G. L., Brooks, E., & Miller, T. (2024). *Identification and performance of stock-recruitment functions in state space assessment models*. NEFSC.
- Buckley, L. J., Caldarone, E. M., & Lough, R. G. (2004). Optimum temperature and food-limited growth of larval Atlantic cod ( *Gadus morhua* ) and haddock ( *Melanogrammus aeglefinus* ) on Georges Bank. *Fisheries Oceanography*, 13(2), 134–140. <https://doi.org/10.1046/j.1365-2419.2003.00278.x>

- 309 Chaikin, S., Riva, F., Marshall, K. E., Lessard, J.-P., & Belmaker, J. (2024). Marine fishes  
310 experiencing high-velocity range shifts may not be climate change winners. *Nature Ecology & Evolution*, 8(5), 936–946. <https://doi.org/10.1038/s41559-024-02350-7>
- 312 Cheng, L., Abraham, J., Trenberth, K. E., Boyer, T., Mann, M. E., Zhu, J., Wang, F., Yu,  
313 F., Locarnini, R., Fasullo, J., Zheng, F., Li, Y., Zhang, B., Wan, L., Chen, X., Wang, D.,  
314 Feng, L., Song, X., Liu, Y., . . . Lu, Y. (2024). New Record Ocean Temperatures and  
315 Related Climate Indicators in 2023. *Advances in Atmospheric Sciences*, 41(6), 1068–  
316 1082. <https://doi.org/10.1007/s00376-024-3378-5>
- 317 Chown, S., Hoffmann, A., Kristensen, T., Angilletta, M., Stenseth, N., & Pertoldi, C. (2010).  
318 Adapting to climate change: a perspective from evolutionary physiology. *Climate Re-*  
319 *search*, 43(1), 3–15. <https://doi.org/10.3354/cr00879>
- 320 De Oliveira, J., & Butterworth, D. (2005). Limits to the use of environmental indices to  
321 reduce risk and/or increase yield in the South African anchovy fishery. *African Journal*  
322 *of Marine Science*, 27(1), 191–203. <https://doi.org/10.2989/18142320509504078>
- 323 Du Pontavice, H., Chen, Z., & Saba, V. S. (2023). A high-resolution ocean bottom tem-  
324 perature product for the northeast U.S. continental shelf marine ecosystem. *Progress in*  
325 *Oceanography*, 210, 102948. <https://doi.org/10.1016/j.pocean.2022.102948>

- 326 Ehrlén, J., & Morris, W. F. (2015). Predicting changes in the distribution and abundance of  
327 species under environmental change. *Ecology Letters*, 18(3), 303–314. [https://doi.org/](https://doi.org/10.1111/ele.12410)  
328 10.1111/ele.12410
- 329 Fabry, V. J., Seibel, B. A., Feely, R. A., & Orr, J. C. (2008). Impacts of ocean acidification on  
330 marine fauna and ecosystem processes. *ICES Journal of Marine Science*, 65(3), 414–432.  
331 <https://doi.org/10.1093/icesjms/fsn048>
- 332 Friedland, K. D., Miles, T., Goode, A. G., Powell, E. N., & Brady, D. C. (2022). The Middle  
333 Atlantic Bight Cold Pool is warming and shrinking: Indices from in situ autumn seafloor  
334 temperatures. *Fisheries Oceanography*, 31(2), 217–223. [https://doi.org/10.1111/fog.](https://doi.org/10.1111/fog.12573)  
335 12573
- 336 Fulton, E. A. (2021). Opportunities to improve ecosystem-based fisheries management by  
337 recognizing and overcoming path dependency and cognitive bias. *Fish and Fisheries*,  
338 22(2), 428–448. <https://doi.org/10.1111/faf.12537>
- 339 García Molinos, J., Burrows, M. T., & Poloczanska, E. S. (2017). Ocean currents modify  
340 the coupling between climate change and biogeographical shifts. *Scientific Reports*, 7(1),  
341 1332. <https://doi.org/10.1038/s41598-017-01309-y>
- 342 Guinotte, J. M., & Fabry, V. J. (2008). Ocean Acidification and Its Potential Effects on

Marine Ecosystems. *Annals of the New York Academy of Sciences*, 1134(1), 320–342.

<https://doi.org/10.1196/annals.1439.013>

Haltuch, M. A., & Punt, A. E. (2011a). The promises and pitfalls of including decadal-scale climate forcing of recruitment in groundfish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences*, 68(5), 912–926. <https://doi.org/10.1139/f2011-030>

Haltuch, M. A., & Punt, A. E. (2011b). The promises and pitfalls of including decadal-scale climate forcing of recruitment in groundfish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences*, 68(5), 912–926. <https://doi.org/10.1139/f2011-030>

Hare, J. A., Morrison, W. E., Nelson, M. W., Stachura, M. M., Teeters, E. J., Griffis, R. B., Alexander, M. A., Scott, J. D., Alade, L., Bell, R. J., Chute, A. S., Curti, K. L., Curtis, T. H., Kircheis, D., Kocik, J. F., Lucey, S. M., McCandless, C. T., Milke, L. M., Richardson, D. E., ... Griswold, C. A. (2016). A Vulnerability Assessment of Fish and Invertebrates to Climate Change on the Northeast U.S. Continental Shelf. *PLOS ONE*, 11(2), e0146756. <https://doi.org/10.1371/journal.pone.0146756>

Hobday, A. J., Spillman, C. M., Paige Eveson, J., & Hartog, J. R. (2016). Seasonal forecasting for decision support in marine fisheries and aquaculture. *Fisheries Oceanography*, 25(S1), 45–56. <https://doi.org/10.1111/fog.12083>

- 360 Hollowed, A., A'mar, T., Barbeaux, S., Bond, N., Ianelli, J., Spncer, P., & Wilderbuer,  
361 T. (2011). *Integrating Ecosystem Aspects and Climate Change Forecasting into Stock*  
362 *Assessments*.
- 363 Hönisch, B., Ridgwell, A., Schmidt, D. N., Thomas, E., Gibbs, S. J., Sluijs, A., Zeebe, R.,  
364 Kump, L., Martindale, R. C., Greene, S. E., Kiessling, W., Ries, J., Zachos, J. C., Royer,  
365 D. L., Barker, S., Marchitto, T. M., Moyer, R., Pelejero, C., Ziveri, P., . . . Williams, B.  
366 (2012). The Geological Record of Ocean Acidification. *Science*, *335*(6072), 1058–1063.  
367 <https://doi.org/10.1126/science.1208277>
- 368 Hyun, S.-Y., Cadrin, S. X., & Roman, S. (2014). Fixed and mixed effect models for fishery  
369 data on depth distribution of Georges Bank yellowtail flounder. *Fisheries Research*, *157*,  
370 180–186. <https://doi.org/10.1016/j.fishres.2014.04.010>
- 371 Johnson, D., Morse, W., Berrien, P., & Vitaliano, J. (1999). *Yellowtail Flounder, Limanda*  
372 *ferruginea, Life History and Habitat Characteristics* (NOAA Technical Memorandum  
373 NMFS-NE-140). US Department of Commerce.
- 374 Kell, L. T., Pilling, G. M., & O'Brien, C. M. (2005). Implications of climate change for the  
375 management of North Sea cod (*Gadus morhua*). *ICES Journal of Marine Science*, *62*(7),  
376 1483–1491. <https://doi.org/10.1016/j.icesjms.2005.05.006>

Kerr, L., Barajas, M., & Wiedenmann, J. (2022). Coherence and potential drivers of stock assessment uncertainty in Northeast US groundfish stocks. *ICES Journal of Marine Science*, 79(8), 2217–2230. <https://doi.org/10.1093/icesjms/fsac140>

Klein, E. S., Smith, S. L., & Kritzer, J. P. (2017). Effects of climate change on four New England groundfish species. *Reviews in Fish Biology and Fisheries*, 27(2), 317–338. <https://doi.org/10.1007/s11160-016-9444-z>

Kleisner, K. M., Fogarty, M. J., McGee, S., Hare, J. A., Moret, S., Perretti, C. T., & Saba, V. S. (2017). Marine species distribution shifts on the U.S. Northeast Continental Shelf under continued ocean warming. *Progress in Oceanography*, 153, 24–36. <https://doi.org/10.1016/j.pocean.2017.04.001>

Laurence, G., & Howell, W. H. (1981). Embryology and influence of temperature and salinity on early development and survival of yellowtail flounder *limanda ferruginea*. *Marine Ecology Progress Series*, 6, 11–18.

Mazur, M. D., Jesse, J., Cadrin, S. X., Truesdell, S. B., & Kerr, L. (2023). Consequences of ignoring climate impacts on New England groundfish stock assessment and management. *Fisheries Research*, 262, 106652. <https://doi.org/10.1016/j.fishres.2023.106652>

McClatchie, S., Goericke, R., Auad, G., & Hill, K. (2010). Re-assessment of the stock–recruit

and temperature–recruit relationships for Pacific sardine (*Sardinops sagax*). *Canadian Journal of Fisheries and Aquatic Sciences*, 67(11), 1782–1790. <https://doi.org/10.1139/F10-101>

Miller, T. J., Applegate, A., Britten, G., Brooks, E. N., Fay, G., Hansell, A., Legault, C. M., Muffley, B., Stock, B. C., & Wiedenmann, J. (2023a). *Factors affecting inferences on natural mortality and associated environmental effects*.

Miller, T. J., Applegate, A., Britten, G., Brooks, E. N., Fay, G., Hansell, A., Legault, C. M., Muffley, B., Stock, B. C., & Wiedenmann, J. (2023b). *Factors affecting inferences on natural mortality and associated environmental effects* [Research {{Track}}]. NEFSC.

Murawski, S. A. (1993). Climate Change and Marine Fish Distributions: Forecasting from Historical Analogy. *Transactions of the American Fisheries Society*, 122(5), 647–658. [https://doi.org/10.1577/1548-8659\(1993\)122%3C0647:CCAMFD%3E2.3.CO;2](https://doi.org/10.1577/1548-8659(1993)122%3C0647:CCAMFD%3E2.3.CO;2)

Nye, J., Link, J., Hare, J., & Overholtz, W. (2009). Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf. *Marine Ecology Progress Series*, 393, 111–129. <https://doi.org/10.3354/meps08220>

Overholtz, W. J., Hare, J. A., & Keith, C. M. (2011). Impacts of Interannual Envi-

ronmental Forcing and Climate Change on the Distribution of Atlantic Mackerel on  
the U.S. Northeast Continental Shelf. *Marine and Coastal Fisheries*, 3(1), 219–232.  
<https://doi.org/10.1080/19425120.2011.578485>

Poloczanska, E. S., Burrows, M. T., Brown, C. J., García Molinos, J., Halpern, B. S.,  
Hoegh-Guldberg, O., Kappel, C. V., Moore, P. J., Richardson, A. J., Schoeman, D. S.,  
& Sydeman, W. J. (2016). Responses of Marine Organisms to Climate Change across  
Oceans. *Frontiers in Marine Science*, 3. <https://doi.org/10.3389/fmars.2016.00062>

Punt, A. E., A’mar, T., Bond, N. A., Butterworth, D. S., De Moor, C. L., De Oliveira, J.  
A. A., Haltuch, M. A., Hollowed, A. B., & Szuwalski, C. (2014a). Fisheries management  
under climate and environmental uncertainty: Control rules and performance simulation.  
*ICES Journal of Marine Science*, 71(8), 2208–2220. <https://doi.org/10.1093/icesjms/fst057>

Punt, A. E., A’mar, T., Bond, N. A., Butterworth, D. S., De Moor, C. L., De Oliveira, J.  
A. A., Haltuch, M. A., Hollowed, A. B., & Szuwalski, C. (2014b). Fisheries management  
under climate and environmental uncertainty: control rules and performance simulation.  
*ICES Journal of Marine Science*, 71(8), 2208–2220. <https://doi.org/10.1093/icesjms/fst057>

Punt, A. E., Butterworth, D. S., De Moor, C. L., De Oliveira, J. A. A., & Haddon, M.  
(2016). Management strategy evaluation: best practices. *Fish and Fisheries*, 17(2),

303–334. <https://doi.org/10.1111/faf.12104>

Saba, V. S., Griffies, S. M., Anderson, W. G., Winton, M., Alexander, M. A., Delworth, T. L., Hare, J. A., Harrison, M. J., Rosati, A., Vecchi, G. A., & Zhang, R. (2016). Enhanced warming of the Northwest Atlantic Ocean under climate change. *Journal of Geophysical Research: Oceans*, 121(1), 118–132. <https://doi.org/10.1002/2015JC011346>

Saba, V., Borggaard, D., Caracappa, J. C., Chambers, R. C., Clay, P. M., Colburn, L. L., Deroba, J., DePiper, G., Du Pontavice, H., Fratantoni, P., Ferguson, M., Gaichas, S., Hayes, S., Hyde, K., Johnson, M., Kocik, J., Keane, E., Kircheis, D., Large, S., . . . Wikfors, G. (2023). NOAA fisheries research geared towards climate-ready living marine resource management in the northeast United States. *PLOS Climate*, 2(12), e0000323. <https://doi.org/10.1371/journal.pclm.0000323>

Stewart, J. S., Hazen, E. L., & Bograd, S. J. (2014a). Combined climate- and prey-mediated range expansion of Humboldt squid (*Dosidicus gigas*), a large marine predator in the California Current System. *Global Change Biology*.

Stewart, J. S., Hazen, E. L., & Bograd, S. J. (2014b). Combined climate- and prey-mediated range expansion of Humboldt squid (*Dosidicus gigas*), a large marine predator in the California Current System. *Global Change Biology*.

- 447 Stock, B. C., & Miller, T. J. (2021). The Woods Hole Assessment Model (WHAM): A general  
448 state-space assessment framework that incorporates time- and age-varying processes via  
449 random effects and links to environmental covariates. *Fisheries Research*, 240, 105967.  
450 <https://doi.org/10.1016/j.fishres.2021.105967>
- 451 Stone, H. H., Gavaris, S., Legault, C. M., Neilson, J. D., & Cadrin, S. X. (2004). Collapse  
452 and recovery of the yellowtail flounder (*Limanda ferruginea*) fishery on Georges Bank.  
453 *Journal of Sea Research*, 51(3-4), 261–270. <https://doi.org/10.1016/j.seares.2003.08.004>
- 454 Stram, D., & Holsman, K. (2022b). *Climate Readiness Synthesis 2022* (p. 28).
- 455 Stram, D., & Holsman, K. (2022a). *Climate Readiness Synthesis 2022* (p. 28). NPFMC.
- 456 Sydeman, W. J., Poloczanska, E., Reed, T. E., & Thompson, S. A. (2015). Climate change  
457 and marine vertebrates. *Oceans and Climate*, 350.
- 458 Tommasi, D., Stock, C. A., Hobday, A. J., Methot, R., Kaplan, I. C., Eveson, J. P., Holsman,  
459 K., Miller, T. J., Gaichas, S., Gehlen, M., Pershing, A., Vecchi, G. A., Msadek, R.,  
460 Delworth, T., Eakin, C. M., Haltuch, M. A., Séférian, R., Spillman, C. M., Hartog, J. R.,  
461 ... Werner, F. E. (2017). Managing living marine resources in a dynamic environment:  
462 The role of seasonal to decadal climate forecasts. *Progress in Oceanography*, 152, 15–49.  
463 <https://doi.org/10.1016/j.pocean.2016.12.011>

- 464 Wang, M., Overland, J. E., & Bond, N. A. (2010). Climate projections for selected large  
465 marine ecosystems. *Journal of Marine Systems*, 79(3-4), 258–266. [https://doi.org/10.](https://doi.org/10.1016/j.jmarsys.2008.11.028)  
466 1016/j.jmarsys.2008.11.028
- 467 Wilson, L. J., Fulton, C. J., Hogg, A. M., Joyce, K. E., Radford, B. T. M., & Fraser,  
468 C. I. (2016). Climate-driven changes to ocean circulation and their inferred impacts  
469 on marine dispersal patterns. *Global Ecology and Biogeography*, 25(8), 923–939. [https:](https://doi.org/10.1111/geb.12456)  
470 [//doi.org/10.1111/geb.12456](https://doi.org/10.1111/geb.12456)