

# Social-ecological models with social hierarchy and space applied to small scale fisheries

SOPHIE WULFING<sup>1\*</sup> AND EASTON R. WHITE<sup>1</sup>

<sup>1</sup>Department of Biological Sciences, University of New Hampshire, 03824, NH, USA

\* Corresponding authors: Sophie Wulfinf (SophieWulfinf@gmail.com) and Dr. Easton White (Easton.White@unh.edu)

## 1 ABSTRACT

Socio-ecological models combine models of ecological systems with human social dynamics in order to better understand human interaction with the environment. One such model of human behavior is replicator dynamics. Replicator dynamics are derived from evolutionary game theory and model how societal influence can change opinions about resource extraction. Previous research on replicator dynamics have shown how evolving opinions on conservation can change how humans interact with their environment and therefore change population dynamics of the harvested species. However, these models have all assumed that human societies are homogenous with no social structure. In this study, we develop a two-patch socio-ecological model with social hierarchy in order to study the effect that social inequity has on decision making. We also analyzed the spatial components of this two-patch model and observed the effect of fish movement on decision making and fish population dynamics. We found that, contrary to our hypothesis, social influences became less significant in our two patch model. Instead, we found that fish movement across patches was a major driver of changes to population dynamics. This indicates the importance of including spatial components to socio-ecological models. Further, this study highlights the importance of understanding species movements when making conservation decisions.

## 2 INTRODUCTION

The study of social ecological models is a growing field in ecology as they treat human behavior as a variable as opposed to a set parameter. This allows for the study of how human decision making can change in response to environmental factors and in turn, change how humans interact with resources and profits (Chris T. Bauch 2005; Innes, Anand, and Bauch 2013; Oraby, Thampi, and Bauch 2014; Chris T. Bauch et al. 2016; Sigdel, Anand, and Bauch 2017; Thampi, Anand, and Bauch 2018). As human societies grow

increasingly intricate and interconnected, these models can help us to analyze how our social structures can influence the environment around us (Liu et al. 2007). These models provide important insight not only into how human decision making can influence the ecological processes but it can also show hidden processes, reveal regime shifts that would otherwise be hidden, and identify vulnerabilities of systems that don't exist within the purely social or ecological models (Liu et al. 2007; Young et al. 2007; Lade et al. 2013). Socio-ecological models have even showed different dynamics at different scales and different amounts of human connectivity (Cumming, Morrison, and Hughes 2017). They can also be utilized in systems where data is difficult to collect, as parameters can be changed in order to analyze different hypothetical scenarios. Socio-ecological models can also inform effective policy decisions. Conservation plans often do not reach their conservation goals, and these setbacks are often attributed to a lack of stakeholder participation (Crona and Bodin 2006; Salas, Barragán-Paladines, and Chuenpagdee 2019; Prince et al. 2021). This can be due to an emergence of conflict for stakeholders, where the conservation plan in place directly hinders their practices, therefore deterring them from participating in the restorative efforts. Socio-ecological models can identify where these areas of potential conflict can arise, compromises that can be made in the system, and alternative conservation practices that encourages participation from all stakeholder groups (Ban et al. 2013). Further, as these models are simulations of human and environmental interactions, they allow flexibility in that they can be adapted to fit the specific system of study and improve place-based management practices (Young et al. 2007; Liu et al. 2007; Felipe-Lucia et al. 2022)

Due to their adaptability, socio-ecological models can use a wide range of strategies to represent human decision making. One such method is replicator dynamics, which model human decision making where an individual makes conservation decisions based on weighing the perceived benefits of conservation with the costs as well as the social pressure to conform to the group stance on conservation. Individuals will therefore “replicate” the behavior of their peers by changing their harvest practices based on the opinion of the majority (Chris T. Bauch and Bhattacharyya 2012). They have been used to show how social learning is a key component to vaccination uptake public health, and preexisting social norma can actually suppress vaccine uptake despite frequent disease outbreaks (Chris T. Bauch and Bhattacharyya 2012; Oraby, Thampi, and Bauch 2014). They can also have conservation applications as pest invasion models have shown ways to simultaneously mitigate pest outbreaks and the cost to address them in the timber industry (Barlow et al. 2014). Further, land use changes have been modeled to have completely different dynamics when human decision making was added to these models (Innes, Anand, and Bauch 2013). However, all previous models of human behavior have assumed that human societies are homogenous, and all people are subject to the same social influence and dynamics. No replicator-dynamics model has incorporated social inequality or

hierarchy, despite the fact the most human societies have varying levels of social influence within them.

Contrary to this assumption made by previous models that human groups are homogenous, the vast majority of real-world societies exhibit some form of hierarchy or inequality. Societies with different social subgroups can often exhibit an “us vs. them” mentality and compete for resources (Borgatti 2003). This is because social status can greatly alter peoples’ interaction with the environment. Competition over resources has been shown to be exacerbated by social hierarchies and ‘top-down’ regulation whereas when social connectivity is considered in management plans, this has been shown to not only improve management outcomes, but reduce costs as well (Krackhardt and Stern 1988; Grafton 2005; Bodin and Crona 2009). Further, members of social networks have been shown to have varying levels of connectivity with others based on attributes such as ethnicity, and this can in turn alter an individual’s relationship with the environment and their views on conservation (Barnes-Mauthe 2013). Barnes-Mauthe (2013) also showed that fishing communities can exhibit homophily, which is the tendency for people to obtain information and opinions from those who are similar to themselves before seeking views from those who are perceived as different. Therefore, people in different social groups may be receiving different information and opinions about conservation and acting accordingly (McPherson, Smith-Lovin, and Cook 2001). For example in Kenya, communication among fishers has been shown to be among fishers using the same gear type which has inhibited successful regulation of the whole fishery (Crona and Bodin 2006). Further, in the Southwest Madagascar octopus fishery, fishing method and location typically falls along gendered lines. When fishing restrictions were imposed on tidal flats, this affected fishing access for women while maintaining this livelihood for male fishers (Baker-Médard 2017). In Thailand, ethnicity has been shown to be a source of fishing conflict which has exacerbated resource depletion (Pomeroy et al. 2007). The existence of social structures is extremely prevalent in human societies and this has been shown to alter how people interact with the environment. However, there has been no previous replicator dynamics study that considers how social hierarchies alters harvest practices.

Small scale fisheries are a particularly relevant system to apply replicator dynamics as fishing practices and policies are often made by communal decision makers. Research on small-scale fisheries is a growing and essential field as they are drastically understudied yet affect many people around the globe. Worldwide, about 32 million fishers make their livelihood in small-scale fisheries, a subsector in which 90 to 95% of fish is distributed for local consumption. These marine products are a vital source of nutrition for these communities (“HIDDEN HARVEST-The Global Contribution of Capture Fisheries” 2012). Due to tight social structures, community decision making, and strong reliance on the environment, small scale fisheries are systems that are well represented by socio ecological models and replicator dynamics (Grafton 2005; Thampi, Anand, and Bauch 2018; Barnes et al. 2019). Conservation efforts in small scale fisheries have

often been unsuccessful, especially when the social and economic components of the industry have been ignored (Salas, Barragán-Paladines, and Chuenpagdee 2019; Prince et al. 2021). However, even when human interactions and decision making have been considered, socio-ecological models have often treated individuals in human societies homogeneous. As human societies are often complex and hierarchical, this simplifying assumption that everyone interacts with the environment and within their community equally can lead to lack of participation in conservation by some groups within a community (Barnes-Mauthe 2013; Cumming, Morrison, and Hughes 2017). Mismanagement of fisheries have even been shown to exacerbate these inequalities (Cinner et al. 2012; Baker-Médard 2017). Further, the specific dynamics of the fishery in question have been shown to be an important component to models, as models with multiple patches can actually mitigate overfishing if there is a high migratory ability of the harvested species (Cressman, Krivan, and Garay 2004).

Instituting effective conservation strategies can be especially difficult if the organism being protected has a migratory pattern that crosses over multiple management jurisdictions such as country borders (Ogburn et al. 2017; Garrone-Neto et al. 2018; Ramírez-Valdez et al. 2021). Borders can also create challenges when gathering population data that requires extensive fieldwork (Cozzi et al. 2020; Hebblewhite and Whittington 2020). The fragmentation of management can also result in a mismatch of conservation strategies that become ineffective when these management bodies do not coordinate efforts (Siddons, Pegg, and Klein 2017). Research on the importance of coordinated research efforts has been conducted on many species of terrestrial animals with large migratory ranges and have consistently shown that cooperation among government bodies is essential to protecting population health of highly migratory species or species whose native ranges expand across multiple countries (Plumptre et al. 2007; Gervasi et al. 2015; Meisingset et al. 2018). Because fish are generally highly migratory, this issue is especially relevant in international waters or waters where different government bodies share jurisdiction (Mchich, Auger, and Raïssi 2000). For this reason, research on two patch fishing models is a commonly used method as different management strategies can be modeled in each patch. Previous research on two-patch fishing models has shown that movement rates between patches can effect population stability when there is different fishing pressures in each patch (Mchich, Auger, and Raïssi 2000; Cai, Li, and Song 2008). Economic output can also be maximized in multi-patch fishing models as high dispersal can result in a higher overall yield of the system than the yield of each patch combined (Auger, Kooi, and Moussaoui 2022). High dispersal across patches is commonly found to be an essential component to maximizing population health and economic gain from fishing (Freedman and Waltman 1977; Moeller and Neubert 2015; Auger, Kooi, and Moussaoui 2022). Two patch models help us to understand better the population dynamics of fish species who face different pressures in each patch and

118 have even resolved conflicts between fishing groups (Mchich, Auger, and Raïssi 2000). However, no previous  
 119 research has combined two patch fishing models with a hierarchical human decision making model in order  
 120 to study how space and social dynamics affect fishery dynamics.

121 In this study, we couple a human-decision replicator dynamics model with social hierarchies with a two-  
 122 patch fishing model in order to understand how decision making. The objectives of this study are: 1) to  
 123 compare the output of this model with that of previous replicator dynamics studies without spatial or social  
 124 hierarchical components, 2) Find the effects of social hierarchies in decision making and how that effects  
 125 fishing dynamics 3) determine the significance of fish dispersal in our two patch model.

### 126 3 METHODS




Figure 1: A conceptual representation of our model as a two-patch extension of Chris T. Bauch et al. (2016). Here, fish populations ( $F_1$  and  $F_2$ ) in each patch increase through natural growth and movement of fish into the patch. Fish populations are decreased through emigration out of the patch and human fishing activity. The number of fishers ( $X_1$  and  $X_2$ ) change in response to fish population levels, the cost of stopping fishing activity, and the opinions of those in the patch and those in the other.

### 3.1 Model Construction

We build on the work of Chris T. Bauch et al. (2016) by extending their approach to a two-patch model (Figure 1). The fish population models are as follows:

$$\frac{dF_1}{dt} = r_1 F_1 (1 - F_1) - \frac{h_1 * F_1}{F_1 + s_1} - i_2 F_1 + i_1 F_2 \quad (1)$$

$$\frac{dF_2}{dt} = r_2 F_2 (1 - F_2) - \frac{h_2 * F_2}{F_2 + s_2} - i_1 F_2 + i_2 F_1 \quad (2)$$

Where the change in fish populations is dependent on  $r$ , the net population growth of each patch, and both populations follow logistic growth. The second term:  $\frac{h * F}{F + s}$ , denotes fish lost to human activity.  $h$  is the harvesting efficiency of the respective human population and  $s$  controls the supply and demand of the fishery. Because we are now working with a two-patch model, the  $i$  parameter denotes the immigration out of each patch and into the other. In this study, we are assuming a closed population between the two patches. Therefore, fish move directly from patch to patch and do not disperse elsewhere.

For the model of human activity and opinion, we used replicator dynamics from evolutionary game theory to simulate societal influence on an individual's opinion. Social dynamics are represented by the proportion of conservationists in a population ( $X$ ) and the proportion of harvesters ( $1 - X$ ). These two groups interact with one another using the term  $X(1 - X)$  which simulates individuals "sampling" other individuals in the population. If one opinion dominates in the population (i.e.  $X \gg (1 - X)$  or  $(1 - X) \gg X$ ), the rate of changing opinions will be slow as the power of societal pressure makes it challenging for the other opinion to gain traction. However, if  $X$  and  $(1 - X)$  are close, the rate of change in opinion will be high as society has a split opinion on conservation versus harvest. In this model, each person holds an opinion (conservation or harvest) by weighing the benefits of conservation ( $U_A$ ) against the benefits of harvest ( $U_B$ ). This gives the replicator equation:

$$X(1 - X)[U_A - U_B] \quad (3)$$

$$X(1 - X)[\Delta U] \quad (4)$$

As individuals "sample" the opinions of others in their group, they can switch from A to B if  $U_B > U_A$  and

147 vice versa. In our model, we adapt  $U_A$  from Chris T. Bauch et al. (2016) with the added influence of the  
 148 other population's opinion.  $U_A$  is therefore given by:

$$U_{A,1} = \frac{1}{(F_1 + c_1)} + d_1 X_1 + \rho_1 X_2 \quad (5)$$

$$U_{A,2} = \frac{1}{(F_2 + c_2)} + d_2 X_2 + \rho_2 X_1 \quad (6)$$

149 Where  $\frac{1}{(F+c)}$  represents the perceived rarity of fish populations within a patch. As  $F$  and  $c$  (the rarity  
 150 valuation parameter) decrease, this term will increase, therefore adding to the perceived benefit of protecting  
 151 fish populations.  $d$  refers to the social influence that each population has on itself, and as an individual  
 152 encounters a conservationist ( $X$ ), the social benefit of also being a conservationist is shown in  $d$ .  $\rho$  has this  
 153 similar effect, but denotes the social effect on the opposite population on decision making. Individuals in  
 154 each population are receiving information about the conservation practices of the other, and the influence  
 155 that this has on each population is encapsulated by  $\rho$ .

156  $U_B$  (the perceived benefits of harvest) is:

$$U_{B,1} = \omega_1 + d_1(1 - X_1) + \rho_1(1 - X_2) \quad (7)$$

$$U_{B,2} = \omega_2 + d_2(1 - X_2) + \rho_2(1 - X_1) \quad (8)$$

157 Where  $\omega$  is the cost of conservation (i.e. revenue lost by not fishing) where now,  $d$  is the social benefit of  
 158 switching to harvesting ( $1 - X$ ) and  $\rho_1$  is the other population's ability to change the opinion of an individual  
 159 to be a harvester.

160 Plugging equations (5), (6) and (7), (8) into equation (3) gives:

$$\frac{dX_1}{dt} = k_1 X_1(1 - X_1) \left[ \frac{1}{F_1 + c_1} - \omega_1 + d_1(2X_1 - 1) + \rho_1(2X_2 - 1) \right] \quad (9)$$

$$\frac{dX_2}{dt} = k_2 X_2(1 - X_2) \left[ \frac{1}{F_2 + c_2} - \omega_2 + d_2(2X_2 - 1) + \rho_2(2X_1 - 1) \right] \quad (10)$$

Table 1: “Parameter values used to simulate sustainable fishing practices in patch 1 and overfishing in patch 2”

Parameter	Population 1	Population 2	Definition
r	0.4	0.35	Fish net growth
s	0.8	0.8	Supply and demand
h	0.25	0.5	Harvesting efficiency
k	1.014	1.014	Social learning rate
<U+03C9>	0.2	0.35	Conservation cost
c	1.5	1.5	Rarity valuation
d	0.5	0.5	Social norm strength (within pop)
i	0.2	0.2	Fish immigration (from opposite patch patch)
<U+03C1>	0.5	0.5	Social norm strength (opposite pop)

Where specifics of the derivation are outlined in the appendix. Coupling the fish population and human opinion models gives:

$$\frac{dF_1}{dt} = r_1 F_1 (1 - F_1) - \frac{h_1 * F_1 (1 - X_1)}{F_1 + s_1} - i_2 F_1 + i_1 F_2 \quad (11)$$

$$\frac{dF_2}{dt} = r_2 F_2 (1 - F_2) - \frac{h_2 * F_2 (1 - X_2)}{F_2 + s_2} - i_1 F_2 + i_2 F_1 \quad (12)$$

$$\frac{dX_1}{dt} = k_1 X_1 (1 - X_1) \left[ \frac{1}{F_1 + c_1} - \omega_1 + d_1 (2X_1 - 1) + \rho_1 (2X_2 - 1) \right] \quad (13)$$

$$\frac{dX_2}{dt} = k_2 X_2 (1 - X_2) \left[ \frac{1}{F_2 + c_2} - \omega_2 + d_2 (2X_2 - 1) + \rho_2 (2X_1 - 1) \right] \quad (14)$$

Where the fishing pressure is now a function of the number of harvesters in a population ( $\frac{hF(1-X)}{F+s}$ ). Further, the opinion of each population will shift based on the perceived fish stock health of their respective patch weighed against the costs and benefits of conservation. As fish stocks decrease, individuals will sway more toward conservation, thereby relieving this fishing pressure. However, we now have an external influence in this model: the opinions of people in the other population. The strength of this external influence is  $\rho$ , and in this study, we plan to simulate inequalities in human societies with this parameter.



## 3.2 Fish Parameters

For our basic analysis, we chose to model a two-patch fishery where the harvested fish species has a mid-range growth rate and regularly diffuses across the two patches, such as the parrotfish modeled in Thampi, Anand, and Bauch (2018). From the human side, the two groups of fishers have the same social influence on one another, representing a non-hierarchical social structure. The default fish growth rate of both patches is 0.35 fish per year. For the harvesting efficiency, we chose a maximal fishing rate of 0.5. These numbers were adapted from a coral reef fishing model Thampi, Anand, and Bauch (2018) where  $r = 0.35$  and  $h = 0.5$  are the mid-level growth rate and max fishing rates analyzed by this paper. For the emigration and immigration parameters, we chose 0.2 for each as these are the values used in the two-patch fishing model described in Cai, Li, and Song (2008). We used the  $s$  parameter described in the Chris T. Bauch et al. (2016) model of  $s = 0.8$ .

## 3.3 Human Parameters

The rate at which humans interact with one another is described by the parameter  $k$ . In our default model, we used  $k = 1.014$  as adapted from the Thampi, Anand, and Bauch (2018) default model. We used the default rarity valuation parameter  $c$  from Thampi, Anand, and Bauch (2018) where  $c = 1.68$ . The cost of conservation default parameter is  $\omega = 0.35$  from Chris T. Bauch et al. (2016). Further, as our default model will have no human social hierarchy, we set  $d = \rho = 0.5$  for our social learning rate as adapted from Chris T. Bauch et al. (2016) which models social decision making regarding deforestation.

Next, we altered these parameters to represent a scenario where patch 1 is fished sustainably, and patch 2 is overfished (Table 1). EASTON HOW DO I EXPLAIN MORE

## 3.4 Analyses

First, we compared the model from Chris T. Bauch et al. (2016) to that of our model by setting our parameters  $i_1 = i_2 = \rho_1 = \rho_2 = 0$  and comparing the results. We then ran different simulations of this model with different parameterizations in order to understand how fish population dynamics are affected by 2-patch socio-ecological models with social hierarchy. Next, we incrementally increased different parameters and simulated this system for 100 years in order to assess how increasing each parameter would affect the overall dynamics of the system. Sensitivity analysis was conducted with the R package FME (CITE). Finally, we created parameter planes of each pair of variables in order to assess how increasing two parameters at a

time would affect the system.

## 4 RESULTS

### 4.1 Model Comparisons

We first compared the output of our model to that of Chris T. Bauch et al. (2016) in order to see how the addition of social hierarchy and two-patch fishing affects the replicator dynamics of a fishery. We found that, while the model in Chris T. Bauch et al. (2016) had multiple states where oscillations occurred (Figure 2 a), there were no scenarios with our model that we tested which resulted in oscillatory behavior (Figure 2 b). This is likely due to the fact that the linear aspects of the fish dispersal term eventually overpowers the non-linear components of the model. Further, because of the outside human influence term,  $\rho$ , people are not responding directly to their respective fishing patch, but also to the conservation opinion of the other group.




Figure 2: a) Oscillatory behavior of the model in Chris T. Bauch et al. (2016) where  $r = 0.06$ ,  $s = 0.8$ ,  $h = 0.1$ ,  $k = 0.17$ ,  $w = 1$ ,  $c = 0.6$ ,  $d = 0.3$ . This is compared to our model, b) which used the same parameter values with the addition of  $i = 0.2$ ,  $\rho = 0.3$ .

### 4.2 Model Analysis

We then modeled a hypothetical scenario where patch 1 is fished sustainably whereas patch 2 is experiencing overfishing (Table 1). Here, the unsustainable practices of human population 2 are so exploitative, that both fish populations eventually collapse (Figure 3). We used this parameterization for the rest of the analysis, however the conclusions from each analyses are consistent with the other parameterizations we tested.




Figure 3: Representation of dynamics with parameters from table 1

213 Next, we ran our model with incrementally higher  $\rho$  values in both populations and observed how this  
 214 affected the final population of the fish patches (Figure 4). We found that under different parameterizations,  
 215  $\rho$  rarely had an effect on the final fish populations. However, there were some instances where  $\rho$  acted as  
 216 a tipping point for population dynamics where instead of continuously changing the final fish populations,  
 217 the  $\rho$  parameter either resulted in stable fish populations or both stocks collapsed once  $\rho$  increased past this  
 218 point.




Figure 4: Final fish populations after 100 years at different  $\rho$  values. Both  $\rho_1$  and  $\rho_2$  were increased simultaneously.

219 We then ran the same analysis with the fish dispersal parameter,  $i$ . Contrary to the effect  $\rho$  had on the  
 220 model, dispersal had a more direct and continuous effect on the final population of fish in each patch. For  
 221 example, as fish movement from patch 2 to patch 1 increased (i.e. from the unsustainable patch to the  
 222 sustainable patch), this actually maintained fish populations in both patches (Figure 5 a). However, if the  
 223 fish movement was increased from patch 1 to patch 2 (from the sustainable fishing to unsustainable), both  
 224 patches eventually collapsed to zero (Figure 5 b).

225 We then conducted sensitivity analysis on the  $i$  and  $\rho$  parameters to confirm the conclusions from the previous




Figure 5: Final fish populations after 100 years at different  $i$  values. a) shows how increases in  $i_1$  affect final populations and b) shows how increases in  $i_2$  affect final populations.

two analyses and further test if the fish diffusion parameter has a higher influence on fish population dynamics than the outside social influence parameter (Table 2). We used the R package FME (CITE) and found that  $F_1$  and  $F_2$  dynamics are consistently more sensitive to perturbations in the  $i$  parameter than to perturbations in the  $\rho$  parameter. This figure shows the L1 sensitivity as it is less sensitive to outliers, however all measures of sensitivity taken showed that the  $F_1$  and  $F_2$  variables are more sensitive to the  $i$  parameters than the  $\rho$

Table 2: Sensitivity analysis of the  $i$  and  $\rho$  parameters on  $F_1$  and  $F_2$  variables

Parameter	Sensitivity
$i_1$	0.0072569
$i_2$	0.0072569
$\rho_1$	0.0010693
$\rho_2$	0.0010693

Finally, to confirm that  $\rho$  has a small effect on the overall dynamics of the fish population model, we generated a series of parameter planes, which show how dynamics change based on variations of two different parameters (Figure 6). Each axis represents a new value for parameter and then the final dynamics of the system were analyzed. If the final fish population was greater than .25, SHOULD I CHANGE THAT CUTOFF????, this was considered stable whereas below .25 was considered population collapse. The corresponding color in the graph represents how each patch fared after 100 years under each parameter combination. The result of this analysis showed that  $\rho$  had no effect on the final fish population's sustainability.

## 5 DISCUSSION

We extended previous models on replicator dynamics to two patch fishing scenarios with social hierarchy in




Figure 6: Parameter plans of perturbations of each parameter compared to perturbations in  $\rho$ . Each Fish population was evaluated after 100 years and if the final population was under 0.25 SHOULD THAT CHANGE, this patch was considered collapsed.

certain environmental conditions. In our model, this shows that the dispersion of fish populations must be well understood in order to institute effective conservation practices. This is because any decision made by one group of people to conserve resources may be rendered ineffective if this species is highly migratory and the other group of harvesters is using unsustainable conservation practices. Further, because of the outside influences from the other human patch, fishers are no longer responding directly to fish levels in their respective patch, but are also influenced by the number of fishers in the other patch. In a scenario where fish is abundant in one patch, this will also encourage fishing in the other patch because this will increase the incentive to fish from the outside influence parameter.

For the remainder of the analyses, we simulated a scenario where patch 1 was being fished sustainably and patch 2 was experiencing overfishing, which resulted in the whole system experiencing collapse (Table 1 and Figure 3). Next, we tested the effect of external social influence on this model and how social hierarchy in decision making would influence dynamics. Contrary to our hypothesis, this parameter often had very little overall effect on the model (Figure 6). Figure 4 showed how fish population dynamics typically crashed when  $\rho$  passed a tipping point, showing that high levels of cooperation between groups actually crashed both populations of fish. Instead of modeling a cohesive system where communication fostered effective conservation, we created a tragedy of the commons scenario, where each community raced to fish each patch as opposed to coming to common understanding of sustainable fishing practices. This is because in our model, the only information being passed on to the outside group is the general opinion regarding fishing versus conservation instead of what sustainable fishing practices were used in order to achieve this benefit. As a result, when one patch is overfished and the other is fished sustainably, the group will continue to overfish their own resources because the opposite patch is encouraging this group to continue fishing. Therefore, this further highlights that the content of the information being disseminated matters in successful conservation (Gray et al. 2012). This confirms previous research on co-management strategies in that despite the fact they are a growing strategy in meeting social and ecological goals in small scale fisheries, they are prone to fail and groups can come to management decisions that are unsuccessful (Cinner et al. 2012). This aligns with previous social-ecological research that shows that the structure of a social network should be taken into consideration when the community manages a resource (Grafton 2005; Newman and Dale 2007; Bodin et al. 2014). This is because people who interact differently with the environment or within a society have to consider different tradeoffs in conservation (Cumming, Morrison, and Hughes 2017).

Instead of social norms controlling a lot of the dynamics of this model, we found that the dispersal of fish between patches was a major driver of population sustainability or collapse (Table 2). As we increased the movement of fish into the sustainable patch (Figure 5 a), this increased populations in that respective patch

because humans in population 1 continued to fish sustainably. Further, as those in population 2 decreased fishing rates, this influenced population 1 to also decrease their number of fishers. As a result, population 1 maintained high fishing levels while population 2 had low stocks. On the contrary, as fish immigrated from the sustainable patch to the unsustainable (Figure 5 b). Here, both fish populations collapsed as  $i_2$  increased. This is because immigration from patch 1 eventually grew to be too great for human population 1 to fish sustainably and human population 2 continued to overfish. High migration has been shown to be an essential part of maximizing economic benefit from fishing in multi-patch models (Moeller and Neubert 2015). Because fish are highly migratory and therefore difficult to track, constraining fishing to one group of people is more challenging (Grafton 2005). This is especially important for fish species that migrate based on lifestage, and requires more management coordination (Siddons, Pegg, and Klein 2017). NEED MORE FISH MOVEMENT SOURCES AND POINTS

The decision to include the external social influence term in our model within the injunctive social norms  $X(1 - X)$  implies that external influence can still change an opinion for or against conservation. However, an individual's willingness to take up a new opinion is still dictated by the overall opinion of their own population. This exemplifies homophily, a concept in sociology where humans tend to take information and the opinions from subgroups similar to them before listening to subgroups of different social standing (Brechwald and Prinstein 2011). Social network based conservation, like in our model, can replace 'top-down' regulation which can exclude stakeholders. However, this method of conservation has been shown to be susceptible to homophily (Newman and Dale 2007). Conservation has been shown to be more effective when human populations are more cohesive and that those with subgroups experience more barriers to effective conservation (Bodin and Crona 2009). Solutions to this issue could be to institute some form of actor that serves as cross-group communicators (Guerrero, Mcallister, and Wilson 2015).

Further research on this model could look at an open system, where fish diffusion doesn't necessarily have to pass between patches and could diffuse into non-fished areas. Also, stronger social networks have been shown to be more adaptable to environmental change (Grafton 2005), therefore further studies could look at the effect of climate change or extreme events on this social system (EASTON IS THIS A CITATION YET). The specific way we chose to incorporate social hierarchy into the model could be changed as well, for example testing a model that includes the outside social influence outside of the  $X(1 - X)$  term. There are many ways to model social systems so another further application of this study would be to compare different models that incorporate social hierarchy to the results of this one. Finally, further work on parameterizing this model to a real-world system could help understand if this model is properly capturing the underlying dynamics of two patch fishing systems with social hierarchy.



- Auger, Pierre, Bob Kooi, and Ali Moussaoui. 2022. “Increase of Maximum Sustainable Yield for Fishery in Two Patches with Fast Migration.” *Ecological Modelling* 467 (May): 109898. <https://doi.org/10.1016/j.ecolmodel.2022.109898>.
- Baker-Médard, Merrill. 2017. “Gendering Marine Conservation: The Politics of Marine Protected Areas and Fisheries Access.” *Society & Natural Resources* 30 (6): 723–37. <https://doi.org/10.1080/08941920.2016.1257078>.
- Ban, Natalie C, Morena Mills, Jordan Tam, Christina C Hicks, Sarah Klain, Natalie Stoeckl, Madeleine C Bottrill, et al. 2013. “A Social–Ecological Approach to Conservation Planning: Embedding Social Considerations.” *Frontiers in Ecology and the Environment* 11 (4): 194–202. <https://doi.org/10.1890/110205>.
- Barlow, Lee-Ann, Jacob Cecile, Chris T. Bauch, and Madhur Anand. 2014. “Modelling Interactions Between Forest Pest Invasions and Human Decisions Regarding Firewood Transport Restrictions.” Edited by Robert Planque. *PLoS ONE* 9 (4): e90511. <https://doi.org/10.1371/journal.pone.0090511>.
- Barnes, Michele L., Örjan Bodin, Tim R. McClanahan, John N. Kittinger, Andrew S. Hoey, Orou G. Gaoue, and Nicholas A. J. Graham. 2019. “Social-Ecological Alignment and Ecological Conditions in Coral Reefs.” *Nature Communications* 10 (1): 2039. <https://doi.org/10.1038/s41467-019-09994-1>.
- Barnes-Mauthe, Michele. 2013. “The Total Economic Value of Small-Scale Fisheries with a Characterization of Post-Landing Trends: An Application in Madagascar with Global Relevance.” *Fisheries Research*, 11.
- Bauch, Chris T. 2005. “Imitation Dynamics Predict Vaccinating Behaviour.” *Proceedings of the Royal Society B: Biological Sciences* 272 (1573): 1669–75. <https://doi.org/10.1098/rspb.2005.3153>.
- Bauch, Chris T., and Samit Bhattacharyya. 2012. “Evolutionary Game Theory and Social Learning Can Determine How Vaccine Scares Unfold.” Edited by Marcel Salathé. *PLoS Computational Biology* 8 (4): e1002452. <https://doi.org/10.1371/journal.pcbi.1002452>.
- Bauch, Chris T., Ram Sigdel, Joe Pharaon, and Madhur Anand. 2016. “Early Warning Signals of Regime Shifts in Coupled Human–Environment Systems.” *Proceedings of the National Academy of Sciences* 113 (51): 14560–67. <https://doi.org/10.1073/pnas.1604978113>.
- Bodin, Örjan, and Beatrice I. Crona. 2009. “The Role of Social Networks in Natural Resource Governance: What Relational Patterns Make a Difference?” *Global Environmental Change* 19 (3): 366–74. <https://doi.org/10.1016/j.gloenvcha.2009.05.002>.
- Bodin, Örjan, Beatrice Crona, Matilda Thyresson, Anna-Lea Golz, and Maria Tengö. 2014. “Conservation Success as a Function of Good Alignment of Social and Ecological Structures and Processes: Social–Ecological Fit and Conservation.” *Conservation Biology* 28 (5): 1371–79. <https://doi.org/10.1111/cobi.12306>.

- Borgatti, S. 2003. "The Network Paradigm in Organizational Research: A Review and Typology." *Journal of Management* 29 (6): 991–1013. [https://doi.org/10.1016/S0149-2063\(03\)00087-4](https://doi.org/10.1016/S0149-2063(03)00087-4).
- Brechwald, Whitney A., and Mitchell J. Prinstein. 2011. "Beyond Homophily: A Decade of Advances in Understanding Peer Influence Processes: BEYOND HOMOPHILY." *Journal of Research on Adolescence* 21 (1): 166–79. <https://doi.org/10.1111/j.1532-7795.2010.00721.x>.
- Cai, Liming, Xuezhi Li, and Xinyu Song. 2008. "MODELING AND ANALYSIS OF A HARVESTING FISHERY MODEL IN A TWO-PATCH ENVIRONMENT." *International Journal of Biomathematics* 01 (03): 287–98. <https://doi.org/10.1142/S1793524508000242>.
- Cinner, Joshua E., Tim R. McClanahan, M. Aaron MacNeil, Nicholas A. J. Graham, Tim M. Daw, Ahmad Mukminin, David A. Feary, et al. 2012. "Comanagement of Coral Reef Social-Ecological Systems." *Proceedings of the National Academy of Sciences* 109 (14): 5219–22. <https://doi.org/10.1073/pnas.1121215109>.
- Cozzi, Gabriele, Dominik M. Behr, Hugh S. Webster, Megan Claase, Caleb M. Bryce, Boitshepho Modise, John W. McNutt, and Arpat Ozgul. 2020. "African Wild Dog Dispersal and Implications for Management." *The Journal of Wildlife Management* 84 (4): 614–21. <https://doi.org/10.1002/jwmg.21841>.
- Cressman, Ross, Vlastimil Krivan, and József Garay. 2004. "Ideal Free Distributions, Evolutionary Games, and Population Dynamics in Multiple-Species Environments."
- Crona, Beatrice, and Örjan Bodin. 2006. "What You Know Is Who You Know? Communication Patterns Among Resource Users as a Prerequisite for Co-Management." *Ecology and Society* 11 (2): art7. <https://doi.org/10.5751/ES-01793-110207>.
- Cumming, Graeme S., Tiffany H. Morrison, and Terence P. Hughes. 2017. "New Directions for Understanding the Spatial Resilience of Social-Ecological Systems." *Ecosystems* 20 (4): 649–64. <https://doi.org/10.1007/s10021-016-0089-5>.
- Felipe-Lucia, María R., Angela M. Guerrero, Steven M. Alexander, Jaime Ashander, Jacopo A. Baggio, Michele L. Barnes, Örjan Bodin, et al. 2022. "Conceptualizing Ecosystem Services Using Social-Ecological Networks." *Trends in Ecology & Evolution* 37 (3): 211–22. <https://doi.org/10.1016/j.tree.2021.11.012>.
- Freedman, H. I., and Paul Waltman. 1977. "Mathematical Models of Population Interactions with Dispersal. I: Stability of Two Habitats with and Without a Predator." *SIAM Journal on Applied Mathematics* 32 (3): 631–48. <https://doi.org/10.1137/0132052>.
- Garrone-Neto, Domingos, Eduardo Antonio Sanches, Felipe Alexandre Lisboa de Miranda Daros, Cleber Mikio Rosa Imanobu, and Pietro Simon Moro. 2018. "Using the Same Fish with Different Rules: A Science-Based Approach for Improving Management of Recreational Fisheries in a Biodiversity Hotspot

of the Western South Atlantic.” *Fisheries Management and Ecology* 25 (4): 253–60. <https://doi.org/10.1111/fme.12288>.

Gervasi, Vincenzo, Henrik Brøseth, Erlend B. Nilsen, Hans Ellegren, Øystein Flagstad, and John D. C. Linnell. 2015. “Compensatory Immigration Counteracts Contrasting Conservation Strategies of Wolverines ( *Gulo Gulo* ) Within Scandinavia.” *Biological Conservation* 191 (November): 632–39. <https://doi.org/10.1016/j.biocon.2015.07.024>.

Grafton, R. Quentin. 2005. “Social Capital and Fisheries Governance.” *Ocean & Coastal Management* 48 (9-10): 753–66. <https://doi.org/10.1016/j.ocecoaman.2005.08.003>.

Gray, Steven, Alex Chan, Dan Clark, and Rebecca Jordan. 2012. “Modeling the Integration of Stakeholder Knowledge in Social–Ecological Decision-Making: Benefits and Limitations to Knowledge Diversity.” *Ecological Modelling* 229 (March): 88–96. <https://doi.org/10.1016/j.ecolmodel.2011.09.011>.

Guerrero, Angela M., Ryan R. J. Mcallister, and Kerrie A. Wilson. 2015. “Achieving Cross-Scale Collaboration for Large Scale Conservation Initiatives: Cross-Scale Collaboration in Conservation.” *Conservation Letters* 8 (2): 107–17. <https://doi.org/10.1111/conl.12112>.

Hebblewhite, Mark, and Jesse Whittington. 2020. “Wolves Without Borders: Transboundary Survival of Wolves in Banff National Park over Three Decades.” *Global Ecology and Conservation* 24 (December): e01293. <https://doi.org/10.1016/j.gecco.2020.e01293>.

“HIDDEN HARVEST-The Global Contribution of Capture Fisheries.” 2012. 66469-GLB. The World Bank. <https://documents1.worldbank.org/curated/en/515701468152718292/pdf/664690ESW0P1210120HiddenHarvest0web.pdf>.

Innes, Clinton, Madhur Anand, and Chris T. Bauch. 2013. “The Impact of Human-Environment Interactions on the Stability of Forest-Grassland Mosaic Ecosystems.” *Scientific Reports* 3 (1): 2689. <https://doi.org/10.1038/srep02689>.

Krackhardt, David, and Robert N. Stern. 1988. “Informal Networks and Organizational Crises: An Experimental Simulation.” *Social Psychology Quarterly* 51 (2): 123. <https://doi.org/10.2307/2786835>.

Lade, Steven J., Alessandro Tavoni, Simon A. Levin, and Maja Schlüter. 2013. “Regime Shifts in a Social-Ecological System.” *Theoretical Ecology* 6 (3): 359–72. <https://doi.org/10.1007/s12080-013-0187-3>.

Liu, Jianguo, Charles L Redman, Stephen H Schneider, Elinor Ostrom, Alice N Pell, Jane Lubchenco, William W Taylor, et al. 2007. “Coupled Human and Natural Systems.”

Mchich, R., P. Auger, and N. Raïssi. 2000. “The Dynamics of a Fish Stock Exploited in Two Fishing Zones.” *Acta Biotheoretica* 48: 201–18.

McPherson, Miller, Lynn Smith-Lovin, and James M Cook. 2001. “Birds of a Feather: Homophily in Social Networks.” *Annual Review of Sociology* 27 (1): 415–44. <https://doi.org/10.1146/annurev.soc.27.1.415>.

- Meisingset, Erling L., Leif E. Loe, Øystein Brekkum, Richard Bischof, Inger M. Rivrud, Unni S. Lande, Barbara Zimmermann, Vebjørn Veiberg, and Atle Mysterud. 2018. "Spatial Mismatch Between Management Units and Movement Ecology of a Partially Migratory Ungulate." Edited by Richard Fuller. *Journal of Applied Ecology* 55 (2): 745–53. <https://doi.org/10.1111/1365-2664.13003>.
- Moeller, Holly V., and Michael G. Neubert. 2015. "ECONOMICALLY OPTIMAL MARINE RESERVES WITHOUT SPATIAL HETEROGENEITY IN A SIMPLE TWO-PATCH MODEL: ECONOMICALLY OPTIMAL MARINE RESERVES." *Natural Resource Modeling* 28 (3): 244–55. <https://doi.org/10.1111/nrm.12066>.
- Newman, Lenore, and Ann Dale. 2007. "Homophily and Agency: Creating Effective Sustainable Development Networks." *Environment, Development and Sustainability* 9 (1): 79–90. <https://doi.org/10.1007/s10668-005-9004-5>.
- Ogburn, Matthew B., Autumn-Lynn Harrison, Frederick G. Whoriskey, Steven J. Cooke, Joanna E. Mills Flemming, and Leigh G. Torres. 2017. "Addressing Challenges in the Application of Animal Movement Ecology to Aquatic Conservation and Management." *Frontiers in Marine Science* 4 (March). <https://doi.org/10.3389/fmars.2017.00070>.
- Oraby, Tamer, Vivek Thampi, and Chris T. Bauch. 2014. "The Influence of Social Norms on the Dynamics of Vaccinating Behaviour for Paediatric Infectious Diseases." *Proceedings of the Royal Society B: Biological Sciences* 281 (1780): 20133172. <https://doi.org/10.1098/rspb.2013.3172>.
- Plumptre, Andrew J., Deo Kujirakwinja, Adrian Treves, Isaiah Owunji, and Helga Rainer. 2007. "Transboundary Conservation in the Greater Virunga Landscape: Its Importance for Landscape Species." *Biological Conservation* 134 (2): 279–87. <https://doi.org/10.1016/j.biocon.2006.08.012>.
- Pomeroy, Robert, John Parks, Richard Pollnac, Tammy Campson, Emmanuel Genio, Cliff Marlessy, Elizabeth Holle, et al. 2007. "Fish Wars: Conflict and Collaboration in Fisheries Management in Southeast Asia." *Marine Policy* 31 (6): 645–56. <https://doi.org/10.1016/j.marpol.2007.03.012>.
- Prince, Jeremy, Watisoni Lalavanua, Jone Tamanitoakula, Laitia Tamata, Stuart Green, Scott Radway, Epeli Loganimoce, et al. 2021. "Spawning Potential Surveys in Fiji: A New Song of Change for <Span Style='font-Variant:small-Caps;'>small-scale</Span> Fisheries in the Pacific." *Conservation Science and Practice* 3 (2). <https://doi.org/10.1111/csp2.273>.
- Ramírez-Valdez, Arturo, Timothy J. Rowell, Katherine E. Dale, Matthew T. Craig, Larry G. Allen, Juan Carlos Villaseñor-Derbez, Andrés M. Cisneros-Montemayor, et al. 2021. "Asymmetry Across International Borders: Research, Fishery and Management Trends and Economic Value of the Giant Sea Bass (*Stereolepis Gigas*)." *Fish and Fisheries* 22 (6): 1392–1411. <https://doi.org/10.1111/faf.12594>.
- Salas, Silvia, María José Barragán-Paladines, and Ratana Chuenpagdee, eds. 2019. *Viability and Sustain-*

ability of Small-Scale Fisheries in Latin America and The Caribbean. Vol. 19. MARE Publication Series.  
Cham: Springer International Publishing. <https://doi.org/10.1007/978-3-319-76078-0>.

Siddons, S. F., M. A. Pegg, and G. M. Klein. 2017. "Borders and Barriers: Challenges of Fisheries Management and Conservation in Open Systems: Challenges of Fisheries Management and Conservation in Open Systems." *River Research and Applications* 33 (4): 578–85. <https://doi.org/10.1002/rra.3118>.

Sigdel, Ram P., Madhur Anand, and Chris T. Bauch. 2017. "Competition Between Injunctive Social Norms and Conservation Priorities Gives Rise to Complex Dynamics in a Model of Forest Growth and Opinion Dynamics." *Journal of Theoretical Biology* 432 (November): 132–40. <https://doi.org/10.1016/j.jtbi.2017.07.029>.

Thampi, Vivek A., Madhur Anand, and Chris T. Bauch. 2018. "Socio-Ecological Dynamics of Caribbean Coral Reef Ecosystems and Conservation Opinion Propagation." *Scientific Reports* 8 (1): 2597. <https://doi.org/10.1038/s41598-018-20341-0>.

Young, Oran R., Gail Osherenko, Julia Ekstrom, Larry B. Crowder, John Ogden, James A. Wilson, Jon C. Day, et al. 2007. "Solving the Crisis in Ocean Governance: Place-Based Management of Marine Ecosystems." *Environment: Science and Policy for Sustainable Development* 49 (4): 20–32. <https://doi.org/10.3200/ENVT.49.4.20-33>.