

Chapter 1

Velocity

In this chapter

By the end of this chapter you should be able to

- explain what a *conceptual model* is, why scientists use them, and what makes a model scientific;
- draw and read *dot diagrams* to compare the speeds of moving objects, and measure a real speed from video frames or from a printed diagram by using an object of known size as a *scale*;
- recognise that positions read off a grid are in *grid units*, not metres, and convert them only with a scale you have measured or been given;
- check a rearranged formula by substituting units, and estimate an instantaneous speed from a dot diagram using a *centred* interval;
- use $\text{distance} = \text{speed} \times \text{time}$ and rearrange it for any one of the three quantities, and convert speeds between m/s, km/h, and mi/h;
- explain why every speed is measured *relative* to something, and say what that something usually is;
- distinguish *average* speed from *instantaneous* speed two ways: as a ratio over a shrinking interval, and as a *derivative*; recover distance from speed two ways: as average speed times time, and as an *integral*;
- read speed from a *position–time graph* as a slope, both of a chord and of a tangent;
- recognize when a model is being *extrapolated* beyond the range where it was tested, and perform a *sanity check* on a result;
- express a measured quantity with its *uncertainty*, and carry that uncertainty through a calculation by recomputing at the ends of the range and by using a derivative;
- convert an answer into units you have direct experience of before judging whether to believe it;
- explain how *velocity* differs from speed, tell a *vector* quantity from a *scalar*, and find the velocity of one object relative to another;
- distinguish an object's *position* from its *coordinate*, say what a change of origin does and does not change, and tell the sign of a coordinate from the sign of a velocity.

How to read this chapter: two perspectives

Every idea in this book can be understood with elementary mathematics alone: ratios, averages, tables, and the areas of rectangles and triangles. Every idea can also be understood with calculus, the mathematics of change, which was invented for exactly these questions. Neither view is the “real” one; each illuminates the other. Throughout, explanations and solutions are labelled

Elementary for the route using only elementary mathematics, and **Calculus** for the route using derivatives and integrals.

This chapter needs only a little calculus: the derivative as the slope of a graph, and the integral as the area under one. If you have not met them yet, read the **Elementary** routes now and return to the **Calculus** routes later; the elementary route is always complete on its own. When a worked example carries both labels, the two routes are independent: read one, then the other, and check that they agree.

1.1 An Ancient Sprinter

In 2003, Mary Pappin Jr., a young Mutthi Mutthi woman from the Willandra Lakes region of south-eastern Australia, noticed a set of footprints pressed into the hard surface of a long-vanished lake bed. The prints were not recent. They had been made in soft, wet clay, then baked, buried, and finally uncovered again by the wind. Hundreds more were soon found nearby: the largest collection of Ice Age human footprints known anywhere in the world.

An archaeologist, Steve Webb, led the scientific study of the site. Among the tracks he described was one that made headlines around the world. Webb calculated that the person who left it had been running barefoot across the mud at more than 10 m/s, about the average speed of an Olympic 100-metre champion. Newspapers ran stories with titles like “Prehistoric hunter could have outrun Usain Bolt.” For a while, the ancient sprinter was famous.

Then, quietly, the story faded. A few years later, other scientists re-examined the same footprints and came to a very different conclusion.

This chapter uses the story of the ancient sprinter to introduce the physics of motion. Along the way we will ask three questions that matter far beyond footprints:

1. How can you measure the speed of something you never saw move?
2. What does it mean to *model* a physical situation, and what can go wrong?
3. How confident should you be in a number that comes out of a calculation?

1.2 Models in Science

Nobody watched the hunter run. There is no stopwatch, no video, and no witness. All that survives is a trail of footprints in dried mud. To turn those footprints into a speed, Webb needed a *model*.

Definition 1.1: Conceptual model

A **conceptual model** is a deliberately simplified description of some part of the world, built so that we can reason about it. A model may combine measurements, diagrams, equations, graphs, tables of data, and computer programs. Every model leaves things out. A good model leaves out the things that do not matter for the question being asked, and keeps the things that do.

You already use models all the time. A subway map is a model of a city’s rail network: it shows which stations connect to which, but it distorts distances and ignores streets entirely. That is exactly right for deciding which train to take, and exactly wrong for estimating how far you will walk. A model is never simply “true” or “false”; it is useful or not useful *for a particular purpose*.

What makes a model scientific? Physicists apply two tests. First, the model must agree with what has already been observed. Second, and more importantly, it must *predict* something that has not yet been checked, so that it can be put at risk. A model whose predictions turn out contrary to what actually happens is refined or, if it cannot be repaired, abandoned. A model that cannot be tested, even in principle, is not doing scientific work at all, however reasonable it sounds. We will meet this test again in Chapter 2, where two rival hypotheses about falling objects are worth arguing about only because they predict different things.

This is a particular case of what might be called the scientific attitude: one of inquiry, integrity, and humility, meaning a willingness to admit error. In science a single verifiable observation that contradicts a claim outweighs any authority behind the claim, no matter how respected the person making it or how many people have repeated it. Keep this in mind as the story of the ancient sprinter unfolds.

Physics is largely the craft of building models that are simple enough to calculate with and faithful enough to trust. The rest of this chapter follows one such model from its construction to its collapse, and then to its repair.

Three models of one set of footprints

Other people studied the same footprints with entirely different models. Webb used a laboratory technique called optically stimulated luminescence, which relies on a model of how buried mineral grains store energy from natural radioactivity, to date the prints to between 19,000 and 23,000 years old. A group of Pintubi trackers, drawing on a lifetime of reading footprints, recognized that most of the trails were left by families walking, while a few belonged to a hunting party chasing an animal. One member of that party was running hard, cutting across to head the animal off. That runner is the “ancient sprinter.” Neither the dating nor the tracking tells us the runner’s speed. For that, Webb needed a third model, and he built it with nothing more than a tape measure.

1.3 From Footprints to Dot Diagrams

1.3.1 What Webb measured

Webb measured two things about the runner’s trail. The first was the **stride length**: the distance from one footprint to the next print made by the *same* foot. Note that a stride is two steps. (A **step** is the distance from a left print to the following right print, or vice versa.) The

second measurement was the **foot length** of the prints themselves. Figure 1.1 shows the idea.




Figure 1.1: A trail of footprints. A stride runs from one right-foot print to the next right-foot print; a step runs from a left print to the following right print. Webb measured stride length and foot length.

Why stride and not step? Because the two feet fall on slightly different lines. Measuring left-to-right-to-left traces a zigzag; measuring right-to-right follows a straight line along the direction of travel, which is what we want.

1.3.2 Simplifying to dots

Suppose the runner planted a right foot once every 0.6 s. (That corresponds to 100 strides per minute, since $60\text{ s}/100 = 0.6\text{ s}$.) Then the right-foot prints mark where the runner was at 0 s, 0.6 s, 1.2 s, and so on. The left-foot prints add nothing new for our purposes, so we drop them. And the shape of each print does not matter either, so we replace each with a dot. Figure 1.2 shows the result.




Figure 1.2: The same trail as a dot diagram. Each dot marks the runner's position at equally spaced instants of time.

Definition 1.2: Dot diagram

A **dot diagram** (also called a *motion diagram*) is a picture that marks the position of a moving object at equally spaced instants of time. The time between dots is the same throughout a single diagram, but it may be any convenient value.

A dot diagram can represent anything that moves. Picture a car with a leaky can of paint dripping onto the road once per second; the trail of drops is a dot diagram. Picture a strobe light flashing in a dark gym while someone runs across; each flash freezes the runner at one dot. And even when nothing physical leaves a mark, we can *imagine* the dots. That is the whole point of a model.

Two rules for dot diagrams


1. Within one diagram, every pair of neighbouring dots is separated by the *same* time interval.

2. To compare the speeds of two objects using their dot diagrams, both diagrams must use the *same* time interval.

1.3.3 Reading speed from spacing


Worked Example 1.1: Ranking speeds from dot diagrams

The dot diagrams below show three objects. The time between dots is the same for all three. Rank the objects from fastest to slowest.



Solution. It is tempting to look at object B, with its many closely packed dots, and think “that one is busy, so it must be fast.” Resist the temptation. The dots are not being laid down faster for B; the time between dots is identical for all three objects. The only thing that differs is how far each object moves during that fixed interval.

Imagine the three objects starting a race together, each laying its first dot at the starting line at the same moment. When the second dot is laid, all three lay it at the same moment too. The object whose second dot is farthest from the start has travelled the greatest distance in the same time, so it is the fastest.



Object A's second dot (highlighted) is farthest along, then C's, then B's. The ranking is $A > C > B$.

Takeaway: widely spaced dots mean high speed; closely spaced dots mean low speed. This works because a fast object covers more ground than a slow one in the same time. ■

Worked Example 1.2: Counting gaps, not dots

The dot diagram in Figure 1.2 has six dots, 0.6 s apart. How much time does the diagram cover?

Solution. The elapsed time is measured by the *gaps* between dots, not by the dots themselves. Six dots have only five gaps between them, so the diagram covers

$$5 \times 0.6 \text{ s} = 3.0 \text{ s}.$$

Be careful: a common error is to answer $6 \times 0.6\text{ s} = 3.6\text{ s}$. This mistake is so common that it has a name: the *fencepost error*, after the observation that a fence with six posts has

only five sections of rail. ■

1.3.4 When speed changes

A dot diagram does more than compare constant speeds. If an object speeds up, each time interval carries it a little farther than the last, so the dots spread apart along the direction of motion. If it slows down, the dots bunch together. Figure 1.3 shows both cases, along with an object moving at constant speed for comparison.




Figure 1.3: Dot diagrams for three kinds of motion. Each row uses the same time between dots.

Notice one thing a dot diagram does *not* tell you: which way the object was going. An object moving left to right and one moving right to left leave exactly the same set of dots. If direction matters, we must add it to the diagram, for example by numbering the dots or drawing an arrow. We return to direction in Section 1.9.

1.3.5 Making dot diagrams in the laboratory

A classic laboratory device for producing dot diagrams is the **ticker-tape timer** (Figure 1.4). A long paper tape is attached to a moving cart and threaded under a small hammer that taps down at a fixed rate, typically 50 times per second. Each tap leaves a carbon dot on the tape. After the run, the tape is a dot diagram of the cart's motion, with 0.02 s between dots. With a ruler, you can then measure the spacing directly.




Figure 1.4: A ticker-tape timer. The cart drags the tape under a hammer that strikes at a steady rate, leaving a dot diagram on the tape. In this sketch the cart is speeding up: the dots nearest the cart, made most recently, are farthest apart.

Check Your Understanding 1.1

A ticker-tape timer is adjusted to tap *less* often, so that more time passes between dots. The cart's motion is unchanged. Will the dots on the tape be closer together or farther apart than before? Explain in one sentence.

Check Your Understanding 1.2

You want to compare a fast cart and a slow cart using ticker tape. Should the timer tap at the same rate for both, faster for the fast cart, or faster for the slow cart? Why?

1.3.6 Measuring speed from a video

A ticker-tape timer makes an excellent dot diagram and is confined to a laboratory bench. The camera in your pocket does the same job for anything you can film, and it is how most measurements in this book can actually be checked.

A video is already a dot diagram in disguise. It is a stack of still frames taken at a fixed rate — commonly 30 or 60 per second, and often several hundred in a phone’s slow-motion mode — so stepping through the frames and marking where the object is gives exactly the equally-spaced-in-time record that Section 1.3 asked for. The frame rate supplies the times for free: if the camera shoots at 30 frames per second, then n frames is $n/30$ seconds, with no clock and no reaction time to worry about.

Distances are the harder half, because a photograph contains no absolute lengths. A cyclist who fills the frame might be two metres away or twenty. What a photograph does contain is *ratios*, so if you know the true size of any one object in the shot, every other length in that shot follows. An object used this way is called a **scale**.

Definition 1.3: Scale

A **scale** is an object of known size appearing in an image, used to convert measured image distances into real ones. Measuring the scale object in image units (usually **pixels**, the small dots a digital image is built from) gives a conversion factor between pixels and metres, which then applies to every distance in that image. A good scale is rigid, of accurately known length, and lies in the same plane, at the same distance from the camera, as the motion being measured.

Objects of known size are easier to find than you might expect: a metre stick laid in the shot, a standard sheet of paper, a doorway, a tennis court’s service line, or a bicycle’s wheelbase.

Worked Example 1.3: A cyclist’s speed from three frames

A cyclist is filmed side-on at 30 frames per second as she rides past a crack in the road. Her bicycle’s wheelbase — the distance between the two axles, quoted by the manufacturer as 1.02 m — measures 96 pixels in the image. Between frame 12 and frame 15, the point where her front wheel meets the road advances 71 pixels. How fast is she going, and how well do you know it?

Solution. *The scale.* The wheelbase gives the conversion factor

$$1.02 \text{ m} \hat{=} 96 \text{ pixels} \quad \implies \quad 1 = \frac{1.02 \text{ m}}{96 \text{ pixels}}.$$

Multiplying by this fraction converts pixels to metres without changing the quantity, exactly as in Section 1.4.4.

The distance.

$$\Delta x = 71 \text{ pixels} \times \frac{1.02 \text{ m}}{96 \text{ pixels}} = 0.754 \text{ m}.$$

The pixels cancel, leaving metres.

The time. From frame 12 to frame 15 is 3 frame intervals, and each lasts $1/30$ of a second, so

$$\Delta t = \frac{3}{30} \text{ s} = 0.100 \text{ s}.$$

The speed.

$$v = \frac{\Delta x}{\Delta t} = \frac{0.754 \text{ m}}{0.100 \text{ s}} = 7.5 \text{ m/s}.$$

How well do we know it? Locating the wheel-road contact point to better than a pixel or two is optimistic, so call the 71 pixels uncertain by about ± 2 , a percentage uncertainty of $2/71 \approx 2.8\%$. The wheelbase measurement and the manufacturer's figure carry their own uncertainty, and so, slightly, does the frame rate. Rounding up to a round 5% to cover all of them,

$$\delta v \approx 0.05 \times 7.5 \text{ m/s} = 0.4 \text{ m/s}, \quad \text{so} \quad v = (7.5 \pm 0.4) \text{ m/s}.$$

Section 1.8 explains why a percentage uncertainty passes straight through a division like this, and why rounding an uncertainty *up* is the honest direction to round.

Sanity check: 7.5 m/s is 27 km/h , a brisk but ordinary cycling speed, and Table 1.1 puts it comfortably above a jog and well below a car. Note also what we have actually measured: her *average* speed over one tenth of a second. That is close to, but not the same as, her speed at any one instant — a distinction Section 1.5 takes up. ■

Why estimating beats guessing

Asked to guess a cyclist's speed from a video, a room full of people will produce answers spread over a factor of three, and the spread does not shrink if you ask them to think harder. The measurement above is not obviously cleverer than a guess — it is just a scale, a frame count, and a division — yet it lands within a few percent, and, more importantly, it comes with a stated uncertainty, so you know how much to trust it. This is the whole method of the subject in miniature: instead of asking the question directly, break it into quantities you can measure, and keep track of how well you measured them.

Check Your Understanding 1.3

You film a friend running past a parked car and want their speed. (a) Name two things in the shot you could use as a scale, and one that would be a poor choice. (b) The car is much closer to the camera than your friend is. Why does using the car as the scale make your answer wrong, and does it make the speed come out too high or too low? (c) You step from frame 40 to frame 46 at 60 frames per second. How much time has passed?

1.3.7 Grid units are not metres

The scale idea has a second home, and it is where most people meet it as a mistake rather than a technique. A dot diagram printed in a book or drawn on a screen sits on some grid, and it is natural to read the dots' positions off that grid as numbers: dot one at 0, dot two at 2, dot three at 8, and so on. Those numbers are real and useful. They are also *not* distances, any more than the pixel counts of Section 1.3.6 were. They are counts of grid squares, and until you know

what one square is worth, they cannot be divided by a time to give a speed.

Three steps, and never fewer

To get a physical speed out of a diagram:

1. **Read** the positions in the diagram's own units — grid squares, pixels, millimetres on the page. Do not call them metres.
2. **Convert**, using a scale you have *measured or been given*: one length in the diagram whose true size you know. This turns the reading into a real distance.
3. **Divide** by the elapsed time, and then **check** the answer against something you can picture.

Skipping step 2 is the same as assuming one grid square equals one metre. That is an assumption, not a reading, and it is almost never true.




Figure 1.5: A dot diagram on a grid. Reading the dots gives the positions 0, 2, 8, 18, 32, 50 — in *grid squares*. One measurement converts the whole diagram: a ruler laid across the last gap, which spans $50 - 32 = 18$ squares, reads 2.7 cm, so one square is 1.5 mm. Without that measurement the grid numbers have no physical size at all.

Worked Example 1.4: Average speed from a printed dot diagram

Figure 1.5 shows a dot diagram whose dots were made $\tau = 0.5$ s apart. Reading off the grid, the dots sit at 0, 2, 8, 18, 32, and 50 squares. A ruler laid across the last gap, from the 32 square to the 50 square, measures 2.7 cm. Find the object's average speed over the whole diagram, and estimate its instantaneous speed at the second-to-last dot.

Solution. *Step 1: read.* The total displacement is $50 - 0 = 50$ squares. There are six dots, so there are *five* intervals (the fencepost count of Section 1.3), and the total time is

$$t = 5 \times 0.5 \text{ s} = 2.5 \text{ s}.$$

Step 2: convert. The ruler measurement is the scale. The last gap spans $50 - 32 = 18$ squares and measures 2.7 cm, so

$$1 \text{ square} = \frac{2.7 \text{ cm}}{18} = 0.15 \text{ cm} = 1.5 \text{ mm}.$$

The total distance is therefore

$$d = 50 \text{ squares} \times \frac{0.15 \text{ cm}}{1 \text{ square}} = 7.5 \text{ cm} = 0.075 \text{ m}.$$

Step 3: divide, and check.

$$\bar{v} = \frac{d}{t} = \frac{0.075 \text{ m}}{2.5 \text{ s}} = 0.030 \text{ m/s} = 3.0 \text{ cm/s}.$$

Three centimetres per second is a slow crawl: a few times a garden snail's pace, and near the top of Table 1.1 where the slowest things live. That is exactly right for something whose whole journey fits inside 7.5 cm of paper in two and a half seconds.

The instantaneous speed. The second-to-last dot is the one at 32. Take the two dots on either side of it, at 18 and 50, so that the interval is centred on the dot we care about:

$$v \approx \frac{(50 - 18) \text{ squares} \times 0.15 \text{ cm}}{2 \times 0.5 \text{ s}} = \frac{4.8 \text{ cm}}{1.0 \text{ s}} = 4.8 \text{ cm/s}.$$

Larger than the average, as it must be: the gaps are growing, so the object is quickest near the end.

Sanity check on the whole approach: had we skipped step 2 and treated squares as metres, we would have got $50/2.5 = 20 \text{ m/s}$ — a highway speed, for a dot diagram a few centimetres long. The error would have been a factor of nearly 700, and the sanity check alone would have caught it. ■

Three habits that catch this error

The grid-unit mistake is easy to make and easy to catch, and each of the three catches is a habit worth building.

Name your units at every step. Write “50 squares,” not “50.” A bare number invites you to divide it by seconds and call the result metres per second. A number with “squares” attached makes the missing conversion visible, because squares/s is not a speed.

Ask where the metres came from. Every physical distance in an answer has to trace back to something actually measured with a real ruler, or given in the problem. If you cannot point at that measurement, you have assumed a scale rather than found one.

Sanity-check the size. This is Chapter 1's recurring theme, and it is the cheapest of the three. A dot diagram you can cover with your hand cannot describe a car on a motorway. When an answer is out by a factor of hundreds, the number itself will usually tell you, if you convert it into units you have a feel for before you write it down.

Check Your Understanding 1.4

A dot diagram on squared paper has dots at grid positions 0, 3, 12, 27, 48, made 0.2 s apart. A ruler shows that the gap from 27 to 48 measures 4.2 cm. (a) How many intervals are there, and what is the total time? (b) What is one grid square worth in centimetres? (c) Find the average speed in cm/s and in m/s. (d) A student answers $48 \text{ m}/(5 \times 0.2 \text{ s}) = 48 \text{ m/s}$. Name the *two* separate mistakes in that line, and say roughly how far out the final answer is.

1.4 Speed: Distance and Time

A dot diagram shows that widely spaced dots mean high speed, but it does not yet give speed a number. To do that we need a second ingredient: the time between dots.

1.4.1 The basic relationship

Before the seventeenth century, people described moving things simply as “slow” or “fast.” Galileo Galilei (1564–1642) is credited with being the first to measure speed by comparing the distance covered with the time it took, and he defined speed as the distance covered per unit of time. Distance was easy for Galileo to measure; short times were not. He sometimes used his own pulse as a clock, and sometimes counted drops from a “water clock” of his own design. The definition he arrived at is the one we still use.

Start with something familiar. If you drive at 30 mi/h for 2 h, you cover 60 mi. Drive twice as long and you cover twice the distance; drive twice as fast and you also cover twice the distance. Distance is proportional to both speed and time, so

$$\text{distance} = \text{speed} \times \text{time}. \quad (1.1)$$

In symbols, with d for distance, v for speed, and t for time,

$$d = vt. \quad (1.2)$$

We usually want speed, so we divide both sides by t :

$$v = \frac{d}{t}. \quad (1.3)$$

Definition 1.4: Speed

The **speed** of an object is the distance it travels divided by the time taken:

$$\text{speed} = \frac{\text{distance travelled}}{\text{time elapsed}}, \quad v = \frac{d}{t}.$$

In SI units, distance is in metres (m) and time in seconds (s), so speed is in metres per second (m/s). The slash in m/s is read “per” and means “divided by.” Any combination of a distance unit and a time unit is a legitimate unit of speed: kilometres per hour and miles per hour suit cars and long journeys, while metres per second suits the shorter distances of most physics problems, and it is the unit this book uses unless there is a good reason not to.

Equation (1.3) packs two ideas into one fraction:

- **Fixed time, bigger distance, bigger speed.** If one object moves 10 m in 2 s and another moves 1000 m in 2 s, the second is faster. The numerator grew.
- **Fixed distance, smaller time, bigger speed.** If one object moves 10 m in 2 s and another moves 10 m in 0.02 s, the second is much faster. The denominator shrank, and shrinking a denominator makes a fraction larger.

Two Views: speed

Elementary Speed is a *ratio*: distance per unit time. The ratio is the same whichever pair of distance and time you take from a constant-speed motion, because doubling the time doubles the distance. Choose a time interval, measure how far the object went, divide. On a dot diagram, the ratio is the spacing of the dots divided by the time between them.

Calculus Speed is a *rate of change*. Let $x(t)$ be the object's position along its line of motion at time t . Constant speed means the position grows in proportion to time, $x(t) = x_0 + vt$, and the speed is the derivative of position with respect to time:


$$v = \frac{dx}{dt}.$$

For $x(t) = x_0 + vt$ the derivative is the constant v : the graph of x against t is a straight line and v is its slope. Written this way the definition does not care whether the speed is constant, and that is its advantage. Section 1.5 uses it for motion whose speed changes.

Worked Example 1.5: Walking speed from stride data

While walking at a steady pace you take one stride every 1.18 s, and each stride is 1.54 m long. How fast are you walking, in metres per second?

Elementary Begin with a labelled dot diagram. It is not strictly needed for the arithmetic, but it organizes the information and makes it obvious which distance goes with which time.



Each stride carries you 1.54 m and takes 1.18 s. By Equation (1.3),

$$v = \frac{d}{t} = \frac{1.54 \text{ m}}{1.18 \text{ s}} = 1.31 \text{ m/s}.$$

Calculus After n strides the walker is at $x = n \times 1.54 \text{ m}$ and the clock reads $t = n \times 1.18 \text{ s}$. Eliminating n gives the position as a function of time,

$$x(t) = \frac{1.54 \text{ m}}{1.18 \text{ s}} t = (1.31 \text{ m/s}) t, \quad \text{so} \quad v = \frac{dx}{dt} = 1.31 \text{ m/s}.$$

The derivative has no t in it, which says the speed is the same at every instant, as “steady pace” promised. Strictly, the walker’s speed wobbles within each stride, and the straight line is a model of the average motion; the dots are on the line, and the wobble is between them.

Is this reasonable? At 1.31 m/s a kilometre takes about 13 minutes and a mile takes about 20 minutes. That is an ordinary walking pace, so the answer passes a quick reality check.

■

Units travel with the numbers

In the example above, metres divided by seconds gave metres per second. Always carry the units through a calculation. If the units of your answer come out wrong, the calculation is wrong, no matter how good the arithmetic looks. The same holds for derivatives: dx/dt has the units of x divided by the units of t , and an integral $\int v dt$ has the units of v times the units of t .

1.4.2 Solving for time or distance

Equation (1.2) can be rearranged to find whichever quantity is unknown:

$$d = vt, \quad v = \frac{d}{t}, \quad t = \frac{d}{v}.$$

You do not need to memorize three formulas. Memorize $d = vt$ and rearrange.

Check a rearrangement with its units

Rearranging is where a formula most often goes wrong, and the error is invisible if you only look at the letters. Suppose you wrote “ $d = v/t$ ” instead of $d = vt$. Nothing about the symbols objects. The units do, immediately:

$$\frac{\text{m/s}}{\text{s}} = \text{m/s}^2,$$

which is not a distance. Whereas vt gives $\text{m/s} \times \text{s} = \text{m}$, as it must.

Make this a reflex. Every time you rearrange, substitute the units in place of the quantities and check that both sides come out the same. It takes a few seconds, needs no understanding of the physics, and catches a large fraction of all algebra slips. It is also the only check available when you have no numbers yet, which is exactly when a wrong rearrangement is most likely to survive into the rest of a solution.

Worked Example 1.6: Paint drops from a wheel

A car travels along a highway at 25 m/s. A blob of wet paint on one tyre touches the road once per rotation, and the wheel turns once every 0.079 s. How far apart are the paint marks, and what does that distance represent physically?

Solution. The marks are a dot diagram with 0.079 s between dots. In that time the car moves

$$d = vt = (25 \text{ m/s})(0.079 \text{ s}) = 1.98 \text{ m}.$$

The car moves this far during exactly one rotation of the wheel, so 1.98 m is the distance the tyre rolls out in one turn: the *circumference* of the wheel. (A tyre about 63 cm across has roughly this circumference, which is sensible for a car.) ■

Worked Example 1.7: Finding the time between strides

A runner moves at 3.2 m/s with a stride length of 2.08 m. How much time passes between successive strides?

Solution. Rearranging $d = vt$ for time,

$$t = \frac{d}{v} = \frac{2.08 \text{ m}}{3.2 \text{ m/s}} = 0.65 \text{ s}.$$

Check the units: metres divided by metres-per-second leaves seconds. Check the size: a bit more than one and a half strides per second is typical of an easy distance run. ■

Check Your Understanding 1.5

Two dot diagrams are made with the same timer. The dots in diagram A are three times as far apart as the dots in diagram B. How many times faster is object A than object B? Now suppose instead that the timer for A ran at half the rate of the timer for B (twice as much time between dots). Which object is faster now, and by what factor?

1.4.3 Speed relative to what?

There is a hidden assumption in every speed we have quoted so far. When we say the walker moves at 1.31 m/s, we mean relative to the ground. The walker's speed relative to a friend walking alongside is zero. If you stroll down the aisle of a moving train, your speed relative to the floor of the carriage is a leisurely 1 m/s, while your speed relative to the countryside outside is that of the train. Everything moves relative to something, and a speed is meaningful only once we know what that something is.

Even “at rest” is relative. As you sit reading this, you are at rest relative to your chair, but Earth carries you around the Sun at about 107 000 km/h, which is 30 km/s, and faster still relative to the centre of our galaxy. When a racing car is said to reach 300 km/h, the speed is relative to the track. Unless stated otherwise, speeds of things in our surroundings are measured relative to the surface of the Earth, and that is the convention this book follows. The ancient sprinter's speed, whatever it was, is a speed relative to the mud of the lake shore. We return to relative motion, with signs and directions, in Section 1.9.5.

Check Your Understanding 1.6

A hungry mosquito sees you resting in a hammock in a steady 3 m/s breeze. How fast, and in which direction, should the mosquito fly in order to hover above you?

1.4.4 Units and conversions

Scientists report speed in metres per second, but everyday life uses kilometres per hour or miles per hour, and you should be able to move fluently between them. The method is to multiply by fractions that equal one. Since 1 km = 1000 m and 1 h = 3600 s,

$$1 \text{ m/s} = 1 \frac{\text{m}}{\text{s}} \times \frac{1 \text{ km}}{1000 \text{ m}} \times \frac{3600 \text{ s}}{1 \text{ h}} = 3.6 \text{ km/h}.$$

Each fraction is equal to one, so multiplying by it changes the units without changing the physical quantity. The metre and second symbols cancel, leaving kilometres per hour.

Where the metre and the second come from

Units are themselves agreements, and they have been refined as measurement improved. The metre was originally defined as one ten-millionth of the distance from the North Pole to the equator, and for a long time it was the distance between two scratches on a platinum–iridium bar kept near Paris. Since 1983 it has been defined as the distance light travels in a vacuum in $1/299\,792\,458$ of a second. The second was once $1/86\,400$ of a mean solar day, but Earth’s rotation is gradually slowing, so since the 1960s the second has been defined by counting $9\,192\,631\,770$ vibrations of a caesium-133 atom. Every speed in this chapter ultimately rests on those two definitions.

Worked Example 1.8: Converting Bolt’s average speed

Usain Bolt ran 100 m in 9.58 s. Express his average speed in metres per second, kilometres per hour, and miles per hour. (1 mi = 1609 m.)

Solution.

$$v = \frac{100 \text{ m}}{9.58 \text{ s}} = 10.44 \text{ m/s.}$$

To kilometres per hour, multiply by 3.6:

$$10.44 \text{ m/s} \times 3.6 = 37.6 \text{ km/h.}$$

To miles per hour, convert metres to miles and seconds to hours:

$$10.44 \frac{\text{m}}{\text{s}} \times \frac{1 \text{ mi}}{1609 \text{ m}} \times \frac{3600 \text{ s}}{1 \text{ h}} = 23.4 \text{ mi/h.}$$

A car doing 37.6 km/h is moving at an ordinary town speed, so Bolt could keep pace with city traffic for about ten seconds. ■

Table 1.1 lists some reference speeds. Keep a few of them in your head; they are the raw material of sanity checks.

1.4.5 Sense-making: a speed that cannot be right

A unit conversion is not busywork. Its real use is to move a number into units where your own experience can judge it, and that judgement is the last line of defence against a wrong answer. Chapter 1 of any physics course should contain at least one number that is obviously absurd once you look at it properly, so here is ours.

Worked Example 1.9: The rental car bill

You hire a car for four days and drive it around a small island. The invoice charges by distance, at \$0.25 per kilometre, and bills you for 48 000 kilometres: a total of \$12,000. The clerk says the figure is taken straight off the car’s odometer, so it cannot be wrong. Show, with a calculation the clerk will understand, that something is.

Solution. The claim is a distance and a time, so turn it into a speed. Over four days,

$$v = \frac{48\,000 \text{ km}}{4 \text{ d}} = 12\,000 \text{ km/d.}$$

Table 1.1: Approximate speeds of familiar things. Values are rounded. Conversions use $1 \text{ h} = 3600 \text{ s}$ and $1 \text{ mi} = 1609.344 \text{ m}$. A handy set of landmarks: 10 m/s is about 36 km/h or 22 mi/h , and 30 m/s is about 108 km/h or 67 mi/h .

Moving object	m/s	km/h	mi/h
Garden snail	0.005	0.02	0.01
Casual walk	1.3	4.7	2.9
Brisk walk	1.8	6.5	4.0
Recreational jog	3	11	6.7
Marathon world-record pace	5.8	21	13
Elite sprinter (average over 100 m)	10.4	37.6	23.4
Elite sprinter (peak)	12.4	44.6	27.7
Highway car	30	108	67
Cruising airliner	250	900	560
Sound in air (room temperature)	343	1235	767

Nobody has any feel for kilometres per day, which is exactly why the number slipped past the clerk. Convert to something familiar. There are 24 h in a day, so

$$12\,000 \text{ km/d} = \frac{12\,000 \text{ km}}{24 \text{ h}} = 500 \text{ km/h},$$

and, using the rough conversion $8 \text{ km} \approx 5 \text{ mi}$,

$$500 \text{ km/h} \times \frac{5 \text{ mi}}{8 \text{ km}} \approx 310 \text{ mi/h}.$$

Now the number can be judged. Table 1.1 puts a highway car at 108 km/h and a cruising airliner at 900 km/h . The bill says you averaged 500 km/h — faster than a Formula One car's top speed — *and* that you did it continuously for four days without stopping to sleep, eat, or refuel.

A second check settles it. Earth's circumference is about $40\,000 \text{ km}$, so $48\,000 \text{ km}$ is more than once around the planet. On a small island.

The resolution is not that the odometer lied. It is that an odometer records the distance a car has travelled *since it was built*, not since you collected it. The company billed you for the car's whole life. What the calculation exposes is not a broken instrument but a wrong assumption about what the instrument measures, and that is the usual way such errors go. Sanity check on the sanity check: a plausible four-day island trip might cover 600 km , at an average of 6.3 km/h over the whole period including nights, or perhaps 60 km/h while actually driving. Those are numbers you can recognise; 500 km/h is not. ■

Convert, then judge

When a calculation produces a number in unfamiliar units, convert it into units you have direct experience of before deciding whether to believe it. 12 000 km/d means nothing to anybody; 310 mi/h is immediately recognisable as impossible. Most errors that survive the algebra do not survive this step.

1.5 Average Speed and Instantaneous Speed

Bolt's 10.44 m/s deserves a closer look. He did not run at that speed for the whole race. He started from rest, took several seconds to reach top speed, ran the middle of the race at roughly 12.4 m/s, and slowed slightly near the finish. The number 10.44 is his **average speed**: total distance divided by total time, with all the speeding up and slowing down smeared together. The number a radar gun would show at any one moment is his **instantaneous speed**.

Two Views: instantaneous speed

Elementary The instantaneous speed at some moment is the average speed over a very short interval around that moment: so short that the object had no time to speed up or slow down appreciably within it. “Very short” depends on the situation. A tenth of a second is short for a sprinter; a whole second is short for a glacier. If shrinking the interval further stops changing the answer, to the precision you care about, you have found the instantaneous speed. A speedometer does exactly this: it measures how far the wheels turn in a small fraction of a second.

Calculus Shrink the interval all the way. The instantaneous speed at time t is the limit of the average speed $\Delta x/\Delta t$ as the interval Δt around t goes to zero, which is the definition of the derivative:

$$v(t) = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}.$$

On a graph of position against time, $\Delta x/\Delta t$ is the slope of the chord joining two points on the curve; as the points slide together the chord becomes the tangent, and the instantaneous speed is the slope of the tangent line. If you have a formula for $x(t)$, the rules of differentiation give you $v(t)$ at every instant at once, with no intervals to shrink.

Definition 1.5: Average and instantaneous speed

The **average speed** over a trip is the total distance travelled divided by the total time:

$$\bar{v} = \frac{d_{\text{total}}}{t_{\text{total}}}.$$

The **instantaneous speed** is the speed at one particular moment: what a speedometer reads. **Elementary** It equals the average speed over a very short time interval around that moment. **Calculus** It is the derivative of position with respect to time, $v = dx/dt$, the slope of the tangent to the position–time graph.

For an object moving at constant speed, average and instantaneous speed are the same, and

every gap in its dot diagram is the same length. For an object that speeds up or slows down, the average speed over a long interval can be quite different from the instantaneous speed at any one point.

There is a practical corollary for dot diagrams, where the shortest interval available is the gap between two dots. If you want the instantaneous speed *at* a particular dot, do not take the gap before it or the gap after it; take the interval from the dot *before* to the dot *after*, and divide by the two intervals of time it spans. The answer is an average, but it is an average over an interval *centred* on the dot you care about, and centring is what makes it a good estimate: the object is running slower than the true value for the first half of the interval and faster for the second, and the two errors largely cancel. Taking the gap on one side alone gives a value that belongs, at best, to the midpoint of that gap rather than to the dot. Worked Example 1.4 uses this.

Estimating an instantaneous speed from a dot diagram

To estimate the speed at dot n , use the dots on either side:

$$v_n \approx \frac{x_{n+1} - x_{n-1}}{2\tau},$$

where τ is the time between dots. Phrase it carefully too: the instantaneous speed does not “occur between” two dots. It is the speed *at* one instant, and the centred average over the surrounding interval is our best *estimate* of it.

Worked Example 1.10: Bolt’s race, twenty metres at a time

In Bolt’s world-record race in Berlin in 2009, timing equipment recorded when he passed each 20 m mark. The times, measured from the starting gun, were 2.88 s, 4.64 s, 6.31 s, 7.92 s, and 9.58 s. Use them to describe how his speed changed during the race.

Elementary Each 20 m segment is a dot-diagram interval in reverse: the distance is fixed and the time varies. The average speed over each segment is 20 m divided by the time spent in it.

mark (m)	time at mark (s)	time in segment (s)	average speed (m/s)
20	2.88	2.88	6.9
40	4.64	1.76	11.4
60	6.31	1.67	12.0
80	7.92	1.61	12.4
100	9.58	1.66	12.0

The first segment is slow because it includes the reaction to the gun (about 0.15 s) and the start from rest. Bolt was still speeding up through 60 m, was fastest between 60 m and 80 m, and slowed slightly at the end. No segment average equals the race average of 10.44 m/s; the race average is pulled down by the slow start. The 12.4 m/s of the fastest segment is itself an average over 1.61 s; his true peak instantaneous speed was a little higher, but the 20 m intervals are too coarse to show it.

Calculus A smooth model for a sprinter's speed is

$$v(t) = v_{\max} (1 - e^{-t/\tau}),$$

which starts at zero, rises quickly, and levels off at $v_{\max}$; the constant τ sets how quickly. Position is recovered by integrating from the start, and the integral of $e^{-t/\tau}$ is $-\tau e^{-t/\tau}$:

$$x(t) = \int_0^t v(t') dt' = v_{\max} [t - \tau (1 - e^{-t/\tau})].$$

With $v_{\max} = 12.4 \text{ m/s}$, the condition $x(9.58 \text{ s}) = 100 \text{ m}$ gives $\tau = 1.52 \text{ s}$. The model then predicts the 20 m marks at 2.9, 4.7, 6.3, 7.9, and 9.6 s, within about a tenth of a second of the measured times. Differentiating $x(t)$ returns $v(t)$, as it must, and the average speed is the integral of v divided by the time,

$$\bar{v} = \frac{1}{T} \int_0^T v(t) dt = \frac{x(T)}{T} = \frac{100 \text{ m}}{9.58 \text{ s}} = 10.44 \text{ m/s},$$

the same number as before, now seen as the average value of a function. The model ignores the reaction time and the slight slowing at the end, which is why it is a model and not a record. ■

1.5.1 From speed back to distance

Average speed is the quantity a driver planning a trip cares about, because it gives the travel time. Rearranging the definition,

$$d_{\text{total}} = \bar{v} t_{\text{total}},$$

which holds for *any* motion, however uneven, provided $\bar{v}$ is the average speed over the whole interval. If your average speed on a four-hour drive is 80 km/h, you covered 320 km, regardless of how many red lights you met along the way. Equation (1.2), $d = vt$, is the special case in which the speed never changed.

Two Views: distance from speed

Elementary On a graph of speed against time, a constant speed is a horizontal line, and $d = vt$ is the area of the rectangle under it: height v , width t . If the speed changes, cut the time into short pieces on which the speed is nearly constant, multiply on each piece, and add. The total is the area under the speed–time graph. Dividing that area by the total time gives the average speed, so $d_{\text{total}} = \bar{v} t_{\text{total}}$ is the statement that a rectangle of height $\bar{v}$ has the same area as the curve.

Calculus “Cut, multiply, add, and let the pieces shrink” is the definition of the definite integral. Because $v = dx/dt$, the fundamental theorem of calculus gives

$$d = \int_{t_1}^{t_2} v(t) dt, \quad \bar{v} = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} v(t) dt. \quad (1.4)$$

The second formula is the average value of the function v on the interval, and the first says the distance is the area under its graph. For a constant v the integral is $v(t_2 - t_1)$ and

Equation (1.2) drops out.

Check Your Understanding 1.7

A car's odometer reads zero at the start of a trip and 40 km half an hour later. (a) What was the car's average speed? (b) Could the car have achieved this average without ever going faster than 80 km/h, if it started from rest? (c) When a police officer writes a speeding ticket, is it for the driver's average speed or instantaneous speed?

Worked Example 1.11: The round-trip trap

You cycle 12 km to a friend's house at 24 km/h, then cycle home along the same road at 12 km/h. What was your average speed for the whole trip?

Elementary The tempting answer is the average of the two speeds, 18 km/h. It is wrong. Average speed is total distance over total time, so we need the time for each leg.

$$t_{\text{out}} = \frac{12 \text{ km}}{24 \text{ km/h}} = 0.50 \text{ h}, \quad t_{\text{back}} = \frac{12 \text{ km}}{12 \text{ km/h}} = 1.00 \text{ h}.$$

Then

$$\bar{v} = \frac{24 \text{ km}}{1.50 \text{ h}} = 16 \text{ km/h}.$$

The slow leg took twice as long, so it counts for twice as much in the average.

Calculus The speed–time graph is two rectangles: height 24 km/h for 0.5 h, then height 12 km/h for 1.0 h. By Equation (1.4),

$$\bar{v} = \frac{1}{1.5} \int_0^{1.5} v \, dt = \frac{24(0.5) + 12(1.0)}{1.5} \text{ km/h} = 16 \text{ km/h}.$$

The time average weights each speed by how long it lasted. In general, for two legs of equal length d at speeds v_1 and v_2 ,

$$\bar{v} = \frac{2d}{d/v_1 + d/v_2} = \frac{2v_1v_2}{v_1 + v_2},$$

the *harmonic* mean of the two speeds, which is always less than their ordinary average unless $v_1 = v_2$ (Exercise 1.37). Whenever the legs of a trip take different times, averaging the speeds directly gives the wrong answer. ■

1.5.2 Position–time graphs

A dot diagram is a picture of *where* an object was; it hides *when* in the spacing between dots. A **position–time graph** puts both on the page. Plot time along the horizontal axis and position along the vertical axis, and mark one point for each dot. Figure 1.6 does this for the walker of Worked Example 1.5.

The points lie on a straight line through the origin. The *slope* of that line, rise over run, is

$$\frac{\Delta x}{\Delta t} = \frac{3.08 \text{ m}}{2.36 \text{ s}} = 1.31 \text{ m/s},$$




Figure 1.6: Position–time graph for a walker with 1.54 m strides every 1.18 s. The points fall on a straight line whose slope is the speed. The red triangle shows a rise of 3.08 m over a run of 2.36 s, a slope of 1.31 m/s.

which is the walker’s speed. This is a general and very useful fact.

Slope of a position–time graph

On a position–time graph, the slope of the straight line (the *chord*) joining two points equals the average speed between those two instants. **Calculus** The slope of the *tangent* at a point equals the instantaneous speed there, dx/dt . A straight graph means constant speed. A graph that bends upward, getting steeper, means the object is speeding up; one that flattens means it is slowing down. A horizontal stretch means the object is not moving at all.

You may notice that the dot diagram of a constant-speed object and its position–time graph carry exactly the same information. They are two views of one model. Which one to draw depends on the question: dot diagrams are quick to sketch and good for comparing objects, while graphs are better for reading off numbers and for handling changing speeds. Figure 1.7 shows what a changing speed looks like.

Worked Example 1.12: Reading a curved position–time graph

From Figure 1.7(a), estimate Bolt’s instantaneous speed 5.0 s into the race, and compare it with his average speed up to that moment.

Elementary Lay a ruler along the curve at $t = 5.0 \text{ s}$ so that it just touches without cutting: that is the tangent, drawn in red. Read two convenient points on it, for example $t = 3.0 \text{ s}$, $x \approx 20 \text{ m}$ and $t = 7.0 \text{ s}$, $x \approx 68 \text{ m}$. Its slope is

$$\frac{\Delta x}{\Delta t} \approx \frac{68 \text{ m} - 20 \text{ m}}{7.0 \text{ s} - 3.0 \text{ s}} = 12 \text{ m/s}.$$

The average speed up to 5.0 s is the slope of the chord from the origin to the point on the curve at $t = 5.0 \text{ s}$, where $x \approx 44 \text{ m}$: about 8.8 m/s. The instantaneous speed is much larger




Figure 1.7: Bolt's world-record race. (a) Position against time: the dots are the measured 20 m split times and the curve is the model of Worked Example 1.10. The graph bends upward as he speeds up, then becomes nearly straight. The dashed chord from the start has slope equal to the average speed; the tangent at 5 s has slope equal to the instantaneous speed then. (b) The model's speed against time. The shaded area under the curve is the distance run, 100 m, and the dashed line is the average speed: a rectangle of that height has the same area as the curve.

than the average because the average still remembers the slow start. Reading a tangent by eye is good to perhaps ten percent; that is usually enough for a sanity check.

Calculus Differentiate the model. Since $d(e^{-t/\tau})/dt = -(1/\tau)e^{-t/\tau}$,

$$v(t) = \frac{dx}{dt} = v_{\max} \left[1 - \tau \cdot \frac{1}{\tau} e^{-t/\tau} \right] = v_{\max} (1 - e^{-t/\tau}),$$

which is the speed function we started from, as it should be. At $t = 5.0$ s, $e^{-5.0/1.52} = 0.037$, so $v = 12.4(1 - 0.037) = 11.9$ m/s. The position at that moment is $x(5.0) = 12.4[5.0 - 1.52(1 - 0.037)] = 43.8$ m, giving an average of $43.8/5.0 = 8.8$ m/s. The ruler estimate of the tangent slope was within one percent of the derivative; the graph and the formula are two views of the same motion. ■

Check Your Understanding 1.8

A cart's position along a track is $x(t) = (1.5 \text{ m/s}^2)t^2$ for the first four seconds. (a) Find its average speed over the four seconds. (b) **Elementary** Estimate its instantaneous speed at $t = 4.0$ s from the average over the interval 3.9 s to 4.1 s. **Calculus** Confirm by differentiating. (c) Why are the answers to (a) and (b) different?

1.6 A Model of Stride Time

We now have the tools to see Webb's problem clearly. He measured the hunter's stride length: 3.73 m. To compute a speed, he also needed the time between strides, and there is no way to read time from a footprint. He could have guessed, but a guess is not science. Instead, he did what scientists usually do: he borrowed a model that someone else had already built and tested.

1.6.1 Why height matters

Watch a small child walking beside an adult. To keep up, the child takes many quick, short steps while the adult takes a few long, unhurried ones. The child's time between strides is shorter. Leg length, and therefore height, affects how quickly a person steps at a given speed.

Webb did not know the hunter's height, but he did know the hunter's foot length, and taller people tend to have longer feet. That opens a chain of inferences:

foot length $\longrightarrow$ height $\longrightarrow$ (with stride length) time between strides $\longrightarrow$ speed.

Each arrow is a step in the model, and each step carries its own uncertainty.

1.6.2 The Charteris model

The equations Webb used came from a 1981 study by three Canadian researchers, John Charteris, Jack Wall, and John Nottrodt. They had been asked a similar question about far older footprints: the 3.6-million-year-old hominin trackway at Laetoli in Tanzania. To answer it, they measured modern volunteers walking at a range of speeds and recorded stride length, foot length, and height. From those data they fitted equations that take stride length and foot length as inputs and return an estimated time between strides.

To make the data from people of different sizes comparable, Charteris and his colleagues worked with ratios rather than raw measurements. One of these is the **relative speed**.

Definition 1.6: Relative speed

The **relative speed** of a walker or runner is their speed divided by their height:

$$\text{relative speed} = \frac{\text{speed}}{\text{height}}.$$

Its unit is m/s/m = 1/s. It tells you how many body-lengths the person covers each second.

A person 1.8 m tall walking at 1.4 m/s has a relative speed of $1.4/1.8 = 0.78/\text{s}$: about three-quarters of a body length per second. A 1.2 m child walking at the same speed has a relative speed of 1.17/s, which is why the child looks so much busier.

1.6.3 The famous result

Feeding the hunter's stride length and foot length into the Charteris equations, Webb obtained a time between strides of 0.360 s. Combined with the measured stride length,

$$v = \frac{d}{t} = \frac{3.73 \text{ m}}{0.360 \text{ s}} = 10.4 \text{ m/s}.$$

Worked Example 1.13: Comparing the hunter to Usain Bolt

Webb's model gives the hunter a speed of 10.4 m/s. Usain Bolt's world-record 100 m time is 9.58 s. Compare the two.

Solution. Bolt's average speed over the race was

$$\bar{v} = \frac{100 \text{ m}}{9.58 \text{ s}} = 10.44 \text{ m/s.}$$

The hunter's modelled speed is within one percent of Bolt's average. Taken at face value, Webb's result says a barefoot hunter on a muddy lake shore 20 000 years ago ran essentially as fast as the fastest human ever timed. That is the claim that made the newspapers. Notice, too, what kind of speed each number is. Bolt's is an average over a race that began from a standstill; the hunter's stride-by-stride figure is closer to an instantaneous speed in mid-run. A fair comparison would set the hunter against Bolt's mid-race 12.4 m/s, which makes the claim less dramatic but does not yet make it wrong. ■

That comparison should make you a little uneasy. Extraordinary results deserve extra scrutiny. Before believing a number, a scientist asks: does the model that produced it hold up?

1.7 Questioning a Model

There are two broad ways to test a model's output. The first is to look at its *other* outputs, the ones nobody put in the headline, and ask whether they make sense. The second is to look inside the model and ask whether it was ever meant for this job. Webb's result fails both tests.

1.7.1 Sanity checks

Definition 1.7: Sanity check

A **sanity check** is a quick, rough test of whether a result is believable. It compares the result, or the intermediate quantities that produced it, against known values, limiting cases, or everyday experience. A sanity check cannot prove a result correct, but it can quickly reveal that something has gone wrong.

Compare the hunter with Bolt in more detail, using numbers that are known for Bolt and predicted by the model for the hunter:

	Usain Bolt (measured)	Hunter (modelled)
Stride length	4.88 m	3.73 m
Time between strides	0.467 s	0.360 s
Height	195 cm	194 cm
Speed	10.44 m/s	10.36 m/s

Bolt's strides are more than a metre longer than the hunter's. On stride length alone, you would expect Bolt to be much faster. The only reason the two speeds come out nearly equal is

that the model assigns the hunter an extraordinarily short stride time, about 0.11 s quicker than Bolt's. Could that be because the hunter was much shorter, with shorter legs that cycle faster? No: the model also predicts that the hunter was 194 cm tall, within a centimetre of Bolt. The model is claiming that a man the same height as Bolt, taking strides a metre shorter than Bolt's, turned his legs over far faster than the greatest sprinter in history. That is hard to believe.

Webb applied the same model to other trackways at the site. None came out as fast as the hunter, but several produced oddly short stride times of their own. When a model produces strange numbers across many inputs, the problem is more likely in the model than in the inputs.

1.7.2 Looking inside the model: extrapolation

Figure 1.8 is a schematic version of one of the plots from the Charteris study. It shows the data the equations were built from. The horizontal axis is relative speed. The vertical axis is stride length measured in foot-lengths, another size-independent ratio.




Figure 1.8: Schematic of the walking data behind the Charteris model, redrawn after Charteris, Wall, and Nottrodt (1981). Points show group means with a bar indicating the spread. The dashed line is a rough straight-line fit. The data stop at a relative speed of about 1.2/s.

Now compute Bolt's relative speed:

$$\frac{10.44 \text{ m/s}}{1.95 \text{ m}} = 5.35/\text{s}.$$

The fastest volunteer in the Charteris data had a relative speed of about 1.2/s. Bolt, and the hunter that Webb's model made Bolt's equal, are more than four times beyond the edge of the data. There is a simpler way to say this: the Charteris volunteers were *walking*. Their model was a model of walking. Webb used it on a sprinter.

Definition 1.8: Interpolation and extrapolation

Using a model to make predictions *inside* the range of conditions it was built and tested on is called **interpolation**. Using it *outside* that range is called **extrapolation**. Interpolation

is usually safe. Extrapolation is a gamble: nothing in the data says how the relationship behaves out there, and the further you go, the worse the gamble.

Two Views: why a fitted line cannot be trusted far from its data

Elementary A straight line fitted to data says: “over the range I was fitted to, each extra unit on the horizontal axis adds the same amount on the vertical axis.” The data cannot say whether that stays true outside the range, because nobody looked there. Real relationships usually bend sooner or later: children stop growing, engines hit a top speed, walkers break into a run. The line does not know about the bend, so the further you follow it past the last data point, the more likely it is to be badly wrong.

Calculus Suppose the true relationship is some smooth curve $y = f(x)$. Near any point x_0 the curve is well approximated by its tangent line,

$$f(x) \approx f(x_0) + f'(x_0)(x - x_0),$$

and a straight-line fit to data clustered near x_0 is essentially an estimate of this tangent: its slope is the derivative $f'(x_0)$ there. But a derivative is a *local* quantity. It describes how f changes right at x_0 , and nothing forces f' to keep the same value far away. Extrapolating a fitted line assumes f' is constant everywhere, which is to assume the curve is a line. The error of the tangent approximation grows with the square of the distance $(x - x_0)^2$ times the curvature f'' , so it is small nearby and can be enormous far off. That is the precise sense in which interpolation is safe and extrapolation is a gamble.

Worked Example 1.14: Extrapolating a straight line

Between the ages of 4 and 10 a child grows from about 100 cm to about 136 cm. Fit a straight line to these figures and use it to “predict” the person’s height at age 40. Then explain what went wrong.

Elementary The growth rate over the six years is

$$\frac{136 \text{ cm} - 100 \text{ cm}}{10 \text{ yr} - 4 \text{ yr}} = 6 \text{ cm per year}.$$

Following the line for a further 30 years adds $30 \times 6 \text{ cm} = 180 \text{ cm}$, for a height of 316 cm at age 40. Nobody is three metres tall. The straight line was a good description of growth between 4 and 10, and would be trustworthy at age 7 (interpolation), but growth slows in the teens and stops around 18. The line has no way of knowing that; only data from those ages could tell it.

Calculus Let $h(t)$ be height as a function of age. Between 4 and 10 the derivative dh/dt is roughly 6 cm per year, and the line we drew is the tangent approximation $h(t) \approx h(4) + h'(4)(t - 4)$. Real growth has $h'(t)$ rising briefly in the growth spurt and then falling to zero, so h'' is strongly negative in the late teens and the tangent approximation fails badly beyond that. The error of a tangent approximation is roughly $\frac{1}{2}h''(t - t_0)^2$; with $(t - t_0) = 36$ years, even a modest curvature produces an error of metres. Webb’s use

of the Charteris line was the same mistake with a different curve: a walking relationship extended, without any data, to a sprint. ■

Extrapolation is not always wrong. Every prediction about the future is one. But a model must be extrapolated with eyes open, and the result treated with suspicion until it is confirmed. Running is not simply fast walking. In a walk, one foot is always on the ground; in a run, both feet leave the ground for part of every stride. The relationship between stride length, stride time, and speed changes when that happens. The Charteris equations know nothing about it.

1.7.3 How the mistake spread

Webb could have caught the problem in several ways: by reading the original study closely enough to notice that it concerned walking; by sanity-checking his output against known sprinters, as we did above; or by asking a colleague who worked with gait models regularly. None of that happened, and the reviewers who checked the paper before publication did not notice either. Once published, the number took on a life of its own. Journalists repeated it, then authors repeated the journalists. Hardly anyone went back to the source.

Lessons from the ancient sprinter

1. **Know what your model was built for.** A model fitted to walking data is a model of walking.
2. **Sanity-check the result.** Compare it, and the intermediate quantities, against things you already know.
3. **Check the other outputs.** A model that gives one spectacular answer and several odd ones is probably wrong about all of them.
4. **Ask for help.** Someone who uses a model daily will spot misuse in seconds.
5. **Check the source.** A number that has been repeated a hundred times is not thereby a hundred times more reliable. In science, an appeal to authority or to popularity carries little weight; a single verifiable check carries a great deal.
6. **Be willing to be wrong.** Scientists, like everyone else, have a large capacity for fooling themselves, and the most widely shared assumptions are often the least questioned. The scientific attitude is one of inquiry, integrity, and humility. Changing your mind in the face of evidence is a strength, not a weakness.

1.8 Uncertainty and a Better Estimate

In 2013 two Spanish researchers, Javier Ruiz and Angélica Torices, revisited the Willandra footprints with a model designed for the right job. Their equations were built from measurements of people actually running, on a running track and on a beach, so that the hunter's conditions fell inside the range of the data rather than far outside it. They also did something Webb had not: they reported how confident they were.

Their estimate for the hunter's speed was

$$v = (7.2 \pm 1.0) \text{ m/s.}$$

1.8.1 Reading an uncertainty

The notation (7.2 ± 1.0) m/s means that the best estimate is 7.2 m/s, and the true value is most likely somewhere between 6.2 m/s and 8.2 m/s. The “ ± 1.0 ” is the **uncertainty**. It is not an admission of sloppiness. It is an honest statement of how precisely the method can pin the answer down.

Definition 1.9: Uncertainty

A measured or calculated quantity is reported as $x \pm \delta x$, where x is the best estimate and δx is the **absolute uncertainty**, in the same units as x . The **percentage uncertainty** is

$$\frac{\delta x}{x} \times 100\%.$$

For the Ruiz and Torices estimate, the percentage uncertainty is $1.0/7.2 \times 100\% \approx 14\%$. That is fairly large, and appropriately so: the estimate rests on a single trackway from one runner who lived twenty thousand years ago, in conditions the model can only approximate.

1.8.2 What the revised number says

A speed of 7.2 m/s is 26 km/h. That is a fast run, roughly the pace of a 100 m race in 14 seconds, sustained barefoot across soft mud while chasing an animal. It is impressive. It is also nowhere near an Olympic sprinter. Even the top of the range, 8.2 m/s, falls well short of Bolt’s 10.44. The headline was wrong; the hunter was fast, not superhuman.

1.8.3 Carrying uncertainty through a calculation

An uncertain input produces an uncertain output. If the speed is known only to within ± 1.0 m/s, then anything computed from the speed, such as the stride time, is uncertain too, and we should say by how much.

Two Views: propagating an uncertainty

Elementary Do the calculation three times: once with the best estimate, and once with each end of the input’s range. The spread of the three outputs is the output’s uncertainty. This always works, and for a single uncertain input it is the method to reach for first. Its only drawback is that the two ends may give unequal deviations, and then you must decide how to report them.

Calculus If the output y is a function of the input x , then a small change δx in the input changes the output by approximately

$$\delta y \approx \left| \frac{dy}{dx} \right| \delta x,$$

the tangent-line approximation again. For $y = d/x$ (a quotient with a fixed numerator), $dy/dx = -d/x^2$, so

$$\delta y \approx \frac{d}{x^2} \delta x = \frac{d}{x} \cdot \frac{\delta x}{x} = y \frac{\delta x}{x}, \quad \text{that is,} \quad \frac{\delta y}{y} \approx \frac{\delta x}{x}.$$

Dividing a fixed number by an uncertain one passes the *percentage* uncertainty through unchanged. The same is true of multiplying: $y = cx$ gives $\delta y/y = \delta x/x$. This one-line rule replaces the three calculations, and it is exact in the limit of small uncertainties.

Worked Example 1.15: Propagating uncertainty to stride time

Using Webb's measured stride length of 3.73 m and the Ruiz–Torices speed of (7.2 ± 1.0) m/s, estimate the hunter's time between strides and its uncertainty. Ignore any uncertainty in the stride length.

Elementary The best estimate comes from the best-estimate speed:

$$t = \frac{d}{v} = \frac{3.73 \text{ m}}{7.2 \text{ m/s}} = 0.52 \text{ s}.$$

To find the uncertainty, repeat the calculation at the two ends of the speed range. A faster speed gives a shorter stride time and vice versa:

$$t_{\text{fast}} = \frac{3.73 \text{ m}}{8.2 \text{ m/s}} = 0.45 \text{ s}, \quad t_{\text{slow}} = \frac{3.73 \text{ m}}{6.2 \text{ m/s}} = 0.60 \text{ s}.$$

The best estimate sits about 0.07 s above the low end and 0.08 s below the high end, so we report $t \approx (0.52 \pm 0.08)$ s, rounding the uncertainty up.

Calculus The percentage uncertainty in the speed is $1.0/7.2 = 14\%$, and $t = d/v$ inherits it:

$$\delta t \approx t \frac{\delta v}{v} = 0.518 \text{ s} \times 0.139 = 0.072 \text{ s},$$

so $t \approx (0.52 \pm 0.07)$ s. The derivative gives a single symmetric figure that sits between the two elementary deviations, 0.066 and 0.083; the difference between them is the curvature of $1/v$, which the tangent approximation ignores. Either report is acceptable, and both say the same thing: the stride time is known to about one part in seven.

Notice that both estimates are longer than Bolt's 0.467 s, as they should be for a shorter-striding, slower runner. The revised model passes the sanity check that Webb's failed.



Three habits of careful modelling

- **Test models against real data** in the conditions where you intend to use them.
- **Remember each model's limits**, and sanity-check anything that comes out of it.
- **Report results with uncertainty**, so that others know how much weight the number can bear.

1.9 From Speed to Velocity

Everything so far has been about *how fast* an object moves. We have said nothing about *which way*. Recall that a dot diagram cannot even tell you whether the object went left or right. For many questions that is fine. For many others, direction is the whole point. A hunter running

toward an animal and one running away from it have the same speed and very different prospects.

1.9.1 Position, displacement, and direction

To keep track of direction, choose a line along the motion, pick a point on it as the origin, and pick one direction as positive. Then every point on the line has a **position** x , a number with a sign. If the hunter runs east and we call east positive, positions ahead of the origin are positive and positions behind it are negative.

Definition 1.10: Displacement

The **displacement** of an object over some time interval is its change in position:

$$\Delta x = x_{\text{final}} - x_{\text{initial}}.$$

Displacement has a sign. Positive displacement means net motion in the positive direction; negative means the opposite. Displacement is *not* the same as distance travelled: an object that goes out and comes back has zero displacement but nonzero distance.

1.9.2 Coordinates are a choice

Setting up that line, origin, and positive direction is called choosing a **coordinate system**, and the number it assigns to an object is the object's **coordinate**. It is worth separating two words that everyday speech runs together.

Definition 1.11: Coordinate and position

An object's **position** is where it actually is: a particular place on the track, the road, or the lake shore. Its **coordinate** is the *number* we assign to that place once we have chosen an origin and a positive direction. The position is a fact about the world; the coordinate is a fact about the world *and* about our bookkeeping. Motion along a single line needs one coordinate and is called **one-dimensional**; motion across a floor needs two; motion through the air needs three.

Nothing forces a particular choice of origin, and different people will make different ones. Figure 1.9 shows a skater, Émilie, on a straight track, measured two ways: x from the left-hand end of the track, and x' from a flag planted 3.2 m along it. The skater is in one place. The two coordinate systems disagree about what to call it.

Which is correct? Both. A coordinate system is a tool, chosen for convenience, and the only rule is to say which one you are using and then stick to it. But this raises a question worth taking seriously: if two physicists can disagree about the numbers, what in the description is actually about the skater?

What a change of origin does and does not change

Move the origin and *every coordinate changes*. Nothing else does. In Figure 1.9 the two systems are related by $x' = x - 3.2$ m, so for any two instants,

$$\Delta x' = x'_{\text{final}} - x'_{\text{initial}} = (x_{\text{final}} - 3.2) - (x_{\text{initial}} - 3.2) = \Delta x.$$




Figure 1.9: One skater, two coordinate systems. Measuring from the left-hand end of the track gives the coordinate $x = 4.3$ m; measuring from the flag gives $x' = 1.1$ m. The skater has not moved. What changed is the origin, and with it the bookkeeping: $x' = x - 3.2$ m at every point of the track.

The shift cancels in the subtraction. So the *displacement* is the same in both systems, and therefore so are the average velocity $\Delta x/\Delta t$, the speed, and the distance travelled. The skater's position, her motion, and every physical question you could ask about her are untouched.

This is the reason displacement earns a symbol of its own. Writing $x_{\text{final}} - x_{\text{initial}}$ keeps two origin-dependent numbers in view; writing Δx keeps only the origin-independent difference, which is the part that is about the skater. Physicists care a great deal about which quantities survive a change of bookkeeping, and you will meet the idea again whenever a new coordinate system appears.

Coordinate, position, and a sentence to avoid

You will hear, and may be tempted to write, “the skater's position is -2 m.” Strictly this mixes the two words: her *coordinate* is -2 m, in some coordinate system that ought to be named, while her *position* is a place on the track. The loose phrasing is common and usually harmless, and this book uses it too. It stops being harmless the moment two coordinate systems are in play at once, which is exactly when problems get interesting, so it is worth being able to say the careful version when you need it.

The skater's namesake

The skater in these figures is called Émilie, after Émilie du Châtelet (1706–1749), who did as much as anyone to give mechanics its modern vocabulary. Her annotated French translation of Newton's *Principia* was published a decade after her death and is still the standard one. She also took the unfashionable side of a long argument about what quantity is conserved when bodies collide, holding that it goes as mv^2 rather than mv ; both turn out to matter, and the first is what we now call kinetic energy. A good deal of what makes this chapter's notation feel obvious was, in her lifetime, contested.

1.9.3 A negative coordinate is not a negative velocity

Signs now appear in two different places, and confusing them is the most common error in this part of the subject. The *coordinate* x carries a sign that says where the object is relative to the origin. The *velocity* v_x carries a sign that says which way the object is going. They are independent: all four combinations occur.




Figure 1.10: A skater at four moments. The dot gives the coordinate, the arrow gives the direction of travel. The sign of x and the sign of v_x are set independently, and all four combinations happen. In (a) the skater is to the left of the origin and moving right: the coordinate is negative and *increasing*, so the velocity is positive.

Case (a) is the one that catches people. The skater is at a negative coordinate, so it feels natural to call the motion negative; but she is heading toward the origin, her coordinate is climbing from -2 m toward zero, and a climbing coordinate is a positive velocity. The test is never where the object is. It is always which way the coordinate is *changing*:

$$v_x > 0 \iff x \text{ is increasing}, \quad v_x < 0 \iff x \text{ is decreasing},$$

whatever the sign of x itself. Notice also that the sign of the velocity, unlike the sign of the coordinate, survives a change of origin: shifting the origin cannot turn rightward motion into leftward motion.

Check Your Understanding 1.9

A skater moves along a straight track with x measured rightward from a flag. Give the sign of her coordinate and the sign of her velocity in each case, or say if it cannot be determined. (a) She is 5 m left of the flag, gliding toward it. (b) She is 5 m left of the flag, gliding away from it. (c) She is level with the flag, gliding rightward. (d) She is 3 m right of the flag and momentarily stationary at the top of a rise. (e) She is 3 m right of the flag, and her speed is 2 m/s. Then say which of your answers would change if the flag were moved 10 m to the left.

1.9.4 Velocity

Definition 1.12: Average velocity

The **average velocity** of an object over a time interval Δt is its displacement divided by the elapsed time:

$$\bar{v}_x = \frac{\Delta x}{\Delta t} = \frac{x_{\text{final}} - x_{\text{initial}}}{t_{\text{final}} - t_{\text{initial}}}.$$

Velocity has both a size and a direction. Along a line, the direction is carried by the sign. The size of the velocity, ignoring its sign, is the speed. The subscript x records that this is the velocity along the x line, with the sign convention chosen for x .

Two Views: velocity, displacement, and distance

Elementary Velocity is speed with a sign. Over an interval, the average velocity is the displacement divided by the time, so it can be positive, negative, or zero; the average speed is the total distance divided by the time and is never negative. The two agree in size only when the object never reverses. On a position–time graph the slope of a chord is now the average velocity: rising lines mean positive velocity, falling lines negative. The instantaneous velocity is the slope over a very short interval, and the instantaneous speed is its size.

Calculus The instantaneous velocity is the derivative of position, sign included:

$$v_x = \frac{dx}{dt}, \quad \text{speed} = |v_x|.$$

Integrating recovers displacement and distance, and the difference between them is exactly the difference between v_x and $|v_x|$:

$$\Delta x = \int_{t_1}^{t_2} v_x \, dt, \quad d = \int_{t_1}^{t_2} |v_x| \, dt.$$

Where the velocity is negative the first integral counts area below the axis as negative; the second counts all area as positive. That is why displacement can vanish while distance does not.

Speed answers “how fast?” Velocity answers “how fast, and which way?” If a car travels at 60 km/h, we know its speed; if it travels at 60 km/h *to the north*, we know its velocity. Two cars passing each other on a highway at 30 m/s have the same speed but opposite velocities: +30 m/s and −30 m/s.

Definition 1.13: Vector and scalar quantities

A quantity that needs both a magnitude and a direction for its complete description is a **vector quantity**. Velocity is a vector quantity; so are force and acceleration, which you will meet in later chapters. A quantity that is fully described by a magnitude alone is a **scalar quantity**. Speed is a scalar, and so are distance, time, and mass.

The symbols, and what the subscript means

A single letter's worth of difference separates three quantities that are easy to say but easy to mix up. The rule is short: **a subscript names an axis and keeps the sign; a bare v means speed and never carries one.**

symbol	name	sign	read it as
x	coordinate along the x axis	either	"where it is, relative to the origin"
Δx	displacement along x	either	"how far and which way it moved"
v_x	velocity along x	either	"how fast and which way it is going"
v	speed	never negative	"how fast it is going"

They are tied together by

$$\Delta x = x_{\text{final}} - x_{\text{initial}}, \quad \bar{v}_x = \frac{\Delta x}{\Delta t}, \quad v = |v_x|.$$

The same pattern runs on every other axis: vertical motion uses y , Δy , and v_y , and in the next chapter accelerations join in as a_x and a_y . Only the bare speed v stands outside the scheme, because it has thrown its direction away and so has no axis to name.

Check Your Understanding 1.10

An object moves along the x axis. At one instant $x = -7\text{ m}$ and $v_x = +4\text{ m/s}$. (a) What is its speed? (b) Is it moving toward or away from the origin? (c) Write the symbol for each of: its distance from the origin; its velocity; the change in its coordinate over the next 2 s. (d) A classmate writes " $v = -4\text{ m/s}$." What is wrong with that, and what did they probably mean?

In this chapter, where motion is along a single line, the direction of a vector is carried by its sign, and that is all the vector machinery we need. In two or three dimensions a velocity is drawn as an arrow whose length gives the speed and whose orientation gives the direction. We return to that picture when we study motion in a plane.

Constant velocity means straight-line motion at constant speed

Constant *speed* means the object neither speeds up nor slows down. Constant *velocity* means constant speed *and* constant direction, and a constant direction is a straight line. A car rounding a curve with its speedometer needle steady has a constant speed but a changing velocity, because its direction changes at every instant. Whenever either the speed or the direction changes, the velocity changes. In the next chapter this distinction becomes the basis of *acceleration*.

Check Your Understanding 1.11

For a certain period of time the speedometer of a car reads a steady 60 km/h. Does this tell you the car has a constant speed? A constant velocity? Explain.

Worked Example 1.16: Speed versus velocity on a return trip

A runner jogs 40 m east along a straight path in 8.0 s, turns around immediately, and jogs 40 m west back to the start in 10.0 s. Find the average speed and the average velocity for the whole trip. Take east as positive.

Elementary The total distance is 80 m and the total time is 18.0 s, so

$$\text{average speed} = \frac{80 \text{ m}}{18.0 \text{ s}} = 4.4 \text{ m/s}.$$

The runner ends where they started, so the displacement is $\Delta x = 0$ and

$$\bar{v}_x = \frac{0}{18.0 \text{ s}} = 0.$$

The runner was certainly moving, but on average they went nowhere. This is not a paradox; it is what velocity is designed to measure. If we instead wanted the velocity on the way back only, it would be $\Delta x/\Delta t = (-40 \text{ m})/(10.0 \text{ s}) = -4.0 \text{ m/s}$, negative because the motion is westward.

Calculus Model each leg at constant velocity: $v_x = +5.0 \text{ m/s}$ for the first 8.0 s and $v_x = -4.0 \text{ m/s}$ for the next 10.0 s. Then

$$\begin{aligned}\Delta x &= \int_0^{18} v_x dt = (5.0)(8.0) + (-4.0)(10.0) = 40 - 40 = 0, \\ d &= \int_0^{18} |v_x| dt = 40 + 40 = 80 \text{ m}.\end{aligned}$$

Dividing by 18.0 s gives $\bar{v}_x = 0$ and an average speed of 4.4 m/s, as before. On the velocity–time graph the first rectangle sits above the axis and the second below it, with equal areas; the displacement is the signed area, zero, and the distance is the unsigned area. ■

1.9.5 Relative velocity

Section 1.4 pointed out that every speed is measured relative to something. With signs in hand we can say precisely how the velocity of an object depends on who is measuring it.

Two Views: relative velocity along a line

Elementary Suppose two objects, A and B, move along the same line, with velocities v_A and v_B measured relative to the ground (signs included). The velocity of A *as seen from* B is

$$v_{A \text{ rel } B} = v_A - v_B.$$

Think of it this way: every second, A moves v_A along the line and B moves v_B , so the gap between them changes by $v_A - v_B$. Two cars driving the same way at 100 km/h and 98 km/h close at 2 km/h; two cars approaching head-on at 100 km/h each, one with velocity +100 and the other −100, close at 200 km/h. The subtraction handles both cases without special rules, provided the signs are kept.

Calculus Let $x_A(t)$ and $x_B(t)$ be the two positions. The position of A relative to B is the difference $x_A - x_B$, and the relative velocity is its derivative:

$$v_{A \text{ rel } B} = \frac{d}{dt}(x_A - x_B) = \frac{dx_A}{dt} - \frac{dx_B}{dt} = v_A - v_B,$$

because the derivative of a difference is the difference of the derivatives. Nothing here required the velocities to be constant; the rule holds instant by instant.

Worked Example 1.17: Closing speeds

You are driving east at 25 m/s. (a) A car ahead of you is driving east at 20 m/s. (b) A car on the other side of the road is driving west at 20 m/s. In each case find the other car's velocity relative to you, and say how fast the gap between you is changing.

Elementary Take east as positive, so your velocity is $v_{\text{you}} = +25$ (all in m/s). (a) The car ahead has $v = +20$; relative to you its velocity is $20 - 25 = -5$ m/s: it drifts toward you (westward, in your frame) at 5 m/s, so you gain on it by 5 m every second. (b) The oncoming car has $v = -20$; relative to you, $-20 - 25 = -45$ m/s. It approaches at 45 m/s, which is 162 km/h: the reason head-on collisions are so much worse than rear-end ones at the same speeds.

Calculus Put your car at the origin at $t = 0$ and the other car a distance D ahead (case a) or ahead on the other side (case b). Then $x_{\text{you}} = 25t$ and $x_{\text{other}} = D + 20t$ or $D - 20t$. The separation $x_{\text{other}} - x_{\text{you}}$ is $D - 5t$ or $D - 45t$, and its derivative is -5 or -45 m/s. The gap closes at those rates and reaches zero (you draw level, or you meet) at $t = D/5$ or $t = D/45$. ■

Check Your Understanding 1.12

Two trains on the same straight track approach each other, one at 30 m/s and the other at 20 m/s, from an initial separation of 5.0 km. Taking the first train's direction as positive, write the velocity of the second train relative to the first, and find how long it takes for the trains to meet if neither brakes.

The word *velocity* gives this chapter its title, and in the chapters ahead velocity, not speed, will be the quantity we work with, because so much of physics depends on direction. For motion along a line, everything you have learned about speed carries over: velocity is the slope of a position–time graph, displacement equals velocity times time when velocity is constant, and dot diagrams work exactly as before, provided you add an arrow. When the motion is one-way along a straight line, physicists often use the words *speed* and *velocity* interchangeably, and we will sometimes do the same; the difference matters the moment direction can change.

It took people nearly two thousand years, from Aristotle to Galileo, to arrive at a clear description of motion. If some of these ideas take you a few hours to absorb, you are doing well.

1.10 Summary

Key ideas

- A **conceptual model** is a simplified description built for a purpose. It is judged by usefulness, not by completeness, and it is scientific only if it makes predictions that could turn out wrong.
- A **dot diagram** marks an object's position at equal time intervals. Within one diagram, wider spacing means higher speed. Comparing diagrams requires the same time interval. Count gaps, not dots.
- **Speed** is distance divided by time. **Elementary** $v = d/t$, equivalently $d = vt$ and $t = d/v$. **Calculus** $v = dx/dt$, the slope of the position–time graph. SI unit: m/s. Multiply m/s by 3.6 to get km/h.
- **Motion is relative.** Every speed is measured relative to something. Unless stated otherwise, speeds are relative to the surface of the Earth.
- **Average speed** is total distance over total time. **Instantaneous speed** is the speed at one moment. **Elementary** The average over a very short interval; the slope of a tangent drawn with a ruler. **Calculus** The derivative dx/dt ; the average speed is $\frac{1}{T} \int_0^T v \, dt$. Average speed is *not* the average of the speeds on different legs unless the legs take equal times.
- **Distance from speed.** **Elementary** Average speed times time; the area under the speed–time graph. **Calculus** $d = \int v \, dt$.
- On a **position–time graph**, the slope of a chord is the average speed (or velocity) between its endpoints, and the slope of the tangent is the instantaneous speed (or velocity).
- A model applied outside the range of its data is being **extrapolated**. **Elementary** Nothing in the data says the relationship stays straight out there. **Calculus** A fitted line is a tangent approximation, and a derivative is local: the error grows like $(x - x_0)^2$. Extrapolated results need independent confirmation.
- A **sanity check** compares a result and its intermediate values against known quantities. Results that fail a sanity check should not be trusted, however impressive they look.
- Report results with **uncertainty**: $x \pm \delta x$. Percentage uncertainty is $\delta x/x \times 100\%$. To propagate it through a calculation: **Elementary** recompute at the ends of the range; **Calculus** $\delta y \approx |dy/dx| \delta x$, and multiplying or dividing by a fixed number preserves the percentage uncertainty.
- **Velocity** is displacement divided by time and carries direction as a sign. Speed is the magnitude of velocity. **Calculus** The velocity is the derivative dx/dt ; displacement is the integral of v_x and distance is the integral of $|v_x|$. Velocity is a **vector** quantity; speed is a **scalar**.
- **Constant velocity** means constant speed *and* constant direction: straight-line motion at a steady speed.

- **Relative velocity** along a line: $v_{A \text{ rel } B} = v_A - v_B$, signs included. Calculus It is the derivative of $x_A - x_B$.
- A number in unfamiliar units cannot be judged. **Convert, then judge:** 12 000 km/d means nothing, while the same speed as 310 mi/h is instantly recognisable as impossible. Most surviving errors die at this step.
- A video is a dot diagram: the **frame rate** supplies equal time intervals, and an object of known size in the shot supplies a **scale** converting pixels to metres. The scale must lie at the same distance from the camera as the motion.
- An object's **position** is where it is; its **coordinate** is the number we assign once an origin and a positive direction are chosen. Moving the origin changes every coordinate and nothing else: displacement, velocity, speed, and distance are all unchanged, which is why Δx deserves a symbol of its own.
- The sign of x and the sign of v_x are independent. $v_x > 0$ means the coordinate is *increasing*, whatever the coordinate's own sign; an object at a negative coordinate moving toward the origin has a positive velocity.
- Numbers read off a diagram are in the **diagram's own units** — grid squares, pixels, millimetres on the page. Turning them into metres takes three steps: *read* in diagram units, *convert* with a measured scale, *divide* and then *check*. Skipping the middle step silently assumes one grid square is one metre.
- **Check a rearrangement with units.** Substituting m and s for the quantities catches a wrong rearrangement in seconds: v/t is m/s², not a distance, while vt is m.
- To estimate an instantaneous speed at a dot, use the **centred** interval from the dot before to the dot after, $v_n \approx (x_{n+1} - x_{n-1})/2\tau$. An average over an interval centred on the point is a far better estimate than one taken to the side, and for steadily changing speed it is exact.

Key equations

	Elementary	Calculus
constant speed	$v = \frac{d}{t}, \quad d = vt, \quad t = \frac{d}{v}$	$x = x_0 + vt, \quad v = \frac{dx}{dt}$
instantaneous speed	$v \approx \frac{\Delta x}{\Delta t}$ (short interval)	$v = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}$
average speed	$\bar{v} = \frac{d_{\text{total}}}{t_{\text{total}}}$	$\bar{v} = \frac{1}{T} \int_0^T v \, dt$
distance from speed	$d = \bar{v} t = \text{area under } v\text{-}t \text{ graph}$	$d = \int_{t_1}^{t_2} v \, dt$
two equal legs	$\bar{v} = \frac{2v_1 v_2}{v_1 + v_2}$	(time-weighted average of v)
unit conversion	1 m/s = 3.6 km/h; 1 mi = 1609 m	
relative speed (gait)	speed/height, unit 1/s	
extrapolation	a fitted line used beyond its data	tangent line $f(x_0) + f'(x_0)(x - x_0)$, local only
uncertainty	$\frac{\delta x}{x} \times 100\%$; recompute at $x \pm \delta x$	$\delta y \approx \left \frac{dy}{dx} \right \delta x$; $\frac{\delta y}{y} = \frac{\delta x}{x}$ for $y = cx$ or c/x
velocity along a line	$\bar{v}_x = \frac{\Delta x}{\Delta t}$; speed = $ \bar{v}_x $ if no reversal	$v_x = \frac{dx}{dt}$; $\Delta x = \int v_x \, dt$, $d = \int v_x \, dt$
relative velocity	$v_{A \text{ rel } B} = v_A - v_B$	$\frac{d}{dt}(x_A - x_B)$

Key terms

conceptual model · scientific attitude · stride · step · dot diagram · fencepost error · ticker-tape timer · frame rate · pixel · scale · grid units · scale bar · units check · centred estimate · speed · rate of change · derivative · motion is relative · average speed · instantaneous speed · integral · area under the graph · position–time graph · chord · tangent · slope · harmonic mean · relative speed · sanity check · interpolation · extrapolation · tangent approximation · uncertainty · percentage uncertainty · propagation of uncertainty · coordinate · coordinate system · origin · one-dimensional motion · position · displacement · velocity · vector quantity · scalar quantity · constant velocity · relative velocity

1.11 Exercises

Exercises marked ★ require more thought or more than one step. Wherever an exercise can be solved both ways, solve it by an **Elementary** route and a **Calculus** route and check that they agree; the answers at the end often show both.

Dot diagrams

- 1.1 Explain in your own words why Webb measured the hunter's *stride* length rather than *step* length.
- 1.2 A dot diagram contains nine dots with 0.5 s between neighbouring dots. How much time does it represent?

- 1.3 Two dot diagrams show dots that are the same distance apart. In diagram A the time between dots is 0.2 s; in diagram B it is 0.4 s. Which object is faster, and by what factor?
- 1.4 Sketch a dot diagram for a car that approaches a red light and slows smoothly to a stop. Mark the direction of motion.
- 1.5 A ticker-tape timer taps 50 times per second. On a piece of tape you count 12 dots, and the distance from the first dot to the last is 33 cm. Find the speed of the cart, assuming it was constant.
- 1.6 A dot diagram cannot tell you the direction of motion. Describe two different ways to add that information to the diagram.
- 1.7 ★ A bicycle wheel has a circumference of 2.1 m. A wet patch on the tyre leaves a mark on the pavement once per revolution, and the marks are 2.1 m apart. A student says, “The marks are evenly spaced, so the bicycle was moving at constant speed.” Is the student right? What, if anything, do the marks tell you about the speed?
- 1.8 ★ A leaky car leaves paint drops on the road. The drops are evenly spaced, yet a passenger insists the car was speeding up the whole time. Can both be true? If so, what must have been happening to the time between drips?
- 1.9 A video is shot at 60 frames per second. How much time passes between frame 15 and frame 27? Between consecutive frames? If instead the clip is a slow-motion recording at 240 frames per second, answer both questions again.
- 1.10 In a photograph, a bicycle whose wheelbase is known to be 1.05 m measures 84 pixels. (a) How many metres does one pixel represent? (b) A crack in the road measures 30 pixels across the same photograph; how wide is it? (c) Why would it be wrong to use this conversion factor on a different photograph taken from twice as far away?
- 1.11 ★ A skateboarder is filmed at 30 frames per second. A doorway of height 2.05 m measures 123 pixels, and between frames 8 and 14 the front wheel advances 62 pixels. (a) Find the skateboarder’s average speed over that interval. (b) If the pixel measurement is uncertain by ± 3 pixels, what is the percentage uncertainty, and what speed would you report? (c) Is the answer an average or an instantaneous speed?
- 1.12 A dot diagram is drawn on squared paper with dots 0.25 s apart. The dots sit at grid positions 0, 4, 16, 36, 64. A ruler laid from the 36 square to the 64 square measures 3.5 cm. (a) What is one grid square worth? (b) Find the average speed over the whole diagram. (c) Estimate the instantaneous speed at the dot at 36.
- 1.13 A classmate reads the same diagram as the previous exercise and reports an average speed of 64 m/s. Explain, in two sentences, exactly which step they omitted, and give one check they could have done in five seconds that would have revealed the error.
- 1.14 ★ The same dot diagram is printed at two sizes, the second an enlargement of the first by a factor of 2.5. (a) Do the dots’ *grid* positions change? (b) Does a ruler measurement of the last gap change? (c) If both readers assume that ruler centimetres are real centimetres, will they get the same speed? (d) What must be printed alongside a diagram before *either* answer can be trusted?

- 1.15** Explain why the quantity “grid squares per second” is not a speed, and what has to be supplied to turn it into one.

Speed, distance, and time

- 1.16** Sound travels through air at about 343 m/s. You see a lightning flash and hear the thunder 4.5 s later. How far away was the lightning? (Light arrives essentially instantly.)
- 1.17** Light travels at 3.00×10^8 m/s, and the Sun is 1.50×10^{11} m from Earth. How long does sunlight take to reach us? Give your answer in seconds and in minutes.
- 1.18** Convert (a) 100 km/h to m/s; (b) 8.0 m/s to km/h; (c) 8.0 m/s to mi/h, using 1 mi = 1609 m.
- 1.19** The men’s marathon world record, set in 2023, is 2 h 0 min 35 s for 42.195 km. Find the average speed in m/s and in km/h.
- 1.20** A walker takes steps 0.75 m long at a rate of 110 steps per minute. Find the walker’s speed in m/s. (Careful: these are steps, not strides.)
- 1.21** Dot diagram A has dots three times as far apart as diagram B, but the time between dots for A is twice that for B. Which object is faster, and by what factor?
- 1.22** A garden snail moves at 0.5 cm/s. How long does it take to cross a path 1.2 m wide?
- 1.23** A cheetah sprints 100 m in 4 s. What is its average speed? What would its average speed be if it sprinted 50 m in 2 s?
- 1.24** A courier company bills you for 9600 km of driving over a two-day hire. Convert this to km/h and to mi/h, and say whether the bill can be right. (Use $8 \text{ km} \approx 5 \text{ mi}$.)
- 1.25** Earth’s circumference is about 40 000 km. At a steady 100 km/h, how many days of non-stop driving would it take to go right around? Use your answer to explain in one sentence why the figure in Worked Example 1.9 is impossible.
- 1.26** ★ A newspaper reports that a glacier is advancing at 3 m/d. Convert this to km/h and to mm/s. Which of the three units gives you the best feel for the motion, and why does the answer depend on what you are comparing it with?
- 1.27** Check each of these rearrangements by substituting units, and say which are wrong: (a) $d = v/t$; (b) $t = d/v$; (c) $v = dt$; (d) $t = v/d$; (e) $d = vt$. For each wrong one, say what physical quantity its units *do* describe, if any.
- 1.28** A student writes “distance = speed / time” and then correctly computes $12 \text{ m}/4 \text{ s} = 3 \text{ m/s}$ for a speed. Their arithmetic is right and their answer is right. What went wrong, and why is it still worth fixing?

Average and instantaneous speed; graphs

- 1.29** You drive to a town 60 km away at 60 km/h and return at 30 km/h. Show that your average speed for the round trip is *not* 45 km/h, and find its correct value.
- 1.30** ★ An object’s position is recorded every two seconds:

t (s)	0	2	4	6	8
x (m)	0	5	10	15	22

- (a) Find the average speed between $t = 0$ and $t = 6$. (b) Find the average speed between $t = 6$ and $t = 8$. (c) Was the object moving at constant speed throughout? Sketch the position–time graph and describe its shape.
- 1.31** ★ Usain Bolt’s average speed over 100 m was 10.44 m/s, but his fastest 20 m segment averaged 12.4 m/s. Explain, using the ideas of average and instantaneous speed, why the two numbers differ and why the race average must be the smaller one.
- 1.32** Using the split times in Worked Example 1.10, find Bolt’s average speed over the first 60 m and over the last 40 m. Which is larger, and why is neither equal to 10.44 m/s?
- 1.33** A car’s position along a straight road is $x(t) = (12 \text{ m/s})t + 50 \text{ m}$. **Elementary** Compute $\Delta x/\Delta t$ between $t = 2 \text{ s}$ and $t = 5 \text{ s}$. **Calculus** Differentiate to find v . Where was the car at $t = 0$?
- 1.34** A particle’s position is $x(t) = (2.0 \text{ m/s}^3)t^3$. (a) Find its average speed between $t = 1.0 \text{ s}$ and $t = 2.0 \text{ s}$. (b) **Calculus** Find its instantaneous speed at $t = 1.0 \text{ s}$ and at $t = 2.0 \text{ s}$ by differentiating. (c) Explain why the average lies between the two instantaneous values.
- 1.35** A cyclist’s speed–time graph is a horizontal line at 8.0 m/s from $t = 0$ to $t = 30 \text{ s}$, then a horizontal line at 4.0 m/s until $t = 90 \text{ s}$. Find the total distance **Elementary** as the area of two rectangles and **Calculus** as $\int v \, dt$, and then the average speed.
- 1.36** ★ A sprinter’s speed is modelled by $v(t) = v_{\max}(1 - e^{-t/\tau})$ with $v_{\max} = 11.0 \text{ m/s}$ and $\tau = 1.4 \text{ s}$. (a) **Calculus** Show by integration that $x(t) = v_{\max}[t - \tau(1 - e^{-t/\tau})]$. (b) Find the time to reach 100 m by trial (a calculator is fine). (c) Find the average speed over the race and the instantaneous speed at the finish.
- 1.37** ★ Show that for two legs of equal length at speeds v_1 and v_2 the average speed $2v_1v_2/(v_1+v_2)$ is never larger than $(v_1+v_2)/2$. (Hint: consider $(v_1 - v_2)^2 \geq 0$.) When are they equal?
- 1.38** A dot diagram has dots at 0, 5, 14, 27, 44, 65 grid squares, $\tau = 0.1 \text{ s}$. Estimate the instantaneous speed, in grid squares per second, at the dot at 27: (a) using the gap before it; (b) using the gap after it; (c) using the centred interval from 14 to 44. Which is the best estimate, and why do (a) and (b) bracket it?
- 1.39** ★ Explain why, for a dot diagram of an object with steadily increasing speed, the centred estimate of Key Idea “Estimating an instantaneous speed from a dot diagram” is exact rather than approximate. (Hint: write the positions as $x_n = \frac{1}{2}a(n\tau)^2$ and compute $(x_{n+1} - x_{n-1})/2\tau$.)

Models and sanity checks

- 1.40** A child 1.2 m tall and an adult 1.8 m tall walk side by side at 1.0 m/s. Find the relative speed of each. Who has the greater relative speed, and what does that mean physically?
- 1.41** ★ The dashed line in Figure 1.8 is approximately

$$\text{stride length (foot-lengths)} \approx 2.2 + 3.5 \times \text{relative speed.}$$

- A walker 1.70 m tall leaves prints with strides 4.3 foot-lengths long. (a) Estimate the walker's relative speed. (b) Estimate the walker's speed in m/s. (c) Is this use of the model an interpolation or an extrapolation? Explain.
- 1.42** The fastest relative speed among the Charteris volunteers was about 1.2/s. Bolt's relative speed was 5.35/s. By what factor was Webb extrapolating beyond the data when he applied the model to a sprinter?
- 1.43** List three distinct sources of uncertainty in Webb's original estimate of the hunter's speed. For each, say whether it is a problem with the *measurement* or with the *model*.
- 1.44** Webb's model output a height of 194 cm and a stride time of 0.360 s for the hunter. Explain, using Bolt's known values, why these two numbers together fail a sanity check.
- 1.45** A gait model predicts that a recreational jogger's time between strides is 0.05 s. Without any further calculation, explain why this result should not be believed.
- 1.46** ★ The temperature in a city rises from 10°C at 8 a.m. to 16°C at 11 a.m. (a) Fit a straight line and use it to "predict" the temperature at 8 p.m. (b) Explain, in the language of Worked Example 1.14, why the prediction is worthless. (c) **Calculus** What does the sign of the second derivative of temperature with respect to time have to be at some point in the afternoon, and why?

Uncertainty

- 1.47** Find the percentage uncertainty in (a) the Ruiz–Torices speed estimate, (7.2 ± 1.0) m/s; (b) Bolt's race time of 9.58 s if the timing system is accurate to ± 0.01 s.
- 1.48** Using the Ruiz–Torices speed and Webb's stride length of 3.73 m, find the hunter's time between strides and the range of values consistent with the uncertainty in the speed.
- 1.49** ★ Suppose Webb's stride-length measurement is uncertain by ± 0.05 m. Holding the speed fixed at 7.2 m/s, find the resulting uncertainty in the stride time **Elementary** by recomputing at both ends and **Calculus** from $\delta t \approx t(\delta d/d)$. Compare it to the uncertainty you found in the previous exercise. Which source of uncertainty dominates?
- 1.50** Express the hunter's revised speed of 7.2 m/s in km/h and mi/h. How long would the hunter take to run 100 m at this speed, and how does that compare to a good high-school sprinter, who might run 100 m in about 11.5 s?
- 1.51** ★ A student times a 100 m run at (14.0 ± 0.3) s. **Elementary** Find the speed and its uncertainty by recomputing at 13.7 s and 14.3 s. **Calculus** Find it from $\delta v \approx |dv/dt| \delta t$ with $v = d/t$. Compare the two answers and explain why they nearly agree.

Velocity and relative velocity

- 1.52** A runner goes 40 m east in 5.0 s, then immediately 40 m west in 5.0 s. Taking east as positive, find the average speed and the average velocity for the whole 10 s.
- 1.53** An object moves along a line from $x = +12$ m to $x = -8$ m in 4.0 s. Find its displacement, its average velocity, and its average speed (assume it did not reverse direction).

- 1.54** ★ Can the average speed of an object over some interval ever be *less* than the magnitude of its average velocity over the same interval? Explain using the definitions of distance and displacement, **Calculus** or using $|\int v_x dt| \leq \int |v_x| dt$.
- 1.55** One airplane flies due north at 300 km/h while another flies due south at 300 km/h. Are their speeds the same? Are their velocities the same? Explain.
- 1.56** You are driving north on a highway at 90 km/h. Without changing your speedometer reading, you round a curve and end up driving east. (a) Did your speed change? (b) Did your velocity change? Explain.
- 1.57** “She moves at a constant speed in a constant direction.” Rephrase this sentence in fewer words, and explain why the shorter version says the same thing.
- 1.58** A particle’s velocity along a line is $v_x = +3.0$ m/s for 4.0 s, then -1.0 m/s for 6.0 s. **Calculus** Find its displacement and the distance it travelled as integrals. **Elementary** Confirm with signed and unsigned rectangle areas.
- 1.59** As you sit in a chair reading this, how fast are you moving relative to the chair? Relative to the Sun? (Use the figure given in the text.)
- 1.60** A car travelling at 100 km/h bumps into the rear of another car travelling in the same direction at 98 km/h. What is the impact speed, that is, the speed of one car relative to the other?
- 1.61** You are driving at the speed limit and see a car coming toward you at the same speed. How fast is the other car approaching you, compared with the speed limit? What would you see if instead it were driving ahead of you in the same direction?
- 1.62** ★ A canoeist can paddle at 8 km/h relative to still water. She sets off upstream on a river that flows at 8 km/h. Describe her motion relative to the riverbank, and relative to the water. What would a dot diagram made by an observer on the bank look like?
- 1.63** ★ Two cyclists start 600 m apart on a straight road and ride toward each other at 7.0 m/s and 5.0 m/s. **Elementary** Use the relative velocity to find when they meet. **Calculus** Write $x_1(t)$ and $x_2(t)$, form $x_2 - x_1$, and find when it is zero. How far has each ridden?
- 1.64** A straight track is marked in metres from its western end. A skater is at the 18 m mark. A second observer sets up a coordinate x' measured from the 25 m mark, positive eastward. (a) What is the skater’s x' coordinate? (b) Write the general relation between x and x' . (c) The skater then moves to the 23 m mark. Find Δx and $\Delta x'$ and comment.
- 1.65** For each case, state whether the coordinate is positive or negative and whether the velocity is positive or negative, taking rightward as positive: (a) an object to the left of the origin moving right; (b) to the left of the origin moving left; (c) to the right of the origin moving left; (d) at the origin moving left.
- 1.66** A student writes: “The ball is at $x = -4$ m, so its velocity is negative.” Identify the error and explain, in one or two sentences, what actually determines the sign of the velocity.
- 1.67** ★ An object’s coordinate is $x(t) = (2 \text{ m/s})t - 6 \text{ m}$. (a) Where is it at $t = 0$, $t = 2$ s, and $t = 5$ s? (b) For what times is its coordinate negative? (c) For what times is its velocity

negative? (d) Now shift the origin 6 m in the negative direction, so that $x' = x + 6$ m. Redo (a), (b), and (c), and say which answers changed and why.

Hands-on

- 1.68** Mark a start line and walk at your normal pace for exactly ten strides, counting one stride each time the same foot lands. Measure the distance covered and the time taken. (a) Find your stride length and the time between strides. (b) Find your walking speed in m/s and in km/h, and compare it with Table 1.1. (c) Compute your relative speed (speed divided by height) and compare it with the range of the Charteris data in Figure 1.8.
- 1.69** Make a real dot diagram. Fill a plastic bottle with water, punch a small hole in the cap so that it drips steadily, and carry it at a constant walking speed along a dry pavement, then again while speeding up from a standstill. Sketch the two trails, and explain in a sentence what each one shows and what it cannot show.
- 1.70** Film a friend sprinting 40 m past cones placed every 10 m, and read the time at each cone from the video. Make a table like the one in Worked Example 1.10, plot position against time, and estimate the top speed from the steepest part of the graph. Report it with an uncertainty based on how precisely you could read the times.
- 1.71** Without using any equations, write a short message to a relative explaining the difference between speed and velocity, and why a car on a curve at a steady 50 km/h does *not* have a constant velocity.
- 1.72** Measure a speed from video. Film someone walking, running, or cycling past, side-on, with the camera held still and an object of known length in the shot and in the same plane as the motion. Step through the frames, use the known object as a scale to convert pixels to metres, and compute the speed. Repeat with a second pair of frames. How much do your two answers differ, and is that difference consistent with the uncertainty you would have quoted?

Answers to Check Your Understanding

- 1.1** Farther apart. With more time between taps, the cart moves farther between dots.
- 1.2** The same rate for both. If the rates differed, dot spacing would no longer measure speed alone; only diagrams with equal time intervals can be compared directly.
- 1.3** (a) Good scales: a metre stick or a sheet of paper laid in the plane your friend runs through, a doorway or paving slab of measured size, or your friend's own height once you have measured it. A poor scale is anything whose true size you do not know (a tree, a cloud) or anything that is not rigid. (b) The number of pixels an object covers depends on how far it is from the camera, so a scale must sit at the same distance as the motion. The nearby car covers *more* pixels for its true length than it would at your friend's distance, which makes the conversion factor (metres per pixel) come out too small. Every distance you then compute, including your friend's displacement, is too small, so the speed comes out *too low*. (c) From frame 40 to frame 46 is 6 frame intervals, so $6/60 = 0.10$ s.
- 1.4** (a) Five dots means *four* intervals, so $t = 4 \times 0.2 \text{ s} = 0.8 \text{ s}$. (b) The gap from 27 to 48 spans 21 squares and measures 4.2 cm, so one square is $4.2/21 = 0.2 \text{ cm}$. (c) The total

- displacement is 48 squares = $48 \times 0.2 = 9.6$ cm, so $\bar{v} = 9.6 \text{ cm}/0.8 \text{ s} = 12 \text{ cm/s} = 0.12 \text{ m/s}$.
- (d) Two mistakes: the student has read grid squares as metres, skipping the conversion in (b); and has counted 5 intervals instead of 4, a fencepost error. The answer 48 m/s is 400 times the correct 0.12 m/s — and it describes a diagram a few centimetres long as moving faster than a motorway car, which is the check that should have stopped it.
- 1.5** Three times faster. In the second case, A covers three times the distance in twice the time, so A is $3/2 = 1.5$ times faster than B.
- 1.6** Into the breeze, toward you, at 3 m/s relative to the air. Its velocity relative to the air then exactly cancels the breeze, so relative to you (and the ground) it is at rest, hovering. This is why a breeze is such an effective deterrent to mosquito bites.
- 1.7** (a) $40 \text{ km}/0.5 \text{ h} = 80 \text{ km/h}$. (b) No. Starting from rest, the car's instantaneous speed was below 80 km/h for some of the time, so it must have exceeded 80 km/h at other times to average 80. (c) Instantaneous speed: a radar gun or speed camera measures how fast the car was going at that moment.
- 1.8** (a) $x(4) = 24 \text{ m}$, so the average speed is $24/4 = 6.0 \text{ m/s}$. (b) **Elementary** $x(4.1) - x(3.9) = 1.5(16.81 - 15.21) = 2.40 \text{ m}$ in 0.2 s: 12.0 m/s. **Calculus** $v = dx/dt = 3.0t = 12.0 \text{ m/s}$ at $t = 4$. (c) The cart is speeding up, so its speed at the end of the interval is greater than its average over the interval; in fact for this motion the final speed is exactly twice the average.
- 1.9** (a) $x < 0$, $v_x > 0$: she is left of the flag but her coordinate is increasing toward zero. (b) $x < 0$, $v_x < 0$. (c) $x = 0$, $v_x > 0$: the coordinate is zero, which is neither positive nor negative, while the velocity is positive. (d) $x > 0$, $v_x = 0$: momentarily at rest. (e) $x > 0$, but the velocity cannot be determined — a *speed* of 2 m/s could be $v_x = +2 \text{ m/s}$ or $v_x = -2 \text{ m/s}$, since speed carries no direction. Moving the flag 10 m left adds 10 m to every coordinate, so the signs in (a) and (b) flip to $x = +5 \text{ m}$ and (c) becomes $x = +10 \text{ m}$; *none* of the velocity answers change, because shifting an origin cannot reverse anybody's direction of travel.
- 1.10** (a) $v = |v_x| = 4 \text{ m/s}$. (b) Toward it: the coordinate is negative and the velocity positive, so x is climbing from -7 m toward zero. (c) Distance from the origin is $|x| = 7 \text{ m}$; the velocity is v_x ; the change in coordinate is Δx . (d) Speed is never negative, so " $v = -4 \text{ m/s}$ " cannot be right as written. They almost certainly meant $v_x = -4 \text{ m/s}$ — and in this case even that is wrong, since the object is moving in the $+x$ direction. Dropping a subscript is the commonest way to turn a right idea into a wrong statement.
- 1.11** A constant speed, yes. A constant velocity, not necessarily: the car may be turning, in which case its direction, and hence its velocity, is changing even though its speed is not.
- 1.12** The second train's velocity is -20 m/s , and relative to the first it is $-20 - 30 = -50 \text{ m/s}$: the gap closes at 50 m/s. They meet after $5000 \text{ m}/(50 \text{ m/s}) = 100 \text{ s}$.

Answers to Selected Exercises

- 1.1** The two feet fall on separate, parallel lines. Measuring from a left print to a right print traces a zigzag; measuring right-to-right follows a straight line along the direction of travel.
- 1.2** 4.0 s. Nine dots have eight gaps: $8 \times 0.5 \text{ s}$.
- 1.3** Object A, twice as fast. It covers the same distance in half the time.

- 1.4** Dots evenly spaced at first, then progressively closer together in the direction of motion, ending in a cluster at the stop line.
- 1.5** 1.5 m/s . Twelve dots span eleven intervals of 0.02 s , so 0.22 s ; $0.33 \text{ m}/0.22 \text{ s} = 1.5 \text{ m/s}$.
- 1.6** Number the dots in time order, or draw an arrow along the direction of motion, or label the first dot.
- 1.7** The student is wrong. The marks are always one circumference apart, whatever the speed, because one mark is made per revolution. Without the time per revolution, the marks say nothing about speed.
- 1.8** Yes. Speed $= d/t$ with d fixed; if speed rises, the time between drips must fall. The can was dripping faster and faster.
- 1.9** At 60 frames per second, frame 15 to frame 27 is 12 intervals, so $12/60 = 0.20 \text{ s}$; consecutive frames are $1/60 = 0.0167 \text{ s}$ apart. At 240 frames per second the same 12 intervals take $12/240 = 0.050 \text{ s}$, and consecutive frames are $1/240 = 0.0042 \text{ s}$ apart. Slow motion buys time resolution, which is why it is the right mode for anything fast.
- 1.10** (a) $1.05 \text{ m}/84 = 0.0125 \text{ m}$, or 12.5 mm , per pixel. (b) $30 \times 0.0125 \text{ m} = 0.38 \text{ m}$. (c) Because the number of pixels a given length covers depends on how far the object is from the camera. At twice the distance the same bicycle would span about half as many pixels, so the conversion factor would be wrong by a factor of two.
- 1.11** (a) The scale is $2.05 \text{ m}/123 = 0.01667 \text{ m}$ per pixel, so the displacement is $62 \times 0.01667 = 1.03 \text{ m}$; six frame intervals at 30 frames per second is 0.20 s , giving $v = 1.03/0.20 = 5.2 \text{ m/s}$. (b) $3/62 \approx 4.8\%$, so round to 5% and report $v = (5.2 \pm 0.3) \text{ m/s}$. (c) An average speed over those 0.20 s , not an instantaneous one.
- 1.12** (a) The last gap spans $64 - 36 = 28$ squares and measures 3.5 cm , so one square is $3.5/28 = 0.125 \text{ cm}$. (b) Five dots give four intervals, so $t = 4 \times 0.25 = 1.0 \text{ s}$; the total distance is $64 \times 0.125 = 8.0 \text{ cm}$, and $\bar{v} = 8.0 \text{ cm/s} = 0.080 \text{ m/s}$. (c) Centred on the dot at 36, use the dots at 16 and 64: $48 \times 0.125 = 6.0 \text{ cm}$ in $2 \times 0.25 = 0.5 \text{ s}$, giving 12 cm/s , or 0.12 m/s .
- 1.13** They omitted the conversion: they divided 64 *grid squares* by the time and wrote the answer in metres per second, which assumes one square is one metre. The five-second check is to convert and compare — 64 m/s is about 230 km/h , and the whole diagram is 8 cm long and records one second of motion.
- 1.14** (a) No. The grid is part of the picture and enlarges with it, so the dots still sit at 0, 4, 16, 36, 64. (b) Yes — it is 2.5 times longer on the page. (c) No. The second reader gets an answer 2.5 times too large, because “ruler centimetres are real centimetres” was true of one printing and not the other. (d) A stated scale: either a ratio such as $1 : 1$, or, better, a scale bar labelled with a real length. That is the only thing that ties a distance on the page to a distance in the world, and it is why a diagram without one cannot be measured at all.
- 1.15** Because a grid square is not a length. It is a subdivision of a particular drawing, and its physical size depends on how that drawing was made and printed. To turn grid squares per second into a speed you need one length in the diagram whose real size you know — a measured gap, a scale bar, or a stated ratio — which converts squares into metres.

- 1.16** About 1.5 km ($343 \times 4.5 \approx 1540$ m).
- 1.17** 500 s, about 8.3 minutes.
- 1.18** (a) 27.8 m/s; (b) 28.8 km/h; (c) 17.9 mi/h.
- 1.19** 5.83 m/s; 21.0 km/h. (Total time 7235 s.)
- 1.20** 1.4 m/s. $0.75 \times 110/60 = 1.375$.
- 1.21** Object A, 1.5 times faster ($3/2$).
- 1.22** 240 s, or 4 minutes.
- 1.23** 25 m/s in both cases: $100/4 = 50/2$.
- 1.24** $9600 \text{ km}/2 \text{ d} = 4800 \text{ km}/\text{d} = 200 \text{ km}/\text{h} \approx 125 \text{ mi}/\text{h}$. The bill cannot be right: it claims racing speed sustained for 48 hours without a single stop.
- 1.25** $40\,000 \text{ km}/(100 \text{ km}/\text{h}) = 400 \text{ h}$, which is about 17 d of non-stop driving. The rental invoice claims more than one circumnavigation in four days, so it would need a speed several times a highway speed, maintained day and night.
- 1.26** $3 \text{ m}/\text{d} = 0.125 \text{ m}/\text{h} = 1.25 \times 10^{-4} \text{ km}/\text{h}$, and $3000 \text{ mm}/86\,400 \text{ s} = 0.035 \text{ mm}/\text{s}$. Metres per day is the most useful here, because the number comes out near 1: it tells you at once that the glacier moves a body-length in a day. The other two units bury the information in zeros. The “best” unit is generally the one that makes the number comparable to something you can picture.
- 1.27** (a) $(\text{m}/\text{s})/\text{s} = \text{m}/\text{s}^2$: wrong; those are the units of an acceleration (Chapter 2). (b) $\text{m}/(\text{m}/\text{s}) = \text{s}$: correct. (c) $(\text{m}/\text{s}) \cdot \text{s}$ written as $d t$ would be $\text{m} \cdot \text{s}$: wrong, and not the units of any quantity we shall meet. (d) $(\text{m}/\text{s})/\text{m} = 1/\text{s}$: wrong; that is a rate or frequency, “so many per second.” (e) $(\text{m}/\text{s}) \cdot \text{s} = \text{m}$: correct.
- 1.28** The written formula contradicts the calculation: the student computed d/t , which is right, while writing “ d/t ” under the name *distance*. It matters for three reasons. A reader — or a grader — can only judge what you wrote. The next time that formula is used, on a problem where the right answer is not obvious, the words will be followed and the answer will be wrong. And a solution is an argument, not just a number: a false statement inside it is a fault even when the final figure survives.
- 1.29** 40 km/h. The trip out takes 1 h and the return 2 h; $120 \text{ km}/3 \text{ h}$. The harmonic-mean formula gives $2(60)(30)/90 = 40$ as well.
- 1.30** (a) 2.5 m/s. (b) 3.5 m/s. (c) No. The graph is a straight line from $t = 0$ to $t = 6$ and then bends upward, becoming steeper: the object sped up in the last interval.
- 1.31** Bolt starts from rest and needs several seconds to reach top speed, so for the early part of the race his instantaneous speed is well below 12.4; he also slows slightly at the end. The average over the whole race blends the slow start with the fast middle, so it must be less than the peak.
- 1.32** First 60 m: $60/6.31 = 9.5 \text{ m}/\text{s}$. Last 40 m: $40/(9.58 - 6.31) = 40/3.27 = 12.2 \text{ m}/\text{s}$. The second is larger because it contains no start. The race average, 10.44, is a time-weighted blend of the two: $(9.5 \times 6.31 + 12.2 \times 3.27)/9.58$.

1.33 **Elementary** $x(5) - x(2) = 110 - 74 = 36$ m in 3 s: 12 m/s. **Calculus** $v = dx/dt = 12$ m/s, constant. At $t = 0$ the car was at $x = 50$ m.

1.34 (a) $x(2) - x(1) = 16 - 2 = 14$ m in 1 s: 14 m/s. (b) $v = 6.0t^2$: 6.0 m/s at $t = 1$ and 24 m/s at $t = 2$. (c) The speed rises steadily through the interval, so its average must lie between its smallest and largest values.

1.35 $8.0(30) + 4.0(60) = 240 + 240 = 480$ m both ways; average speed $480/90 = 5.3$ m/s.

1.36 (a) $\int_0^t v_{\max}(1 - e^{-t'/\tau}) dt' = v_{\max} [t' + \tau e^{-t'/\tau}]_0^t = v_{\max}[t - \tau(1 - e^{-t/\tau})]$. (b) About 10.5 s: $x(10.5) = 11.0[10.5 - 1.4(1 - e^{-7.5})] = 11.0 \times 9.10 = 100.1$ m. (c) Average 9.5 m/s; at the finish $v = 11.0(1 - e^{-7.5}) \approx 11.0$ m/s.

1.37 $\frac{v_1 + v_2}{2} - \frac{2v_1v_2}{v_1 + v_2} = \frac{(v_1 + v_2)^2 - 4v_1v_2}{2(v_1 + v_2)} = \frac{(v_1 - v_2)^2}{2(v_1 + v_2)} \geq 0$, with equality only when $v_1 = v_2$.

1.38 (a) $(27 - 14)/0.1 = 130$ squares per second. (b) $(44 - 27)/0.1 = 170$ squares per second. (c) $(44 - 14)/0.2 = 150$ squares per second. The centred value is the best estimate. Since the object is speeding up steadily, the interval before the dot lies wholly in slower motion and the one after wholly in faster motion, so (a) and (b) fall on opposite sides of the truth; the centred estimate is their average, and the two errors cancel.

1.39 With $x_n = \frac{1}{2}a(n\tau)^2$,

$$\frac{x_{n+1} - x_{n-1}}{2\tau} = \frac{\frac{1}{2}a\tau^2[(n+1)^2 - (n-1)^2]}{2\tau} = \frac{\frac{1}{2}a\tau^2(4n)}{2\tau} = an\tau,$$

and $an\tau$ is exactly $v(t_n) = at_n$ at the instant of dot n . The two errors do not merely largely cancel here; they cancel completely, because the position is a quadratic in t and the centred difference of a quadratic is its exact derivative at the midpoint. This is the same special property of parabolas noted in Worked Example 2.3.

1.40 Child: 0.83/s; adult: 0.56/s. The child has the greater relative speed: at the same speed the child covers more of their own body length each second, which is why the child appears to hurry.

1.41 (a) $(4.3 - 2.2)/3.5 = 0.60$ /s. (b) $0.60 \times 1.70 \approx 1.0$ m/s. (c) Interpolation: 0.60 lies within the data range of roughly 0.4 to 1.2, and 1.0 m/s is an ordinary walking speed.

1.42 About 4.5 times $(5.35/1.2)$.

1.43 Examples: stride length measured imperfectly and strides vary (measurement); footprints may have eroded or deformed over 20 000 years (measurement); foot length only loosely predicts height (model); the height-and-stride-to-stride-time equations have their own scatter (model); the model was built for walking, not running (model, and the largest problem).

1.44 The model makes the hunter the same height as Bolt but with a shorter stride and a much shorter stride time. A person of the same height taking strides a metre shorter than Bolt's and cycling their legs 0.11 s faster than Bolt is not credible; the output contradicts what we know about the fastest runner ever measured.

1.45 0.05 s per stride means 20 strides per second. Bolt, at full sprint, manages about two strides per second (0.467 s each). No jogger turns their legs over ten times faster than Bolt.

- 1.46** (a) 2°C per hour, so 34°C at 8 p.m. (b) The line was fitted to three morning hours and says nothing about the afternoon; temperature peaks and then falls, so the relationship bends. (c) Negative: the temperature rises in the morning and falls in the evening, so its rate of change must decrease somewhere in between, and a decreasing first derivative means a negative second derivative.
- 1.47** (a) About 14%. (b) About 0.1%.
- 1.48** 0.52 s, with a range of about 0.45 s to 0.60 s; roughly (0.52 ± 0.08) s.
- 1.49** **Elementary** $3.68/7.2 = 0.511$ s and $3.78/7.2 = 0.525$ s, so about ± 0.007 s. **Calculus** $\delta t \approx 0.518 \times (0.05/3.73) = 0.007$ s. This is roughly ten times smaller than the ± 0.08 s from the speed uncertainty, so the speed uncertainty dominates.
- 1.50** 26 km/h; 16 mi/h. At 7.2 m/s, 100 m takes about 14 s, slower than a good high-school sprinter's 11.5 s (8.7 m/s), though the hunter was barefoot, on mud, and not running a race.
- 1.51** **Elementary** $100/14.0 = 7.14$ m/s; $100/13.7 = 7.30$ and $100/14.3 = 6.99$, so (7.14 ± 0.15) m/s. **Calculus** $|dv/dt| = d/t^2 = 100/196 = 0.51$, so $\delta v \approx 0.51 \times 0.3 = 0.15$ m/s; equivalently $\delta v/v = \delta t/t = 2.1\%$. They agree because a 2% change is small enough for the tangent approximation to be accurate.
- 1.52** Average speed 8.0 m/s; average velocity 0.
- 1.53** Displacement -20 m; average velocity -5.0 m/s; average speed 5.0 m/s.
- 1.54** No. Distance travelled is always at least as large as the magnitude of displacement, since any back-tracking adds to distance but subtracts from displacement. Dividing both by the same time interval preserves the inequality, so average speed $\geq |\bar{v}_x|$, with equality only when the object never reverses direction. In integral form, $|\int v_x dt| \leq \int |v_x| dt$ because cancellation can only reduce a signed sum.
- 1.55** Same speed, 300 km/h. Different velocities, because velocity includes direction and the directions are opposite.
- 1.56** (a) No; the speedometer reading, which is the speed, stayed at 90 km/h. (b) Yes; the direction changed from north to east, so the velocity changed. (In the next chapter you will see that this means the car accelerated.)
- 1.57** "She moves at a constant velocity." Constant velocity means constant speed and constant direction.
- 1.58** Displacement $3.0(4.0) + (-1.0)(6.0) = 6.0$ m; distance $12 + 6 = 18$ m. The first rectangle (area 12) is above the axis, the second (area 6) below; signed sum 6, unsigned sum 18.
- 1.59** Zero relative to the chair. About 107 000 km/h, or 30 km/s, relative to the Sun.
- 1.60** 2 km/h. Relative to the slower car, the faster one closes in at the difference of their velocities, $100 - 98$.
- 1.61** Twice the speed limit: with velocities $+v$ and $-v$, the relative velocity is $-2v$. A car ahead of you at the same speed would appear to stand still: relative velocity zero.

- 1.62** Relative to the bank she does not move at all: her 8 km/h upstream relative to the water is cancelled by the water's 8 km/h downstream. Relative to the water she is moving at 8 km/h. An observer on the bank would see all the dots of her dot diagram piled on top of one another at a single point.
- 1.63** Closing speed $7.0 + 5.0 = 12.0$ m/s, so they meet after $600/12 = 50$ s. With $x_1 = 7.0t$ and $x_2 = 600 - 5.0t$, $x_2 - x_1 = 600 - 12t = 0$ at $t = 50$ s. They ride 350 m and 250 m.
- 1.64** (a) $x' = 18 - 25 = -7$ m. (b) $x' = x - 25$ m. (c) $\Delta x = 23 - 18 = +5$ m and $\Delta x' = (-2) - (-7) = +5$ m: identical, because the -25 m cancels in the subtraction. The coordinates disagree; the displacement does not.
- 1.65** (a) $x < 0$, $v_x > 0$. (b) $x < 0$, $v_x < 0$. (c) $x > 0$, $v_x < 0$. (d) $x = 0$ (neither positive nor negative), $v_x < 0$.
- 1.66** The student has read the sign of the *coordinate* as though it were the sign of the *velocity*. Where the ball is says nothing about which way it is going. The velocity is negative only if the coordinate is *decreasing*; a ball at $x = -4$ m heading toward the origin has a positive velocity.
- 1.67** (a) -6 m, -2 m, $+4$ m. (b) $x < 0$ for $t < 3$ s. (c) Never: $v_x = dx/dt = +2$ m/s at every instant. (d) With $x' = x + 6$ m $= (2 \text{ m/s})t$, the positions become 0, 4, and 10 m; x' is never negative for $t \geq 0$; and the velocity is still $+2$ m/s. The answers to (a) and (b) changed because they are about coordinates; the answer to (c) did not, because velocity is a difference and the shift cancels.
- 1.68** Answers depend on your own measurements. Typical values: stride about 1.4 to 1.6 m, time between strides about 1.0 to 1.2 s, speed about 1.2 to 1.5 m/s (4.5 to 5.5 km/h), relative speed roughly 0.7 to 0.91/s, comfortably inside the Charteris range of about 0.4 to 1.2.
- 1.69** The constant-speed trail has evenly spaced drops; the speeding-up trail has drops that spread farther apart along the direction of motion. Neither trail, by itself, tells you which way you were walking or how fast the drips were falling.
- 1.70** Typical school sprinters reach 7 to 9 m/s. Reading times from ordinary video (about 30 frames per second) gives an uncertainty of a few hundredths of a second per cone, which for 10 m segments of about 1.3 s is a few percent in speed.
- 1.72** Answers depend on your own measurements. Two frame pairs from the same walk-past typically differ by a few percent, which should sit comfortably inside a quoted uncertainty of around 5%; if they differ by much more, the usual culprits are a scale object at a different distance from the camera than the motion, a camera that moved, or a subject whose speed genuinely changed between the two pairs.

Sources and Further Reading

- P. G. Hewitt, *Conceptual Physics*, 12th ed. (Pearson, 2014). Chapter 1 (“About Science”) on scientific methods, the scientific attitude, and testable hypotheses; Chapter 3 (“Linear Motion”), Sections 3.1–3.3, on relative motion, speed, instantaneous and average speed, velocity, and vector and scalar quantities; Appendix A on the definitions of the metre and the second. The reference speeds in Table 1.1 and several of the Check Your Understanding

questions and exercises are adapted from this book. Hewitt’s treatment corresponds to the **Elementary** route throughout; the **Calculus** route is our addition.

- S. Webb, M. L. Cupper, and R. Robins, “Pleistocene human footprints from the Willandra Lakes, southeastern Australia,” *Journal of Human Evolution* 50 (2006), 405–413.
- S. Webb, “Further research of the Willandra Lakes fossil footprint site, southeastern Australia,” *Journal of Human Evolution* 52 (2007), 711–715.
- J. Charteris, J. C. Wall, and J. W. Nottrodt, “Functional reconstruction of gait from the Pliocene hominid footprints at Laetoli, northern Tanzania,” *Nature* 290 (1981), 496–498.
- J. Ruiz and A. Torices, “Humans running at stadiums and beaches and the accuracy of speed estimations from fossil trackways,” *Ichnos* 20 (2013), 31–35.
- International Association of Athletics Federations and Deutscher Leichtathletik-Verband, *Biomechanics Report, World Championships Berlin 2009: Sprint Men* (2009). The source of the 20 m split times and the reaction time used in Worked Example 1.10 and Figure 1.7. The stride count and stride length quoted for Bolt are from the same analysis.
- “Prehistoric Aussie aboriginals could have outrun Usain Bolt,” *Metro* (UK), 14 October 2009. An example of how the original claim was reported.
- Scholars High School Physics 1: Mechanics (AoPS course 4926), Lesson 1 class transcript, 11 September 2026. The rental-car sense-making problem, the video-frame method of measuring a cyclist’s speed with a bicycle wheelbase as a scale, and the treatment of coordinate systems and the coordinate–position distinction follow the arc of that lesson. All numbers, examples, and prose here are our own.
- Émilie du Châtelet, *Principes mathématiques de la philosophie naturelle* (Paris, 1759), her translation of and commentary on Newton’s *Principia*, still the standard French edition. She also argued, against the prevailing view, that the quantity conserved in collisions goes as mv^2 rather than mv — an early step toward the concept of kinetic energy we meet later in this book.
- The Mungo National Park visitor site, <https://www.visitmungo.com.au>, has photographs of the trackways and background on the Willandra Lakes World Heritage Area and its Traditional Owners.