

(a)
29,761
Survey respondents, all ages
Excluded: aged < 18 years (n = 8,317)
21,444
Adults aged ≥ 18 years
Excluded: no GP or family physician visit in the past 12 months (n = 7,768)
13,676
GP or family physician visit, past 12 months
Excluded: psychologist, psychotherapist or psychiatrist visit, past 12 months (n = 833)
12,843
Eligible population
Excluded: missing household income (n = 1)
12,842
Analysis frame (7,725 households)
PHQ-8 ≥ 10 (n = 581)
Bar widths are proportional to unweighted counts.
(b)
Information available before assessment (no depressive-symptom questions)
M0 Routine characteristics
age, sex, education, household income, chronic conditions
M1 + Two diet questions
fruit and vegetable/salad consumption frequency
M2 + Activity and social support
walking, cycling and sport days, physical demand of daily activities, social support (3 items)
Rank adults by predicted probability of PHQ-8 ≥ 10
Invite the top share c (c = assessment capacity)
Detailed assessment
Detection yield: selected PHQ-8–positive adults per 1,000 eligible adults
Comparators at the same capacity
Random selection
Perfect ranking (upper bound)
PHQ-2 ranking: symptom-based comparator sharing two items with the PHQ-8; not an upper bound