

EMERGENCY DEPARTMENT WOMEN AND CHILDREN
HOSPITAL KUALA LUMPUR
Acute Gastroenteritis Order Set

DATE :

TIME SEEN:

PATIENT DETAILS

NAME		ADDRESS	
IC/PASSPORT		GENDER	MALE/FEMALE
AGE		RACE	MALAY/CHINESE/INDIAN/OTHERS _____
MRN		NEXT OF KIN CONTACT NO	

Weight (kg) :

PMHX :

Allergies :

ASSESS

Look at child's general condition	Well, alert		Restless or irritable		Lethagic or unconscious	
Look for sunken eyes	No sunken eyes		Sunken eyes		Sunken eyes	
Offer the child fluid	Drinks normally		Drinks eagerly, thirsty		Not able to drink or drinks poorly	
Pinch skin of abdomen	Skin goes back immediately		Skin goes back slowly		Skin goes back slowly (> 2 secs)	
Classify	Mild Dehydration IMCI :no signs of Dehydration		≥ 2 above signs Moderate Dehydration 5-10% Dehydrated IMCI :Some signs of dehydration		≥ 2 above signs Severe Dehydration > 10% Dehydration	

INVESTIGATIONS

**Mild / Moderate AGE
(If clinically indicated)**

- ☐ FBC
- ☐ RP
- ☐ Dxt

Severe AGE

- ☐ FBC (if clinically indicated)
- ☐ RP
- ☐ VBG
- ☐ Blood C&S (if clinically indicated)
- ☐ Dxt

Others (When Indicated)

- ☐ Urine FEME
- ☐ Urine C&S
- ☐ Stool C&S

ORS

Dissolve 1 sachet completely in 250mls of cooled boiled water.

Ordered by: Provider, Time, Signature	Orders	Noted by SN/MA Time & Initials
MILD DEHYDRATION		
	☐ Give Extra Fluid (as much as the child will take) Breastfeed frequently and longer ☐ Start Oral Rehydration Therapy if observation is required 10mls/kg after each loose stool : _____ mls /purge ☐ Continue Feeding usual food ☐ I/O Charting if observation is required ☐ V/s Monitoring if observation is required ☐ Consider discharge if child is able to take orally well	
MODERATE DEHYDRATION		
	☐ Give Extra Fluid (as much as the child will take) ☐ Start Oral Rehydration Therapy and observation 75mls/kg over 4 hours : _____ mls/ 4 hours ☐ Reassess frequently within 4 hours ☐ I/O Charting ☐ V/s Monitoring	
SEVERE DEHYDRATION		
	☐ Call for help from senior MO / EP ☐ ABC ☐ Insert IV/IO Line ☐ Fluid resuscitation 20mls/kg 0.9% NS or Hartmann's rapid IV bolus* : _____ mls bolus Repeat if necessary ☐ Replace Deficit % Dehydration x BW (g) given over 4 -6 hours % /100 x BW (g) = _____ mls/4-6 H (depending on Sr Na levels, may need to prolong the duration of correction) ☐ Fluid Maintenance Holliday & Segar : 1 st 10kg : 100mls 2 nd 10kg : 50mls Ever kg thereafter : 20mls ☐ Continuous V/s Monitoring ☐ I/O Charting ☐ Admit	

OUTCOME

** Consider reduction in bolus volume and rate in orang Asli, malnutrition, medical conditions requiring fluid restriction*

☐ Discharge

☐ Admit

Discharge advice given

- ✓ Return if child is not able to drink or breastfeed properly
- ✓ Becomes sicker
- ✓ Develops fever
- ✓ Blood in stool
- ✓ Lethargic

ATTENDING DOCTOR
SIGNATURE AND STAMP

ATTENDING SN/MA
SIGNATURE AND STAMP