ADR Reporting: Task 2 Report

This is my ADR reporting work for the 18 fictional intake cases in the pack. For each case I filled one standard ADR report form (pages 3 onwards) with the patient details, the reaction, the suspect drug, concomitant medicines, relevant medical history and the reporter. The five mandatory fields are marked with an asterisk (*) on every form: patient initials, age, suspect drug, reaction start date and outcome.

The intake file does not give patient initials, any calendar dates or any medical history. As the README says, unknown facts must stay unknown, so I wrote **Unknown** for these fields and did not make up initials or dates. The only timing given is the onset in days after treatment began, and I recorded this next to the reaction start date. Doses and indications are written exactly as reported, and anything that looks wrong is listed in the notes box of that case. All cases are fictional and this work is for training only.

1. Mandatory field check

Case	Patient initials*	Age*	Suspect drug*	Reaction start date*	Outcome*
SYN-PV-007	Unknown (not provided)	63 years	Omeprazole 500 mg twice daily	Unknown (onset 16 days after start)	Recovered
SYN-PV-008	Unknown (not provided)	36 years	Losartan 5 mg once daily	Unknown (onset 2 days after start)	Recovering
SYN-PV-009	Unknown (not provided)	78 years	Amoxicillin 5 mg once daily	Unknown (onset 7 days after start)	Recovering
SYN-PV-010	Unknown (not provided)	57 years	Atorvastatin 5 mg once daily	Unknown (onset 21 days after start)	Recovered
SYN-PV-011	Unknown (not provided)	23 years	Metformin 5 mg once daily	Unknown (onset 19 days after start)	Recovering
SYN-PV-012	Unknown (not provided)	24 years	Amlodipine 5 mg once daily	Unknown (onset 4 days after start)	Not recovered (at follow-up)
SYN-PV-013	Unknown (not provided)	75 years	Ciprofloxacin 500 mg twice daily	Unknown (onset 5 days after start)	Not recovered (at follow-up)
SYN-PV-014	Unknown (not provided)	75 years	Sertraline 5 mg once daily	Unknown (onset 9 days after start)	Recovering
SYN-PV-015	Unknown (not provided)	33 years	Omeprazole 20 mg once daily	Unknown (onset 16 days after start)	Recovered
SYN-PV-016	Unknown (not provided)	66 years	Losartan 20 mg once daily	Unknown (onset 10 days after start)	Not recovered (at follow-up)
SYN-PV-017	Unknown (not provided)	57 years	Amoxicillin 5 mg once daily	Unknown (onset 21 days after start)	Recovering
SYN-PV-018	Unknown (not provided)	56 years	Atorvastatin 500 mg twice daily	Unknown (onset 17 days after start)	Not recovered (at follow-up)
SYN-PV-019	Unknown (not provided)	29 years	Metformin 5 mg once daily	Unknown (onset 9 days after start)	Recovering
SYN-PV-020	Unknown (not provided)	56 years	Amlodipine 5 mg once daily	Unknown (onset 7 days after start)	Recovering
SYN-PV-021	Unknown (not provided)	33 years	Ciprofloxacin 5 mg once daily	Unknown (onset 9 days after start)	Recovered
SYN-PV-022	Unknown (not provided)	43 years	Sertraline 5 mg once daily	Unknown (onset 21 days after start)	Not recovered (at follow-up)
SYN-PV-023	Unknown (not provided)	26 years	Omeprazole 500 mg twice daily	Unknown (onset 4 days after start)	Recovering
SYN-PV-024	Unknown (not provided)	79 years	Losartan 500 mg twice daily	Unknown (onset 3 days after start)	Recovering

Age, suspect drug and outcome are available for all 18 cases. Patient initials and reaction start date are not provided in any case.

2. Reporting template (filled for the whole pack)

Field	Required	Student entry	Source evidence
Identifiable patient	Yes	Present in all 18. Age is given for every case. Sex is given in 10 cases and unknown in 8. Initials are not provided.	Age, Sex columns
Identifiable reporter	Yes	Present as reporter type only: 10 physician, 5 pharmacist, 3 consumer. Name and contact are not provided.	Reporter column
Suspect product	Yes	Present in all 18 (drug, dose and route given).	SuspectProduct, Dose, Route
Adverse event	Yes	Present in all 18 (reaction term and onset day given).	Reaction, OnsetDays
Chronology	No	Onset day only. Treatment start and stop dates are unknown.	OnsetDays, narrative
Seriousness	No	Given for all cases: 7 medically important, 7 hospitalisation, 4 not serious.	Seriousness column
Outcome	No	Stated in the narrative (recovered, recovering or not recovered).	Narrative

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-007** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	63 years
Sex	Male
B. Adverse reaction	
Reaction / event	Headache
Description	A 63-year-old patient took Omeprazole by mouth (500 mg twice daily) for hypertension. Headache was reported 16 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovered.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 16 day(s) after treatment began.
Outcome*	Recovered
Dechallenge / rechallenge	Dechallenge: Negative. Rechallenge: Negative.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Omeprazole
Dose and frequency	500 mg twice daily
Route	Oral
Indication	Hypertension
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Consumer
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (500 mg twice daily, 1000 mg/day) is outside the usual adult range for Omeprazole (20 to 40 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Hypertension' does not match the usual use of Omeprazole. The real indication should be confirmed.
• Dechallenge is negative but the outcome is recovered. When recovery happened compared with stopping the drug is not stated.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-008** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	36 years
Sex	Male
B. Adverse reaction	
Reaction / event	Fatigue
Description	A 36-year-old patient took Losartan by mouth (5 mg once daily) for infection. Fatigue was reported 2 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 2 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Negative. Rechallenge: Not done.
Seriousness (as reported)	Not serious
Severity	Not stated
Expectedness (as given)	Unexpected
C. Suspect drug	
Suspect drug*	Losartan
Dose and frequency	5 mg once daily
Route	Oral
Indication	Infection
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Pharmacist
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Losartan (25 to 100 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Infection' does not match the usual use of Losartan. The real indication should be confirmed.
• Dechallenge is negative but the outcome is recovering. When recovery happened compared with stopping the drug is not stated.
• Expectedness is given as Unexpected. It should be checked against the reference safety information.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-009** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	78 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Rash
Description	A 78-year-old patient took Amoxicillin by mouth (5 mg once daily) for anxiety. Rash was reported 7 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 7 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Positive. Rechallenge: Not done.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Amoxicillin
Dose and frequency	5 mg once daily
Route	Oral
Indication	Anxiety
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Vitamin D. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Amoxicillin (about 750 to 3000 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Anxiety' does not match the usual use of Amoxicillin. The real indication should be confirmed.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-010** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	57 years
Sex	Female
B. Adverse reaction	
Reaction / event	Myalgia
Description	A 57-year-old patient took Atorvastatin by mouth (5 mg once daily) for infection. Myalgia was reported 21 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovered.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 21 day(s) after treatment began.
Outcome*	Recovered
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Negative.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Atorvastatin
Dose and frequency	5 mg once daily
Route	Oral
Indication	Infection
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Pharmacist
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Atorvastatin (10 to 80 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Infection' does not match the usual use of Atorvastatin. The real indication should be confirmed.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-011** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	23 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Nausea
Description	A 23-year-old patient took Metformin by mouth (5 mg once daily) for hypertension. Nausea was reported 19 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 19 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Positive.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Metformin
Dose and frequency	5 mg once daily
Route	Oral
Indication	Hypertension
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Consumer
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Metformin (500 to 3000 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Hypertension' does not match the usual use of Metformin. The real indication should be confirmed.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-012** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	24 years
Sex	Female
B. Adverse reaction	
Reaction / event	Peripheral oedema
Description	A 24-year-old patient took Amlodipine by mouth (5 mg once daily) for hypertension. Peripheral oedema was reported 4 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: not recovered (at follow-up).
Reaction start date*	Unknown (calendar date not provided). Reported onset: 4 day(s) after treatment began.
Outcome*	Not recovered (at follow-up)
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Positive.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Amlodipine
Dose and frequency	5 mg once daily
Route	Oral
Indication	Hypertension
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Pharmacist
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-013** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	75 years
Sex	Female
B. Adverse reaction	
Reaction / event	Tendon pain
Description	A 75-year-old patient took Ciprofloxacin by mouth (500 mg twice daily) for diabetes. Tendon pain was reported 5 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: not recovered (at follow-up).
Reaction start date*	Unknown (calendar date not provided). Reported onset: 5 day(s) after treatment began.
Outcome*	Not recovered (at follow-up)
Dechallenge / rechallenge	Dechallenge: Negative. Rechallenge: Positive.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Ciprofloxacin
Dose and frequency	500 mg twice daily
Route	Oral
Indication	Diabetes
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Vitamin D. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The indication 'Diabetes' does not match the usual use of Ciprofloxacin. The real indication should be confirmed.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-014** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	75 years
Sex	Female
B. Adverse reaction	
Reaction / event	Dizziness
Description	A 75-year-old patient took Sertraline by mouth (5 mg once daily) for diabetes. Dizziness was reported 9 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 9 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Negative. Rechallenge: Negative.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Sertraline
Dose and frequency	5 mg once daily
Route	Oral
Indication	Diabetes
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Sertraline (25 to 200 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Diabetes' does not match the usual use of Sertraline. The real indication should be confirmed.
• Dechallenge is negative but the outcome is recovering. When recovery happened compared with stopping the drug is not stated.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-015** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	33 years
Sex	Male
B. Adverse reaction	
Reaction / event	Headache
Description	A 33-year-old patient took Omeprazole by mouth (20 mg once daily) for hypertension. Headache was reported 16 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovered.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 16 day(s) after treatment began.
Outcome*	Recovered
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Positive.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Omeprazole
Dose and frequency	20 mg once daily
Route	Oral
Indication	Hypertension
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The indication 'Hypertension' does not match the usual use of Omeprazole. The real indication should be confirmed.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-016** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	66 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Fatigue
Description	A 66-year-old patient took Losartan by mouth (20 mg once daily) for diabetes. Fatigue was reported 10 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: not recovered (at follow-up).
Reaction start date*	Unknown (calendar date not provided). Reported onset: 10 day(s) after treatment began.
Outcome*	Not recovered (at follow-up)
Dechallenge / rechallenge	Dechallenge: Positive. Rechallenge: Negative.
Seriousness (as reported)	Not serious
Severity	Not stated
Expectedness (as given)	Unexpected
C. Suspect drug	
Suspect drug*	Losartan
Dose and frequency	20 mg once daily
Route	Oral
Indication	Diabetes
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (20 mg once daily, 20 mg/day) is outside the usual adult range for Losartan (25 to 100 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Diabetes' is vague for Losartan. The exact reason for use should be confirmed.
• Dechallenge is positive but the patient had not recovered at follow-up. This should be clarified.
• Expectedness is given as Unexpected. It should be checked against the reference safety information.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-017** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	57 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Rash
Description	A 57-year-old patient took Amoxicillin by mouth (5 mg once daily) for hypertension. Rash was reported 21 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 21 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Positive.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Amoxicillin
Dose and frequency	5 mg once daily
Route	Oral
Indication	Hypertension
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Vitamin D. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Amoxicillin (about 750 to 3000 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Hypertension' does not match the usual use of Amoxicillin. The real indication should be confirmed.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-018** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	56 years
Sex	Female
B. Adverse reaction	
Reaction / event	Myalgia
Description	A 56-year-old patient took Atorvastatin by mouth (500 mg twice daily) for anxiety. Myalgia was reported 17 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: not recovered (at follow-up).
Reaction start date*	Unknown (calendar date not provided). Reported onset: 17 day(s) after treatment began.
Outcome*	Not recovered (at follow-up)
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Positive.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Atorvastatin
Dose and frequency	500 mg twice daily
Route	Oral
Indication	Anxiety
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (500 mg twice daily, 1000 mg/day) is outside the usual adult range for Atorvastatin (10 to 80 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Anxiety' does not match the usual use of Atorvastatin. The real indication should be confirmed.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-019** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	29 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Nausea
Description	A 29-year-old patient took Metformin by mouth (5 mg once daily) for anxiety. Nausea was reported 9 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 9 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Positive. Rechallenge: Negative.
Seriousness (as reported)	Not serious
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Metformin
Dose and frequency	5 mg once daily
Route	Oral
Indication	Anxiety
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Pharmacist
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Metformin (500 to 3000 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Anxiety' does not match the usual use of Metformin. The real indication should be confirmed.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-020** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	56 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Peripheral oedema
Description	A 56-year-old patient took Amlodipine by mouth (5 mg once daily) for diabetes. Peripheral oedema was reported 7 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 7 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Unknown. Rechallenge: Not done.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Amlodipine
Dose and frequency	5 mg once daily
Route	Oral
Indication	Diabetes
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Pharmacist
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The indication 'Diabetes' does not match the usual use of Amlodipine. The real indication should be confirmed.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-021** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	33 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Tendon pain
Description	A 33-year-old patient took Ciprofloxacin by mouth (5 mg once daily) for infection. Tendon pain was reported 9 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovered.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 9 day(s) after treatment began.
Outcome*	Recovered
Dechallenge / rechallenge	Dechallenge: Positive. Rechallenge: Not done.
Seriousness (as reported)	Serious (other medically important condition)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Ciprofloxacin
Dose and frequency	5 mg once daily
Route	Oral
Indication	Infection
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Ciprofloxacin (500 to 1500 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• Seriousness is reported as 'serious (other medically important condition)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-022** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	43 years
Sex	Female
B. Adverse reaction	
Reaction / event	Dizziness
Description	A 43-year-old patient took Sertraline by mouth (5 mg once daily) for infection. Dizziness was reported 21 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: not recovered (at follow-up).
Reaction start date*	Unknown (calendar date not provided). Reported onset: 21 day(s) after treatment began.
Outcome*	Not recovered (at follow-up)
Dechallenge / rechallenge	Dechallenge: Negative. Rechallenge: Positive.
Seriousness (as reported)	Not serious
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Sertraline
Dose and frequency	5 mg once daily
Route	Oral
Indication	Infection
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Consumer
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (5 mg once daily, 5 mg/day) is outside the usual adult range for Sertraline (25 to 200 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Infection' does not match the usual use of Sertraline. The real indication should be confirmed.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-023** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	26 years
Sex	Male
B. Adverse reaction	
Reaction / event	Headache
Description	A 26-year-old patient took Omeprazole by mouth (500 mg twice daily) for anxiety. Headache was reported 4 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 4 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Negative. Rechallenge: Negative.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Known
C. Suspect drug	
Suspect drug*	Omeprazole
Dose and frequency	500 mg twice daily
Route	Oral
Indication	Anxiety
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	None reported.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (500 mg twice daily, 1000 mg/day) is outside the usual adult range for Omeprazole (20 to 40 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• The indication 'Anxiety' does not match the usual use of Omeprazole. The real indication should be confirmed.
• Dechallenge is negative but the outcome is recovering. When recovery happened compared with stopping the drug is not stated.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.

ADVERSE DRUG REACTION (ADR) REPORT FORM

Case ID: **SYN-PV-024** | Fictional training case, not for regulatory submission | * = mandatory field

A. Patient information	
Patient initials*	Unknown (not provided; case is de-identified)
Age*	79 years
Sex	Unknown
B. Adverse reaction	
Reaction / event	Fatigue
Description	A 79-year-old patient took Losartan by mouth (500 mg twice daily) for hypertension. Fatigue was reported 3 day(s) after treatment began. The reporter said treatment was stopped and the outcome is: recovering.
Reaction start date*	Unknown (calendar date not provided). Reported onset: 3 day(s) after treatment began.
Outcome*	Recovering
Dechallenge / rechallenge	Dechallenge: Positive. Rechallenge: Positive.
Seriousness (as reported)	Serious (hospitalisation)
Severity	Not stated
Expectedness (as given)	Unexpected
C. Suspect drug	
Suspect drug*	Losartan
Dose and frequency	500 mg twice daily
Route	Oral
Indication	Hypertension
Treatment start / stop dates	Unknown (dates not provided)
Action taken	Drug stopped (as reported)
D. Concomitant medications	
Medicines	Paracetamol. Dose, dates and indication are unknown.
E. Relevant medical history	
History	Unknown (not provided in the report)
F. Reporter	
Reporter type	Physician
Name and contact	Unknown (not provided)

G. Notes and follow-up needed
• The reported dose (500 mg twice daily, 1000 mg/day) is outside the usual adult range for Losartan (25 to 100 mg/day). Check for a unit or entry error, or a medication error if the dose is correct.
• Rechallenge is recorded as positive, but the report does not say the drug was restarted. This needs checking with the reporter.
• Seriousness is reported as 'serious (hospitalisation)', but the reason is not stated.
• Expectedness is given as Unexpected. It should be checked against the reference safety information.