

Table 1: **Cross-tabulation of Combined Healthcare Source (4 Groups) and Antibiotic Use**

Healthcare Source (4 Groups)	Antibiotic Use, <i>n</i> (%)		Total, <i>n</i> (%)
	No (0)	Yes (1)	
Public health facility	410 (44.42)	513 (55.58)	923 (100.00)
Private health facility	448 (44.23)	565 (55.77)	1,013 (100.00)
Pharmacy	932 (57.42)	691 (42.58)	1,623 (100.00)
Other informal/other	838 (48.78)	880 (51.22)	1,718 (100.00)
Total	2,628 (49.80)	2,649 (50.20)	5,277 (100.00)

Pearson $\chi^2(3) = 61.74$, $p < 0.001$

Table 2: Cross-tabulation of Detailed Healthcare Source (11 Categories) and Antibiotic Use

Detailed Healthcare Source	Antibiotic Use, <i>n</i> (%)		Total, <i>n</i> (%)
	No (0)	Yes (1)	
Public hospital	226 (40.57)	331 (59.43)	557 (100.00)
Private hospital/clinic	58 (29.74)	137 (70.26)	195 (100.00)
Public clinic/health centre	130 (49.06)	135 (50.94)	265 (100.00)
Private physician	390 (47.68)	428 (52.32)	818 (100.00)
Pharmacy	932 (57.42)	691 (42.58)	1,623 (100.00)
Community health worker	48 (51.06)	46 (48.94)	94 (100.00)
Mobile/outreach clinic	6 (85.71)	1 (14.29)	7 (100.00)
Shop/market/street	77 (41.18)	110 (58.82)	187 (100.00)
Traditional practitioner	646 (55.17)	525 (44.83)	1,171 (100.00)
Relative/friend	3 (60.00)	2 (40.00)	5 (100.00)
Other	112 (31.55)	243 (68.45)	355 (100.00)
Total	2,628 (49.80)	2,649 (50.20)	5,277 (100.00)
Pearson $\chi^2(10) = 159.84, p < 0.001$			

Table 3: **Factors Associated with Antibiotic Use Among Children Under 5 Years (Combined Source Model)**

Variable	Crude OR (95% CI)		Adjusted OR (95% CI)	
	COR	95% CI	AOR	95% CI
Source of care (Ref: Pharmacy)				
Public health facility	1.74	(1.43–2.11)	1.63	(1.33–1.98)
Private health facility (non-pharmacy)	1.65	(1.37–1.98)	1.51	(1.25–1.83)
Other informal/other	1.46	(1.25–1.70)	1.41	(1.20–1.66)
Child sex (Ref: Female)				
Male	–	–	1.07	(0.94–1.22)
Child age group in months (Ref: 0–5)				
6–11	–	–	1.17	(0.91–1.50)
12–23	–	–	1.10	(0.88–1.38)
24–35	–	–	1.22	(0.97–1.54)
36–47	–	–	1.04	(0.82–1.32)
48–59	–	–	1.26	(0.98–1.63)
Child illness status (Ref: Single/No)				
Multiple illnesses	–	–	2.67	(2.34–3.05)
Mother's education (Ref: No education)				
Primary	–	–	0.89	(0.69–1.14)
Secondary	–	–	0.93	(0.73–1.18)
Higher	–	–	1.00	(0.75–1.33)
Residence (Ref: Rural)				
Urban	–	–	0.95	(0.79–1.14)
Wealth quintile (Ref: Poorest)				
Poor	–	–	1.02	(0.84–1.23)
Middle	–	–	1.02	(0.83–1.25)
Richer	–	–	1.30	(1.05–1.61)
Richest	–	–	1.25	(0.98–1.60)
Family size (Ref: Small 1–3)				
Medium (4–6)	–	–	0.84	(0.69–1.02)
Large (7+)	–	–	0.88	(0.70–1.12)
Crowding category (Ref: Low 1–2)				
Medium (3–4)	–	–	1.18	(1.01–1.37)
High (5+)	–	–	1.14	(0.88–1.48)

Table 4: **Factors Associated with Antibiotic Use Among Children Under 5 Years (Detailed Source Model)**

Variable	Crude OR (95% CI)		Adjusted OR (95% CI)	
	COR	95% CI	AOR	95% CI
Detailed source of care (Ref: Pharmacy)				
Public hospital	1.86	(1.48–2.35)	1.72	(1.35–2.19)
Private hospital/clinic	2.99	(2.05–4.38)	2.70	(1.81–4.03)
Public clinic/health centre	1.54	(1.14–2.07)	1.43	(1.06–1.93)
Private physician	1.45	(1.19–1.76)	1.34	(1.10–1.64)
Community health worker	1.83	(1.17–2.86)	1.86	(1.18–2.92)
Mobile/outreach clinic	0.15	(0.02–1.29)	0.14	(0.02–1.23)
Shop/market/street	2.00	(1.42–2.83)	1.88	(1.34–2.64)
Traditional practitioner	1.14	(0.96–1.35)	1.06	(0.89–1.27)
Relative/friend	0.56	(0.07–4.30)	0.44	(0.06–3.28)
Other	3.01	(2.28–3.96)	3.13	(2.34–4.19)
Child sex (Ref: Female)				
Male	–	–	1.09	(0.96–1.23)
Child age group in months (Ref: 0–5)				
6–11	–	–	1.17	(0.91–1.51)
12–23	–	–	1.12	(0.89–1.41)
24–35	–	–	1.27	(1.01–1.61)
36–47	–	–	1.09	(0.85–1.38)
48–59	–	–	1.29	(1.00–1.68)
Child illness status (Ref: Single/No)				
Multiple illnesses	–	–	2.75	(2.40–3.15)
Mother's education (Ref: No education)				
Primary	–	–	0.91	(0.70–1.16)
Secondary	–	–	0.95	(0.74–1.21)
Higher	–	–	0.99	(0.74–1.32)
Residence (Ref: Rural)				
Urban	–	–	0.90	(0.75–1.09)
Wealth quintile (Ref: Poorest)				
Poor	–	–	1.02	(0.84–1.24)
Middle	–	–	1.02	(0.83–1.26)
Richer	–	–	1.28	(1.03–1.59)
Richest	–	–	1.21	(0.94–1.56)
Family size (Ref: Small 1–3)				
Medium (4–6)	–	–	0.83	(0.68–1.01)
Large (7+)	–	–	0.88	(0.69–1.11)
Crowding category (Ref: Low 1–2)				
Medium (3–4)	–	–	1.16	(1.00–1.35)
High (5+)	–	–	1.17	(0.90–1.51)