

Linear classifiers and Support Vector Machines (SVM)

Sylvain Le Corff

1. Reminder on Logistic regression
2. Linear Support Vector Machine (SVM)

1. Reminder on Logistic regression

2. Linear Support Vector Machine (SVM)

Setting

- Historical data about **individuals** $i = 1, \dots, n$.
- **Features** vector $X_i \in \mathbb{R}^d$ for each individual i .
- For each i , X_i **belongs to a group** ($Y_i = 0$) or not ($Y_i = 1$).
- $Y_i \in \{0, 1\}$ is the **label** of i .

Objective

- Given a new feature vector, **predict a label in $\{0, 1\}$** .
- Use data $\mathcal{D}_n = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$ **to construct a classifier**.

The best solution f^* (which is independent of \mathcal{D}_n) is

$$f^* = \operatorname{argmin}_{f: \mathbb{R}^d \rightarrow \{0,1\}} \mathbb{E}[\mathbb{1}_{Y \neq f(X)}] = \operatorname{argmin}_{f: \mathbb{R}^d \rightarrow \{0,1\}} \mathbb{P}(Y \neq f(X)).$$

Bayes Predictor (explicit solution)

Binary classification with 0 – 1 loss:

$$f^*(X) = \begin{cases} +1 & \text{if } \mathbb{P}(Y = 1|X) \geq \mathbb{P}(Y = 0|X) \\ & \Leftrightarrow \mathbb{P}(Y = 1|X) \geq 1/2, \\ 0 & \text{otherwise.} \end{cases}$$

The explicit solution requires to know the conditional law of Y given X ...

Fully parametric modeling.

Estimate the law of (X, Y) and use the **Bayes formula** to deduce an estimate of the conditional law of Y : *LDA/QDA, Naive Bayes...*

Parametric conditional modeling.

Estimate the conditional law of Y by a **parametric** law: *linear regression, logistic regression, Feed Forward Neural Networks...*


Nonparametric conditional modeling.

Estimate the conditional law of Y by a **non parametric** estimate: *kernel methods, nearest neighbors...*

In the LDA case, the classification rule is of the form:

$$f^*(x) = 1 \Leftrightarrow \langle w, x \rangle + b \geq 0,$$

where w and b depends on the model parameters.



- Relax the Gaussian assumption ? (logistic model, SVM).
- Design nonlinear classification rules ? (kernels, neural networks).

- ▶ One of the most widely used classification algorithm.
- ▶ Logistic model is **generalized linear model** of the linear model in the context of binary classification ($\mathcal{Y} = \{0, 1\}$).
- ▶ It **models the distribution of Y given X** . For $y \in \{0, 1\}$

$$\mathbb{P}(Y = 1|X) = \sigma(X^\top w + b)$$

where $\sigma : z \mapsto (1 + e^{-z})^{-1}$, $w \in \mathbb{R}^d$ is a vector of model weights and $b \in \mathbb{R}$ is the intercept, and where σ is the **sigmoid** function:

- ▶ The sigmoid is a modelling choice to map $\mathbb{R} \rightarrow [0, 1]$ (to model a probability).
- ▶ We could also consider

$$\mathbb{P}(Y = 1|X) = F(X^\top w + b)$$

for any **distribution function** F .

- ▶ Another popular choice is the Gaussian distribution

$$F(z) = \mathbb{P}(\mathcal{N}(0, 1) \leq z),$$

which leads to the **probit model**.

- In the case of the sigmoid (**logistic regression**),

$$\mathbb{P}(Y = 1|X) = \frac{\exp(b + w^\top X)}{1 + \exp(b + w^\top X)} = \frac{1}{1 + \exp(-(b + w^\top X))}$$

$$\mathbb{P}(Y = 0|X) = \frac{1}{1 + \exp(b + w^\top X)}$$

- Log-odd ratio:

$$\log \left(\frac{\mathbb{P}(Y = 1|X)}{\mathbb{P}(Y = 0|X)} \right) = X^\top w + b.$$

Compute \hat{w}_n and \hat{b}_n as follows:

$$(\hat{w}_n, \hat{b}_n) \in \operatorname{argmin}_{w \in \mathbb{R}^d, b \in \mathbb{R}} \frac{1}{n} \sum_{i=1}^n \left(-Y_i (X_i^\top w + b) + \log(1 + e^{X_i^\top w + b}) \right).$$

→ It is an **average of losses**, one for each sample point.

→ It is a **convex and smooth problem**.

Using the **logistic loss** function

$$\ell : (y, y') \mapsto \log(1 + e^{-yy'})$$

yields

$$(\hat{w}_n, \hat{b}_n) \in \operatorname{argmin}_{w \in \mathbb{R}^d, b \in \mathbb{R}} \frac{1}{n} \sum_{i=1}^n \ell(Y_i, \langle w, X_i \rangle + b).$$

1. Reminder on Logistic regression

2. Linear Support Vector Machine (SVM)


- ▶ Training dataset of pairs (X_i, Y_i) for $1 \leq i \leq n$.
- ▶ Features $X_i \in \mathbb{R}^d$ and labels $Y_i \in \{-1, 1\}$.
- ▶ Given a features vector $x \in \mathbb{R}^d$, we want to predict its associated label.
- ▶ Focus on linear classification, i.e. classifiers defined by $w \in \mathbb{R}^d$ and $b \in \mathbb{R}$ such that

$$h(x) = \text{sign}(x^\top w + b).$$

Preliminary definitions

A dataset is **linearly separable** if there exists an hyperplane H (linear classification rule) such that the following assumptions hold.

- Points $X_i \in \mathbb{R}^d$ such that $Y_i = 1$ are on one side of the hyperplane.
- Points $X_i \in \mathbb{R}^d$ such that $Y_i = -1$ are on the other side.
- H does not pass through any point X_i .



A **hyperplane** is a translation of a set of vectors orthogonal to w .

$$H_{w,b} = \{x \in \mathbb{R}^d : w^\top x + b = 0\}.$$

→ $w \in \mathbb{R}^d$ is a **non-zero vector normal** to the hyperplane.

→ $b \in \mathbb{R}$ is a scalar.

Following for instance the results obtained for linear discriminant analysis and logistic regression, a **hyperplane $H_{w,b}$ may be used as a classifier** by defining

$$h_{w,b} : x \mapsto \begin{cases} 1 & \text{if } \langle w; x \rangle + b > 0, \\ -1 & \text{otherwise.} \end{cases}$$

If H do not pass through any sample point x_i , we can scale w and b so that

$$\min_{(x,y) \in D_n} |w^T x + b| = 1$$

For such w and b , we call H the **canonical** hyperplane




Figure: The marginal hyperplanes are the hyperplanes parallel to the separating hyperplane and passing through the closest points on the negative or positive sides.

The distance of any point $x \in \mathbb{R}^d$ to H is given by

$$d(x, H_{w,b}) = \frac{|\langle w, x \rangle + b|}{\|w\|}$$

So, if H is a canonical hyperplane, its **margin** is given by

$$\min_{(x,y) \in D_n} \frac{|w^\top x + b|}{\|w\|} = \frac{1}{\|w\|}.$$

If \mathcal{D}_n is strictly **linearly separable**, we can find a **canonical separating hyperplane**

$$H_{w,b} = \{x \in \mathbb{R}^d : w^\top x + b = 0\},$$

that satisfies

$$|\langle w, X_i \rangle + b| \geq 1 \text{ for any } i = 1, \dots, n,$$

which entails that a point X_i is correctly classified if

$$Y_i(\langle X_i, w \rangle + b) \geq 1.$$

The **margin** of H is equal to $1/\|w\|$.

Hard Support Vector Machines is a classification procedure which aims at building a linear classifier with the largest possible margin, i.e. **the largest minimal distance between a point in the training set and the hyperplane.**

The hyperplane which **correctly separates all training data sets with the largest margin** is obtained with:

$$(\hat{w}_n, \hat{b}_n) \in \underset{\substack{(w,b) \in \mathbb{R}^d \times \mathbb{R}; \|w\|=1, \\ \forall i \in \{1, \dots, n\}, Y_i(\langle w; X_i \rangle + b) > 0}}{\operatorname{argmax}} \left\{ \min_{1 \leq i \leq n} |\langle w; X_i \rangle + b| \right\} .$$

The **hard Support Vector Machines** procedure is equivalent to solving the following optimization problem:

$$(\hat{w}_n, \hat{b}_n) \in \underset{(w,b) \in \mathbb{R}^d \times \mathbb{R}; \|w\|=1}{\operatorname{argmax}} \left\{ \min_{1 \leq i \leq n} Y_i (\langle w; X_i \rangle + b) \right\},$$

A **solution to the hard Support Vector Machines optimization** problem is obtained by setting $(\hat{w}_n, \hat{b}_n) = (w_\star / \|w_\star\|, b_\star / \|w_\star\|)$ where

$$(w_\star, b_\star) \in \underset{\substack{(w,b) \in \mathbb{R}^d \times \mathbb{R} \\ \forall i \in \{1, \dots, n\}, Y_i (\langle w; X_i \rangle + b) \geq 1}}{\operatorname{argmin}} \|w\|^2.$$

Proof on blackboard !

Maximum margin problem

In the **Hard SVM case**, a way of classifying \mathcal{D}_n with maximum margin is to solve the following problem:

$$(w_*, b_*) \in \operatorname{argmin}_{w \in \mathbb{R}^d, b \in \mathbb{R}} \left\{ \frac{1}{2} \|w\|_2^2 \right\},$$

under **the constraints**:

$$Y_i(\langle X_i, w \rangle + b) \geq 1,$$

for all $1 \leq i \leq n$.

- ▶ This problem admits a **unique** solution.
- ▶ It is a **'quadratic programming'** problem.
- ▶ Dedicated optimization algorithms can solve this on a large scale very efficiently

Consider a **constrained optimization problem**:

$$P^{\star} = \min_{x \in \mathbb{R}^d} f(x)$$

under **the constraints**, for all $1 \leq i \leq p$, $1 \leq j \leq q$,

$$h_i(x) = 0 \quad \text{and} \quad g_j(x) \leq 0,$$

where $f, h_1, \dots, h_p, g_1, \dots, g_q$ are defined on \mathbb{R}^d .

Lagrangian

The **Lagrangian** is the function defined on $\mathbb{R}^d \times \mathbb{R}^p \times \mathbb{R}_+^q$ by

$$\mathcal{L}(x, \lambda, \mu) := f(x) + \sum_{i=1}^p \lambda_i h_i(x) + \sum_{j=1}^q \mu_j g_j(x)$$

$\lambda \in \mathbb{R}^p$, $\mu \in \mathbb{R}_+^q$ are the **Lagrange** or **dual** variables.

The **Lagrange dual** function is defined by:

$$D : (\lambda, \mu) \mapsto \inf_{x \in \mathbb{R}^d} \mathcal{L}(x, \lambda, \mu).$$

Let \mathcal{D} be the subset of \mathbb{R}^d of feasible points. Using

$$\sup_{\mu \geq 0, \lambda} \inf_{x \in \mathbb{R}^d} \mathcal{L}(x, \lambda, \mu) \leq \inf_{x \in \mathbb{R}^d} \sup_{\mu \geq 0, \lambda} \mathcal{L}(x, \lambda, \mu) = \inf_{x \in \mathcal{D}} f(x)$$

yealds the **weak duality relation**:

$$D^* = \sup_{\mu \geq 0, \lambda} D(\lambda, \mu) \leq \inf_{x \in \mathcal{D}} f(x) = P^*.$$

Equality, known as **strong duality relation** requires some additional assumptions.

Strong duality holds under

- ▶ **convexity** of the problem
- ▶ **constraint qualifications**

A simple way to have constraint qualification (sufficient but not necessary)

Slater's conditions

There is some strictly feasible point $x \in \mathbb{R}^d$ such that

$$h_i(x) = 0 \quad \text{for all } i = 1, \dots, p$$

$$g_j(x) < 0 \quad \text{for all } j = 1, \dots, q$$

Assume that (i) f, g_1, \dots, g_q are **differentiable** and **convex**, (ii) that h_1, \dots, h_p are **affine** functions and that (iii) **Slater's condition** holds.

Then $x^* \in \mathbb{R}^d$ is a solution of the primal problem if and only if there is $(\lambda^*, \mu^*) \in \mathbb{R}^p \times \mathbb{R}_+^q$ such that

$$\nabla_x \mathcal{L}(x^*, \lambda^*, \mu^*) = \nabla f(x^*) + \sum_{i=1}^p \lambda_i^* \nabla h_i(x^*) + \sum_{j=1}^q \mu_j^* \nabla g_j(x^*) = 0,$$

with

$$\begin{aligned} h_i(x^*) &= 0 && \text{for any } i = 1, \dots, p, \\ g_j(x^*) &\leq 0 && \text{for any } j = 1, \dots, q, \\ \mu_j^* g_j(x^*) &= 0 && \text{for any } j = 1, \dots, q. \end{aligned}$$

- ▶ These are known as the **KKT conditions**
- ▶ The last one is called **complementary slackness**

Take-home message: Lagrangian duality

If

- primal problem is **convex** and
- constraint functions satisfy the **Slater's** conditions

then

- ▶ **strong duality** holds.

If in addition we have that

- functions f, g_1, \dots, g_n are **differentiable**

then

- ▶ KKT conditions are **necessary and sufficient** for optimality

In the **Hard SVM case**, a way of classifying \mathcal{D}_n with maximum margin is to solve the following problem:

$$\min_{w \in \mathbb{R}^d, b \in \mathbb{R}} f(w),$$

under **the constraints**:

$$g_i(w) \leq 0,$$

for all $1 \leq i \leq n$, where

- ▶ $f(w) = \|w\|_2^2 / 2$ is **strongly convex**, since

$$\nabla^2 f(w) = I_d \succ 0$$

- ▶ Constraints are $g_i(w, b) \leq 0$ with **affine** functions

$$g_i(w, b) = 1 - Y_i(\langle X_i, w \rangle + b).$$

The **KKT conditions** allows to obtain the dual formulation of the problem.

Lagrangian

- ▶ Introduce dual variables $\mu_i \geq 0$ for $i = 1, \dots, n$ corresponding to the constraints $g_i(w, b) \leq 0$.
- ▶ For $w \in \mathbb{R}^d$, $b \in \mathbb{R}$ and $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{R}_+^n$, define the Lagrangian

$$\mathcal{L}(w, b, \mu) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^n \mu_i (1 - y_i (\langle w, x_i \rangle + b)) .$$

$$\mathcal{L}(w, b, \mu) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^n \mu_i (1 - y_i(\langle w, x_i \rangle + b))$$

KKT conditions

Set the gradient to zero

$$\nabla_w L(w, b, \mu) = w - \sum_{i=1}^n \mu_i y_i x_i = 0 \quad \text{namely} \quad w = \sum_{i=1}^n \mu_i y_i x_i$$

$$\nabla_b L(w, b, \mu) = - \sum_{i=1}^n \mu_i y_i = 0 \quad \text{namely} \quad \sum_{i=1}^n \mu_i y_i = 0$$

Write the complementary slackness condition: $\forall i = 1, \dots, n$

$$\mu_i (1 - y_i(\langle w, x_i \rangle + b)) = 0 \quad \text{namely} \quad \mu_i = 0 \quad \text{or} \quad y_i(\langle w, x_i \rangle + b) = 1$$

At the optimum,

- ▶ There are **dual** variables $\mu_i \geq 0$ such that the **primal** solution (w, b) satisfies

$$w = \sum_{i=1}^n \mu_i y_i x_i$$

- ▶ We have that

$$\mu_i \neq 0 \quad \text{iff} \quad y_i(\langle w, x_i \rangle + b) = 1$$

This means that

- ▶ w writes as a linear combination of the features vectors x_i that belong to the marginal hyperplanes $\{x \in \mathbb{R}^d : w^T x + b = \pm 1\}$
- ▶ These vectors x_i are called **support vectors**

The support vectors fully define the maximum-margin hyperplane, hence the name **Support Vector Machine**

Linearly separable dataset

31 / 52

```
X, y = make_blobs(n_samples = 200, centers = 2, random_state = 0, cluster_std = 0.50)
simulated_data = pd.DataFrame(columns = ["X1", "X2", "Label"])


simulated_data["X1"] = X[:,0]
simulated_data["X2"] = X[:,1]
simulated_data["Label"] = y

# Use the 'Label' argument to provide a factor variable
sns.set_style("whitegrid")
sns.lmplot(x = "X1", y = "X2", data = simulated_data, fit_reg = False, hue = 'Label', legend = True)

slope = 1.1
offset = 0.85
margin = 0.2

xfit = np.linspace(-1, 3.5)
yfit = m * xfit + b

plt.plot(xfit, yfit, '--k')
plt.fill_between(xfit, yfit - d, yfit + d, color = 'AAAAAA', alpha = 0.2)
```



```


# Classification based on a support vector classifier
model = SVC(kernel="linear", C=10)
model.fit(X, y)
sns.set_style("whitegrid")
sns.lmplot(x = "x", y = "y", data = simulated_data, fit_reg = False, hue = 'Label', legend = True)

xlim = [np.min(X[:,0]), np.max(X[:,0])]
ylim = [np.min(X[:,1]), np.max(X[:,1])]
xplot = np.linspace(xlim[0], xlim[1], 30)
yplot = np.linspace(ylim[0], ylim[1], 30)


Vplot, Xplot = np.meshgrid(yplot, xplot)
xy = np.vstack([Xplot.ravel(), Vplot.ravel()]).T
P = model.decision_function(xy).reshape(Xplot.shape)

# plot decision boundary and margins
plt.contour(Xplot, Vplot, P, colors = 'k', levels = [-1, 0, 1], alpha = 0.8,
            linestyle = ['--', '-', '--'])

```




```
sns.set_style("whitegrid")
sns.lmplot(x = "x", y = "y", data = simulated_data, fit_reg = False, hue = 'Label', legend = True)
# plot decision boundary and margins
plt.contour(Xplot, Yplot, P, colors = 'k', levels = [-1, 0, 1], alpha = 0.8,
            linestyle = ['--', '-', '--'])
plt.scatter(model.support_vectors_[0], model.support_vectors_[1], s = 5, c = 'k');
```



Under **strong duality**, primal and dual problems are strongly related, and one can be used to solve the other.

- Recall that the Lagrangian is

$$\mathcal{L}(w, b, \mu) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^n \mu_i (1 - y_i (\langle w, x_i \rangle + b))$$

- Plug $w = \sum_{i=1}^n \mu_i y_i x_i$ in this equation to obtain

$$\begin{aligned} \mathcal{L}(w, b, \mu) = & \frac{1}{2} \left\| \sum_{i=1}^n \mu_i y_i x_i \right\|_2^2 + \sum_{i=1}^n \mu_i - b \sum_{i=1}^n \mu_i y_i \\ & - \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle. \end{aligned}$$

- ▶ Recalling that $\sum_{i=1}^n \mu_i y_i = 0$ and doing some algebra provides the dual formulation.

Dual formulation

The dual problem amounts to solve:

$$\max_{\mu \in \mathbb{R}^n} \quad \sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle ,$$

under the constraints

$$\mu_i \geq 0 \quad \text{and} \quad \sum_{i=1}^n \mu_i y_i = 0 \quad \text{for all } i = 1, \dots, n .$$

- ▶ As in the primal formulation, it is again a **quadratic programming problem**.
- ▶ At optimum, we have (using KKT conditions) that the decision function is expressed using the dual variables as

$$x \mapsto \text{sign}(w^\top x + b) = \text{sign}\left(\sum_{i=1}^n \mu_i y_i \langle x, x_i \rangle + b\right)$$

- ▶ The intercept b can be expressed for any support vector x_i as

$$b = y_i - \sum_{j=1}^n \mu_j y_j \langle x_i, x_j \rangle$$

This allows to write the margin as a function of the dual variables

- ▶ Multiplying the last equality by $\mu_i y_i$ and summing entails

$$\sum_{i=1}^n \mu_i y_i b = \sum_{i=1}^n \mu_i y_i^2 - \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle$$

- ▶ Namely recalling that at optimum $\sum_{i=1}^n \mu_i y_i = 0$ and $w = \sum_{i=1}^n \mu_i y_i x_i$ we get

$$0 = \sum_{i=1}^n \mu_i = \|w\|_2^2, \quad \text{namely}$$
$$\text{margin} = \frac{1}{\|w\|_2^2} = \frac{1}{\sum_{i=1}^n \mu_i} = \frac{1}{\|\mu\|_1}$$

→ Restricting the problem to linearly separable training data sets is a **somehow strong assumption**.

→ Inequality constraints in the quadratic optimization problem **can be relaxed**.

Replace the constraints

$$Y_i(\langle w, X_i \rangle + b) \geq 1 \quad \text{for all } i = 1, \dots, n,$$

→ Restricting the problem to linearly separable training data sets is a **somehow strong assumption**.

→ Inequality constraints in the quadratic optimization problem **can be relaxed**.

Replace the constraints

$$Y_i(\langle w, X_i \rangle + b) \geq 1 \quad \text{for all } i = 1, \dots, n,$$

that are too strong, by the **relaxed** ones

$$Y_i(\langle w, X_i \rangle + b) \geq 1 - s_i \quad \text{for all } i = 1, \dots, n,$$

for **slack variables** $s_1, \dots, s_n \geq 0$

The original problem

$$\min_{w \in \mathbb{R}^d, b \in \mathbb{R}} \frac{1}{2} \|w\|_2^2,$$

under the constraints

$$Y_i(\langle X_i, w \rangle + b) \geq 1 \text{ for all } i = 1, \dots, n.$$

is replaced by the relaxation using slack variables

$$\min_{w \in \mathbb{R}^d, b \in \mathbb{R}, s \in \mathbb{R}^n} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n s_i,$$

under the constraints

$$Y_i(\langle X_i, w \rangle + b) \geq 1 - s_i \text{ and } s_i \geq 0 \forall i = 1, \dots, n.$$

- ▶ The slack $s_i \geq 0$ measures the distance by which x_i violates the desired inequality $Y_i(\langle X_i, w \rangle + b) \geq 1$
- ▶ A vector x_i with $0 < Y_i(\langle X_i, w \rangle + b) < 1$ is correctly classified but is an outlier, since $s_i > 0$
- ▶ If we omit outliers, training data is correctly classified by the hyperplane $\{x \in \mathbb{R}^d : \langle x, w \rangle + b = 0\}$ with a margin $1/\|w\|_2^2$
- ▶ The margin $1/\|w\|_2^2$ is called a **soft-margin** (in the non-separable case), while it is a **hard-margin** in the separable case

Relaxed margin problem

$$\min_{w \in \mathbb{R}^d, b \in \mathbb{R}, s \in \mathbb{R}^n} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n s_i,$$

under the constraints

$$Y_i(\langle X_i, w \rangle + b) \geq 1 - s_i \text{ and } s_i \geq 0 \quad \forall i = 1, \dots, n.$$

Once again:

- ▶ This problem admits a **unique** solution.
- ▶ It is a quadratic programming problem.

The constant $C > 0$ is chosen using V -fold cross-validation.

Lagrangian

$$\begin{aligned}\mathcal{L}(w, b, s, \mu, \beta) = & \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n s_i \\ & + \sum_{i=1}^n \mu_i (1 - s_i - y_i(\langle w, x_i \rangle + b)) - \sum_{i=1}^n \beta_i s_i\end{aligned}$$

with $\mu_i \geq 0$ and $\beta_i \geq 0$.

At optimum:

- ▶ set the **gradients** ∇_w , ∇_b and ∇_s to zero ;
- ▶ write **the complementary conditions**.

$$\nabla_w L(w, b, s, \mu, \beta) = w - \sum_{i=1}^n \mu_i y_i x_i = 0 \quad \text{i.e.} \quad w = \sum_{i=1}^n \mu_i y_i x_i$$

$$\nabla_b L(w, b, s, \mu, \beta) = - \sum_{i=1}^n \mu_i y_i = 0 \quad \text{i.e.} \quad \sum_{i=1}^n \mu_i y_i = 0$$

$$\nabla_s L(w, b, s, \mu, \beta) = C - \mu_i - \beta_i = 0 \quad \text{i.e.} \quad \mu_i + \beta_i = C$$

and the complementary condition

$$\mu_i (1 - s_i - y_i (\langle w, x_i \rangle + b)) = 0 \quad \text{i.e.} \quad \mu_i = 0 \quad \text{or} \quad y_i (\langle w, x_i \rangle + b) = 1 - s_i$$

$$\beta_i s_i = 0 \quad \text{i.e.} \quad \beta_i = 0 \quad \text{or} \quad s_i = 0$$

for all $i = 1, \dots, n$

- ▶ $w = \sum_{i=1}^n \mu_i y_i x_i$
- ▶ If $\mu_i \neq 0$ we say that x_i is a **support vector** and in this case $y_i(\langle w, x_i \rangle + b) = 1 - s_i$.
 - ▶ If $s_i = 0$ then x_i belongs to a margin hyperplane.
 - ▶ If $s_i \neq 0$ then x_i is an outlier and $\beta_i = 0$ and then $\mu_i = C$.

Support vectors either belong to a marginal hyperplane, or are outliers with $\mu_i = C$

- ▶ Plugging $w = \sum_{i=1}^n \mu_i y_i x_i$ in $L(w, b, s, \mu, \beta)$ leads to the same formula as before

$$\sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle$$

- ▶ Plugging $w = \sum_{i=1}^n \mu_i y_i x_i$ in $L(w, b, s, \mu, \beta)$ leads to the same formula as before

$$\sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle$$

- ▶ with the constraints

$$\mu_i \geq 0, \quad \beta_i \geq 0, \quad \sum_{i=1}^n \mu_i y_i = 0, \quad \mu_i + \beta_i = C$$

that can be rewritten for as

$$0 \leq \mu_i \leq C, \quad \sum_{i=1}^n \mu_i y_i = 0$$

for all $i = 1, \dots, n$

Dual problem

$$\begin{aligned} \max_{\mu \in \mathbb{R}^n} \quad & \sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle \\ \text{subject to} \quad & 0 \leq \mu_i \leq C \text{ and } \sum_{i=1}^n \mu_i y_i = 0 \text{ for all } i = 1, \dots, n \end{aligned}$$

- ▶ This is the same problem as before, but with the extra constraint

$$\mu_i \leq C$$

- ▶ It is again a convex quadratic program

As in the linearly separable case, the **label** prediction is expressed using the **dual variables**.

Labels given by

$$x \mapsto \text{sign}(w^T x + b) = \text{sign}\left(\sum_{i=1}^n \mu_i y_i \langle x, x_i \rangle + b\right)$$

The intercept b can be expressed for a support vector x_i such that $0 < \mu_i < C$ as

$$b = y_i - \sum_{j=1}^n \mu_j y_j \langle x_i, x_j \rangle$$

The dual problem

$$\max_{\mu \in \mathbb{R}^n} \quad \sum_{i=1}^n \mu_i - \frac{1}{2} \sum_{i,j=1}^n \mu_i \mu_j y_i y_j \langle x_i, x_j \rangle$$

subject to $0 \leq \mu_i \leq C$ and $\sum_{i=1}^n \mu_i y_i = 0$ for all $i = 1, \dots, n$

and the label prediction (using dual variables)

$$x \mapsto \text{sign}(w^T x + b) = \text{sign}\left(\sum_{i=1}^n \mu_i y_i \langle x, x_i \rangle + b\right)$$

depends only on the features x_i via their **inner products** $\langle x_i, x_j \rangle$!

► This will be particularly important later: **kernel methods**

Going back to the primal problem

$$\min_{w \in \mathbb{R}^d, b \in \mathbb{R}, s \in \mathbb{R}^n} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n s_i$$

subject to $y_i(\langle x_i, w \rangle + b) \geq 1 - s_i$ and $s_i \geq 0$ for all $i = 1, \dots, n$

Going back to the primal problem

$$\min_{w \in \mathbb{R}^d, b \in \mathbb{R}, s \in \mathbb{R}^n} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n s_i$$

subject to $y_i(\langle x_i, w \rangle + b) \geq 1 - s_i$ and $s_i \geq 0$ for all $i = 1, \dots, n$

We remark that it can be rewritten as follows.

Reformulation of the primal problem

$$\operatorname{argmin}_{w \in \mathbb{R}^d, b \in \mathbb{R}} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n \max \left(0, 1 - y_i(\langle x_i, w \rangle + b) \right).$$

The hinge loss function

$$\ell(y, y') = \max(0, 1 - yy') = (1 - yy')_+,$$

the problem can be written as

Reformulation of the primal problem

$$\operatorname{argmin}_{w \in \mathbb{R}^d, b \in \mathbb{R}} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n \ell(y_i, \langle x_i, w \rangle + b).$$

Leads to an alternative understanding of the linear SVM.

Recall that the natural loss is the 0/1 one given by

$$\ell_{0/1}(y, z) = \mathbb{1}_{yz \leq 0}.$$

Instead of the Linear SVM, it would be nice to consider

$$\operatorname{argmin}_{w \in \mathbb{R}^d, b \in \mathbb{R}} \frac{1}{2} \|w\|_2^2 + C \sum_{i=1}^n \mathbb{1}_{y_i(\langle x_i, w \rangle + b) \leq 0},$$

but impossible numerically (NP-hard)

Hinge loss is a **convex surrogate** for the 0/1 loss

LDA/QDA

- ▶ Model: $X|Y \sim \mathcal{N}$

Logistic regression

- ▶ Logistic regression has a nice probabilistic interpretation
- ▶ Model $\text{logit}(\mathbb{P}(Y = 1|X))$ is linear in X
- ▶ Relies on the choice of the logit link function
- ✗ does not work on separable dataset

SVM

- ▶ No model, only aims at separating points
- ✓ Thought for separable case
- ✓ But can be relaxed for the non-separable case