

LSPU Self-paced Learning Module (SLM)

Course	SCI 1- Chemistry for Engineers
Sem/AY	First Semester/2023-2024
Module No.	5
Lesson Title	ELECTROCHEMICAL ENERGY
Week Duration	1
Date	October 2-6, 2023
Description of the Lesson	This module discusses the oxidation reduction, the functions of voltaic and electrolytic cells, calculate the standard cell potential generated by a voltaic cell and the properties and types of batteries.

Learning Outcomes

Intended Learning Outcomes	<p>Students should be able to meet the following intended learning outcomes:</p> <ul style="list-style-type: none"> • Recognize oxidation reduction (redox) reactions and be able to assign oxidation numbers to atoms in molecules and ions. • Describe the functions of the various components of simple voltaic and electrolytic cells. • Diagram electrochemical cells, labeling the anode, cathode, and directions of ion and electron movement. • Given appropriate reduction potentials, calculate the cell voltage (standard cell potential, E_{cell}°) generated by a voltaic cell. • Discuss how a spontaneous redox reaction can be used to create a battery and recognize the connection between the components of the cell and the properties of the battery.
Targets/ Objectives	<p>At the end of the lesson, students should be able to:</p> <ul style="list-style-type: none"> • Assign oxidation numbers to atoms in molecules and ions. • Write the oxidation and reduction half reactions. • Identify the compounds being oxidized and reduced. • Identify the oxidizing agent and reducing agent. • Balance a redox reaction by the half-reaction method. • Identify the various components of simple voltaic and electrolytic cells.



- Calculate the cell voltage (standard cell potential, E_{cell}°) generated by a voltaic cell.
- Recognize the connection between the components of the cell and the properties of the battery.
- Recognize the chemical reaction used in a lead-acid storage battery and lithium ion battery.
- Describe the basic components and processes in the H₂/O₂ fuel cell.



Student Learning Strategies

Online Activities (Synchronous/ Asynchronous)

- A. Online Discussion via Google Meet
You will be directed to attend in a One-Hour class discussion on Electrochemical energy. To have access to the Online Discussion, refer to the announcement that will be posted in the Google Classroom.

(For further instructions, refer to your Google Classroom and see the schedule of activities for this module)

- B. Learning Guide Questions:
1. Define Electrochemistry.
 2. What is redox reaction?
 3. What are the rules in assigning oxidation states?
 4. What are the rules in half-reaction method?
 5. What is the difference between an electrolytic cell and a galvanic cell?

***Note:** The insight that you will post on online discussion forum using Learning Management System (LMS) will receive additional scores in class participation.*

Offline Activities (e-Learning/Self- Paced)

Lecture Guide

Electrochemistry

- Is the study of the interconversion of electrical and chemical energy. This conversion takes place in an electrochemical cell that may be a/an
 - a. Voltaic (galvanic) cell in which a spontaneous reaction generates electrical energy.
 - b. Electrolytic cell in which electrical energy is used to bring about a nonspontaneous reaction.

Electrochemistry is the study of chemical processes that cause electrons to move. This movement of electrons is called electricity, which can be generated



ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna


by movements of electrons from one element to another in a reaction known as an oxidation-reduction ("redox") reaction.

A **redox reaction** is a reaction that involves a change in oxidation state of one or more elements. An **oxidation-reduction reaction** is any chemical reaction in which the oxidation number of a molecule, atom, or ion change by gaining or losing an electron.


Redox reactions are common and vital to some of the basic functions of life, including photosynthesis, respiration, combustion, and corrosion or rusting.

- In an electrochemical cell, these two half-reactions occur at two different electrodes, which most often consist of metal plates or wires.

Reduction occurs at the **cathode**



Oxidation takes place at the **anode**, where a species such as zinc metal produces electrons:



It is always true that in an electrochemical cell, **anions move to the anode; cations move to the cathode**

Oxidation-Reduction Reactions

Terminology Concepts

A **redox reaction** involves a transfer of electrons between two species.

Oxidation occurs when a species loses electrons and increases its oxidation number (O.N.). An equation that shows this loss is called an oxidation half-reaction. In this half-reaction, the electrons are in the product side.

Reduction occurs when a specie gain electron and decreases its oxidation number (O.N.). An equation that shows this gain is called a reduction half-reaction. In this half-reaction, the electrons are in the reactant side.

- The ion or molecule that donates the electrons (i.e. oxidized) is called the **reducing agent**.
- The ion or molecule that accepts the electrons (i.e. reduced) is called the **oxidizing agent**.
- Oxidation and reduction occur together in the same reaction. There is no net change in the number of electrons, just an exchange of electrons


Example 1: Iron-Vanadium Couple

Given the redox reaction



ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna



which species is oxidized? Which is reduced? Identify the reducing agent and the oxidizing agent.

Solution:

Fe^{3+} is reduced into Fe^{2+} and

V^{2+} is oxidized into V^{3+} .

This is because the oxidized species loses electrons, and the reduced species gains electrons.

Iron gains an electron ($Fe^{3+} \rightarrow Fe^{2+}$), and vanadium loses an electron ($V^{2+} \rightarrow V^{3+}$).

Thus, Fe^{3+} is the oxidizing agent and V^{2+} is the reducing agent.

Rules for Assigning Oxidation States

1. Free elements have an oxidation state of 0. (e.g., He, N_2 , O_2 has an oxidation state of 0)
2. The oxidation state of one atom ion must equal the net charge. (e.g., F^- oxidation state is -1, K^+ oxidation state is +1).
3. The sum of the oxidation states are equal to the total net charge for a compound. (e.g., MnO_4^- has a net charge of -1, Mn (+7) O_4 (-8) = -1)
4. The alkali metals (Group I elements) have an oxidation state of +1. (e.g., Li_2O , Li = +1)
5. The alkaline earth metals (Group II elements) always have an oxidation state of +2. (e.g., CaO, Ca = +2)
6. Oxygen has an oxidation state of -2 in a compound
7. Fluorine has an oxidation state of -1 in a compound
8. Hydrogen has an oxidation state of +1 in a compound.
9. Transition metals and other metals may have more than one common ionic charge. (e.g., Chromium's common ionic charges are Cr^{2+} and Cr^{3+})

REMEMBER!!!

The sum of the oxidation states is equal to zero for neutral compounds and equal to the charge for polyatomic ion species

Example 2. What is the oxidation state of magnesium in MgF_2 ?

Answer:

MgF_2 total charge = 0

Total Charge = (+2) + (-1 x 2) = 0



Using rule 4 and 7.

Example 3. What is the oxidation state of hydrogen in H_2O ?


Answer:

H_2O total charge = 0

Total Charge = $(+1 \times 2) + (-2) = 0$

Using rule 6 and 8

Example 4: Assigning Oxidation State. Determine the Oxidation States of each element in the following reactions:



Answer:


Fe and O_2 are free elements therefore they each have an oxidation state of 0.

O has an oxidation state of -2, which means Fe has an oxidation state of +3.



Answer:


The oxidation state of Fe corresponds to its charge; therefore, the oxidation state is +2.



Answer:

Ag has an oxidation state of 0, H has an oxidation state of +1, S has an oxidation state of -2 and hence Ag in $Ag_2S_{(g)}$ has an oxidation state of +1.

Example 5. Determine the Oxidation State of the bold element in each of the following:




The oxidation numbers of Na and O are +1 and -2. Because sodium phosphite is neutral, the sum of the oxidation numbers must be zero.

Letting x be the oxidation number of phosphorus,

$$3(+1) + x + 3(-2) = 0$$

x=oxidation number of P= +3.



Hydrogen and oxygen have oxidation numbers of +1 and -2. The ion has a charge of -1, so the sum of the oxidation numbers must be -1.

Letting y be the oxidation number of phosphorus,

$$y + 2(+1) + 4(-2) = -1$$




ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna


y= oxidation number of P= +5

Example 6. Identifying Reduced and Oxidized Elements. Determine which element is oxidized and which element is reduced in the following reactions (be sure to include the oxidation state of each):



Answer:

Zn is oxidized (Oxidation number: $0 \rightarrow +2$); H^+ is reduced (Oxidation number: $+1 \rightarrow 0$)



Answer:

Al is oxidized (Oxidation number: $0 \rightarrow +3$); Cu^{2+} is reduced ($+2 \rightarrow 0$)



Answer:

This is not a redox reaction because each element has the same oxidation number in both reactants and products: O= -2, H= +1, C= +4.

Balancing Redox Reactions

Method 1: Oxidation Number Method

Step 1: Assign oxidation numbers to each atom.

Step 2: Determine the net change in charge to determine the ratio of atoms

Step 3: Use the ratio to eliminate the net charge change

Step 4: Use the ratio as coefficients for the elements

Method 2: Half-Reaction Method

Step 1: Determine oxidation numbers for each atom

Step 2: Use oxidation numbers to determine what is oxidized and what is reduced.

Step 3: Write a half-reaction for reduction

Step 4: Write a half-reaction for oxidation

Step 5: Balance all elements except H and O

- if have acid redox reaction: Balance the O using H_2O , balance the H using protons
- if have base redox reaction: Balance O using OH^-

Step 6: Add up the charge on each side




Step 7: Balance the charges by adding electrons

Step 8: Multiply the half-reactions by factors that cancel out electrons

Step 9: Add the two half-reactions back together to eliminate out intermediates


Balancing Half-Equations (Oxidation or Reduction)

Given the oxidation half-equation




O.N. Fe: +2 → +3

It is clear that mass and charge balance can be achieved by adding an electron to the right:



****Loss of electron and increase in oxidation number**

Given a reduction half-equation




O.N. Cl: 0 → -1

Mass balance is obtained by writing a coefficient of 2 for Cl^{-} ; charge is then balanced by adding two electrons to the left. The balanced half-equation is



Example 7. Balance the given half-equations.




Solution:

Oxidation numbers: N: +5 → +2; O: -2 → -2; N is reduced

Atom balance: 1N on each side; no adjustment is required

Total oxidation number: N: 5(1) → N: 2(1)


Add electrons: The oxidation number for N goes from +5 to +2. It is reduced by 3. add 3 electrons to the reactant side.



Balance charge:

$$\text{reactants: } -1 + 3(-1) = -4 \quad \text{products: } 0$$

basic medium: add OH^{-} . To balance, add $4OH^{-}$ on the right.




Reactants: -1 + 3(-1) = -4 products: 4(-1) = -4

Balance H: reactants: 0 H products: 4 H




To balance, add $2H_2O$ on the left.



Reactants: 4 H products: 4 H

Check O: reactants: $3 + 2 = 5$ products: $4 + 1 = 5$

The half-equation balanced:




b. $Cl_2(g) \rightarrow ClO_3^-(aq)$ (acidic solution)

Oxidation numbers: Cl: $0 \rightarrow +5$; O: $-2 \rightarrow -2$

Cl is oxidized


Atom balance: reactant: 2 Cl product: 1 Cl

Multiply ClO_3^- by 2




"Total oxidation" number: Cl: $0(2)=0 \rightarrow Cl: 5(2)=10$

Add electrons: The oxidation number for Cl goes from 0 to 10 The oxidation number increases by 10. Add 10 electrons to the product side.



Balance charge: reactants: 0 products: $2(-1)+10(-1)=-12$


Acidic medium: add H^+ . To balance, add $12H^+$ on the right.



Reactants: 0 Products: $2(-1) + 10(-1) + 12(+1) = 0$

Balance H: reactant: 0 H products: 12 H


To balance, add $6H_2O$ on the left.



Reactants: 12 H products: 12 H

Check O: reactants: 6 products: $2(3)=6$

The half-equation is balanced:





Example 8. Balance the following redox reactions



1. $Fe_2^+(aq) + NO_3^-(aq) \rightarrow Fe_3^+(aq) + NO(g)$ (basic solution)









Split into two half-equations:





Balance the half-equations





Multiply by 3:








Split into two half-equations:



Balance the half-equations




Multiply by 2:



Example 9. Balance the reaction $MnO_4^- + I^- \rightarrow I_2 + Mn^{2+}$

Steps to balance:


1. Separate the half-reactions that undergo oxidation and reduction



2. Balance the half-reactions




** add 2 in front of the I on the left-hand side so there is an equal number of atoms on both sides




** balance oxygen atoms by adding $4H_2O$ on the right side and adding $8H^+$ on the left side to balance the H atoms

3. Balance the charges in each half-reaction




** right hand side has an overall charge of 0, adding $2e^-$ so both sides of the equation have equal charges of -2



** left hand side there are 8 hydrogen atoms with a +1 charge, MnO_4^- ion has a charge of -1, overall is +7. Right hand side Mn^{2+} is +2, H_2O is 0, overall is +2. add 5 electrons on the left side to make sure both sides have equal charges of +2

4. Multiply both sides of both reactions by the least common multiple that will allow the half-reactions to have the same number of electrons and cancel each other out.








ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna

5. Add the two half-reactions in order to obtain the overall equation by cancelling out the electrons and any H_2O and H^+ ions that exist on both sides of the equation.




Example 10. Balance the following reaction in acidic aqueous solution






Solution:



1. Split into two half-equations:





2. Balance the half-reactions



3. Balance the charges in each half-reaction






** left hand side has an overall charge of 0 and right hand side has an overall charge of +2. Adding $2e^-$ to right hand side, so both sides of the equation have equal charges of -2





** left hand side there are 8 hydrogen atoms with a +1 charge, MnO_4^- ion has a charge of -1, overall is +7. Right hand side Mn^{2+} is +2, H_2O is 0, overall is +2. add 5 electrons on the left side to make sure both sides have equal charges of +2



4. Multiply both sides of both reactions by the least common multiple that will allow the half-reactions to have the same number of electrons and cancel each other out.



5. Add the two half-reactions in order to obtain the overall equation by cancelling out the electrons and any H_2O and H^+ ions that exist on both sides of the equation.



An apparatus that is used to generate electricity from a spontaneous redox reaction or, conversely, that uses electricity to drive a nonspontaneous redox reaction is called an electrochemical cell.

There are two types of electrochemical cells:

- a. galvanic cells and
- b. electrolytic cells.

Galvanic cells are named for the Italian physicist and physician Luigi Galvani (1737–1798), who observed that dissected frog leg muscles twitched when a small electric shock was applied, demonstrating the electrical nature of nerve impulses. A galvanic (voltaic) cell uses the energy released during a spontaneous redox reaction ($\Delta G < 0$) to generate electricity. This type of electrochemical cell is often called a voltaic cell after its inventor, the Italian physicist Alessandro Volta (1745–1827). In contrast, an electrolytic cell consumes electrical energy from an external source, using it to cause a nonspontaneous redox reaction to occur ($\Delta G > 0$).



ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna




Figure 1. Electrochemical cells. galvanic cells (left) and electrolytic cell (right)

A galvanic cell transforms the energy released by a spontaneous redox reaction into electrical energy that can be used to perform work. The oxidative and reductive half-reactions usually occur in separate compartments that are connected by an external electrical circuit; in addition, a second connection that allows ions to flow between the compartments (shown here as a vertical dashed line to represent a porous barrier) is necessary to maintain electrical neutrality. The potential difference between the electrodes (voltage) causes electrons to flow from the reductant to the oxidant through the external circuit, generating an electric current.

In an electrolytic cell, an external source of electrical energy is used to generate a potential difference between the electrodes that forces electrons to flow, driving a nonspontaneous redox reaction; only a single compartment is employed in most applications. In both kinds of electrochemical cells, the anode is the electrode at which the oxidation half-reaction occurs, and the cathode is the electrode at which the reduction half-reaction occurs.

**** A galvanic (voltaic) cell converts the energy released by a spontaneous chemical reaction to electrical energy. An electrolytic cell consumes electrical energy from an external source to drive a nonspontaneous chemical reaction.*


Voltaic (Galvanic) Cells

Consider the reaction of metallic zinc with cupric ion (Cu^{2+}) to give copper metal and Zn^{2+} ion. The balanced chemical equation is as follows:




ISO 9001:2015 Certified
Level I Institutionally Accredited


Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna



Galvanic or voltaic cells involve spontaneous electrochemical reactions in which the half-reactions are separated (Figure 2) so that current can flow through an external wire.



Oxidation half-reaction:



Reduction half-reaction:





Figure 2. In this standard galvanic cell, the half-cells are separated; electrons can flow through an external wire and become available to do electrical work. As the reaction proceeds, the zinc rod dissolves, and a mass of metallic copper forms (right). These changes occur spontaneously.

The beaker on the left side of the figure is called a half-cell, and contains a 1 M solution of copper (II) nitrate $[\text{Cu}(\text{NO}_3)_2]$ with a piece of copper metal partially submerged in the solution. The copper metal is an electrode. The copper is undergoing oxidation; therefore, the copper electrode is the anode. The anode is connected to a voltmeter with a wire and the other terminal of the voltmeter is connected to a silver electrode by a wire. The silver is undergoing reduction; therefore, the silver electrode is the cathode. The half-cell on the right side of the figure consists of the silver electrode in a 1 M solution of silver nitrate $[\text{AgNO}_3]$. At this point, no current flows—that is, no significant movement of electrons through the wire occurs because the circuit is open. The circuit is closed using a salt bridge, which transmits the current with moving ions. The electrolyte in the salt bridge serves two purposes: it completes the circuit by



carrying electrical charge and maintains electrical neutrality in both solutions by allowing ions to migrate between them. The salt bridge consists of a concentrated, nonreactive, electrolyte solution such as the sodium nitrate [$NaNO_3$] solution used in this example. As electrons flow from left to right through the electrode and wire, nitrate ions (anions) pass through the porous plug on the left into the copper (II) nitrate solution. This keeps the beaker on the left electrically neutral by neutralizing the charge on the copper (II) ions that are produced in the solution as the copper metal is oxidized. At the same time, the nitrate ions are moving to the left, sodium ions (cations) move to the right, through the porous plug, and into the silver nitrate solution on the right. These added cations “replace” the silver ions that are removed from the solution as they were reduced to silver metal, keeping the beaker on the right electrically neutral. Without the salt bridge, the compartments would not remain electrically neutral and no significant current would flow. However, if the two compartments are in direct contact, a salt bridge is not necessary. The instant the circuit is completed, the voltmeter reads +0.46 V, this is called the cell potential. The cell potential is created when the two dissimilar metals are connected, and is a measure of the energy per unit charge available from the oxidation-reduction reaction. The volt is the derived SI unit for electrical potential

$$volt = V = \frac{J}{C}$$

In this equation, A is the current in amperes and C the charge in coulombs. Note that volts must be multiplied by the charge in coulombs (C) to obtain the energy in joules (J).

When the electrochemical cell is constructed in this fashion, a positive cell potential indicates a spontaneous reaction and that the electrons are flowing from the left to the right. There is a lot going on in Figure 2, so it is useful to summarize things for this system:

- Electrons flow from the anode to the cathode: left to right in the standard galvanic cell in the figure.
- The electrode in the left half-cell is the anode because oxidation occurs here. The name refers to the flow of anions in the salt bridge toward it.
- The electrode in the right half-cell is the cathode because reduction occurs here. The name refers to the flow of cations in the salt bridge toward it.
- Oxidation occurs at the anode (the left half-cell in the figure).
- Reduction occurs at the cathode (the right half-cell in the figure).




ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna


- The cell potential, +0.46 V, in this case, results from the inherent differences in the nature of the materials used to make the two half-cells.
- The salt bridge must be present to close (complete) the circuit and both an oxidation and reduction must occur for current to flow.

The cell diagram (or cell notation) is a shorthand notation to represent the redox reactions of an electrical cell. For the cell described, the cell diagram is as follows:




- A double vertical line (||) is used to separate the anode half reaction from the cathode half reaction. This represents the salt bridge.
 - The anode (where oxidation occurs) is placed on the left side of the (|).
 - The cathode (where reduction occurs) is placed on the right side of the (|).


A single vertical line (|) is used to separate different states of matter on the same side, and a comma is used to separate like states of matter on the same side. For example:




Example 11. Write the cell diagram for this reaction








Solution:



Example 12. Write cell reactions for this cell diagram



Solution:







ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines

Laguna State Polytechnic University

Province of Laguna



Determining Standard State Cell Potentials

A cell's standard state potential is the potential of the cell under standard state conditions, which is approximated with concentrations of 1 mole per liter (1 M) and pressures of 1 atmosphere at 25°C.

To calculate the standard cell potential for a reaction


- Write the oxidation and reduction half-reactions for the cell.
- Look up the reduction potential, $E^{\circ}_{\text{reduction}}$, for the reduction half-reaction in a table of reduction potentials
- Look up the reduction potential for the reverse of the oxidation half-reaction and reverse the sign to obtain the oxidation potential. For the oxidation half-reaction,

$$E^{\circ}_{\text{oxidation}} = -E^{\circ}_{\text{reduction}}$$

- Add the potentials of the half-cells to get the overall standard cell potential.

$$E^{\circ}_{\text{cell}} = E^{\circ}_{\text{reduction}} + E^{\circ}_{\text{oxidation}}$$

Example 13. Find the standard cell potential for an electrochemical cell with the following cell reaction.



Solution:

Write the half-reactions for each process.



Look up the standard potentials for the reduction half-reaction.

$$E^{\circ}_{\text{reduction } Cu^{2+}} = +0.3419$$

Look up the standard reduction potential for the reverse of the oxidation reaction and change the sign.

$$E^{\circ}_{\text{reduction } Zn^{2+}} = -0.7618 V$$






ISO 9001:2015 Certified
Level I Institutionally Accredited


Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna

$$E_{\text{oxidation Zn}^{2+}}^{\circ} = +0.7618 \text{ V}$$

Add the cell potentials together to get the overall standard cell potential.







Example 14. Calculate E_{cell}° for the following redox reaction under standard conditions:




Solution: Write the half-reactions for each process.



Look up the standard potentials for the reduction half-reaction.




Add the cell potentials together to get the overall standard cell potential.



****If the stoichiometric coefficient is multiplied by 2, the standard potential does not change**

Example 15. Calculate E_{cell}° for the following redox reaction under standard conditions:




Solution:

Write the half-reactions for each process.



ISO 9001:2015 Certified
Level I Institutionally Accredited


Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna



Look up the standard potentials for the reduction half-reaction.



Add the cell potentials together to get the overall standard cell potential.




Batteries: Chemistry and Reactions

A **battery** is an electrochemical cell or series of cells that produces an electric current. It is used in many day-to-day devices such as cellular phones, laptop computers, clocks, and cars. An ideal battery would never run down, produce an unchanging voltage, and be capable of withstanding environmental extremes of heat and humidity. Real batteries strike a balance between ideal characteristics and practical limitations.


There are two basic types of batteries: primary and secondary

Primary Batteries


Primary batteries are single-use batteries because they cannot be recharged. A common primary battery is the dry cell (Figure 3). The dry cell is a zinc-carbon battery. The zinc can serve as both a container and the negative electrode. The positive electrode is a rod made of carbon that is surrounded by a paste of manganese (IV) oxide, zinc chloride, ammonium chloride, carbon powder, and a small amount of water. The reaction at the anode can be represented as the ordinary oxidation of zinc:



The reaction at the cathode is more complicated, in part because more than one reaction occurs. The series of reactions that occurs at the cathode is approximately



The overall reaction for the zinc-carbon battery can be represented as:





with an overall cell potential which is initially about 1.5 V, but decreases as the battery is used. It is important to remember that the voltage delivered by a battery is the same regardless of the size of a battery. For this reason, D, C, A, AA, and AAA batteries all have the same voltage rating. However, larger batteries can deliver more moles of electrons. As the zinc container oxidizes, its contents eventually leak out, so this type of battery should not be left in any electrical device for extended periods.







Figure 3. A cross section of a zinc-carbon dry cell.

Alkaline batteries (Figure 4) were developed in the 1950s partly to address some of the performance issues with zinc-carbon dry cells. They are manufactured to be exact replacements for zinc-carbon dry cells. As their name suggests, these types of batteries use alkaline electrolytes, often potassium hydroxide. The reactions are

Anode:



Cathode:



An alkaline battery can deliver about three to five times the energy of a zinc-carbon dry cell of similar size. Alkaline batteries are prone to leaking potassium hydroxide, so these should also be removed from devices for long-term storage. While some alkaline batteries are rechargeable, most are not. Attempts to recharge an alkaline battery that is not rechargeable often leads to rupture of the battery and leakage of the potassium hydroxide electrolyte.



ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna







Figure 4. Alkaline batteries were designed as direct replacements for zinc-carbon (dry cell) batteries.


Secondary Batteries

Secondary batteries are rechargeable. These are the types of batteries found in devices such as smartphones, electronic tablets, and automobiles.

Nickel-cadmium, or NiCd, batteries (Figure 5) consist of a nickel-plated cathode, cadmium-plated anode, and a potassium hydroxide electrode. The positive and negative plates, which are prevented from shorting by the separator, are rolled together and put into the case. This is a “jelly-roll” design and allows the NiCd cell to deliver much more current than a similar-sized alkaline battery. The reactions are:



The voltage is about 1.2 V to 1.25 V as the battery discharges. When properly treated, a NiCd battery can be recharged about 1000 times. Cadmium is a toxic heavy metal so NiCd batteries should never be opened or put into the regular trash.








ISO 9001:2015 Certified
Level I Institutionally Accredited

Republic of the Philippines
Laguna State Polytechnic University
Province of Laguna

Figure 5. NiCd batteries use a “jelly-roll” design that significantly increases the amount of current the battery can deliver as compared to a similar-sized alkaline battery.

Lithium ion batteries (Figure 6) are among the most popular rechargeable batteries and are used in many portable electronic devices. The reactions are:



With the coefficients representing moles, x is no more than about 0.5 moles. The battery voltage is about 3.7 V. Lithium batteries are popular because they can provide a large amount current, are lighter than comparable batteries of other types, produce a nearly constant voltage as they discharge, and only slowly lose their charge when stored.







Figure 6. In a lithium ion battery, charge flows between the electrodes as the lithium ions move between the anode and cathode.

Lead Acid Battery (Figure 7) is the type of secondary battery used in your automobile. It is inexpensive and capable of producing the high current required by automobile starter motors. The reactions for a lead acid battery are:



Each cell produces 2 V, so six cells are connected in series to produce a 12-V car battery. Lead acid batteries are heavy and contain a caustic liquid electrolyte, but are often still the battery of choice because of their high current density. Since these batteries contain a significant amount of lead, they must always be disposed of properly.




Figure 7. The lead acid battery in your automobile consists of six cells connected in series to give 12 V. Their low cost and high current output make these excellent candidates for providing power for automobile starter motors.

Fuel Cells

A **fuel cell** is a device that converts chemical energy into electrical energy. Fuel cells are similar to batteries but require a continuous source of fuel, often hydrogen. They will continue to produce electricity as long as fuel is available. Hydrogen fuel cells have been used to supply power for satellites, space capsules, automobiles, boats, and submarines (Figure 8).





Figure 8. In this hydrogen fuel-cell schematic, oxygen from the air reacts with hydrogen, producing water and electricity.

In a hydrogen fuel cell, the reactions are:



The voltage is about 0.9 V. The efficiency of fuel cells is typically about 40% to 60%, which is higher than the typical internal combustion engine (25% to 35%) and, in the case of the hydrogen fuel cell, produces only water as exhaust.



	Currently, fuel cells are rather expensive and contain features that cause them to fail after a relatively short time.
	Engaging Activities I. Determine the oxidation number of the elements in each of the following compounds; a. H_2CO_3 b. N_2 c. $Zn(OH)_4^{2-}$ II. Identify the species being oxidized and reduced in each of the following reactions: a. $Cr^{+} + Sn^{4+} \rightarrow Cr^{3+} + Sn^{2+}$ b. $3Hg^{2+} + 2Fe_{(s)} \rightarrow 3Hg_2 + 2Fe^{3+}$




Performance Tasks

III. Balance the following redox reactions.


- $VO_4^{3-} + Fe^{2+} \rightarrow VO^{2+} + Fe^{3+}$ in acidic solution
- $Fe(OH)_3 + OCl^{-} \rightarrow FeO_4^{2-} + Cl^{-}$ in acidic solution

IV. Evaluate the following.

- Calculate the standard cell potential, E_{cell}° for a silver-copper galvanic cell given the following reaction:



- Regarding the following reaction:



- List the species being oxidized:_____ List the species being reduced:_____
- Calculate E_{cell}° for this cell.



Learning Resources

- Brown, Holme, Peterson, Sack and Gabler. Chemistry for Engineering Students Philippine Edition. Cengage Learning. 2018
- Masterton, Hurley, Petersen, Sack and Gabler. Principles and Reaction: Chemistry for Engineering Students Philippine Edition. Cengage Learning. 2018
- Brown and Holme. Chemistry for Engineering Students 2nd Edition. Cengage Learning. 2012
- Brown, LeMay, Bursten. Chemistry the Central Science. Pearson Education Limited. 2015
- Electrochemistry Basics. (2020, August 16). Retrieved from [https://chem.libretexts.org/Bookshelves/Analytical_Chemistry/Supplemental_Modules_\(Analytical_Chemistry\)/Electrochemistry/Basics_of_Electrochemistry](https://chem.libretexts.org/Bookshelves/Analytical_Chemistry/Supplemental_Modules_(Analytical_Chemistry)/Electrochemistry/Basics_of_Electrochemistry)
- Electrochemistry - Cells and Batteries. (2019, August 22). Retrieved from [https://chem.libretexts.org/Courses/Sacramento_City_College/SCC%3A_CHEM_330 - Adventures in Chemistry \(Alviar-Agnew\)/08%3A_Oxidation_and_Reduction/8.03%3A_Electrochemistry - Cells and Batteries](https://chem.libretexts.org/Courses/Sacramento_City_College/SCC%3A_CHEM_330_-_Adventures_in_Chemistry_(Alviar-Agnew)/08%3A_Oxidation_and_Reduction/8.03%3A_Electrochemistry_-_Cells_and_Batteries)
- Galvanic Cells. (n.d.). Retrieved October 31, 2020 from <https://opentextbc.ca/chemistry/chapter/17-2-galvanic-cells/>
- Batteries and Fuel Cells. (n.d.). Retrieved October 31, 2020 from <https://opentextbc.ca/chemistry/chapter/17-5-batteries-and-fuel-cells/>
- Lead Storage Batteries. (n.d.). Retrieved October 31, 2020 from <https://courses.lumenlearning.com/introchem/chapter/lead-storage-battery/>
- Batteries: Using Chemistry to Generate Electricity. (2019, July 2). Retrieved from https://chem.libretexts.org/Courses/can/intro/16%3A_Oxidation_and_Reduction/16.6%3A_Batteries%3A_Using_Chemistry_to_Generate_Electricity
- Battery Basics. (n.d.). Retrieved October 31, 2020 from <https://www.pveducation.org/pvcdrom/batteries/battery-basics>



Elaine L. Ching
Instructor

This module is for educational purpose only. Under section Sec. 185 of RA 8293, which states, "The fair use of a copyrighted work for criticism, comment, news reporting, teaching including multiple copies for classroom use, scholarship, research, and similar purposes is not an infringement of copyright. The unauthorized reproduction, use, and dissemination of this module without joint consent of the authors is strictly prohibited and shall be prosecuted to the full extent of the law, including appropriate administrative sanctions, civil, and criminal"