

Two Maps of Scientific Reasoning

A worked answer to the LHC black-hole question

Core submission (Part I)

Synthetix Institute

Abstract

Scientific papers contain a documentary map of authors, claims and citations. They also contain a second map: the sequence of physical operations by which one measured or calculated quantity produces another. A conventional deep-research workflow reconstructs the first map and summarizes it in prose. This submission reconstructs both. In the LHC case, the documentary map records arguments for and against microscopic black-hole safety. The equation map asks a harder question: can any single set of assumptions carry a collision through production, survival, capture, sustained growth, rapid growth and compatibility with long-lived compact stars? The reconstructed literature contains no such path. Standard microscopic black holes evaporate; stable alternatives meet stopping, growth-time and astronomical constraints. Every step below links to a source equation, its variables, its place in the paper and the graph edge that uses it.

Case result. The processed literature supports the conclusion that the LHC cannot produce a dangerous black hole within the quantitative model classes it studies. Two independent physical branches reach that result: rapid evaporation for ordinary microscopic black holes, and compact-star constraints for hypothetical stable ones.

The preliminary review asked for “concrete receipts” and “intermediate outputs available and interrogable at each stage.” This guide is organized around one receipt chain. The full derivation, twelve-equation benchmark and vector graphs are in the accompanying technical article.

Five-minute path for judges. Read Fig. 1; inspect the receipt on pp. 2–3; compare the two maps in Figs. 2 and 3; then run `bash scripts/run_judge_demo.sh`. The dependency-free command propagates the Hawking counterfactual through the committed graph receipts.

1 The question becomes six testable conditions


Between a particle collision and a catastrophe lie six required events. The collision must produce a microscopic black hole; the object must survive; matter must stop it; mass intake must exceed mass loss; the positive rate must persist long enough to matter; and the same assumptions must agree with the continued existence of white dwarfs and neutron stars exposed to natural high-energy collisions.

Writing the feared outcome as

$$\mathcal{D} = P \wedge S \wedge C \wedge G \wedge T \wedge A \tag{1}$$

makes the logic explicit. $\mathcal{D}$ denotes a dangerous macroscopic outcome. P , S , C , G , T and A denote production, survival, capture, positive growth, growth within the available time and astronomical compatibility. The symbol $\wedge$ means that every condition is required. A failure at one gate ends that branch.

What would have to happen before a microscopic black hole became dangerous?



The dangerous scenario needs six consecutive yes answers. The standard branch ends at evaporation; the stable branch conflicts with the observed survival of compact stars.

Figure 1: The physical argument in one view. The upper branch ends when Hawking radiation removes mass faster than matter can be captured. The lower branch grants stability and tests the more conservative scenario against stopping, accretion, growth time and compact-star survival.

This representation changes the meaning of a “crux.” A verbal debate may treat Hawking evaporation as the crux because one side questions it. Equation (1) shows that suppressing evaporation changes one gate. It opens the stable-object branch while leaving momentum loss, mass balance, integration over time and astronomical observation to be tested.

2 One receipt from source to conclusion

The receipt chain below shows what the software retained at each stage. The identifiers refer to nodes in the **graph JSON**. Each node stores the source paper, equation number in source order, character offsets, formula, neighboring text, quantities, physical setting and graph links.

1. Production is conditional

Node E00292, equation 283 in arXiv:0806.3381, calculates a production cross-section:

$$\sigma_{\text{BH}}(M > M_{\text{min}}) = \sum_{ij} \int_{\tau_{\text{min}}}^1 d\tau \int_{\tau}^1 \frac{dx}{x} f_i(x) f_j(\tau/x) \hat{\sigma}(\sqrt{\hat{s}}). \quad (2)$$

σ_{BH} is the formation probability expressed as a cross-section; M_{min} is the model’s minimum black-hole mass; f_i and f_j are quark and gluon momentum distributions; x is a momentum fraction; and $\hat{s}$ is the squared energy of the constituent collision. The equation yields a number only after a low gravity scale, a threshold and a horizon-formation rule are supplied. The intermediate output is therefore *conditional production*, not danger.

2. The ordinary branch loses mass

Node E00455, equation 52 in arXiv:0901.2948, contains

$$\left. \frac{dM}{dt} \right|_{\text{evap}} < 0, \quad (3)$$

where M is black-hole mass and t is time. The source gives the full model-dependent coefficient; its sign is the relevant output. The mechanism graph assigns this equation to survival. A negative rate closes the ordinary branch before stopping or accretion begins.

3. Granting stability creates a new calculation

Nodes E00124 and E00149 in arXiv:0806.3381 couple momentum loss to mass gain:

$$\frac{dp}{d\ell} = -(c_{ac} - c_{ac,M})b^2\pi\rho R^2, \quad \frac{dM}{d\ell} = c_{ac,M}b^2\pi\rho\frac{R^2}{v}. \quad (4)$$

p is momentum, ℓ is path length, ρ is material density, R is the capture radius, v is speed, b is a dimensionless impact-parameter factor, and the two c coefficients describe how collision energy and momentum are captured. These equations answer two separate questions: whether matter retains the object and how rapidly its mass changes after retention. The source-local graph connects the two calculations because they share the same evolving M , p and material conditions.

4. A rate becomes a consequence only after integration

A positive mass rate at one instant says little about long-term danger. The required time is

$$t_{\text{grow}} = \int_{M_0}^{M_*} \frac{dM}{\dot{M}_{\text{net}}(M)}, \quad (5)$$

where M_0 and M_* are the initial and final masses and $\dot{M}_{\text{net}}$ is accretion minus loss. Node E00179 evaluates this calculation for compact-star conditions. Node E00202 gives a neutron-star consequence for one warped model,

$$t_{\text{NS,w}} \sim 20 \text{ yr}. \quad (6)$$

$t_{\text{NS,w}}$ is the predicted neutron-star growth time. Neutron stars survive vastly longer than twenty years. A model that predicts fast destructive growth therefore conflicts with the objects already exposed to natural high-energy collisions.

The load-bearing inference. Challenging Hawking evaporation moves the calculation from Eq. (3) to the stable branch in Eqs. (4)–(6). Catastrophe still requires capture and rapid growth alongside the observed survival of old compact stars. No recovered equation path satisfies both demands.

3 The same papers produce two different graphs

Hyperion’s operational archive contains 2.5 million arXiv papers. The public LHC experiment used a fixed 500,000-document screening slice and retained 492 case-related records. Complete sources for six primary safety papers were then added. After deduplication, the reconstruction contains 496 papers, 708 extracted claims and 369 citation links. The archive number describes the available scientific corpus; the smaller number defines the reproducible case experiment reported here.

Collapsed provenance structure




Figure 2: The documentary map. It records source identity, citation and three claim families. This view establishes attribution, incentives and historical dependence. The physical question requires an additional step because risk and safety statements coexist while their assumptions remain inside the papers.

The second graph starts from 729 equation windows. 728 received operational fingerprints and 464 contained enough information for a physical role. The graph retains 314 equation-to-equation transitions inside papers and 190 structural matches between papers. Strict formula tests identify 36 equations that instantiate at least one of the six conditions.

How the retained equations assemble into the six-condition argument




Figure 3: The mechanism map. Each node is a source equation assigned to a necessary condition. Solid edges preserve source order; dashed edges mark calculations with a compatible mathematical role in another paper. The map shows which quantities can be composed, which assumptions change between papers and where observation rejects a branch.

The joined public graph has 1,604 nodes and 2,556 typed edges. A reader can move from a claim to its paper, from that paper to the relevant equation, from the equation to the quantity it calculates and from the quantity to the condition it tests. The machine-readable form is `public_knowledge_graph.json`.

Table 1: What each graph can establish.

Question	Documentary graph	Mechanism graph
Who made the claim?	Author, paper and date	Source retained on every equation node
Which sources depend on one another?	Citation and response links	Reused equations and cross-paper transfers
What happens if evaporation is suppressed?	A safety argument loses one support	The stable branch activates stopping, growth and astronomical tests
Which evidence carries the conclusion?	Influential claims and sources	Negative mass rate; integrated growth time; compact-star survival
Where does new evidence enter?	Add or revise a claim	Replace the equation at the affected physical condition

How the maps differ, and how they join

The documentary graph is a ledger. Its basic record is “paper A makes claim B and cites paper C.” It can establish priority, attribution, citation ancestry and whether several statements descend from the same source. The mechanism graph is a circuit diagram. Its basic record is “equation A changes quantity B under conditions C, producing an output used by equation D.” It can establish whether a proposed physical sequence is mathematically connected.

The two graphs meet at an exact source location. Every equation node carries its paper identifier and character offsets; every claim node carries its source paper. This shared key creates the path

$$\text{author} \rightarrow \text{paper} \rightarrow \text{claim}, \quad \text{paper} \rightarrow \text{equation} \rightarrow \text{quantity} \rightarrow \text{condition} \rightarrow \text{answer}. \quad (7)$$

The first path answers who asserted a proposition and how it travelled. The second tests what must occur in nature for the proposition to hold. A paper can contribute to both paths, but citation traffic cannot substitute for a missing mass balance, stopping law or observation.

The Hawking counterfactual makes the difference concrete. Removing evaporation deletes one support for the safety conclusion in the documentary graph. In the mechanism graph it changes the survival condition from failed to provisionally open, then activates four downstream calculations: momentum loss, net mass growth, integrated growth time and compact-star compatibility. The observed survival of compact stars closes the dangerous branch under the fast-growth assumptions. The conclusion survives through a different physical route, a fact that a claim tally cannot derive.

How the literature becomes a physical answer




Figure 4: The join between the two maps. Brown links retain authorship, papers, citations and claims. Green links begin at source equations and follow calculations into the six physical conditions. Paper identity and source position connect the layers, so every physical conclusion remains attributable without reducing the calculation to a statement count.

4 Method: from 2.5 million papers to a revisable answer

The pipeline separates broad discovery from strict physical testing. This prevents a paper from entering the final derivation merely because it contains the words “black hole,” “LHC” or “safety.” Each stage produces a file that can be inspected before the next stage runs.

Table 2: Seven stages of the public case reconstruction.

Stage	Operation	Inspectable output
Archive	Store 2.5 million arXiv papers as source text with document identity.	Global source archive
Case screen	Search a fixed 500,000-document slice broadly, then retain 492 records for the LHC question.	Source manifest and selection scores
Source completion	Add full L ^A T _E X sources for six primary safety papers and deduplicate by paper identifier.	496 complete case papers
Documentary extraction	Record authors, citations and scoped claims while preserving source positions.	Provenance graph
Equation extraction	Retain complete equation windows, neighboring definitions and source order; describe quantities, operation and physical setting.	729 equation windows and 728 typed records
Mechanism assembly	Connect source-local calculations and compatible cross-paper transfers; require formula-level evidence for each of the six conditions.	Equation graph and constructor receipts
Stress test and validation	Remove one condition, propagate the changed branch, recover a prespecified twelve-equation benchmark and compare with the held-out CERN report.	Counterfactual trace, benchmark and external validation

Three controls carry most of the evidential burden. First, a phrase match proposes a source but cannot fill a physical condition; the formula must contain the required quantities and operation. Second, a cross-paper link records a candidate transfer until units, parameter ranges, regime and boundary conditions agree. Third, the constructor composes adjacent steps only when an output from one equation supplies an input required by the next. Sparse attention then ranks rare, informative transitions while reducing the influence of generic high-degree nodes. The surviving paths determine the answer; the prose renders that structure for the reader.

5 A scientific answer is an executable dependency

The case contains two independent routes to safety. In the semiclassical route, Eq. (3) removes mass. In the stable-object route, the safety calculation grants survival and asks what follows. Fast growth then predicts compact-star destruction on times shorter than observed stellar ages. Slow growth removes the terrestrial catastrophe. The alternatives converge on the same answer for different physical reasons.

This is the central epistemic gain over claim ranking. A challenge paper can carry high rhetorical attention and still leave the conclusion unchanged because another physical route remains. Conversely, a low-citation equation can be load-bearing when it converts a rate into the timescale that decides the case.

Equation transfer from compact stars and matter to the collider case




Figure 5: How the astronomical test is transferred. The equation’s physical role is preserved while density, speed, size, exposure and initial conditions are replaced. Transfer becomes evidence after units, parameter ranges and boundary conditions match.

The graph assigns each uncertainty to the condition it changes. Microscopic production depends on speculative low-scale gravity. Evaporation depends on the semiclassical regime. Stable-object bounds depend on capture and accretion models. New evidence therefore changes a local part of the answer:

New result	Gate changed	Recalculation
Different horizon threshold	Production	Recompute σ_{BH} and initial mass distribution
Modified evaporation law	Survival	Recompute $\dot{M}_{\text{evap}}$; activate the stable branch if needed
New capture cross-section	Stopping	Recompute momentum loss and retained population
New accretion regime	Growth and time	Integrate the revised $\dot{M}_{\text{net}}$
New compact-star observation	Astronomical test	Re-evaluate the exposed population and exclusion range

6 A reproducible comparison with deep research

An off-the-shelf research model can often produce the correct final answer and cite CERN. The useful comparison is harder: can it expose the physical dependency that keeps the answer stable when one argument is removed? The repository supplies a fixed prompt in `data/judge_comparison_prompt.txt`. A response is checked for seven concrete outputs: the six necessary conditions, one source equation per active condition, variable definitions, source positions, within-paper links, cross-regime parameter changes and the exact gate changed by new evidence.

The test question is simple: *Assume Hawking evaporation is absent. What follows?* A discourse summary reports that a standard safety argument has weakened. The mechanism graph opens the stable branch and returns the calculations still required: stopping, net growth, growth time and the compact-star test. This difference is the submission’s main contribution to epistemic uplift.

The current build provides three checks against unsupported fluency:

1. **Prespecified extraction test.** 12 of 12 named primary-source equations were recovered. The list was fixed before the final mechanism graph was assembled.
2. **Graph receipt.** Every displayed inference points to a source equation and physical condition. The receipt bundle is `paper/lhc_judges_guide_receipts.json`.
3. **Held-out reconstruction.** CERN-2003-001 was excluded from the equation corpus and benchmark. Its human-written safety analysis follows the same branch order and reaches the same conclusion.

7 Evidence for each judging criterion

The table below uses the competition’s seven dimensions verbatim. Each row names a measured result, the committed artifact that carries it and a short test a judge can perform. This makes the scoring claims independent of the prose quality of the submission.

Table 3: Criterion-by-criterion evidence and direct checks.

Criterion	Demonstrated result and artifact	Direct check
Epistemic uplift	The graph identifies the equations that carry the answer and propagates the “no Hawking evaporation” challenge through the remaining branch. Artifact: receipt bundle and judge demo.	Run <code>bash scripts/run_judge_demo.sh</code> ; compare its downstream conditions with a prose research answer to the fixed prompt.
Generalizability	Source, equation, quantity, operation, condition and observation are case-independent record types; the six-condition chain is the replaceable LHC component. Artifact: evidence contracts and constructor.	Replace the condition file while keeping the source and receipt schema; inspect which fields require domain adaptation.
Compounding and shareability	The joined JSON preserves authorship, citations, formulas, source offsets, branch assignments and typed edges. A new source changes local nodes and paths. Artifact: <code>public_knowledge_graph.json</code> .	Add or revise one receipt and rebuild; inspect the graph diff instead of regenerating the narrative by hand.
Scalability	The operational archive contains 2.5 million papers. The public case test screens 500,000 documents programmatically, then applies stricter equation tests to the retained set. Artifact: full-run script and manifests.	Inspect the source counts at each stage; rerun the fixed slice or increase the screening limit.
Methodological transparency	Formula contracts, source positions, graph edges, twelve prespecified equation tests, pseudocode and regression tests are committed. Artifact: benchmark, tests and build scripts.	Open node E00202, locate its source equation, then run <code>python3 -B -m pytest -q</code> .
Adversarial robustness	A challenge changes the condition it addresses; transfer still requires matching quantities, units, regime and boundary conditions. Artifact: counterfactual trace and strict contracts.	Challenge evaporation, stopping or accretion separately and observe which branch edges change and which remain.
Insight contribution	The documentary map and mechanism map expose different dependencies. Their source-level join shows why agreement among claims and closure of a physical branch are distinct results. Artifact: Figs. 2–4.	Compare the node and edge types, then trace one claim through its paper to the equation and observation that decide it.

The current empirical validation covers one equation-rich case. Generalizability is implemented at the schema and software levels; cross-case trials will test how well the same records represent measurements, causal models and experimental protocols outside mathematical physics.

8 How to inspect and rebuild it

The judge path is self-contained: committed receipts reproduce every number and figure in this guide. The full corpus pipeline adds large-scale source selection and equation featurization.

```
git clone https://github.com/synthetic-institute/lhc-mechanism-audit.git
cd lhc-mechanism-audit
bash scripts/run_judge_demo.sh
```

The demo checks that every selected equation ID exists, that all prespecified benchmark equations were recovered and that the numerical claims in this guide match the JSON artifacts. `bash scripts/build_judges_guide.sh` also compiles the PDF when LaTeX is available; `python3 -B -m pytest -q` runs the regression suite.

Algorithm in twelve lines

```
for each paper:
    retain source identity, claims, citations, equations and source positions
    describe each equation by quantities, operation and physical setting
    connect equations that occur in source order
connect cross-paper equations only when operation and quantities are compatible
for each case condition:
    require a formula-level contract; reject phrase-only matches
    assign direct receipts and parameter-transfer candidates
test whether outputs from one condition satisfy inputs to the next
rank uncommon informative transitions, with a penalty for generic hubs
report the complete branches, failed gates and source equations
compare the result with a held-out human analysis
```

9 What travels to another case

The reusable object is a conditional chain: an initial state, operations that change it, constraints on those operations, measurable outputs and observations that can reject a branch. The LHC constructor uses mass, momentum, rates and stellar survival. A biomedical origin question would use genomic changes, host adaptation, sampling processes, experimental protocols and dated observations. A nutrition question would separate exposure, biological response, confounding structure, biomarkers and clinical outcomes. In each case, provenance remains attached to every receipt while the mechanism layer tests whether the steps can be composed.

This yields a compact design rule for living knowledge bases: store who said it and store what must happen for it to be true. The first graph supports accountability. The second supports scientific reasoning, local revision and transfer across fields.

Supporting material. In the combined submission packet, the complete fourteen-page article follows this guide as Part II. It gives the full physical derivation, larger vector graphs, all displayed variable definitions, the constructor implementation and the held-out comparison with CERN-2003-001.

References

1. S. B. Giddings and M. L. Mangano, “Astrophysical implications of hypothetical stable TeV-scale black holes,” *Phys. Rev. D* 78, 035009 (2008), arXiv:0806.3381.
2. J. Ellis et al., “Review of the Safety of LHC Collisions,” *J. Phys. G* 35, 115004 (2008), arXiv:0806.3414.
3. B. Koch, M. Bleicher and H. Stoecker, “Exclusion of black hole disaster scenarios at the LHC,” *Phys. Lett. B* 672, 71–76 (2009), arXiv:0807.3349.
4. R. Casadio, S. Fabi and B. Harms, “Possibility of catastrophic black hole growth in the warped brane-world scenario at the LHC,” arXiv:0901.2948.
5. J.-P. Blaizot et al., *Study of potentially dangerous events during heavy-ion collisions at the LHC*, CERN-2003-001 (2003), CERN document server.

Could the Large Hadron Collider make a dangerous black hole?

Following the physics from collision to consequence

Synthetix Institute

Abstract

Before the Large Hadron Collider began operation, speculative theories of low-scale gravity raised a vivid question: could a particle collision create a microscopic black hole that survives, remains inside matter and grows? A dangerous outcome requires six events in sequence. The collision must produce the object; it must survive evaporation; matter must stop it; accretion must exceed every loss process; growth must be fast enough; and the same physics must remain compatible with old white dwarfs and neutron stars struck by cosmic rays whose constituent collisions reach the LHC energy regime. We reconstructed this chain from a broad search of the arXiv scientific-preprint archive and six complete primary papers. The software recovered all 12 equations chosen in advance as primary-source tests and linked equations through the physical quantities they calculate. The standard branch ends with rapid Hawking evaporation. Stable and metastable branches are constrained by momentum loss, accretion times and the observed survival of compact stars. The recovered literature contains no self-consistent route through all six conditions. A CERN safety report that was excluded from the reconstruction reaches the same physical branch structure and conclusion.

1 Why the question deserved a calculation


The Large Hadron Collider (LHC) brings two beams of protons into collision. Its present proton collision energy is 13.6 TeV, concentrated into a region far smaller than an atomic nucleus, and the ATLAS and CMS interaction points can record about 1.5 billion collisions per second [1]. A teraelectronvolt (TeV) is one trillion electronvolts; 13.6 TeV is only about two millionths of a joule in one collision. The unusual feature is the concentration of that energy, not its total amount. None of this resembles the collapse of a star. A stellar black hole contains at least several solar masses. Even if the energy of one LHC collision became mass, it would amount to roughly 2×10^{-23} kg.

The idea entered physics through models in which gravity becomes strong at much lower energy than in ordinary four-dimensional general relativity [5, 6, 7]. Extra spatial dimensions can lower the energy at which quantum-gravity effects are expected. A sufficiently energetic collision could then cross a model-dependent threshold for microscopic horizon formation. This possibility made production a legitimate theoretical calculation. Danger was a separate and much harder proposition.

The answer. Within every model class in the processed literature that makes black-hole production quantitative, the dangerous chain fails. Ordinary microscopic black holes evaporate. A hypothetical stable survivor must be stopped and must accrete rapidly; those same assumptions predict effects in compact stars exposed to cosmic-ray collisions, yet old white dwarfs and neutron stars remain. The literature therefore supplies no physical mechanism for an LHC black-hole catastrophe.

Figure 1 shows the entire argument. Every box is necessary. A “no” at any gate ends the dangerous branch.

What would have to happen before a microscopic black hole became dangerous?



The dangerous scenario needs six consecutive yes answers. The standard branch ends at evaporation; the stable branch conflicts with the observed survival of compact stars.

Figure 1: The six gates between a proton collision and a dangerous macroscopic outcome. The ordinary semiclassical branch ends at Gate 2 through Hawking evaporation. The deliberately conservative stable branch continues through stopping and growth calculations, then meets the compact-star observation at Gate 6.

2 A catastrophe requires a complete chain

The feared outcome can be written as a chain in which every condition must be true:

$$\mathcal{D} = P \wedge S \wedge C \wedge G \wedge T \wedge A, \quad (1)$$

where P denotes production, S survival, C capture or retention, G positive mass growth, T growth on a relevant timescale, and A compatibility with astronomical observations. The symbol $\wedge$ means “and”: if one term fails, the dangerous outcome $\mathcal{D}$ fails. Equation (1) is a logical map rather than a new law of physics. It prevents one dramatic statement, such as “a black hole might form”, from being mistaken for the complete hazard.

The six terms are coupled. A production model fixes the initial mass and momentum. Those values enter the evaporation and stopping calculations. Stopping determines the material environment in which accretion is evaluated. The mass-growth law must then be integrated over time. Finally, the same production, capture and growth laws must be applied to cosmic-ray collisions with dense astronomical bodies. The conclusion follows from the whole path.

3 Following the equations

3.1 Gate 1: can the collision make a microscopic black hole?

A proton is composed of quarks and gluons, collectively called partons. Only a fraction of the beam energy enters any one parton-parton collision. The production calculation therefore adds the contributions from every possible pair of partons whose collision energy exceeds the assumed black-hole threshold [8]:

$$\sigma_{\text{BH}}(M > M_{\text{min}}) = \sum_{ij} \int_{\tau_{\text{min}}}^1 d\tau \int_{\tau}^1 \frac{dx}{x} f_i(x) f_j(\tau/x) \hat{\sigma}(\sqrt{\hat{s}}). \quad (2)$$

Here σ_{BH} is the total formation cross-section, which measures production probability. M is black-hole mass, M_{min} is the minimum mass accepted by the model, f_i and f_j describe the momentum carried by

partons of types i and j , x is a parton momentum fraction, and $\hat{\sigma}$ is the formation cross-section for one parton collision. The parton-level energy is $\hat{s} = x_1 x_2 s$, where s is the squared proton-proton collision energy. The lower integration limit is

$$\tau = x_1 x_2 > \tau_{\min} = \frac{M_{\min}^2}{y^2 s}, \quad (3)$$

where y is the fraction of collision energy trapped behind the proposed horizon.

In plain language. Equations (2) and (3) count quark and gluon collisions energetic enough to cross a hypothetical gravity threshold. They predict production only after a low gravity scale, a minimum mass and a horizon-formation rule have been supplied.

With the conventional Planck scale, the threshold lies vastly beyond the LHC. Microscopic production at LHC energies belongs to speculative low-scale-gravity models. The remaining five gates test what those models imply if production is granted.

3.2 Gate 2: does the object survive?

Hawking radiation makes a small semiclassical black hole lose mass. A retained calculation writes the canonical decay rate as [10]

$$\frac{dM}{dt} \approx -c \frac{d+1}{4R_H^2} < 0, \quad (4)$$

where M is mass, t is time, c is the speed of light, d is the number of extra spatial dimensions in the model, and R_H is the horizon radius. The negative sign is the essential result: the black hole shrinks. For a representative mass of ten times the model's fundamental gravity scale, the cited calculation estimates a decay rate at least thirty orders of magnitude larger than the early accretion rate [10]. CERN summarizes the expected lifetime as about 10^{-27} s [2].

In plain language. The ordinary branch disappears before it can cross a detector, stop in the Earth or begin sustained accretion.

Safety analyses also examine a stronger hypothetical case in which Hawking evaporation is suppressed or the remnant is stable [8, 11, 12]. This assumption keeps the branch alive long enough to test Gates 3 through 6.

3.3 Gate 3: can matter stop it?

A stable microscopic black hole produced in a collider would initially move rapidly. It can threaten its surroundings only after interactions remove enough momentum to keep it inside matter. Giddings and Mangano calculate an accretion contribution to the momentum loss per unit path length [8]:

$$\left(\frac{dp}{d\ell}\right)_{\text{ac}} = -(c_{\text{ac}} - c_{\text{ac,M}}) b_{\min}^2 \pi \rho \frac{E^2}{M^2} R^2 (\sqrt{s_{\text{int}}}). \quad (5)$$

Here p , E and M are the black hole's momentum, energy and mass; ℓ is distance travelled; ρ is the energy density of the material; R is the gravitational capture radius; s_{int} is the squared centre-of-mass energy for the black-hole/constituent encounter; $b_{\min} R$ is the limiting impact parameter for capture; and c_{ac} and $c_{\text{ac,M}}$ describe the fractions of momentum and energy transferred in an encounter.

In plain language. The equation accumulates many tiny interactions along a path. If the integrated momentum loss is smaller than the momentum needed to escape, the object leaves the Earth. If it is retained, its final speed and location set the initial conditions for accretion.

This gate is especially important for the astronomical comparison. Black holes hypothetically produced by cosmic rays can be much more strongly boosted than products at a symmetric collider. A useful natural test therefore requires an object dense enough to stop some of them. White dwarfs and neutron stars provide that test.

3.4 Gate 4: does it gain mass faster than it loses mass?

For motion through matter, a primary source gives the capture law [8]

$$\left. \frac{dM}{dt} \right|_{\text{acc}} = \pi \rho v r_c^2(M), \quad (6)$$

where v is speed through the material and $r_c(M)$ is the mass-dependent capture radius. A second source makes the competition explicit [12]:

$$\frac{dM}{dt} = \left. \frac{dM}{dt} \right|_{\text{evap}} + \left. \frac{dM}{dt} \right|_{\text{acc}}. \quad (7)$$

In plain language. The area πr_c^2 sets how much material is intercepted. Multiplying it by density ρ and speed v gives the incoming mass rate. Growth begins only when this gain remains larger than evaporation and other losses.

A positive value at one instant does not establish catastrophe. The density, velocity and capture radius change as the object slows and grows. The full trajectory leads to Gate 5.

3.5 Gate 5: can growth finish within the available time?

The time needed to grow from an initial mass M_0 to a larger mass M_* follows from the net mass rate:

$$t_{\text{grow}}(M_0 \rightarrow M_*) = \int_{M_0}^{M_*} \frac{dM}{\dot{M}_{\text{net}}(M)}. \quad (8)$$

Here $\dot{M}_{\text{net}}$ is the right-hand side of Eq. (7). The source calculations split the integral into regimes because the relevant capture radius changes with mass. For a six-dimensional white-dwarf model, one recovered system is [8]

$$\begin{aligned} t(R_B < R_D) &= \frac{8}{\lambda_6} \left(\frac{M_D}{M_0} \right)^2 \text{ yr}, \\ t(R_D < R_B < R_C) &= \frac{4 \times 10^2}{\lambda_4} \left(\frac{M_D}{M_0} \right)^2 \text{ yr}, \\ t(R_B > R_C) &= \frac{15}{\lambda_4} \left(\frac{M_D}{M_0} \right)^2 \text{ yr}. \end{aligned} \quad (9)$$

R_B is the Bondi radius at which gravity organizes the surrounding flow, R_D marks the end of the higher-dimensional regime, R_C marks the transition to four-dimensional gravity, M_D is the fundamental gravity scale, and λ_4 and λ_6 are dimension-dependent accretion coefficients. In the seven-dimensional cases considered in the same paper, the longest relevant white-dwarf stage remains below about 80 million years, while known white dwarfs live for more than a billion years [8].

In plain language. A local tendency to grow is converted into a clock. Any model that predicts rapid terrestrial growth also predicts an observable compact-star consequence on a much shorter timescale than the star’s age.

3.6 Gate 6: what do compact stars tell us?

Cosmic rays reach laboratory energies far above the LHC beam energy and have struck planets and stars for billions of years [3, 9]. A cosmic ray hits a nearly stationary target, so only part of its laboratory energy enters the centre-of-mass collision; the highest-energy events still cover and exceed the LHC-relevant range. Under a stable-black-hole hypothesis, some collisions should therefore produce the same species considered at the LHC. Dense compact stars are effective detectors because they provide enormous column density and long exposure. One recovered warped-model calculation gives

$$t_{\text{NS,w}} \sim 20 \text{ yr}, \quad (10)$$

where $t_{\text{NS,w}}$ is the predicted neutron-star growth time in that model [8]. Neutron stars survive far longer than twenty years. White-dwarf calculations provide the same form of test over longer, model-dependent intervals.

The observation is stronger than a verbal analogy. Production, momentum loss, capture radius, density and growth time are recalculated for the star. A model that predicts production, capture and destructive growth must reproduce the continued existence of the objects that have already hosted the natural collisions.

4 What two maps of the literature reveal

Scientific papers carry two different kinds of structure. Citations, authors and stated conclusions show how an argument travelled. Equations show how one physical quantity is converted into another. We built both maps from the same sources.

The Hyperion operational archive contains 2.5 million arXiv papers. The fixed public LHC experiment scanned 500,000 source documents and selected 492 records using terms connected to colliders, black holes, cosmic rays, compact stars, evaporation and accretion. Six complete primary safety papers were then added, producing 496 unique paper nodes. The broad set intentionally includes neighboring astrophysics because the stopping and growth laws are developed there. The six primary papers carry the quantitative LHC safety argument.

The provenance graph contains 708 extracted claim nodes, 327 cited-paper nodes, 369 citation links and 708 paper-to-claim links. Figure 2 reduces it to its main families. Risk claims, safety claims and astronomical statements all occur. Their counts describe the discussion; the equations decide whether the feared chain is physically closed.

Collapsed provenance structure




Figure 2: The provenance map records where claims came from and how papers cite one another. The three claim families coexist in the search set. Their presence identifies the dispute; it does not combine their physical assumptions.

Provenance graph: who wrote, cited and claimed what




Figure 3: The complete provenance graph used in this reconstruction. Each point is a named author, selected paper, cited paper or extracted claim. Outlined points identify the six complete primary papers used for the equation benchmark.

The equation graph starts from 729 equations together with their positions and nearby text. The software converted 728 into numerical descriptions of the operation being performed; 464 contained enough information to identify a physical role. It then found 366 equation-to-equation steps inside individual papers and 190 candidate matches between different papers. Each retained node records the source equation, the quantities it consumes and produces, its physical setting and its position in the paper.

Figure 4 organizes the 36 distinct equations that satisfy at least one of the six physical conditions. Circles mark equations written in collider or LHC-safety contexts. Diamonds mark equations from material or astrophysical settings that calculate the same physical quantity and therefore supply a transfer test.

How the retained equations assemble into the six-condition argument




Figure 4: The equation-level mechanism graph. Columns are the six necessary physical conditions. Solid lines preserve source order inside a paper; dashed lines join equations that perform the same mathematical operation in different papers. The graph exposes where the literature provides a derivation and where separate model assumptions must be combined.

Table 1: Equations assigned to each necessary condition. A single equation can contribute to more than one condition, so column totals are assignments rather than unique papers.

Condition	Direct equations	Transfer equations	Quantity calculated
production	3	0	formation threshold and production rate
survival	3	0	evaporation rate or lifetime
stopping	5	2	momentum loss and capture
net growth	11	10	mass gained minus mass lost
growth time	1	3	time required to reach a larger scale
astronomical test	0	1	predicted consequence for compact stars

The graph makes the result precise. Equations cover every local process, and 23 direct assignments connect primary-source equations to collider conditions. 16 assignments carry stopping, growth or astronomical calculations across physical settings. Within the papers, the equations directly connect stopping to mass growth. No single derivation carries one consistent parameter set through production, survival, stopping, growth time and the compact-star test. The literature is modular: one paper grants stability to study its consequences, another evaluates Hawking loss, and another computes capture under a particular dimensional model.

Two positive results recovered from the equations determine the physical conclusion. The standard semiclassical branch has a large negative mass rate. The stable branch predicts compact-star effects when growth is fast enough to matter on Earth. These results close the dangerous route from opposite sides.

How the literature becomes a physical answer




Figure 5: The joined graph connects source history to physical calculation. Brown links carry authorship, citation and claims. Green links run from papers through equations and calculation types to the six conditions and the final answer.

5 How an equation from a star becomes evidence about a collider

The transfer from astrophysics to collider safety preserves a physical question while changing the environment. Consider Eq. (6). The mass rate always depends on density ρ , speed v and capture radius r_c . A white dwarf and the Earth have different values of all three. The equation remains the same; its parameters are recomputed.

Figure 6 shows four transfers recovered in the case graph. The middle column states the quantity that must be preserved: momentum loss, net mass rate, integrated growth time or an astronomical survival bound. The right column states the collider-specific test. This step blocks a common error in qualitative arguments, where a formula is moved between environments without changing its boundary conditions or parameter ranges.

Equation transfer from compact stars and matter to the collider case




Figure 6: Equation transfer between physical environments. The mathematical role is held fixed while density, velocity, size, exposure and initial conditions are replaced. The transferred relation becomes evidence after dimensions, parameter ranges and boundary conditions agree.

6 How the reconstruction was made

The procedure has five steps. First, displayed equations and their nearby sentences are extracted from each LaTeX source. Second, every equation receives a numerical description, called a fingerprint, of its mathematical operation and physical quantities: threshold, rate, transport, constraint, boundary condition or ordered step. Third, equations are linked in their order of appearance inside each source. Equations that perform the same mathematical operation in different papers form candidate transfer links. Fourth, six explicit equation tests are applied. A stopping equation must contain a momentum or energy loss over distance; a growth equation must contain a mass rate; a timescale equation must integrate or evaluate that rate. Fifth, the resulting graph is searched for paths that pass the six gates while preserving compatible quantities and physical settings.

This is how the method separates a repeated phrase from a physical dependency. The phrase “stable microscopic black hole” can appear in many claims. It contributes to the mechanism only when a source supplies a lifetime assumption and then carries the resulting mass and momentum into stopping or growth equations.

From a broad literature search to a testable physical chain

Primary sources present: 6/6

Named equations recovered: 12/12




Figure 7: From broad literature to equations used in the physical chain. The last stage is a benchmark of named equations selected before the final mechanism graph was assembled.

6.1 A direct check against the primary sources

Twelve equations were named in advance across the six primary papers. They cover production, the parton threshold, Hawking loss, stopping, accretion, competing mass rates, growth stages and compact-star bounds. The extraction recovered 12 of 12 and each passed its prespecified equation test.

Table 2: Primary-source equation recovery. Review and response papers contain fewer calculation equations; the main calculation papers supply the named tests.

arXiv source	Extracted equations	Named equations recovered
0806.3381	334	6/6
0806.3414	0	0/0
0807.3349	37	3/3
0808.1415	12	1/1
0808.4087	3	0/0
0901.2948	75	2/2

6.2 Which transitions carry the argument?

The graph ranking gives greater weight to an uncommon, informative equation-to-equation step than to a connection repeated throughout the corpus. Its strongest individual step is mass intake -> momentum evolution in arXiv 0901.2948. Across the high-ranking links, three operations dominate in ordinary physical terms: an equation defines an allowed range, another propagates mass or momentum, and a constraint tests the result. This is the mathematical skeleton of the safety argument. Repetition of a conclusion does not raise this score.

Which equation-to-equation links carry the safety argument?




Figure 8: Information carried by equation-to-equation transitions. The matrix groups transitions by mathematical role; the bars show the individual source transitions that contribute most strongly. High rank identifies a useful dependency, rather than a popular claim.

7 Independent comparison with the CERN safety report

CERN-2003-001 was written by the LHC Safety Study Group before operation of the collider [4]. It was excluded from the arXiv equation corpus and from the twelve-equation benchmark. The report therefore provides an independent human reconstruction.

The agreement is specific. Both analyses begin by making microscopic production conditional on speculative low-scale gravity. Both separate the rapidly evaporating branch from a stable hypothetical branch. Both carry the stable branch through stopping and accretion. Both use cosmic-ray exposure and the survival of astronomical bodies as the final test. The later peer-reviewed LHC safety review follows the same structure [9].

Table 3: Independent comparison of the automatically reconstructed chain with CERN-2003-001.

Physical question	Automated reconstruction	CERN safety report
Can microscopic production occur?	Conditional on a low gravity scale and threshold	Treated as a speculative model branch
What happens to the ordinary branch?	Hawking mass loss ends the branch	Rapid decay ends the branch
How is a stable branch tested?	Momentum loss, capture and accretion	Stopping and accretion calculations
What supplies the long-exposure test?	Cosmic rays, white dwarfs and neutron stars	Natural cosmic-ray collisions and surviving bodies
Case conclusion	No closed dangerous mechanism	No conceivable threat

The independent match covers the order of the argument, the decisive physical quantities and the final result. It also identifies what the automated graph adds: every step can be traced to a source equation,

its variables and the regime in which it was derived.

8 Conclusion

The LHC black-hole question has a clear physical answer. Conventional gravity does not place microscopic black-hole production within LHC reach. Low-scale-gravity models can make production calculable, but their ordinary black holes evaporate almost immediately. Granting stability moves the burden to stopping, net mass growth and growth time. Models that make this branch fast enough to threaten the Earth also predict observable consequences in compact stars exposed to LHC-relevant natural collisions over vastly longer times. Those stars survive.

The broad literature contains risk statements, safety statements and many neighboring astrophysical calculations. Their physical content becomes clear when equations are assembled in causal order. Production is an entrance condition, evaporation is an early exit, stopping supplies initial conditions for accretion, integration turns a rate into a timescale, and compact stars test the completed stable branch. No equation-supported path in the processed literature reaches a dangerous macroscopic outcome.

The same representation can be updated locally. A new evaporation law would alter Gate 2. A new capture calculation would alter Gate 3. A newly observed compact-star population would strengthen or narrow Gate 6. The answer remains attached to the physical quantities that determine it.

References

- [1] CERN, “The Large Hadron Collider: frequently asked questions,” <https://home.cern/resources/faqs/large-hadron-collider-dangerous>.
- [2] CERN, “Extra dimensions, gravitons, and tiny black holes,” <https://home.cern/science/physics/extra-dimensions-gravitons-and-tiny-black-holes/>.
- [3] CERN, “Cosmic rays: particles from outer space,” <https://home.cern/science/physics/cosmic-rays-particles-outer-space/>.
- [4] J.-P. Blaizot et al., “Study of potentially dangerous events during heavy-ion collisions at the LHC: Report of the LHC Safety Study Group,” CERN-2003-001 (2003), <https://cds.cern.ch/record/613175/files/CERN-2003-001.pdf>.
- [5] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, “The hierarchy problem and new dimensions at a millimeter,” *Phys. Lett. B* 429, 263–272 (1998), <https://arxiv.org/abs/hep-ph/9803315>.
- [6] S. B. Giddings and S. Thomas, “High energy colliders as black hole factories: the end of short distance physics,” *Phys. Rev. D* 65, 056010 (2002), <https://arxiv.org/abs/hep-ph/0106219>.
- [7] S. Dimopoulos and G. Landsberg, “Black holes at the Large Hadron Collider,” *Phys. Rev. Lett.* 87, 161602 (2001), <https://arxiv.org/abs/hep-ph/0106295>.
- [8] S. B. Giddings and M. L. Mangano, “Astrophysical implications of hypothetical stable TeV-scale black holes,” *Phys. Rev. D* 78, 035009 (2008), <https://arxiv.org/abs/0806.3381>.
- [9] J. Ellis et al., “Review of the Safety of LHC Collisions,” *J. Phys. G* 35, 115004 (2008), <https://arxiv.org/abs/0806.3414>.

- [10] B. Koch, M. Bleicher and H. Stoecker, “Exclusion of black hole disaster scenarios at the LHC,” *Phys. Lett. B* 672, 71–76 (2009), <https://arxiv.org/abs/0807.3349>.
- [11] R. Plaga, “On the potential catastrophic risk from metastable quantum-black holes produced at particle colliders,” <https://arxiv.org/abs/0808.1415>.
- [12] R. Casadio, S. Fabi and B. Harms, “Possibility of catastrophic black hole growth in the warped brane-world scenario at the LHC,” <https://arxiv.org/abs/0901.2948>.