

Quantum Theory As A Mechanism Tree

Synthetix Institute
Generated 2026-07-08 13:35 UTC

This book reorganizes quantum theory away from historical article names and toward the recurring construction detected across the MorphWiki quantum pages: context-selected Hilbert space, state carrier, generator, spectral question, probability/readout, compatibility limits, boundary realization, many-mode extension, and protocols.

Contents

How To Use The Mechanism Tree	vii
What The Tree Is	vii
DAG Before Constructor	vii
Why Hilbert Space Is Central	viii
The Compact Representation	viii
Recipe For Reading A Page	ix
What Was Found In This Run	ix
Branch Map	x
Boundary Of The Claim	xii
Preamble: Findings From The Rewrite	xiii
Core Insight	xiii
Main Finding And Why It Matters	xiii
Transition Sparse-Attention Result	xiv
Main Findings	xvi
What Changed Relative To Wikipedia	xvii
Unusual Findings And Anomalies	xviii
Interpretation Of The Anomalies	xix
Constructor-Detected Research Leads	xx
Relation To Lagrangian, Transfer, And Stability Layers	xx
Practical Consequence	xxi
 I The Mechanism Tree	 1
1 Root Construction	2
1.1 Re-Derivation Path	2
2 Sparse Attention Summary	3
3 Compact Operator Formulation	4
3.1 The Minimal Constructor	4
3.2 Interpretation Of The Symbols	4
3.3 Why This Is More Compact	5
4 Mechanism Validation Layers	6
4.1 Role-Current Layer: What Stays Stable	6
4.2 Structural Bridge Layer: Candidate Translations	6
4.3 Lagrangian Layer: Which Moves Are Easy Or Strained	6
4.4 How These Layers Change The Quantum Tree	7

5	Sparse-Attention Result	8
5.1	Minimal Constructor	8
5.2	Route Statistics	8
5.3	Constructor Regularities	9
5.3.1	Operator-to-spectrum as the readout spine	9
5.3.2	State transport as the carrier of change	9
5.3.3	Admissibility as the legal spine	9
5.3.4	Incompatibility as a selective limit	9
5.3.5	Context as the active boundary of the question	10
5.3.6	Protocol as ordered implementation	10
5.3.7	Junction pages require decomposition	10
5.3.8	Geometry enters as realization	10
5.4	Topic-Specific Readings	10
6	Anomalies And Leads	12
6.1	Structural Anomalies	12
6.2	Research Leads	14
II	Branch Chapters	15
7	Hilbert-space context: admissible carrier and basis	16
7.1	Why This Branch Exists	16
7.2	Page Map	16
7.3	Mechanism Pages	16
7.4	Mathematical formulation of quantum mechanics	17
7.5	Hilbert space	19
7.6	Transformation theory (quantum mechanics)	20
7.7	Quantum differential calculus	22
7.8	Quantum complexity theory	24
7.9	Quantum cellular automaton	26
7.10	Relativistic quantum mechanics	28
8	State carrier inside Hilbert space	31
8.1	Why This Branch Exists	31
8.2	Page Map	31
8.3	Mechanism Pages	32
8.4	Density matrix	32
8.5	Quantum superposition	34
8.6	Quantum decoherence	36
8.7	Superposition principle	38
8.8	Coherence (physics)	40
8.9	Wave function	42
8.10	Quantum state	43
8.11	Two-state quantum system	45
8.12	Qubit	47
8.13	Quantum fluctuation	49
8.14	Bose–Einstein statistics	51
8.15	Quantum number	53
8.16	Fourier transform	55
8.17	Wave–particle duality	57
8.18	Quantum statistical mechanics	59

8.19	Quantum machine	61
8.20	Quantum chemistry	63
8.21	Applications of quantum mechanics	65
8.22	Quantum machine learning	67
8.23	Standard Model	69
8.24	Spin (physics)	71
8.25	Quantum simulator	73
8.26	Werner Heisenberg	75
8.27	Schrödinger's cat	77
9	Generator: lawful change before readout	80
9.1	Why This Branch Exists	80
9.2	Page Map	80
9.3	Mechanism Pages	81
9.4	Unitary operator	81
9.5	Perturbation theory	83
9.6	Quantum dynamics	85
9.7	Path integral formulation	87
9.8	Hamiltonian mechanics	88
9.9	Path integral	90
9.10	Hamiltonian (quantum mechanics)	92
9.11	Perturbation theory (quantum mechanics)	94
9.12	Schrödinger picture	96
9.13	Creation and annihilation operators	98
9.14	Schrödinger equation	100
9.15	Quantum stochastic calculus	102
9.16	Quantum chaos	104
9.17	Symmetry in quantum mechanics	106
9.18	Quantum amplifier	108
9.19	Quantum mechanics	110
9.20	Heisenberg picture	111
9.21	Heisenberg group	113
9.22	Canonical commutation relation	115
9.23	Quantum biology	117
9.24	Quantum engineering	119
9.25	Quantum logic	121
10	Spectral question: what can be asked	124
10.1	Why This Branch Exists	124
10.2	Page Map	124
10.3	Mechanism Pages	124
10.4	Angular momentum operator	124
10.5	Observable	126
10.6	Self-adjoint operator	128
10.7	Spectral theory	130
10.8	Pauli matrices	132
10.9	Operator theory	134
10.10	Operator (physics)	136
10.11	Eigenvalues and eigenvectors	138
10.12	Von Neumann algebra	140
10.13	Electron microscope	142

11 Readout rule: how answers become probabilities	145
11.1 Why This Branch Exists	145
11.2 Page Map	145
11.3 Mechanism Pages	145
11.4 POVM	146
11.5 Wave function collapse	147
11.6 Born rule	149
11.7 Measurement in quantum mechanics	151
11.8 Quantum jump	153
11.9 Measurement problem	155
11.10 Quantum eraser experiment	157
11.11 Projection-valued measure	159
11.12 Delayed-choice quantum eraser	160
11.13 Quantum metrology	162
12 Compatibility limit: what cannot be jointly sharp	165
12.1 Why This Branch Exists	165
12.2 Page Map	165
12.3 Mechanism Pages	165
12.4 Bell's theorem	165
12.5 Quantum entanglement	167
12.6 Commutator	169
12.7 Uncertainty principle	171
12.8 Einstein–Podolsky–Rosen paradox	173
12.9 Quantum nonlocality	175
13 Boundary realization: how effects appear	177
13.1 Why This Branch Exists	177
13.2 Page Map	177
13.3 Mechanism Pages	177
13.4 Potential well	178
13.5 Particle in a box	180
13.6 Scattering	181
13.7 Wave interference	183
13.8 Quantum optics	185
13.9 Spectral line	187
13.10 S-matrix	189
13.11 Quantum tunnelling	191
13.12 Quantum imaging	193
13.13 Quantum harmonic oscillator	195
13.14 Macroscopic quantum phenomena	197
13.15 Quantum metamaterial	199
14 Many-mode extension: fields, particles, and scaling	202
14.1 Why This Branch Exists	202
14.2 Page Map	202
14.3 Mechanism Pages	203
14.4 Fermi–Dirac statistics	203
14.5 Dirac equation	205
14.6 Quantum electrodynamics	207
14.7 Renormalization	209
14.8 Gauge theory	211

14.9 Photon	212
14.10Quantum field theory	214
14.11Fermion	216
14.12Boson	218
14.13Quantum chromodynamics	219
14.14AdS/CFT correspondence	221
14.15Klein–Gordon equation	223
14.16Quantum gravity	225
14.17Spin network	227
14.18Electron	229
14.19Loop quantum gravity	231
14.20Spin foam	233
14.21String theory	235
14.22Quantum geometry	237
14.23Fock space	239
14.24Quantum spacetime	241
15 Protocol layer: engineered transformations	243
15.1 Why This Branch Exists	243
15.2 Page Map	243
15.3 Mechanism Pages	244
15.4 Quantum information science	244
15.5 Quantum network	246
15.6 Quantum algorithm	248
15.7 Quantum error correction	250
15.8 Quantum channel	252
15.9 Quantum logic gate	253
15.10Quantum neural network	255
15.11Quantum circuit	257
15.12Quantum information	259
15.13Quantum computing	261
15.14Quantum key distribution	263
15.15Quantum sensor	265
15.16Quantum teleportation	267
15.17Quantum bus	269
15.18Quantum image processing	271
15.19Quantum finite automaton	273
15.20Quantum cryptography	275
15.21Quantum programming	277
16 Annotations: history, interpretations, and popular frames	280
16.1 Why This Branch Exists	280
16.2 Page Map	280
16.3 Mechanism Pages	281
16.4 Introduction to quantum mechanics	281
16.5 Old quantum theory	283
16.6 History of quantum mechanics	285
16.7 Introduction to Quantum Mechanics (book)	287
16.8 Quantum mind	289
16.9 History of quantum field theory	291
16.10Erwin Schrödinger	293
16.11Interpretations of quantum mechanics	295

16.12Quantum mysticism	297
16.13David Hilbert	299
16.14Quantum Computing: A Gentle Introduction	301
16.15QBism	303
16.16Relational quantum mechanics	305
16.17Modern Quantum Mechanics	307
16.18The Quantum Universe	309
16.19Quantum Theory: Concepts and Methods	311
Source Boundary	313

How To Use The Mechanism Tree

What The Tree Is

This book is a derivation map for quantum topics. Each named topic is treated as a contribution to one prediction-making mechanism. First specify the context: preparation, basis, boundary condition, gauge, detector arrangement, or domain. That context selects the Hilbert space and operator domain. A state is then transported by a generator, an observable defines the possible spectral answers, the Born or trace rule assigns probabilities to those answers, and compatibility constraints record which questions cannot be made jointly sharp. The page title is therefore not the root of the explanation. The root is the role the page plays in that construction.

Mechanism-first reading: identify the operation a named topic performs in the quantum construction.

Constructor sequence: context $\rightarrow$ Hilbert-space carrier and operator domain $\rightarrow$ generator/evolution $\rightarrow$ observable spectrum $\rightarrow$ probability readout $\rightarrow$ compatibility constraint $\rightarrow$ boundary/protocol realization.

DAG Before Constructor

The mechanism tree uses two distinct layers. The directed acyclic graph gives the order in which a mechanism becomes complete: context and admissibility appear before state transport; state transport appears before spectral readout; readout and compatibility appear before boundary, field, detector, or protocol realization. The constructor then fills each ordered node with the local quantum objects needed to make a prediction. This distinction matters. The DAG is the assembly order and prevents a page from being treated as a finished mechanism when it supplies only one role. The constructor is the role contract: it asks which carrier, operator, map, question, readout, closure condition, and realization are present. A topic becomes a usable mechanism when it has both a place in the DAG and a sufficiently filled constructor frame.

$$\begin{aligned} \text{DAG : } & C \prec (\mathcal{H}_C, \mathcal{D}_C) \prec \rho \prec H_C, U_C, \Phi_C \prec A_C, E_{A_C} \prec \text{Pr} \prec \mathcal{K}, \mathcal{R} \\ \text{Constructor : } & (C, \mathcal{H}_C, \mathcal{D}_C, \rho, H_C, A_C, E_{A_C}, \text{Pr}, \mathcal{K}, \mathcal{R}). \end{aligned}$$

The nonstandard result of this run is empirical rather than axiomatic. Standard quantum mechanics already contains Hilbert spaces, operators, spectra, Born probabilities, and commutators. What the rewrite adds is that a large topic system built from ordinary article names collapses into a stable constructor order. Particles, fields, protocols, interpretations, and boundary effects no longer appear as equal roots; they become later roles in one state-to-spectrum mechanism.

Why Hilbert Space Is Central

Hilbert space is central because quantum mechanics requires a space of admissible states before it requires a coordinate geometry of positions. Euclidean space labels positions, domains, detectors, or boundary conditions. Hilbert space carries phase, superposition, spin, entanglement, operator domains, and probability amplitudes. It supplies the inner product, norm, linear superposition, operator action, spectral projectors, and unitary maps required for quantum prediction. The usual position wave function illustrates the distinction. In the expression $\langle \psi(x) |$, the coordinate $x \in \mathbb{R}^3$ labels a representation. The state itself is $|\psi\rangle \in L^2(\mathbb{R}^3)$, and adding spin, identical particles, fields, or open-system mixtures moves the carrier to tensor products, Fock spaces, density operators, or operator algebras. Thus Euclidean space is often a basis or realization domain; Hilbert space is the admissibility selector.

$$\begin{aligned}
 |\psi\rangle \in \mathcal{H}, \quad \rho \geq 0, \quad \text{Tr } \rho = 1 & \quad \text{state carrier} \\
 p_i = |\langle e_i | \psi \rangle|^2, \quad \text{Pr}(\Delta) = \text{Tr}(\rho E_A(\Delta)) & \quad \text{probability structure} \\
 A = A^\dagger, \quad A = \int_{\sigma(A)} \lambda dE_A(\lambda) & \quad \text{operator-to-spectrum conversion} \\
 \rho_t = U(t)\rho U(t)^\dagger, \quad U^\dagger U = I & \quad \text{identity-preserving evolution.}
 \end{aligned}$$

In this sense Hilbert space is the selector chosen by the first branch. A context selects $(\mathcal{H}_C, \mathcal{D}_C)$: the admissible carrier and the operator domain. The later branches explain what is done with that selected carrier: states inhabit it, generators move states through it, observables resolve spectra on it, Born rules read probabilities from it, and commutators identify when two questions cannot share one sharp basis. Hilbert space selects the legal arena; it does not select the outcome.

The Compact Representation

The compact representation is the minimal formal record needed to say what a quantum mechanism is doing. It is a compressed version of standard quantum mechanics:

Compact tree: selector $\rightarrow$ carrier $\rightarrow$ map $\rightarrow$ question $\rightarrow$ readout.

Compatibility constrains which questions can be jointly sharp. Realization adds boundaries, fields, detectors, protocols, and scaling limits.

- **Context** means the experimental or mathematical setting in which a quantum question is posed: preparation, basis, boundary, gauge, detector, domain, or representation.
- **Hilbert-space carrier and domain** means the Hilbert space, operator domain, and constraints selected by that context. Examples include normalization, positivity of a density operator, boundary conditions, gauge constraints, and domain conditions for an unbounded operator.
- **Generator/evolution** means the Hamiltonian, unitary map, quantum channel, action, or other lawful map that transports the state before readout.
- **Observable spectrum** means the allowed answer set of a measurable question, represented by a self-adjoint operator, projection-valued measure, POVM, or channel readout.
- **Probability readout** means the Born or trace rule that turns the state and spectral channels into probabilities for observed outcomes.

- **Compatibility constraint** means a restriction on which questions can be jointly resolved, usually expressed by commutators, uncertainty relations, contextuality tests, conservation laws, or admissibility checks.
- **Boundary/protocol realization** means the physical or engineered implementation: a potential well, scattering boundary, cavity, field mode, detector, circuit, channel, or measurement protocol.

$$\mathfrak{M}_Q = (C, \mathcal{H}_C, \mathcal{D}_C, \rho, H_C, A_C, E_{A_C}, \text{Pr}, \mathcal{K}, \mathcal{R})$$

C : context, preparation, basis, boundary, gauge, or detector arrangement

$\mathcal{H}_C, \mathcal{D}_C$: admissible state space and operator domain

$$\rho_t = U_C(t)\rho U_C(t)^\dagger, \quad U_C(t) = \exp(-iH_C t/\hbar)$$

$$A_C = \int_{\sigma(A_C)} \lambda dE_{A_C}(\lambda)$$

$$\text{Pr}(\Delta \mid C, \rho, A) = \text{Tr}(\rho_t E_{A_C}(\Delta))$$

$\mathcal{K}$: compatibility tests such as commutators, constraints, and admissibility checks

$\mathcal{R}$: realization layer: boundary, field mode, detector, circuit, or scaling limit.

This representation explains the tree. Context selects the Hilbert-space carrier and operator domain. The wave function and density matrix are states on that carrier. The Hamiltonian and unitary operator contribute the generator. Observables and projection-valued measures contribute the spectral question. The Born rule contributes the probability readout. Commutators, uncertainty, Bell-type pages, and EPR-type pages contribute compatibility tests. Tunnelling, particle-in-a-box, scattering, fields, gauge theory, circuits, and channels contribute realization or protocol layers.

Recipe For Reading A Page

1. **Locate the page in the tree.** The branch tells which part of $\mathfrak{M}_Q$ the page mainly modifies.
2. **Fill the compact tuple.** Identify C , $\mathcal{H}_C$, ρ , H_C or the relevant map, A_C , the spectral measure or readout, and any compatibility condition.
3. **Separate topic-specific pages from core-derived pages.** Topic-specific pages have native equations or explicit constructor skeletons. Core-derived pages inherit the shared constructor spine and specialize it with branch and topic evidence.
4. **Read anomalies as diagnostics.** An anomaly is a junction where the topic uses several roles at once; the diagnosis is decomposition, not rejection.
5. **Use transfer carefully.** A mechanism can be transferred only when the state carrier, operator role, readout, and compatibility test survive the move; field-specific nouns may change.

What Was Found In This Run

The current export contains 147 quantum pages. Of these, 145 have topic-specific equation skeletons or explicit constructor overrides, and 1 are expanded from the compact constructor as core-derived mechanisms. The sparse-attention profile is strongly operator/spectral: 145 pages

exceed the operator/spectrum threshold, with mean signal 0.328. State evolution is also broad (145 pages), while boundary/context (54), incompatibility (59), and explicit protocol (32) are more selective. The practical result is a compact reading of quantum theory as operator-to-spectrum conversion under context and compatibility constraints. This is the main structural finding of the rewrite. Quantum theory can be introduced as a constructor in which particles, waves, fields, detectors, boundaries, circuits, and interpretations are roles attached to a smaller state-operator-readout spine. The portable unit is the legal transformation from context and state to spectral readout.

Branch Map

Role	Use in the derivation	Pages
Hilbert-space context: admissible carrier and basis	A quantum calculation first fixes the Hilbert space, operator domain, basis, preparation context, representation, gauge, or boundary condition. This is not the measured answer; it is the legal carrier on which states, transformations, observables, and probabilities can be defined.	7
State carrier inside Hilbert space	A state is the probability-bearing element of the selected Hilbert space or its density-operator state space. Wave functions, density matrices, superpositions, and coherent states are different representations of this predictive carrier.	24
Generator: lawful change before readout	Hamiltonians, unitary maps, equations of motion, and path weights describe the lawful transport of the state before a question is resolved.	22
Spectral question: what can be asked	An observable is a permitted question whose operator form determines the possible numerical answers.	10
Readout rule: how answers become probabilities	Measurement connects a state and an observable to recorded frequencies. Projection, POVMs, Born weights, and collapse language are alternative ways of presenting this state-to-spectrum readout step.	10
Compatibility limit: what cannot be jointly sharp	The non-classical part of the theory appears when two otherwise legal questions do not compose into one common sharp question. Commutators, uncertainty relations, contextuality, Bell tests, and entanglement live here.	6
Boundary realization: how effects appear	Many named quantum effects are boundary realizations of the same construction. A potential, barrier, box, cavity, detector, or medium changes the allowed spectral channels without changing the basic prediction problem.	12
Many-mode extension: fields, particles, and scaling	Quantum field theory, gauge theory, renormalization, photons, fermions, and related topics extend the same state-operator-spectrum logic to variable particle number, local fields, and scale-dependent descriptions.	21

Role	Use in the derivation	Pages
Protocol layer: engineered transformations	Quantum computing, channels, circuits, algorithms, networks, sensors, and error correction turn the same formal machinery into controlled sequences of operations.	18
Annotations: history, interpretations, and popular frames	Some pages help readers navigate the subject but do not form steps in the mechanism. They are kept as annotations so books, historical figures, interpretations, and popular frames do not distort the constructive tree.	16

Boundary Of The Claim

The book is a mechanism-indexed synthesis generated from Wikipedia topic scaffolds, MorphWiki mechanism summaries, and ranked source-equation witness profiles. Its uses are reading, teaching, hypothesis generation, and constructor-target selection. Formal derivation, domain review, and experimental validation are the tests that promote a mechanism reading into a physical claim.

Preamble: Findings From The Rewrite

Core Insight

The rewrite reveals a compact operational reading of quantum theory. The standard formalism has a simpler explanatory order than the usual topic list. The order recovered by the sparse-attention pass is:

Constructor sequence: context $\rightarrow$ Hilbert-space carrier and operator domain $\rightarrow$ generator/evolution $\rightarrow$ observable spectrum $\rightarrow$ probability readout $\rightarrow$ compatibility constraint $\rightarrow$ boundary/protocol realization.

Here, context means the preparation, basis, boundary, gauge, domain, detector, or representation in which a question is posed. It selects the Hilbert space and operator domain allowed by that context, together with normalization, positivity, boundary, gauge, or domain constraints. This carrier is central: it defines which states are legal, how amplitudes and norms are compared, which operators may act, and how spectral projectors can be read probabilistically. The generator is the Hamiltonian, unitary map, channel, or action that carries the state before readout. The observable spectrum is the allowed answer set of the question being asked. The probability readout is the Born or trace-rule assignment to those answers. The compatibility constraint records which otherwise valid questions cannot be resolved together, for example through a non-zero commutator, uncertainty relation, contextuality test, conservation law, or admissibility condition. Boundary and protocol realization specify how the same construction is embodied as a potential well, scattering boundary, cavity, field mode, detector, circuit, or measurement sequence.

Main Finding And Why It Matters

The main finding is that the quantum corpus can be reorganized away from named topics into a smaller operational constructor: context-selected Hilbert space, state carrier, generator, observable spectrum, probability readout, compatibility limits, and realization or protocol. This changes the explanatory root. In the usual presentation, quantum theory appears as a collection of named effects and puzzles: particles, waves, measurement, tunnelling, entanglement, collapse, fields, and interpretations. In the mechanism-tree view, these become roles inside one compact state-operator-spectrum construction.

The importance is practical. The tree gives a recipe for constructing and comparing theories. A page becomes useful when one can specify what state is carried, what operator or map acts, what spectrum or readout is produced, what compatibility constraint applies, and what changes when the realization changes. This turns topic similarity into mechanism similarity: two subjects may use different nouns while preserving the same state-operator-readout role.

This organization of existing quantum theory has direct consequences for physical AI. A constructor-oriented AI can search over compact mechanisms, test whether the operator/spectral role survives changes of representation, and use anomalies as places where several roles

collide. The result is a compact, falsifiable, and transferable representation of quantum theory.

Transition Sparse-Attention Result

A second sparse-attention pass treated the rewrite itself as the object of study: Wikipedia/topic view to mechanism-tree view. This comparison asks what becomes visible only after the same quantum topics are reorganized by constructor role rather than by article title.

The transition run covered 147 pages. Of these, 145 currently have topic-specific equation skeletons or explicit constructor overrides, while 2 are expanded by the shared compact constructor. The distinction is evidential, not structural: both page types are read through the same context-state-generator-spectrum-readout sequence.

The transition signal is asymmetric: the rewrite adds operational roles more strongly than it adds object names. The mechanism view exposes the state, operator, readout, compatibility, boundary, and protocol roles that are implicit in the topic view.

Route role	Mean signal	Interpretation
operator-to-spectrum readout	0.3272	constructor role preserved in the rewrite
state evolution	0.2321	constructor role preserved in the rewrite
normalization or admissibility	0.1646	constructor role preserved in the rewrite
preparation, basis, or boundary context	0.0943	constructor role preserved in the rewrite
non-commuting compatibility limits	0.0861	constructor role preserved in the rewrite
controlled update protocol	0.0753	constructor role preserved in the rewrite

New Structural Information

- The rewrite converts a noun-indexed encyclopedia into a derivation graph.**
The new object is not a better summary of each topic; it is an ordering relation: which topic plays context, state, generator, observable, readout, compatibility, boundary, field, protocol, or annotation.
- The dominant stable role is operator-to-spectrum readout, not object naming.**
The rewrite makes explicit that many named quantum topics become different ways of asking a legal spectral question of a state.
- Particles become stable role-realizations inside field/mode/readout machinery.**
The particle pages are not discarded; they are relocated as field/mode/statistics/readout constructions. This is a more precise statement than 'particles are not fundamental'.
- Interpretations mostly act on readout semantics rather than replacing the**

formal constructor. QBism, relational quantum mechanics, collapse language, and popular frames can be kept without letting them become false roots of the derivation tree.

5. **Boundary pages are realization gates: they change allowed spectra without changing the core prediction problem.** Tunnelling, particle-in-a-box, scattering, cavities, and spectral lines become one family: boundary-shaped spectra.
6. **Protocol pages are an engineering layer over the constructor, not the root of the theory.** Quantum computing is reorganized as controlled composition of states, operators, readouts, and error constraints rather than a separate ontology of qubits.
7. **Anomalies identify where several constructor roles collide.** Anomalies are not errors in the tree. They are research handles: EPR, measurement problem, quantum gravity, quantum biology, and related pages require several roles at once.

Highest-Attention Transition Pages

- **Quantum logic** (generators, constructed): dominant roles after rewriting are measurement, context, state. Mechanism reading: This page sits between generators and annotations. The ambiguity is useful: it marks a place where two constructor roles meet and should be separated before the page is used as a derivation.
- **Quantum biology** (generators, constructed): dominant roles after rewriting are context, observable, measurement. Mechanism reading: Quantum biology is an open-system transfer problem. The anomaly is that the biological environment is not background noise only; it is part of the boundary that may preserve, destroy, or select coherence. A rigorous constructor must name the state carrier, the environmental coupling, the coherence or transport observable, and the classical control that would remove the quantum contribution.
- **Schrödinger’s cat** (states, constructed): dominant roles after rewriting are measurement, state, context. Mechanism reading: Schrödinger’s cat is a macroscopic readout protocol, not a spectrum-first topic. It couples microscopic unitary evolution to a macroscopic boundary and forces the reader to separate three steps: coherent state transport, decoherence or apparatus coupling, and the rule by which one record is selected or conditioned.
- **Measurement problem** (measurement, constructed): dominant roles after rewriting are measurement, context, state. Mechanism reading: The measurement problem is a readout junction. It sits where unitary state transport, detector context, probability assignment, and state update meet. Use four parts: state evolution before measurement, coupling to an apparatus or environment, POVM or projection readout, and the rule used to condition the state after the record.
- **Delayed-choice quantum eraser** (measurement, constructed): dominant roles after rewriting are measurement, context, state. Mechanism reading: The delayed-choice eraser is a protocol-order stress test. Its mechanism is the arrangement of which-path information, later measurement choice, and conditional correlation readout. The anomaly is not retrocausality by default; it is that the relevant statistics are defined only after the full measurement protocol is specified.
- **Introduction to quantum mechanics** (annotations, constructed): dominant roles after rewriting are measurement, context, state. Mechanism reading: An introductory page is anomalous because pedagogy compresses the whole constructor into one narrative. It mixes states, operators, spectra, measurement, examples, and interpretations. It should

be used as a map, then decomposed into mechanism branches before making technical claims.

- **Einstein–Podolsky–Rosen paradox** (incompatibility, constructed): dominant roles after rewriting are measurement, context, state. Mechanism reading: EPR is a compatibility test, not a page about a peculiar object. Its mechanism is a bipartite state, separated measurement contexts, and a correlation readout that cannot be reduced to pre-existing local values. The construction should therefore start from the joint state and the allowed local observables, then ask which correlation constraints fail.
- **Quantum nonlocality** (incompatibility, constructed): dominant roles after rewriting are measurement, context, state. Mechanism reading: This incompatibility page mixes operator-to-spectrum readout, state evolution, controlled update protocol. Its mechanism should state which otherwise legal questions fail to share a single sharp representation, and what experiment or inequality exposes that failure.

Operational Use

- **Teach quantum theory as a derivation tree.** The reader sees the required assembly order: context, state, generator, observable spectrum, probability readout, compatibility limit, and realization. This avoids presenting interpretations, particles, and protocols as equal primitives.
- **Build constructor targets for the decoder.** The 145 topic-specific pages are seed targets. The 2 core-derived pages become a supervised specialization set: each needs a topic-native state, operator/map, spectrum/readout, compatibility condition, and realization.
- **Find transfer candidates across fields.** A page can be transferred only if the role survives. This lets FieldBridge search for analogues of a mechanism rather than semantic analogues of words.
- **Use anomalies as research prompts.** Multi-role pages such as EPR, measurement problem, quantum gravity, and quantum biology are not clean branches. They mark places where compatibility, boundary, protocol, and transport signals collide.
- **Separate interpretation from machinery.** Interpretive pages can be retained as readout/probability annotations without allowing them to rewrite the Hamiltonian/operator/spectral core.

The practical consequence is a derivation tree rather than a topic list: topic-specific pages show local equations, while core-derived pages show how the same compact constructor specializes to the page role.

Main Findings

1. **Quantum theory collapses to a smaller constructor.** The constructor order is context, state space, generator/evolution, observable spectrum, Born readout, and compatibility constraint. This is not a new postulate of quantum mechanics; it is a cleaner order in which to present the existing formalism.
2. **Operator/spectral structure is the spine.** 145 of 147 pages exceed the operator/spectrum threshold, with mean signal 0.328. This is the strongest and most universal signal in the rewrite.

3. **Evolution is secondary but broad.** 145 pages carry state-evolution or transport signal. Evolution is not absent; it is the motion of the predictive carrier before a readout.
4. **Context and boundary are realization gates.** 54 pages emphasize preparation, boundary, or domain context. Tunnelling, cavities, scattering, and particle-in-a-box become boundary-shaped spectra rather than separate conceptual primitives.
5. **Incompatibility is the non-classical junction.** 59 pages carry explicit incompatibility signal. Entanglement, Bell phenomena, commutators, and uncertainty are grouped by failure of joint sharpness.
6. **Protocol is rare and therefore informative.** Only 32 pages exceed the protocol threshold. Quantum computing and information are therefore best treated as an engineering layer over the state-operator-spectrum machinery.
7. **The tree compresses named topics into mechanism roles.** Hilbert-space context: admissible carrier and basis: 7, State carrier inside Hilbert space: 24, Generator: lawful change before readout: 22, Spectral question: what can be asked: 10, Readout rule: how answers become probabilities: 10, Compatibility limit: what cannot be jointly sharp: 6, Boundary realization: how effects appear: 12, Many-mode extension: fields, particles, and scaling: 21, Protocol layer: engineered transformations: 18, Annotations: history, interpretations, and popular frames: 16. This is the mechanism-level table of contents produced by the rewrite.
8. **Particles are stable role-realizations.** Particle pages resolve into field modes, spectra, statistics, and readout-stable excitations. The claim is not that particles are fake. The claim is that particle identity is a stable role inside a state-operator-readout construction.
9. **String theory and AdS/CFT sit on the field/geometry translation frontier.** String theory appears as a spectral many-mode constructor. AdS/CFT is the sharper geometry-translation case because its boundary and geometry signals are stronger in the current export.
10. **Black holes are not yet a root in this export.** The sparse-attention script flags black-hole physics as a boundary/information-closure target rather than as a finished claim. A serious black-hole reading requires an explicit rerun on black holes, horizons, Hawking radiation, entropy, holography, and information-loss topics.
11. **Interpretations sit mostly on the readout layer.** Interpretations change the meaning assigned to state, probability, collapse, update, or observer language. They do not, in this tree, replace the Hamiltonian, operator, spectrum, or Born-rule machinery.
12. **Geometry appears as realization and reconstruction.** Geometry is necessary for embodiment, boundary conditions, and physical interpretation, but source-equation stability checks preserve structure and spectral/operator roles more strongly than geometry.

What Changed Relative To Wikipedia

A conventional encyclopedia organizes quantum theory by names and historical articles. The MorphWiki rewrite organizes the same material by the operations needed to construct a prediction. The page named measurement no longer becomes the origin of the theory; it becomes the readout junction. The page named particle becomes a realization. The page named tunnelling becomes a boundary spectral channel. The page named quantum field theory becomes a scaling and many-mode extension. The tree therefore gives a new order of explanation rather than a new interpretation of consciousness, observation, or ontology.

Unusual Findings And Anomalies

The most useful anomalies are not errors. They are pages that sit between branches or violate the dominant operator/spectrum pattern. These pages mark conceptual junctions where a theory-builder should look for missing assumptions, hidden boundary conditions, or possible transfers to other fields.

The anomaly labels describe a page's role in the mechanism tree, not the literal physical object named by the page. For example, a page can be structurally anomalous because context, protocol, or compatibility carries the explanation before spectra are read out.

- **pre-spectral admissibility:** another construction step fixes the admissible question before eigenvalues become meaningful.
- **context participates in the law:** preparation, apparatus, context, representation, or boundary conditions are part of the mechanism rather than surrounding detail.
- **admissibility meets incompatibility:** the page joins the rules that make a quantum state legal with the rules that restrict jointly resolvable questions.
- **operation order matters:** the order of operations matters; the topic cannot be reduced to a static state, operator, and spectrum.
- **several mechanisms meet:** several construction steps meet in one topic, so the page is a junction rather than a clean branch leaf.
- **branch interface:** the topic belongs at an interface between two explanatory roles and should be read as a bridge before being assigned to one branch.
- **Schrödinger's cat.** Primary reading: State carrier inside Hilbert space; neighboring reading: Annotations: history, interpretations, and popular frames. Schrödinger's cat is a macroscopic readout protocol, not a spectrum-first topic. It couples microscopic unitary evolution to a macroscopic boundary and forces the reader to separate three steps: coherent state transport, decoherence or apparatus coupling, and the rule by which one record is selected or conditioned.
- **Einstein–Podolsky–Rosen paradox.** Primary reading: Compatibility limit: what cannot be jointly sharp; neighboring reading: Annotations: history, interpretations, and popular frames. EPR is a compatibility test, not a page about a peculiar object. Its mechanism is a bipartite state, separated measurement contexts, and a correlation readout that cannot be reduced to pre-existing local values. The construction should therefore start from the joint state and the allowed local observables, then ask which correlation constraints fail.
- **Quantum biology.** Primary reading: Generator: lawful change before readout; neighboring reading: Hilbert-space context: admissible carrier and basis. Quantum biology is an open-system transfer problem. The anomaly is that the biological environment is not background noise only; it is part of the boundary that may preserve, destroy, or select coherence. A rigorous constructor must name the state carrier, the environmental coupling, the coherence or transport observable, and the classical control that would remove the quantum contribution.
- **Introduction to quantum mechanics.** Primary reading: Annotations: history, interpretations, and popular frames; neighboring reading: State carrier inside Hilbert space. An introductory page is anomalous because pedagogy compresses the whole constructor

into one narrative. It mixes states, operators, spectra, measurement, examples, and interpretations. It should be used as a map, then decomposed into mechanism branches before making technical claims.

- **Measurement problem.** Primary reading: Readout rule: how answers become probabilities; neighboring reading: Generator: lawful change before readout. The measurement problem is a readout junction. It sits where unitary state transport, detector context, probability assignment, and state update meet. Use four parts: state evolution before measurement, coupling to an apparatus or environment, POVM or projection readout, and the rule used to condition the state after the record.
- **Quantum gravity.** Primary reading: Many-mode extension: fields, particles, and scaling; neighboring reading: Generator: lawful change before readout. Quantum gravity is a field/boundary junction. It asks whether geometry itself becomes part of the quantum state carrier or remains a realization layer for an operator theory. The missing constructor is explicit: a state of geometry, a constraint or evolution operator, a boundary or semi-classical readout, and a test showing which geometric quantities survive quantization.
- **Scattering.** Primary reading: Boundary realization: how effects appear; neighboring reading: Generator: lawful change before readout. Scattering is a boundary-to-spectrum mechanism. The relevant object is the map from asymptotic in-states to out-states. The constructor should specify the Hamiltonian or interaction region, the boundary/asymptotic channels, the S-matrix or cross-section readout, and the conservation constraints.
- **Quantum state.** Primary reading: State carrier inside Hilbert space; neighboring reading: Generator: lawful change before readout. Quantum state is the carrier, not the final prediction. It is anomalous because it precedes several downstream roles: admissibility, evolution, observable choice, and probability readout. A clean derivation should state whether the carrier is a vector, density operator, field state, or register, and which transformations preserve its legality.

Interpretation Of The Anomalies

The sparse-attention interpretation is aggregated from role labels assigned by the mechanism pass. It summarizes recurring junction types in the mechanism tree rather than presenting a list of physical anomalies.

- **several mechanisms meet.** several construction steps meet in one topic, so the page is a junction rather than a clean branch leaf. Mechanism implication: one topic name covers several formal roles, so the page is a junction rather than a single branch leaf.
- **pre-spectral admissibility.** another construction step fixes the admissible question before eigenvalues become meaningful. Mechanism implication: the eigenvalue problem is downstream of another role that first defines the admissible question.
- **operation order matters.** the order of operations matters; the topic cannot be reduced to a static state, operator, and spectrum. Mechanism implication: the order of preparation, transformation, and readout is part of the mechanism.
- **context participates in the law.** preparation, apparatus, context, representation, or boundary conditions are part of the mechanism rather than surrounding detail. Mechanism implication: preparation, apparatus, domain, detector, or boundary conditions participate in the observed law.

- **admissibility meets incompatibility.** the page joins the rules that make a quantum state legal with the rules that restrict jointly resolvable questions. Mechanism implication: admissibility conditions and jointly resolvable questions meet in the same topic.
- **branch interface.** the topic belongs at an interface between two explanatory roles and should be read as a bridge before being assigned to one branch. Mechanism implication: the topic lies at an interface between explanatory roles and should be read as a bridge.

Constructor-Detected Research Leads

The useful anomalies in this book are constructor junctions: places where a topic requires several formal roles at once. Some are established foundational problems; others are under-formulated mechanism questions that can guide derivation and experiment.

- **Measurement as a four-map junction.** The established measurement problem becomes operationally sharper when split into pre-measurement evolution, apparatus or environment coupling, POVM or projection readout, and post-record state conditioning. The useful signal is the missing local map between probability readout and state update.
- **Geometry as a possible state carrier.** Quantum gravity is a recognized frontier. The constructor reading isolates the specific unresolved choice: whether geometry remains a substrate on which operators act, or becomes part of the quantum state carrier itself. The corresponding test is which geometric quantities survive as admissible observables after quantization.
- **Boundary-selected spectra across different phenomena.** Scattering, tunnelling, cavities, particle-in-a-box systems, spectral lines, and optical boundary problems share a boundary-to-spectrum construction. The research lead is to treat them as one transferable mechanism family rather than as separate examples tied to different physical nouns.
- **Environment as an active selector in quantum biology.** In quantum biology, the useful constructor signal is that the environment can be part of the mechanism: it may preserve, destroy, or select coherence. A useful derivation must name the carrier, coupling, coherence or transport observable, and classical control.
- **Hamiltonian dual role.** The Hamiltonian is both a generator of time evolution and an observable with an energy spectrum. This is standard formalism. The constructor lead is to keep those roles distinct when deriving mechanisms that use energy conservation, spectral measurement, and unitary transport in the same page.
- **Quantum simulation as a target-carrier-validation triangle.** A simulator is an engineered realization of another Hamiltonian, channel, or field theory. The anomaly is practical: the physical carrier, encoded target, and validation observable are distinct roles. A simulator claim is incomplete until all three are specified.

Relation To Lagrangian, Transfer, And Stability Layers

The later validation chapter explains how the global evidence layers check this tree. Stability tests ask which part of a formalism remains the same after a rewrite. Typed transfer and bridge-family layers ask which constructor roles can move between contexts. The Lagrangian navigation layer asks whether a proposed move through the atlas is a low-resistance continuation, a strained bridge, or a void-boundary target. In ordinary quantum language: the invariant is mostly operator/structural role, while geometry, boundary, and representation are where the role becomes embodied.

Practical Consequence

A reader, teacher, or AI constructor should not start from named quantum objects. The constructive order is: define the admissible context, define the predictive carrier, define lawful change, define the legal question, resolve its spectrum, assign probabilities, test incompatibility, then dress the construction in the relevant physical realization. That is the main structural result of the rewrite.

Part I

The Mechanism Tree

Chapter 1

Root Construction

Given a context-selected Hilbert space, an admissible state, and a permitted observable, quantum theory determines which numerical outcomes can occur, assigns probabilities to those outcomes, and marks which questions cannot be made simultaneously sharp.

1.1 Re-Derivation Path

1. **Context and Hilbert space.** Fix the Hilbert space, operator domain, basis, preparation condition, boundary, gauge, or representation.
2. **State carrier.** Write a normalized vector, wave function, density operator, or field/register state on that carrier.
3. **Generator.** Let a Hamiltonian, unitary map, constraint, or action move that carrier before readout.
4. **Spectral question.** Represent a measurable question by an operator with allowed answers.
5. **Readout.** Resolve the operator into outcome channels and assign probabilities.
6. **Compatibility.** Mark when two legal questions cannot be made jointly sharp.
7. **Realization and extension.** Add boundaries, fields, scaling limits, detectors, or protocols as later embodiments of the same construction.

$$\begin{aligned} B &\mapsto \rho_B \\ \rho_t &= U_t \rho_B U_t^\dagger \\ O &= \sum_i \lambda_i P_i \\ p_i &= \text{Tr}(P_i \rho_t) \\ [O_1, O_2] &\neq 0. \end{aligned}$$

Chapter 2

Sparse Attention Summary

Operation	Mean signal	Pages above 0.10
State evolution	0.232	145
Normalization and admissibility	0.165	136
Observables and spectra	0.328	145
Preparation and boundary context	0.095	54
Incompatible questions	0.086	59
Controlled update protocol	0.076	32

The dominant recurrent signal is observables and spectra. Evolution, context, incompatibility, closure, and protocol are branch-forming operations. This is why the book begins with admissible context, state carrier, generator, and legal question rather than with particles or measurement folklore.

Chapter 3

Compact Operator Formulation

The compact form of nonrelativistic quantum theory is a map from a context and a state to a probability measure over an operator spectrum.

This chapter is the formal spine of the book. It does not introduce new physics. It rewrites familiar quantum mechanics as a sequence of necessary construction steps. Each later page is placed by asking which step it changes: admissible context, state carrier, generator, observable, readout, compatibility, realization, many-mode extension, or protocol.

3.1 The Minimal Constructor

$$\begin{aligned} C &\longmapsto (\mathcal{H}_C, \mathcal{D}_C) \\ \rho &\in \mathcal{S}(\mathcal{H}_C), \quad \rho \geq 0, \quad \text{Tr } \rho = 1 \\ \rho_t &= U_C(t) \rho U_C(t)^\dagger, \quad U_C(t) = \exp(-iH_C t/\hbar) \\ A_C &= \int_{\sigma(A_C)} \lambda dE_{A_C}(\lambda) \\ \text{Pr}(\Delta \mid C, \rho, A) &= \text{Tr}(\rho_t E_{A_C}(\Delta)) \\ [A_C, B_C] \neq 0 &\Rightarrow \text{no generic common sharp spectral refinement.} \end{aligned}$$

3.2 Interpretation Of The Symbols

- C is the context: preparation, basis, boundary condition, gauge choice, detector arrangement, domain, or representation.
- $\mathcal{H}_C$ is the admissible state space selected by that context.
- $\mathcal{D}_C$ is the domain or admissibility condition for the generator and observables.
- ρ is the predictive state carrier. It may be a state vector, density operator, field state, or register state.
- H_C is the generator of lawful change before readout.
- A_C is the question being asked. Its spectral measure E_{A_C} supplies the outcome channels.
- $\text{Pr}(\Delta \mid C, \rho, A)$ is the Born probability of an outcome set Δ .
- A non-zero commutator marks a compatibility limit: two legal questions may not share one sharp readout basis.

3.3 Why This Is More Compact

The usual presentation introduces particles, waves, measurement, operators, interpretations, and fields as separate conceptual blocks. The compact constructor shows that many of these are changes of role rather than separate roots. A particle is a stable state/readout role. A barrier is a context that changes the operator domain. A field is a many-mode extension of the state space. A quantum circuit is a protocolized composition of maps. An interpretation changes the meaning of state, probability, or update, but usually leaves the constructor intact.

Chapter 4

Mechanism Validation Layers

The mechanism tree is evaluated by three independent evidence layers: role-current stability asks what survives a rewrite, Gromov-Wasserstein (GW) neighborhoods measure relational bridge candidates, and the learned Lagrangian asks whether a proposed move is an easy continuation, a strained bridge, or a void-boundary design target.

4.1 Role-Current Layer: What Stays Stable

The role-current layer measures stability of formal roles under rewrite and translation. When a theory is rewritten, translated, or moved through the corpus, the test asks which part of the formal apparatus keeps its role. For a mechanism tree, this is the identity test: two formulations belong together when the same role contract survives the move.

- The run promoted ? strict currents and ? near currents. These mark role-preserving directions in the learned representation.

4.2 Structural Bridge Layer: Candidate Translations

The Gromov-Wasserstein layer compares relational neighborhoods. It contributes a structural bridge score: two formulations are stronger bridge candidates when their local neighborhoods preserve comparable relations among roles. In the current constructor, this layer is combined with typed transfer edges, bridge-family candidates, source-local transitions, and source-equation validation.

- The run scanned ? Gromov-Wasserstein candidates, hardened ? bridge records, and retained ? artifact-validated method bridges.
- GW nominates relationally compatible bridge candidates; typed transfer edges, directed source transitions, source equations, and residual checks rank them for use.

4.3 Lagrangian Layer: Which Moves Are Easy Or Strained

The learned Lagrangian is the navigation layer of the atlas. It is a representation-space action surrogate over source-equation fingerprints, separate from the physical action of a quantum system. Given a construction-role state, it asks which next states are cheap continuations, which require a strained bridge, and which absences sit on a meaningful void front. In this sense it finds roads through the operational grammar: low-action corridors where a formal role can be continued, translated, or tested. In the current quantum tree this layer supplies the

global road-map constraint. A full page-coordinate export will allow direct page-level action scoring.

- For the quantum tree, the Lagrangian is therefore a construction constraint rather than a leaf selector. It tells how open construction steps should be interpreted in the atlas, but the present export cannot yet say that a particular Wikipedia page lies on a particular low-action road. That requires full fingerprints or witness transition vectors for the page.

4.4 How These Layers Change The Quantum Tree

A quantum concept is not placed by name alone. It is placed by the role it plays in the mechanism, checked against source equation witnesses, and interpreted through stability and transfer layers. If a page preserves structure and spectral role, it belongs near the spine. If it changes the admissible domain, it belongs in boundary realization. If it introduces many modes or scale flow, it belongs in the field/scaling extension. If it describes a sequence of controlled operations, it belongs in the protocol layer. This is the human-readable reason for the tree: it unfolds quantum theory as a construction, then asks which parts survive when notation, representation, geometry, or field changes.

Chapter 5

Sparse-Attention Result

Quantum theory, in this sparse-attention reading, is a constructor that turns admissible state descriptions into spectra and probabilities under compatibility and boundary constraints.

Particles, strings, fields, measurements, and interpretations collapse into roles in the same constructor rather than separate root categories.

5.1 Minimal Constructor

The sparse-attention run returns the same constructor as a role sequence. Rendered in standard quantum notation, the sequence is:

$$\begin{aligned} B &\mapsto (\mathcal{H}_B, \mathcal{D}_B) \\ \rho_B(t) &= U_B(t)\rho_B(0)U_B(t)^\dagger \\ O_B &= \sum_i \lambda_i P_i \quad \text{or, more generally,} \quad O_B = \int_{\sigma(O_B)} \lambda dE_{O_B}(\lambda) \\ p_i &= \text{Tr}(P_i \rho_B(t)), \quad \text{Pr}(\Delta) = \text{Tr}(\rho_B(t) E_{O_B}(\Delta)) \\ [O_1, O_2] \neq 0 &\Rightarrow \text{no generic common sharp readout} \\ \mathcal{R} &: \text{boundary, field, detector, circuit, or scaling realization.} \end{aligned}$$

- B denotes the context: preparation, basis, boundary, domain, gauge, or detector arrangement.
- $(\mathcal{H}_B, \mathcal{D}_B)$ is the admissible state space and operator domain selected by that context.
- $\rho_B(t)$ is the propagated state carrier.
- O_B , P_i , and E_{O_B} express the observable question and its spectral readout channels.
- The probabilities p_i or $\text{Pr}(\Delta)$ are Born-rule readouts of the state against those channels.
- Non-commuting observables define compatibility limits; realization layers specify how the same role is embodied.

5.2 Route Statistics

Role	Mean signal	Pages > 0.10
state evolution	0.232	146
admissibility and closure	0.165	137
observable spectra	0.327	146
preparation and boundary context	0.094	54
incompatible questions	0.086	59
controlled update protocol	0.075	32

The strongest recurrent signal is observable spectra. In the current export it is active above threshold on nearly every page, while protocol/update is much sparser. This is the quantitative reason the tree begins with state, generator, observable, spectrum, and readout before it introduces algorithms, experiments, or interpretations.

5.3 Constructor Regularities

The regularities below are the strongest reusable patterns in the sparse-attention export. They are public reading rules for the quantum book, not internal diagnostics or a fixed ontology. Title-token rules and other lexical artifacts are omitted here because they are useful for auditing but too weak for the main exposition.

5.3.1 Operator-to-spectrum as the readout spine

Reading. The stable readout of a quantum construction is usually spectral. A state becomes experimentally or mathematically predictive when an allowed operator question exposes eigenvalues, projectors, or a spectral measure.

Reading consequence. Introduce observables, self-adjointness, spectral projectors, and the Born rule as the common readout interface before presenting particles, fields, or protocols as separate subjects.

5.3.2 State transport as the carrier of change

Reading. Quantum topics repeatedly require a lawful carrier of change: a state vector, density operator, field mode, or channel is propagated before it is read.

Reading consequence. Derive state evolution as the map between preparation and later readout, using unitary, semigroup, or channel language depending on whether the system is closed, open, or operational.

5.3.3 Admissibility as the legal spine

Reading. Closure and admissibility form the legal spine of the theory: states must be normalizable or positive, maps must preserve the allowed state set, and operators must have a domain on which the question is well-defined.

Reading consequence. Every derivation should state what makes the state, operator, map, or probability assignment legal before it discusses physical interpretation.

5.3.4 Incompatibility as a selective limit

Reading. Incompatibility is selective. It becomes central when two questions, bases, or transformations cannot be made sharp at the same time.

Reading consequence. Use commutators, non-common eigenbases, Bell-type constraints, or contextuality tests only where the page is actually about jointly unavailable readouts.

5.3.5 Context as the active boundary of the question

Reading. Context is sometimes part of the law. Preparation, boundary conditions, detector arrangement, basis choice, or representation can change which state and operator are admissible.

Reading consequence. When context is active, write the context before the state: specify the Hilbert space, domain, boundary, apparatus, or channel that makes the later spectral question meaningful.

5.3.6 Protocol as ordered implementation

Reading. Protocols are a sparse but real layer. They become central when an ordered sequence of operations, measurements, controls, or updates determines what can be inferred.

Reading consequence. Place quantum algorithms, circuits, finite automata, Bell tests, and delayed-choice experiments in this layer: their content depends on operation order, not only on a static Hamiltonian.

5.3.7 Junction pages require decomposition

Reading. Some pages are junctions rather than leaves. EPR, wave function, delayed-choice eraser, entanglement, quantum biology, and similar topics join state, evolution, readout, boundary, protocol, and compatibility in one place.

Reading consequence. Such pages should be read as junctions of several formal roles: state carrier, transformation, readout, context, and compatibility condition. Their position marks where a single topic name covers more than one construction step.

5.3.8 Geometry enters as realization

Reading. Geometry enters mainly through realization-heavy branches. In this export, geometric language most often specifies boundary, field, context, or reconstruction layers rather than replacing the operator-spectral core.

Reading consequence. Introduce geometry when the theory needs a domain, boundary, metric, representation, field realization, or reconstruction map. Keep the operator and admissibility structure visible underneath.

5.4 Topic-Specific Readings

- **Particle:** In this reading, a particle is not the root unit. It is a stable field/mode/readout realization of the state-operator-spectrum constructor. Photon, electron, fermion, and boson do not share one identical route profile; 'particle' decomposes into spectral, boundary, closure, and statistics roles. Relevant pages: Boson, Fermion, Photon, Fock space, Creation and annihilation operators, Electron.
- **Black-hole direction:** Black holes do not appear as a primary constructor node in this export. They enter as a boundary/geometry stress test through quantum gravity, string, AdS/CFT, and horizon-like witnesses. This suggests a useful rerun: black-hole topics

should be processed explicitly to test whether horizon physics is a boundary readout, an information-closure problem, or an geometry-free operator-kernel gap. Relevant pages: Quantum gravity, AdS/CFT correspondence, Quantum geometry, Loop quantum gravity, Electron, Fermion.

- **String-theory direction:** String theory appears as a many-mode field/scale extension with strong spectral content and geometry/boundary pressure, not as a separate noun ontology. The string is best read as a carrier of spectra under constraints; AdS/CFT is the sharper geometry-translation page because its boundary signal is higher. Relevant pages: String theory, AdS/CFT correspondence, Renormalization, Quantum mechanics, Quantum geometry, Loop quantum gravity.
- **Measurement and interpretations:** Measurement is a readout/probability layer, and interpretations modify the meaning of that layer rather than the constructor equations. QBism and relational pages can have strong operator/spectrum evidence while still being annotations, because they narrate the readout layer instead of adding new dynamics. Relevant pages: QBism, Wave function collapse, POVM, Interpretations of quantum mechanics, Born rule, Quantum mechanics.
- **Geometry:** Geometry behaves as realization and reconstruction: it makes the operator construction legal on domains, boundaries, gauges, or holographic duals, but it is not the strongest preserved current. Quantum gravity, AdS/CFT, and spin foam cluster near field/boundary roles; this matches the current source-equation stability checks. Relevant pages: Spin foam, Loop quantum gravity, Quantum geometry, AdS/CFT correspondence, Electron, Fermion.
- **Void leads:** The best discovery leads are missing gates: contexts where a state, operator, boundary, or compatibility relation is implied but not closed by evidence. The black-hole/holography area and geometry-free operator kernels are natural tests for this, They require explicit topic reruns and typed equations. Relevant pages: Quantum mechanics, Quantum geometry, Fock space, Photon, Electron, Fermion.

These statements are role hypotheses produced by deterministic sparse attention over the MorphWiki quantum pages and existing source-equation findings. They are not final physics claims. The next test is to rerun the same analysis on explicit black-hole, horizon, Hawking-radiation, entropy, holography, and information-loss topics with typed equation witnesses.

Chapter 6

Anomalies And Leads

6.1 Structural Anomalies

The labels below describe why a page is structurally interesting in the mechanism tree. They do not describe the literal object named by the page.

- **pre-spectral admissibility:** another construction step fixes the admissible question before eigenvalues become meaningful.
- **context participates in the law:** preparation, apparatus, context, representation, or boundary conditions are part of the mechanism rather than surrounding detail.
- **admissibility meets incompatibility:** the page joins the rules that make a quantum state legal with the rules that restrict jointly resolvable questions.
- **operation order matters:** the order of operations matters; the topic cannot be reduced to a static state, operator, and spectrum.
- **several mechanisms meet:** several construction steps meet in one topic, so the page is a junction rather than a clean branch leaf.
- **branch interface:** the topic belongs at an interface between two explanatory roles and should be read as a bridge before being assigned to one branch.
- **Schrödinger’s cat.** Schrödinger’s cat is a macroscopic readout protocol, not a spectrum-first topic. It couples microscopic unitary evolution to a macroscopic boundary and forces the reader to separate three steps: coherent state transport, decoherence or apparatus coupling, and the rule by which one record is selected or conditioned.
- **Einstein–Podolsky–Rosen paradox.** EPR is a compatibility test, not a page about a peculiar object. Its mechanism is a bipartite state, separated measurement contexts, and a correlation readout that cannot be reduced to pre-existing local values. The construction should therefore start from the joint state and the allowed local observables, then ask which correlation constraints fail.
- **Quantum biology.** Quantum biology is an open-system transfer problem. The anomaly is that the biological environment is not background noise only; it is part of the boundary that may preserve, destroy, or select coherence. A rigorous constructor must name the state carrier, the environmental coupling, the coherence or transport observable, and the classical control that would remove the quantum contribution.

- **Introduction to quantum mechanics.** An introductory page is anomalous because pedagogy compresses the whole constructor into one narrative. It mixes states, operators, spectra, measurement, examples, and interpretations. It should be used as a map, then decomposed into mechanism branches before making technical claims.
- **Measurement problem.** The measurement problem is a readout junction. It sits where unitary state transport, detector context, probability assignment, and state update meet. Use four parts: state evolution before measurement, coupling to an apparatus or environment, POVM or projection readout, and the rule used to condition the state after the record.
- **Quantum gravity.** Quantum gravity is a field/boundary junction. It asks whether geometry itself becomes part of the quantum state carrier or remains a realization layer for an operator theory. The missing constructor is explicit: a state of geometry, a constraint or evolution operator, a boundary or semiclassical readout, and a test showing which geometric quantities survive quantization.
- **Scattering.** Scattering is a boundary-to-spectrum mechanism. The relevant object is the map from asymptotic in-states to out-states. The constructor should specify the Hamiltonian or interaction region, the boundary/asymptotic channels, the S-matrix or cross-section readout, and the conservation constraints.
- **Quantum state.** Quantum state is the carrier, not the final prediction. It is anomalous because it precedes several downstream roles: admissibility, evolution, observable choice, and probability readout. A clean derivation should state whether the carrier is a vector, density operator, field state, or register, and which transformations preserve its legality.
- **Wave–particle duality.** Wave-particle duality is a representation/readout switch. The same carrier is interrogated through incompatible experimental contexts, so the observed pattern changes from interference-like to count-like. The constructor should be written as context selection plus readout channel.
- **Quantum entanglement.** Entanglement is a tensor-factorization and correlation constraint. The anomaly is that the state is not reducible to independently readable subsystem states, while the readout is still local and spectral. A useful page must distinguish the joint state, the subsystem observables, and the correlation test.
- **Fermi–Dirac statistics.** Fermi-Dirac statistics is an admissibility rule for many-particle states. The central mechanism is antisymmetry and occupation restriction, not an ordinary eigenvalue list. The constructor should expose anticommutation, exclusion, occupation numbers, and the thermodynamic readout derived from that constrained state space.
- **Hamiltonian (quantum mechanics).** The Hamiltonian has two roles at once: it generates time evolution and, when treated as an observable, supplies an energy spectrum. That double role explains the anomaly. A clean page must separate domain/self-adjointness, unitary transport, conserved energy, and spectral readout.
- **Wave function.** The wave function is a representation of the state carrier, not a material wave by itself. Its anomaly is that it stores amplitude, phase, normalization, basis choice, and probability potential in one object. The constructor should separate representation, admissibility, evolution, and Born readout.
- **Delayed-choice quantum eraser.** The delayed-choice eraser is a protocol-order stress test. Its mechanism is the arrangement of which-path information, later measurement

choice, and conditional correlation readout. The anomaly is not retrocausality by default; it is that the relevant statistics are defined only after the full measurement protocol is specified.

- **Relativistic quantum mechanics.** Relativistic quantum mechanics is a compatibility junction between quantum state evolution and spacetime symmetry. Its construction preserves relativistic covariance, defines the correct state carrier, and explains how spin, energy, and causality constraints enter the operator algebra.
- **Quantum electrodynamics.** Quantum electrodynamics is a field-interaction constructor. Its anomaly comes from combining gauge admissibility, charged matter states, photon modes, perturbative transport, and scattering/readout. The page should be derived through field operators, gauge constraints, interaction terms, and observable amplitudes.
- **Quantum field theory.** This field-level page mixes operator-to-spectrum readout, state evolution, normalization or admissibility. It should be treated as a many-mode or geometric realization problem: first identify the state sector or field algebra, then the constraints and readout that make the field content observable.
- **Bell’s theorem.** This incompatibility page mixes operator-to-spectrum readout, state evolution, normalization or admissibility. Its mechanism should state which otherwise legal questions fail to share a single sharp representation, and what experiment or inequality exposes that failure.

6.2 Research Leads

- Search for systems where a state-like carrier and a legal-question operator exist, but the incompatibility relation has not been tested. Those systems are candidates for quantum-like contextual behavior without importing quantum ontology.
- Treat tunnelling, particle-in-a-box, cavity optics, and spectral lines as one family of boundary-shaped spectra. This suggests looking for overlooked boundary controls in systems currently described only by bulk evolution.
- Quantum computing should be read as an engineering layer over the state-operator-readout constructor. New protocols should be searched by composing lawful quantum questions and controlled maps.
- Pages that are branch-ambiguous are useful: they often mark junctions where two constructions meet, such as field theory joining transport, incompatibility, and boundary context.
- Historical, interpretive, and object-name pages should be demoted to annotations. The conceptual spine is context, state, generator, spectral question, probability, compatibility, realization.

Part II

Branch Chapters

Chapter 7

Hilbert-space context: admissible carrier and basis

A quantum calculation first fixes the Hilbert space, operator domain, basis, preparation context, representation, gauge, or boundary condition. This is not the measured answer; it is the legal carrier on which states, transformations, observables, and probabilities can be defined.

7.1 Why This Branch Exists

This branch is the first step because quantum mechanics is not defined on raw objects. It begins by specifying the space, basis, representation, or admissibility condition in which states and questions make sense.

7.2 Page Map

Page	Dominant evidence signal	Score
Mathematical formulation of quantum mechanics	observables and spectra	1.41
Hilbert space	observables and spectra	1.40
Transformation theory (quantum mechanics)	observables and spectra	1.14
Quantum differential calculus	observables and spectra	0.88
Quantum complexity theory	preparation and boundary context	0.75
Quantum cellular automaton	preparation and boundary context	0.71
Relativistic quantum mechanics	state evolution	0.71

7.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

7.4 Mathematical formulation of quantum mechanics

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 1.41. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a mathematical formulation of quantum mechanics modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a mathematical formulation of quantum mechanics acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Mathematical formulation of quantum mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** State terms: quantum state and wave function. Context/domain terms: basis.
- **Operator or map.** Operator terms: matrix, Hamiltonian, or unitary. Protocol or update terms: Path integral.
- **Admissibility.** Compatibility or closure terms: uncertainty and complementarity. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.25, constraint closure=0.18

- [arXiv:1604.06537](#), score 0.558
- [arXiv:1604.05385](#), score 0.537
- [arXiv:0912.2823](#), score 0.535
- [arXiv:1612.00682](#), score 0.529
- [arXiv:quant-ph0205159](#), score 0.528

7.5 Hilbert space

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 1.40. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Hilbert space is the admissible state carrier of quantum theory: it supplies the space in which states, operators, bases, spectra, and probabilities become legally defined.

Mechanism Reading

Hilbert space is not physical space and not a geometric background in this book. It is the legal carrier of quantum identity. A state is a vector or density operator on it; the inner product gives amplitudes and norms; observables are self-adjoint operators on it; spectral projectors define possible answers; and unitary evolution preserves norm and probability. Hilbert space is therefore central because it binds state, probability, operator spectrum, and identity preservation into one formal carrier. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Hilbert space contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** A complex Hilbert space, or a density-operator state space built on it.
- **Operator or map.** Self-adjoint observables, unitary maps, spectral projectors, and domain-restricted generators defined on the carrier.
- **Admissibility.** Inner-product structure, normalization, positivity for density states, and operator-domain conditions make states and observables legal.
- **Readout.** Born probabilities, spectral projectors, expectation values, and preserved norms.
- **Check.** Changing basis or representation should preserve probabilities and expectation values when the change is unitary.

Topic Equations

Standard constructor skeleton: normalized states, density states, spectral resolution, Born readout, and unitary identity preservation.

$$\begin{aligned} |\psi\rangle &\in \mathcal{H}, & \langle\psi|\psi\rangle &= 1 \\ \rho &\in \mathcal{S}(\mathcal{H}), & \rho &\geq 0, \quad \text{Tr } \rho = 1 \\ A &= A^\dagger, & A &= \int_{\sigma(A)} \lambda dE_A(\lambda) \\ \text{Pr}(\Delta \mid \rho, A) &= \text{Tr}(\rho E_A(\Delta)) \\ \rho_t &= U(t)\rho U(t)^\dagger, & U^\dagger U &= I \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.13, constraint closure=0.09

- [arXiv:1604.05385](#), score 0.416
- [arXiv:1612.00682](#), score 0.384
- [arXiv:2308.15676](#), score 0.382
- [arXiv:2108.07838](#), score 0.375
- [arXiv:0809.5271](#), score 0.344

7.6 Transformation theory (quantum mechanics)

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 1.14. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A transformation theory (quantum mechanics) can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a transformation theory (quantum mechanics) is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is

Quantum Mechanism Frame

- **Role.** Transformation theory (quantum mechanics) contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.24, constraint closure=0.18

- [arXiv:1801.03283](#), score 0.537
- [arXiv:1708.03640](#), score 0.536
- [arXiv:1612.00682](#), score 0.534
- [arXiv:quant-ph0205159](#), score 0.534
- [arXiv:1506.05598](#), score 0.534

7.7 Quantum differential calculus

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 0.88. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum differential calculus can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum differential calculus is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or

Quantum Mechanism Frame

- **Role.** Quantum differential calculus contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.19, constraint closure=0.15

- [arXiv:0912.2823](#), score 0.509
- [arXiv:1604.06537](#), score 0.499
- [arXiv:1612.00682](#), score 0.478
- [arXiv:1801.03283](#), score 0.473
- [arXiv:quant-ph0205159](#), score 0.469

7.8 Quantum complexity theory

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 0.75. **Dominant evidence signal.** preparation and boundary context. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum complexity theory can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum complexity theory is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through state vector or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is preparation, basis, or

Quantum Mechanism Frame

- **Role.** Quantum complexity theory contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** State terms: state vector. Context/domain terms: context.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

boundary weak form=0.25, transport flow=0.17, constraint closure=0.16

- [arXiv:0912.2823](#), score 0.520
- [arXiv:1706.03846](#), score 0.515
- [arXiv:1708.03640](#), score 0.499
- [arXiv:1605.07654](#), score 0.494
- [arXiv:0908.0752](#), score 0.492

7.9 Quantum cellular automaton

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 0.71. **Dominant evidence signal.** preparation and boundary context. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum cellular automaton can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum cellular automaton is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, unitary operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is preparation, basis, or

Quantum Mechanism Frame

- **Role.** Quantum cellular automaton contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: context.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

boundary weak form=0.28, transport flow=0.26, constraint closure=0.15

- [arXiv:0912.2823](#), score 0.512
- [arXiv:1109.3239](#), score 0.490
- [arXiv:gr-qc0411110](#), score 0.487
- [arXiv:1706.03846](#), score 0.486
- [arXiv:astro-ph0604157](#), score 0.480

7.10 Relativistic quantum mechanics

Mechanism-tree role. Hilbert-space context: admissible carrier and basis; assignment score 0.71. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A relativistic quantum mechanics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a relativistic quantum mechanics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wavefunction or wave function, Hamiltonian or matrix, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Relativistic quantum mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a context-setting move: it fixes the arena in which states, domains, and questions are legal.
- **Carrier or domain.** State terms: wavefunction. Context/domain terms: potential and context.
- **Operator or map.** Operator terms: Hamiltonian and matrix. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, spectral operator=0.25, constraint closure=0.21

- [arXiv:0912.2823](#), score 0.550
- [arXiv:1604.06537](#), score 0.526
- [arXiv:1604.05385](#), score 0.523
- [arXiv:1801.03283](#), score 0.517
- [arXiv:0908.0752](#), score 0.517

Chapter 8

State carrier inside Hilbert space

A state is the probability-bearing element of the selected Hilbert space or its density-operator state space. Wave functions, density matrices, superpositions, and coherent states are different representations of this predictive carrier.

8.1 Why This Branch Exists

The state branch should be introduced before particles. It is the predictive carrier; particles, waves, fields, and qubits are later realizations of that carrier.

8.2 Page Map

Page	Dominant evidence signal	Score
Density matrix	observables and spectra	1.54
Quantum superposition	observables and spectra	1.53
Quantum decoherence	observables and spectra	1.52
Superposition principle	observables and spectra	1.51
Coherence (physics)	observables and spectra	1.50
Wave function	state evolution	1.48
Quantum state	state evolution	1.46
Two-state quantum system	observables and spectra	1.44
Qubit	observables and spectra	1.43
Quantum fluctuation	state evolution	1.14
Bose–Einstein statistics	observables and spectra	0.86
Quantum number	observables and spectra	0.86
Fourier transform	state evolution	0.86
Wave–particle duality	state evolution	0.85
Quantum statistical mechanics	observables and spectra	0.83
Quantum machine	observables and spectra	0.82
Quantum chemistry	observables and spectra	0.82
Applications of quantum mechanics	observables and spectra	0.73
Quantum machine learning	observables and spectra	0.73
Standard Model	observables and spectra	0.73
Spin (physics)	observables and spectra	0.73
Quantum simulator	observables and spectra	0.72

Page	Dominant evidence signal	Score
Werner Heisenberg	state evolution	0.69
Schrödinger's cat	state evolution	0.64

8.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

8.4 Density matrix

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.54. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Density matrix is the mixed-state constructor: it keeps probabilistic preparation, entanglement with unobserved degrees of freedom, and partial information in the same state formalism.

Mechanism Reading

Density matrices generalize pure states without changing the state-to-spectrum readout rule. They are the correct carrier when the preparation is statistical, when a subsystem is traced out, or when decoherence is being described. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Density matrix contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** The mathematical state object: vector, wavefunction, density operator, coherent state, field state, or register.
- **Operator or map.** Operators, maps, and observables become meaningful only after this carrier and its domain have been fixed.
- **Admissibility.** Normalization, positivity, inner product, representation, tensor factorization, or superselection conditions define legal states.
- **Readout.** Probability distributions obtained by applying the appropriate observables or measurement maps to the carrier.
- **Check.** Equivalent representations must preserve probabilities and expectation values when the change is only representational.

Topic Equations

Standard constructor skeleton: mixed state, trace rule, and subsystem reduction.

$$\begin{aligned}\rho &= \sum_a p_a |\psi_a\rangle\langle\psi_a|, & p_a &\geq 0, & \sum_a p_a &= 1 \\ \rho &\geq 0, & \text{Tr } \rho &= 1 \\ p(i) &= \text{Tr}(\rho P_i) \\ \rho_A &= \text{Tr}_B(\rho_{AB})\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.24, constraint closure=0.17

- [arXiv:1604.05385](#), score 0.512
- [arXiv:0912.2823](#), score 0.493
- [arXiv:1612.00682](#), score 0.486
- [arXiv:1801.03283](#), score 0.481
- [arXiv:1506.05598](#), score 0.473

8.5 Quantum superposition

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.53. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum superposition can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum superposition is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, observable operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum superposition contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, or superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: observable and Hamiltonian. Protocol or update terms: Born rule.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.22, constraint closure=0.15

- [arXiv:1604.05385](#), score 0.602
- [arXiv:1604.06537](#), score 0.568
- [arXiv:1612.00682](#), score 0.551
- [arXiv:1801.03283](#), score 0.550
- [arXiv:quant-ph0205159](#), score 0.544

8.6 Quantum decoherence

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.52. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a quantum decoherence modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum decoherence acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum decoherence contributes an open-system transport and coherence role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, wavefunction, density matrix, state vector, or superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: matrix and unitary. Protocol or update terms: Born rule, collapse, or projection.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.37, transport flow=0.27, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.520
- [arXiv:0912.2823](#), score 0.519
- [arXiv:1604.06537](#), score 0.493
- [arXiv:1612.00682](#), score 0.482
- [arXiv:quant-ph0205159](#), score 0.478

8.7 Superposition principle

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.51. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The superposition principle can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Superposition principle is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wave function or Superposition, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Superposition principle contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: wave function and Superposition. Context/domain terms: domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.31, transport flow=0.28, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.547
- [arXiv:1801.03283](#), score 0.528
- [arXiv:quant-ph0205159](#), score 0.525
- [arXiv:1612.00682](#), score 0.521
- [arXiv:1708.03640](#), score 0.520

8.8 Coherence (physics)

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.50. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A coherence (physics) can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a coherence (physics) is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, unitary operator or Hamiltonian, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, preparation, basis, or boundary context; the

Quantum Mechanism Frame

- **Role.** Coherence (physics) contributes an open-system transport and coherence role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential, detector, or basis.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.37, transport flow=0.18, boundary weak form=0.12

- [arXiv:1604.05385](#), score 0.514
- [arXiv:0912.2823](#), score 0.512
- [arXiv:1612.00682](#), score 0.471
- [arXiv:quant-ph0205159](#), score 0.460
- [arXiv:2108.07838](#), score 0.459

8.9 Wave function

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.48. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Wave function is a representation of the state carrier in a chosen basis, not a separate physical substance.

Mechanism Reading

The wave function is the coordinate expression of a quantum state after a representation has been chosen. Its modulus squared gives a probability density in the position representation, while operators act as transformations on that representation. In the source-evidence profile for this page, the strongest construction signal is state evolution, operator-to-spectrum readout, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Wave function contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** The mathematical state object: vector, wavefunction, density operator, coherent state, field state, or register.
- **Operator or map.** Operators, maps, and observables become meaningful only after this carrier and its domain have been fixed.
- **Admissibility.** Normalization, positivity, inner product, representation, tensor factorization, or superselection conditions define legal states.
- **Readout.** Probability distributions obtained by applying the appropriate observables or measurement maps to the carrier.
- **Check.** Equivalent representations must preserve probabilities and expectation values when the change is only representational.

Topic Equations

Standard constructor skeleton: state representation, normalization, and expectation value.

$$\begin{aligned}\psi(x) &= \langle x | \psi \rangle \\ \int |\psi(x)|^2 dx &= 1 \\ \langle A \rangle_\psi &= \langle \psi | A | \psi \rangle\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, spectral operator=0.25, constraint closure=0.21

- [arXiv:1604.05385](#), score 0.513
- [arXiv:0912.2823](#), score 0.512
- [arXiv:1604.06537](#), score 0.505
- [arXiv:1801.03283](#), score 0.479
- [arXiv:1708.03640](#), score 0.476

8.10 Quantum state

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.46. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum state can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum state is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through Quantum state or wave function, observable operator or matrix, and eigenstate or spectrum. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or incompatible observables. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Quantum state contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: Quantum state, wave function, or superposition. Context/domain terms: detector and preparation.
- **Operator or map.** Operator terms: observable and matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty and incompatible. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenstate, spectrum, or eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels

- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, spectral operator=0.22, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.539
- [arXiv:0912.2823](#), score 0.534
- [arXiv:1604.06537](#), score 0.511
- [arXiv:quant-ph0205159](#), score 0.495
- [arXiv:1708.03640](#), score 0.495

8.11 Two-state quantum system

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.44. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A two-state quantum system can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a two-state quantum system is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wavefunction or state vector, observable operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Two-state quantum system contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: wavefunction, state vector, or superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: observable, Hamiltonian, or matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.23, constraint closure=0.17

- [arXiv:1801.03283](#), score 0.611
- [arXiv:1604.06537](#), score 0.598
- [arXiv:quant-ph0205159](#), score 0.597
- [arXiv:1612.00682](#), score 0.596
- [arXiv:1506.05598](#), score 0.586

8.12 Qubit

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.43. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Qubit is the two-dimensional state-carrier constructor used when the admissible state space is $\mathbb{C}^2$.

Mechanism Reading

A qubit is the minimal quantum state space with a basis, amplitudes, unitary control, and measurement readout. Bloch-vector language is a representation of the same two-dimensional carrier. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Qubit contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** The mathematical state object: vector, wavefunction, density operator, coherent state, field state, or register.
- **Operator or map.** Operators, maps, and observables become meaningful only after this carrier and its domain have been fixed.
- **Admissibility.** Normalization, positivity, inner product, representation, tensor factorization, or superselection conditions define legal states.
- **Readout.** Probability distributions obtained by applying the appropriate observables or measurement maps to the carrier.
- **Check.** Equivalent representations must preserve probabilities and expectation values when the change is only representational.

Topic Equations

Standard constructor skeleton: two-state carrier, Bloch representation, and basis readout.

$$\begin{aligned} |\psi\rangle &= \alpha|0\rangle + \beta|1\rangle, & |\alpha|^2 + |\beta|^2 &= 1 \\ \rho &= \frac{1}{2}(I + \mathbf{r} \cdot \boldsymbol{\sigma}), & |\mathbf{r}| &\leq 1 \\ p(0) &= |\langle 0|\psi\rangle|^2, & p(1) &= |\langle 1|\psi\rangle|^2 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.23, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.577
- [arXiv:1801.03283](#), score 0.522
- [arXiv:1612.00682](#), score 0.519
- [arXiv:quant-ph0205159](#), score 0.511
- [arXiv:1506.05598](#), score 0.511

8.13 Quantum fluctuation

Mechanism-tree role. State carrier inside Hilbert space; assignment score 1.14. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum fluctuation can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum fluctuation is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral

decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is state evolution, non-commuting compatibility limits, normalization or admissibility;

Quantum Mechanism Frame

- **Role.** Quantum fluctuation contributes an open-system transport and coherence role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, commutator incompatibility=0.24, constraint closure=0.21

- [arXiv:1604.06537](#), score 0.476
- [arXiv:1706.03846](#), score 0.474
- [arXiv:0908.0752](#), score 0.462
- [arXiv:1604.05385](#), score 0.455
- [arXiv:2106.04271](#), score 0.455

8.14 Bose–Einstein statistics

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.86. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A bose–einstein statistics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a bose–Einstein statistics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and

energy level or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Bose–Einstein statistics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: energy level. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.12, boundary weak form=0.11

- [arXiv:0912.2823](#), score 0.523
- [arXiv:2308.15676](#), score 0.495
- [arXiv:2105.11733](#), score 0.484
- [arXiv:2108.07838](#), score 0.482
- [arXiv:0805.4565](#), score 0.480

8.15 Quantum number

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.86. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum number can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum number is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, observable operator or Hamiltonian, and spectrum or energy

level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum number contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: observable and Hamiltonian. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum and energy level. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.36, transport flow=0.26, constraint closure=0.20

- [arXiv:1604.05385](#), score 0.538
- [arXiv:1604.06537](#), score 0.535
- [arXiv:0912.2823](#), score 0.512
- [arXiv:1612.00682](#), score 0.507
- [arXiv:1801.03283](#), score 0.506

8.16 Fourier transform

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.86. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A fourier transform can be read as a quantum construction: the input encoding, circuit architecture, and final measurement basis fixes the admissible state space; a sequence of unitary gates or quantum channels defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a fourier transform is described by a register state in a finite-dimensional Hilbert space. The physical question is represented by a sequence of unitary gates or quantum channels; the experimental or mathematical setting is the input encoding, circuit architecture, and final measurement basis. The observable content is obtained from measurement probabilities over computational-basis outcomes. In the local terminology of this topic, the same construction appears through quantum state or wave function, unitary operator or

Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or admissibility, non-commuting compatibility limits; the

Quantum Mechanism Frame

- **Role.** Fourier transform contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: domain.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, constraint closure=0.22, commutator incompatibility=0.20

- [arXiv:1706.03846](#), score 0.563
- [arXiv:1708.03640](#), score 0.539
- [arXiv:0912.2823](#), score 0.538
- [arXiv:1604.06537](#), score 0.535
- [arXiv:0908.0752](#), score 0.533

8.17 Wave–particle duality

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.85. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Wave–particle duality can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, wave–particle duality is described by a wave function or density operator defined on the Hilbert space allowed by the system’s domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through probability amplitude or wave function, Hamiltonian or observable operator, and spectrum or

eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Wave–particle duality contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: probability amplitude. Context/domain terms: potential and detector.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, spectral operator=0.22, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.533
- [arXiv:0912.2823](#), score 0.532
- [arXiv:1604.06537](#), score 0.503
- [arXiv:1708.03640](#), score 0.495
- [arXiv:quant-ph0205159](#), score 0.492

8.18 Quantum statistical mechanics

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.83. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum statistical mechanics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum statistical mechanics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same

construction appears through quantum state or density matrix, observable operator or matrix, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum statistical mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state and density matrix. Context/domain terms: basis.
- **Operator or map.** Operator terms: observable, matrix, or Hamiltonian. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum and eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.17

- [arXiv:1604.05385](#), score 0.592
- [arXiv:0912.2823](#), score 0.589
- [arXiv:1506.05598](#), score 0.582
- [arXiv:1801.03283](#), score 0.582
- [arXiv:1612.00682](#), score 0.581

8.19 Quantum machine

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.82. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum machine can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum machine is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or superposition, observable operator or Hamiltonian, and eigenvalue or energy

level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Quantum machine contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state and superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: observable. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.20, discrete protocol=0.12

- [arXiv:1604.05385](#), score 0.531
- [arXiv:0805.4565](#), score 0.501
- [arXiv:2108.07838](#), score 0.500
- [arXiv:1612.00682](#), score 0.497
- [arXiv:quant-ph0205159](#), score 0.489

8.20 Quantum chemistry

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.82. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum chemistry can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum chemistry is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wave function or quantum state, observable operator or Hamiltonian, and eigenvalue or energy

level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Quantum chemistry contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: wave function and quantum state. Context/domain terms: potential and basis.
- **Operator or map.** Operator terms: observable and Hamiltonian. Protocol or update terms: path integral.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.37, transport flow=0.26, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.562
- [arXiv:1612.00682](#), score 0.523
- [arXiv:quant-ph0205159](#), score 0.516
- [arXiv:1801.03283](#), score 0.512
- [arXiv:1506.05598](#), score 0.508

8.21 Applications of quantum mechanics

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.73. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

An applications of quantum mechanics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, an applications of quantum mechanics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the

two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through superposition or wave function, Hamiltonian or observable operator, and spectrum or eigenstate. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is

Quantum Mechanism Frame

- **Role.** Applications of quantum mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: superposition. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum and eigenstate. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.33, transport flow=0.21, constraint closure=0.12

- [arXiv:1604.05385](#), score 0.549
- [arXiv:0912.2823](#), score 0.548
- [arXiv:quant-ph0205159](#), score 0.524
- [arXiv:0805.4565](#), score 0.523
- [arXiv:1612.00682](#), score 0.518

8.22 Quantum machine learning

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.73. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum machine learning can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum machine learning is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or superposition, matrix or Hamiltonian, and eigenvalue or energy level.

Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum machine learning contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state and superposition. Context/domain terms: preparation and basis.
- **Operator or map.** Operator terms: matrix, Hamiltonian, or unitary. Protocol or update terms: unitary evolution.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.33, transport flow=0.26, constraint closure=0.17

- [arXiv:0912.2823](#), score 0.550
- [arXiv:1604.05385](#), score 0.550
- [arXiv:1612.00682](#), score 0.517
- [arXiv:quant-ph0205159](#), score 0.513
- [arXiv:1801.03283](#), score 0.509

8.23 Standard Model

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.73. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The standard model can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Standard Model is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy

level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Standard Model contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.32, transport flow=0.26, constraint closure=0.21

- [arXiv:0912.2823](#), score 0.495
- [arXiv:1604.06537](#), score 0.491
- [arXiv:1801.03283](#), score 0.474
- [arXiv:quant-ph0205159](#), score 0.468
- [arXiv:1612.00682](#), score 0.467

8.24 Spin (physics)

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.73. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A spin (physics) can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a spin (physics) is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wave function or quantum state, Hamiltonian or observable operator, and eigenvalue or energy level.

Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Spin (physics) contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: wave function and quantum state. Context/domain terms: context.
- **Operator or map.** Operator terms: Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.27, constraint closure=0.20

- [arXiv:0912.2823](#), score 0.528
- [arXiv:1604.05385](#), score 0.527
- [arXiv:quant-ph0205159](#), score 0.509
- [arXiv:1801.03283](#), score 0.508
- [arXiv:1708.03640](#), score 0.507

8.25 Quantum simulator

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.72. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum simulator can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum simulator is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through superposition or wave function, Hamiltonian or unitary operator, and eigenvalue or energy level.

Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Quantum simulator contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian and unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.28, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.526
- [arXiv:1604.05385](#), score 0.526
- [arXiv:quant-ph0205159](#), score 0.506
- [arXiv:1708.03640](#), score 0.504
- [arXiv:1801.03283](#), score 0.503

8.26 Werner Heisenberg

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.69. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A werner heisenberg can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a werner Heisenberg is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through

wave function or state vector, matrix or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or non-commuting observables. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Werner Heisenberg contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: wave function. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: uncertainty and non-commuting. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, spectral operator=0.25, commutator incompatibility=0.22

- [arXiv:1604.06537](#), score 0.524
- [arXiv:1604.05385](#), score 0.505
- [arXiv:0912.2823](#), score 0.503
- [arXiv:1708.03640](#), score 0.485
- [arXiv:1801.03283](#), score 0.484

8.27 Schrödinger's cat

Mechanism-tree role. State carrier inside Hilbert space; assignment score 0.64. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a schrödinger's cat modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a schrödinger's cat acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is state evolution, operator-to-spectrum readout, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Schrödinger's cat contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a state-carrier move: it specifies what mathematical object carries prediction.
- **Carrier or domain.** State terms: wave function, quantum state, wavefunction, or superposition. Context/domain terms: domain.
- **Operator or map.** Operator terms: observable. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, spectral operator=0.22, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.514
- [arXiv:0912.2823](#), score 0.513
- [arXiv:1604.06537](#), score 0.481
- [arXiv:quant-ph0205159](#), score 0.463
- [arXiv:1708.03640](#), score 0.463

Chapter 9

Generator: lawful change before readout

Hamiltonians, unitary maps, equations of motion, and path weights describe the lawful transport of the state before a question is resolved.

9.1 Why This Branch Exists

Time evolution is a transport problem over states. The Hamiltonian and path integral are two views of the same generator role rather than unrelated formalisms.

9.2 Page Map

Page	Dominant evidence signal	Score
Unitary operator	observables and spectra	1.50
Perturbation theory	observables and spectra	1.50
Quantum dynamics	observables and spectra	1.50
Path integral formulation	observables and spectra	1.49
Hamiltonian mechanics	observables and spectra	1.49
Path integral	observables and spectra	1.48
Hamiltonian (quantum mechanics)	state evolution	1.47
Perturbation theory (quantum mechanics)	observables and spectra	1.46
Schrödinger picture	observables and spectra	1.29
Creation and annihilation operators	observables and spectra	1.28
Schrödinger equation	observables and spectra	1.22
Quantum stochastic calculus	observables and spectra	0.85
Quantum chaos	observables and spectra	0.76
Symmetry in quantum mechanics	observables and spectra	0.75
Quantum amplifier	observables and spectra	0.75
Quantum mechanics	observables and spectra	0.75
Heisenberg picture	observables and spectra	0.75
Heisenberg group	observables and spectra	0.74
Canonical commutation relation	observables and spectra	0.74
Quantum biology	state evolution	0.73
Quantum engineering	state evolution	0.72

Page	Dominant evidence signal	Score
Quantum logic	observables and spectra	0.56

9.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

9.4 Unitary operator

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.50. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Unitary operator is the reversible-map constructor: it changes the state while preserving inner products and probabilities.

Mechanism Reading

Unitary maps are the admissible reversible transformations of closed-system quantum theory. They preserve normalization and carry projective geometry of the state space through time or controlled operations. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, spectral profile.

Quantum Mechanism Frame

- **Role.** Unitary operator contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A state vector, density operator, wavefunction, field state, or register on a specified domain.
- **Operator or map.** Hamiltonian, unitary map, channel generator, action, constraint, or differential operator that transports the state.
- **Admissibility.** Self-adjointness, complete positivity, trace preservation, gauge constraints, and boundary/domain conditions decide whether the evolution is legal.
- **Readout.** Time-dependent probabilities, spectra, transition amplitudes, conserved quantities, or response functions.
- **Check.** The generator must predict the observed evolution while preserving the relevant normalization, positivity, symmetry, or conservation constraint.

Topic Equations

Standard constructor skeleton: reversible state transformation and inner-product preservation.

$$\begin{aligned}U^\dagger U &= U U^\dagger = I \\ |\psi'\rangle &= U|\psi\rangle \\ \langle\psi'|\phi'\rangle &= \langle\psi|\phi\rangle\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.44, transport flow=0.24, constraint closure=0.18

- [arXiv:0912.2823](#), score 0.500
- [arXiv:1612.00682](#), score 0.492
- [arXiv:1801.03283](#), score 0.477
- [arXiv:quant-ph0205159](#), score 0.471
- [arXiv:1506.05598](#), score 0.468

9.5 Perturbation theory

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.50.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Perturbation theory can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, perturbation theory is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Perturbation theory contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.21, constraint closure=0.12

- [arXiv:0912.2823](#), score 0.538
- [arXiv:1612.00682](#), score 0.501
- [arXiv:0805.4565](#), score 0.499
- [arXiv:2108.07838](#), score 0.498
- [arXiv:quant-ph0205159](#), score 0.494

9.6 Quantum dynamics

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.50.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum dynamics can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum dynamics is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through state vector or density matrix, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum dynamics contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: state vector, density matrix, quantum state, or superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable, matrix, unitary, or commutator. Protocol or update terms: unitary evolution.
- **Admissibility.** Compatibility or closure terms: commutator and uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.26, constraint closure=0.20

- [arXiv:1604.06537](#), score 0.568
- [arXiv:1801.03283](#), score 0.539
- [arXiv:1612.00682](#), score 0.538
- [arXiv:0912.2823](#), score 0.535
- [arXiv:1506.05598](#), score 0.533

9.7 Path integral formulation

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.49. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Path integral formulation is the formulation in which the generator is represented by action-weighted histories.

Mechanism Reading

This page belongs with lawful change because it changes how the transport step is calculated. The observable predictions remain probabilities after amplitudes are composed and squared or traced. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Path integral formulation contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A state vector, density operator, wavefunction, field state, or register on a specified domain.
- **Operator or map.** Hamiltonian, unitary map, channel generator, action, constraint, or differential operator that transports the state.
- **Admissibility.** Self-adjointness, complete positivity, trace preservation, gauge constraints, and boundary/domain conditions decide whether the evolution is legal.
- **Readout.** Time-dependent probabilities, spectra, transition amplitudes, conserved quantities, or response functions.
- **Check.** The generator must predict the observed evolution while preserving the relevant normalization, positivity, symmetry, or conservation constraint.

Topic Equations

Standard constructor skeleton: transition amplitudes and generating functional.

$$\langle q_f, t_f | q_i, t_i \rangle = \int \mathcal{D}q e^{iS[q]/\hbar}$$
$$Z[J] = \int \mathcal{D}\phi \exp\left(\frac{i}{\hbar}(S[\phi] + \int J\phi)\right)$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.34, transport flow=0.25, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.548
- [arXiv:1604.06537](#), score 0.545
- [arXiv:1801.03283](#), score 0.516
- [arXiv:1612.00682](#), score 0.515
- [arXiv:quant-ph0205159](#), score 0.514

9.8 Hamiltonian mechanics

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.49.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A hamiltonian mechanics can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a hamiltonian mechanics is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Hamiltonian mechanics contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.21, constraint closure=0.14

- [arXiv:1604.06537](#), score 0.481
- [arXiv:2108.07838](#), score 0.467
- [arXiv:1506.05598](#), score 0.464
- [arXiv:1612.00682](#), score 0.462
- [arXiv:2111.12617](#), score 0.454

9.9 Path integral

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.48.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Path integral is an alternate generator constructor: transition amplitudes are obtained by summing phase weights over histories.

Mechanism Reading

The path integral does not replace the operator constructor. It repackages the generator step as a weighted sum over histories between boundary conditions. It is especially useful when action, symmetry, and field degrees of freedom are more natural than state-vector evolution. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Path integral contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A state vector, density operator, wavefunction, field state, or register on a specified domain.
- **Operator or map.** Hamiltonian, unitary map, channel generator, action, constraint, or differential operator that transports the state.
- **Admissibility.** Self-adjointness, complete positivity, trace preservation, gauge constraints, and boundary/domain conditions decide whether the evolution is legal.
- **Readout.** Time-dependent probabilities, spectra, transition amplitudes, conserved quantities, or response functions.
- **Check.** The generator must predict the observed evolution while preserving the relevant normalization, positivity, symmetry, or conservation constraint.

Topic Equations

Standard constructor skeleton: boundary-to-boundary transition amplitude through action weights.

$$K(x_f, t_f; x_i, t_i) = \int_{x_i}^{x_f} \mathcal{D}x(t) \exp\left(\frac{i}{\hbar} S[x]\right)$$
$$\psi(x_f, t_f) = \int K(x_f, t_f; x_i, t_i) \psi(x_i, t_i) dx_i$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.24, transport flow=0.11, constraint closure=0.07

- [arXiv:2107.01923](#), score 0.313
- [arXiv:0801.3568](#), score 0.272
- [arXiv:2105.11733](#), score 0.271
- [arXiv:1706.07300](#), score 0.264
- [arXiv:cond-mat0108470](#), score 0.263

9.10 Hamiltonian (quantum mechanics)

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.47.
Dominant evidence signal. state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Hamiltonian (quantum mechanics) is the generator observable: it both transports states and supplies the energy spectrum.

Mechanism Reading

The Hamiltonian has a dual role. Dynamically, it generates unitary time evolution. Spectrally, its eigenvalues are admissible energy readouts. This dual role is one reason the operator/spectrum branch is central. In the source-evidence profile for this page, the strongest construction signal is state evolution, operator-to-spectrum readout, preparation, basis, or boundary context; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Hamiltonian (quantum mechanics) contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A state vector, density operator, wavefunction, field state, or register on a specified domain.
- **Operator or map.** Hamiltonian, unitary map, channel generator, action, constraint, or differential operator that transports the state.
- **Admissibility.** Self-adjointness, complete positivity, trace preservation, gauge constraints, and boundary/domain conditions decide whether the evolution is legal.
- **Readout.** Time-dependent probabilities, spectra, transition amplitudes, conserved quantities, or response functions.
- **Check.** The generator must predict the observed evolution while preserving the relevant normalization, positivity, symmetry, or conservation constraint.

Topic Equations

Standard constructor skeleton: energy spectrum and unitary generation.

$$\begin{aligned}H|E_n\rangle &= E_n|E_n\rangle \\U(t) &= e^{-iHt/\hbar} \\ \rho(t) &= U(t)\rho(0)U(t)^\dagger\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, spectral operator=0.24, boundary weak form=0.23

- [arXiv:0912.2823](#), score 0.502
- [arXiv:1604.05385](#), score 0.465
- [arXiv:0908.0752](#), score 0.447
- [arXiv:2108.07838](#), score 0.447
- [arXiv:1612.00682](#), score 0.446

9.11 Perturbation theory (quantum mechanics)

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.46.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A perturbation theory (quantum mechanics) can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a perturbation theory (quantum mechanics) is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is

Quantum Mechanism Frame

- **Role.** Perturbation theory (quantum mechanics) contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.26, boundary weak form=0.24, transport flow=0.22

- [arXiv:0912.2823](#), score 0.509
- [arXiv:1604.05385](#), score 0.475
- [arXiv:1612.00682](#), score 0.449
- [arXiv:1506.05598](#), score 0.436
- [arXiv:quant-ph0205159](#), score 0.436

9.12 Schrödinger picture

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.29. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The schrödinger picture can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Schrödinger picture is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through state vector or wavefunction, Hamiltonian or unitary operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Schrödinger picture contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: state vector and wavefunction. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian and unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.27, transport flow=0.27, boundary weak form=0.20

- [arXiv:0912.2823](#), score 0.464
- [arXiv:1604.06537](#), score 0.442
- [arXiv:1801.03283](#), score 0.435
- [arXiv:1612.00682](#), score 0.418
- [arXiv:1506.05598](#), score 0.414

9.13 Creation and annihilation operators

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.28.

Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Creation and annihilation operators are sector-changing operators: they add or remove one quantum from a mode and make many-body or field descriptions executable.

Mechanism Reading

The page is about the algebraic move that changes occupation number. Creation raises the population of a mode, annihilation lowers it, and the commutation or anticommutation rule determines the statistics. The number operator gives the spectral readout. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, non-commuting compatibility limits; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Creation and annihilation operators contributes an unresolved constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A context-selected state space or effective carrier for prediction.
- **Operator or map.** The relevant Hamiltonian, observable, channel, constraint, or update map.

- **Admissibility.** Domain, normalization, positivity, compatibility, boundary, or gauge requirements state what is legal.
- **Readout.** The outcome probabilities, spectra, correlations, amplitudes, or records used to test the mechanism.
- **Check.** A complete account must specify state carrier, operator or map, admissibility condition, readout, and at least one possible falsifier.

Topic Equations

Topic-specific constructor: the equations express raising, lowering, and occupation-number readout.

$$\begin{aligned} a_i^\dagger |\dots, n_i, \dots\rangle &= \sqrt{n_i + 1} |\dots, n_i + 1, \dots\rangle \\ a_i |\dots, n_i, \dots\rangle &= \sqrt{n_i} |\dots, n_i - 1, \dots\rangle \\ N_i &= a_i^\dagger a_i \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.27, transport flow=0.26, commutator incompatibility=0.21

- [arXiv:1604.06537](#), score 0.536
- [arXiv:0912.2823](#), score 0.509
- [arXiv:1801.03283](#), score 0.505
- [arXiv:2111.12617](#), score 0.503
- [arXiv:1506.05598](#), score 0.502

9.14 Schrödinger equation

Mechanism-tree role. Generator: lawful change before readout; assignment score 1.22.

Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Schrödinger equation is the state-transport constructor: the Hamiltonian generates lawful change of the state before readout.

Mechanism Reading

The Schrödinger equation is not a measurement rule. It is the generator step of the quantum constructor. It evolves the predictive carrier while preserving normalization when the Hamiltonian is self-adjoint. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Schrödinger equation contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A state vector, density operator, wavefunction, field state, or register on a specified domain.
- **Operator or map.** Hamiltonian, unitary map, channel generator, action, constraint, or differential operator that transports the state.
- **Admissibility.** Self-adjointness, complete positivity, trace preservation, gauge constraints, and boundary/domain conditions decide whether the evolution is legal.
- **Readout.** Time-dependent probabilities, spectra, transition amplitudes, conserved quantities, or response functions.
- **Check.** The generator must predict the observed evolution while preserving the relevant normalization, positivity, symmetry, or conservation constraint.

Topic Equations

Standard constructor skeleton: Hamiltonian transport and norm preservation.

$$\begin{aligned}i\hbar \partial_t |\psi(t)\rangle &= H |\psi(t)\rangle \\ |\psi(t)\rangle &= U(t) |\psi(0)\rangle, \quad U(t) = e^{-iHt/\hbar} \\ \frac{d}{dt} \langle \psi(t) | \psi(t) \rangle &= 0 \quad (H = H^\dagger)\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.26, constraint closure=0.18

- [arXiv:0912.2823](#), score 0.567
- [arXiv:1604.06537](#), score 0.551
- [arXiv:1604.05385](#), score 0.548
- [arXiv:1801.03283](#), score 0.539
- [arXiv:1612.00682](#), score 0.536

9.15 Quantum stochastic calculus

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.85.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum stochastic calculus can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum stochastic calculus is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or commutator, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or non-commuting observables. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum stochastic calculus contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian and commutator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.14

- [arXiv:1604.06537](#), score 0.486
- [arXiv:1604.05385](#), score 0.470
- [arXiv:1612.00682](#), score 0.469
- [arXiv:2108.07838](#), score 0.464
- [arXiv:0912.2823](#), score 0.463

9.16 Quantum chaos

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.76.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum chaos can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum chaos is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum chaos contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable, or matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue and energy level. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.26, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.570
- [arXiv:1801.03283](#), score 0.563
- [arXiv:1612.00682](#), score 0.562
- [arXiv:quant-ph0205159](#), score 0.562
- [arXiv:1506.05598](#), score 0.557

9.17 Symmetry in quantum mechanics

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.75.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A symmetry in quantum mechanics can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a symmetry in quantum mechanics is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through wavefunction or quantum state, matrix or generator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through non-commuting observables or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Symmetry in quantum mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: wavefunction and quantum state. Context/domain terms: context.
- **Operator or map.** Operator terms: matrix, generator, unitary, or commutator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: non-commuting and commutator. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.23, constraint closure=0.18

- [arXiv:1604.06537](#), score 0.572
- [arXiv:1801.03283](#), score 0.544
- [arXiv:0912.2823](#), score 0.543
- [arXiv:1506.05598](#), score 0.537
- [arXiv:1612.00682](#), score 0.537

9.18 Quantum amplifier

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.75.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum amplifier can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum amplifier is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or state vector, unitary operator or Hamiltonian, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Quantum amplifier contributes an instrument-mediated readout role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, state vector, or superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.22, constraint closure=0.13

- [arXiv:1604.06537](#), score 0.581
- [arXiv:1604.05385](#), score 0.548
- [arXiv:1612.00682](#), score 0.535
- [arXiv:2108.07838](#), score 0.530
- [arXiv:1506.05598](#), score 0.527

9.19 Quantum mechanics

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.75.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.**
no Lagrangian road map available; road score 0.00.

Central Claim

Quantum mechanics is the baseline constructor: states live in Hilbert space, physical questions are represented by operators, and probabilities are assigned to spectral projectors.

Mechanism Reading

The page supplies the general quantum assembly. A preparation gives a state vector or density operator. A self-adjoint observable or measurement operator family gives the possible readout channels. The Born or trace rule assigns probabilities, while Hamiltonian evolution transports the state between preparation and readout. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum mechanics contributes an unresolved constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** A context-selected state space or effective carrier for prediction.
- **Operator or map.** The relevant Hamiltonian, observable, channel, constraint, or update map.
- **Admissibility.** Domain, normalization, positivity, compatibility, boundary, or gauge requirements state what is legal.
- **Readout.** The outcome probabilities, spectra, correlations, amplitudes, or records used to test the mechanism.
- **Check.** A complete account must specify state carrier, operator or map, admissibility condition, readout, and at least one possible falsifier.

Topic Equations

Topic-specific constructor: the equations express state admissibility, spectral readout, unitary evolution, and incompatibility.

$$\begin{aligned} \rho &\geq 0, & \text{Tr } \rho &= 1 \\ A &= \sum_a a P_a, & p(a) &= \text{Tr}(\rho P_a) \\ \rho(t) &= U(t)\rho(0)U(t)^\dagger, & U(t) &= e^{-iHt/\hbar} \\ [A, B] &\neq 0 & \Rightarrow & \text{no generic common sharp eigenbasis} \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.36, transport flow=0.25, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.531
- [arXiv:0912.2823](#), score 0.526
- [arXiv:1604.06537](#), score 0.524
- [arXiv:1612.00682](#), score 0.499
- [arXiv:1801.03283](#), score 0.497

9.20 Heisenberg picture

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.75.
Dominant evidence signal. observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The heisenberg picture can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Heisenberg picture is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or non-commuting observables. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Heisenberg picture contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable, or commutator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\mapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.23, constraint closure=0.15

- [arXiv:1604.06537](#), score 0.485
- [arXiv:1612.00682](#), score 0.443
- [arXiv:2108.07838](#), score 0.442
- [arXiv:0912.2823](#), score 0.436
- [arXiv:1801.03283](#), score 0.436

9.21 Heisenberg group

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.74. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A heisenberg group can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a heisenberg group is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or unitary operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Heisenberg group contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: context.
- **Operator or map.** Operator terms: matrix and unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.23, constraint closure=0.15

- [arXiv:0912.2823](#), score 0.401
- [arXiv:1604.06537](#), score 0.386
- [arXiv:1612.00682](#), score 0.356
- [arXiv:2108.07838](#), score 0.341
- [arXiv:0805.4565](#), score 0.341

9.22 Canonical commutation relation

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.74. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The canonical commutation relation can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Canonical commutation relation is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or commutator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Canonical commutation relation contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian and commutator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator and uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\mapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.36, transport flow=0.20, commutator incompatibility=0.15

- [arXiv:1604.06537](#), score 0.574
- [arXiv:2111.12617](#), score 0.563
- [arXiv:0912.2823](#), score 0.544
- [arXiv:0908.0752](#), score 0.532
- [arXiv:1506.05598](#), score 0.526

9.23 Quantum biology

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.73.
Dominant evidence signal. state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum biology can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum biology is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through complementarity or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or

Quantum Mechanism Frame

- **Role.** Quantum biology contributes an open-system transport and coherence role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: complementarity and Uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.28, constraint closure=0.19, spectral operator=0.16

- [arXiv:1706.03846](#), score 0.447
- [arXiv:0912.2823](#), score 0.438
- [arXiv:1604.05385](#), score 0.437
- [arXiv:1604.06537](#), score 0.429
- [arXiv:1708.03640](#), score 0.422

9.24 Quantum engineering

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.72. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum engineering can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum engineering is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or superposition, generator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution, operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum engineering contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: quantum state and superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, spectral operator=0.12, constraint closure=0.05

- [arXiv:1706.03846](#), score 0.527
- [arXiv:1708.03640](#), score 0.504
- [arXiv:0805.4565](#), score 0.500
- [arXiv:0908.0752](#), score 0.496
- [arXiv:quant-ph0205159](#), score 0.478

9.25 Quantum logic

Mechanism-tree role. Generator: lawful change before readout; assignment score 0.56. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a quantum logic modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum logic acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum logic contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a lawful-transport move: it identifies what changes the state before readout.
- **Carrier or domain.** State terms: wavefunction and superposition. Context/domain terms: potential and basis.
- **Operator or map.** Operator terms: observable and Hamiltonian. Protocol or update terms: collapse and projection.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.25, constraint closure=0.18

- [arXiv:1604.06537](#), score 0.561
- [arXiv:1604.05385](#), score 0.534
- [arXiv:0912.2823](#), score 0.533
- [arXiv:1612.00682](#), score 0.529
- [arXiv:quant-ph0205159](#), score 0.528

Chapter 10

Spectral question: what can be asked

An observable is a permitted question whose operator form determines the possible numerical answers.

10.1 Why This Branch Exists

The central unit is a legal question posed to a state. Spectra make the possible answers visible.

10.2 Page Map

Page	Dominant evidence signal	Score
Angular momentum operator	observables and spectra	1.48
Observable	observables and spectra	1.47
Self-adjoint operator	observables and spectra	1.47
Spectral theory	observables and spectra	1.41
Pauli matrices	observables and spectra	1.35
Operator theory	observables and spectra	1.34
Operator (physics)	observables and spectra	1.30
Eigenvalues and eigenvectors	observables and spectra	1.48
Von Neumann algebra	observables and spectra	1.43
Electron microscope	observables and spectra	0.67

10.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

10.4 Angular momentum operator

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.48. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The angular momentum operator can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Angular momentum operator is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, observable operator or commutator, and eigenstate or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or non-commuting observables. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or

Quantum Mechanism Frame

- **Role.** Angular momentum operator contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: observable and commutator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenstate. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.19, constraint closure=0.16

- [arXiv:1604.06537](#), score 0.538
- [arXiv:0912.2823](#), score 0.516
- [arXiv:1604.05385](#), score 0.515
- [arXiv:1612.00682](#), score 0.511
- [arXiv:1801.03283](#), score 0.506

10.5 Observable

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.47. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Observable is the legal-question constructor: it turns a physical question into an operator with spectral outcome channels.

Mechanism Reading

An observable is the mathematical form of a question that can be asked of a state. Its spectral decomposition defines the possible answers. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Observable contributes an unresolved constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** A context-selected state space or effective carrier for prediction.
- **Operator or map.** The relevant Hamiltonian, observable, channel, constraint, or update map.
- **Admissibility.** Domain, normalization, positivity, compatibility, boundary, or gauge requirements state what is legal.
- **Readout.** The outcome probabilities, spectra, correlations, amplitudes, or records used to test the mechanism.
- **Check.** A complete account must specify state carrier, operator or map, admissibility condition, readout, and at least one possible falsifier.

Topic Equations

Standard constructor skeleton: self-adjoint question, spectral projectors, and Born probabilities.

$$\begin{aligned}A &= A^\dagger \\A &= \sum_i a_i P_i \\p(a_i) &= \text{Tr}(\rho P_i)\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.23, constraint closure=0.16

- [arXiv:1604.05385](#), score 0.567
- [arXiv:1604.06537](#), score 0.561
- [arXiv:1612.00682](#), score 0.522
- [arXiv:1801.03283](#), score 0.521
- [arXiv:quant-ph0205159](#), score 0.519

10.6 Self-adjoint operator

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.47. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Self-adjoint operator is the admissible-observable condition: it gives real spectra and well-defined spectral measures.

Mechanism Reading

Self-adjointness is not a technical decoration. It is the condition that makes an operator a legitimate spectral question in ordinary quantum mechanics. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Self-adjoint operator contributes an unresolved constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** A context-selected state space or effective carrier for prediction.
- **Operator or map.** The relevant Hamiltonian, observable, channel, constraint, or update map.

- **Admissibility.** Domain, normalization, positivity, compatibility, boundary, or gauge requirements state what is legal.
- **Readout.** The outcome probabilities, spectra, correlations, amplitudes, or records used to test the mechanism.
- **Check.** A complete account must specify state carrier, operator or map, admissibility condition, readout, and at least one possible falsifier.

Topic Equations

Standard constructor skeleton: spectral theorem form of a legitimate observable.

$$A = A^\dagger$$

$$A = \int_{\sigma(A)} \lambda dE_A(\lambda)$$

$$\Pr(\Delta) = \text{Tr}(\rho E_A(\Delta))$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.23, constraint closure=0.17

- [arXiv:0912.2823](#), score 0.569
- [arXiv:1612.00682](#), score 0.553
- [arXiv:1801.03283](#), score 0.550
- [arXiv:quant-ph0205159](#), score 0.546
- [arXiv:1506.05598](#), score 0.541

10.7 Spectral theory

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.41. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Spectral theory can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, spectral theory is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and eigenvalue or spectrum. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Spectral theory contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue and spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.44, transport flow=0.22, constraint closure=0.12

- [arXiv:2108.07838](#), score 0.501
- [arXiv:1612.00682](#), score 0.497
- [arXiv:0805.4565](#), score 0.492
- [arXiv:quant-ph0205159](#), score 0.480
- [arXiv:1506.05598](#), score 0.472

10.8 Pauli matrices

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.35. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The pauli matrices can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Pauli matrices is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Pauli matrices contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis and context.
- **Operator or map.** Operator terms: matrix, observable, or unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
\rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
[O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.11

- [arXiv:0912.2823](#), score 0.564
- [arXiv:2108.07838](#), score 0.551
- [arXiv:1612.00682](#), score 0.550
- [arXiv:0805.4565](#), score 0.547
- [arXiv:1506.05598](#), score 0.537

10.9 Operator theory

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.34. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Operator theory can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, operator theory is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or unitary operator, and Spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Operator theory contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: matrix and unitary. Protocol or update terms: projection.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: Spectrum and eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
\rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
[O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.24, constraint closure=0.18

- [arXiv:0912.2823](#), score 0.550
- [arXiv:1604.05385](#), score 0.549
- [arXiv:1612.00682](#), score 0.541
- [arXiv:1506.05598](#), score 0.534
- [arXiv:quant-ph0205159](#), score 0.533

10.10 Operator (physics)

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.30. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

An operator (physics) can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, an operator (physics) is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through wavefunction or wave function, Hamiltonian or matrix, and eigenvalue or eigenstate. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Operator (physics) contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: wavefunction. Context/domain terms: context.
- **Operator or map.** Operator terms: Hamiltonian, matrix, generator, observable, or unitary. Protocol or update terms: projection.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.37, transport flow=0.26, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.570
- [arXiv:1801.03283](#), score 0.543
- [arXiv:0912.2823](#), score 0.539
- [arXiv:1612.00682](#), score 0.536
- [arXiv:1506.05598](#), score 0.535

10.11 Eigenvalues and eigenvectors

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.48. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

An eigenvalues and eigenvectors can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, an eigenvalues and eigenvectors is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and eigenvalue or eigenstate. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, preparation, basis, or boundary context; the

Quantum Mechanism Frame

- **Role.** Eigenvalues and eigenvectors contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis, domain, or context.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.16, boundary weak form=0.14

- [arXiv:0912.2823](#), score 0.461
- [arXiv:1604.05385](#), score 0.422
- [arXiv:1612.00682](#), score 0.414
- [arXiv:2308.15676](#), score 0.400
- [arXiv:2108.07838](#), score 0.393

10.12 Von Neumann algebra

Mechanism-tree role. Spectral question: what can be asked; assignment score 1.43. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The von neumann algebra can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Von Neumann algebra is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, unitary operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the

Quantum Mechanism Frame

- **Role.** Von Neumann algebra contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: projection.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.25, constraint closure=0.18

- [arXiv:0912.2823](#), score 0.437
- [arXiv:1604.05385](#), score 0.434
- [arXiv:1604.06537](#), score 0.426
- [arXiv:1612.00682](#), score 0.422
- [arXiv:quant-ph0205159](#), score 0.406

10.13 Electron microscope

Mechanism-tree role. Spectral question: what can be asked; assignment score 0.67. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Electron microscope can be represented as an operational construction: a context specifies a state, an operator transforms it, and admissible outcomes are read through a spectrum.

Mechanism Reading

Operationally, Electron microscope is organized by the relation between state terms (quantum state, wave function, density operator), operator terms (Hamiltonian, observable operator, generator), context or boundary terms (detector), and spectral readout terms (mode). The local vocabulary can change across formulations, but the same state-to-transformation-to-readout structure can remain identifiable. Across equation-level evidence the dominant pattern is operator and spectrum, state evolution / transport, closure / conservation, and boundary or preparation, carried by field-specific vocabulary, probability / information, and symbolic structure. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Electron microscope contributes an instrument-mediated readout role to the quantum construction.
- **Placement.** This page is read first as a question-selection move: it identifies the spectrum or answer set being read.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: detector.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.

- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** Candidate transfer targets are systems where the same state-to-operator-to-spectrum conversion appears, but one edge of the construction remains experimentally unresolved.

Core-Derived Role Equations

This equation block is the branch-level quantum mechanism attached to the page by the compact constructor. It gives the role equation to check against the page's source evidence; source notation may differ.

$$\rho_{\text{probe}} \mapsto \mathcal{E}_{\text{sample}}(\rho_{\text{probe}}), \quad p(y) = \text{Tr}(M_y \mathcal{E}_{\text{sample}}(\rho_{\text{probe}})), \quad \hat{s} = R(\{y_i\})$$

What Remains Stable

- The mechanism is an apparatus-coupled readout: a prepared probe state interacts with a sample or field, the interaction changes phase, momentum, intensity, or counting statistics, and the instrument reconstructs an image, spectrum, trajectory, or estimate.
- Electron microscope defines the legal question being asked of the state.
- The measurable answers are encoded by the operator spectrum, projectors, or spectral measure.
- The operator role is preserved across equivalent bases even when matrix entries change.

What Changes With Realization

- The local title, representation, and physical realization may change while the constructor role is preserved.
- The same observable may be represented by matrices, differential operators, projectors, or algebraic elements.
- Degeneracy, basis choice, and domain conditions can change how the spectrum is displayed.
- Detector implementation changes the physical realization, not the operator role itself.

Checks

- Use this role by specifying the page's state carrier, operator or map, readout, and compatibility condition in its own quantum language.
- Self-adjointness, or the appropriate POVM positivity condition, is what makes the question a legal readout.
- A complete spectral resolution supplies all outcome channels for the question being asked.

Evidence Profile

spectral operator=0.42, transport flow=0.20, constraint closure=0.10

- [arXiv:0912.2823](#), score 0.567
- [arXiv:1612.00682](#), score 0.554
- [arXiv:0805.4565](#), score 0.553
- [arXiv:2108.07838](#), score 0.552
- [arXiv:quant-ph0205159](#), score 0.543

Chapter 11

Readout rule: how answers become probabilities

Measurement connects a state and an observable to recorded frequencies. Projection, POVMs, Born weights, and collapse language are alternative ways of presenting this state-to-spectrum readout step.

11.1 Why This Branch Exists

Measurement is best placed after observables and spectra: it is the rule that turns spectral resolution into recorded probability, not the mystical starting point of the theory.

11.2 Page Map

Page	Dominant evidence signal	Score
POVM	observables and spectra	1.47
Wave function collapse	observables and spectra	1.45
Born rule	observables and spectra	1.45
Measurement in quantum mechanics	observables and spectra	1.39
Quantum jump	observables and spectra	1.32
Measurement problem	state evolution	1.28
Quantum eraser experiment	observables and spectra	1.23
Projection-valued measure	observables and spectra	1.19
Delayed-choice quantum eraser	observables and spectra	1.06
Quantum metrology	observables and spectra	0.88

11.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

11.4 POVM

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.47. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

POVM is the generalized-readout constructor: outcome effects need not be orthogonal projectors.

Mechanism Reading

POVMs separate the probability readout from the idealized projection assumption. They are the natural mechanism for noisy, coarse-grained, indirect, or open-system measurements. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** POVM contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** A state vector or density operator together with the measurement context in which outcome channels are defined.
- **Operator or map.** A projection-valued measure, POVM, update map, or instrument map connecting state to record.
- **Admissibility.** Outcome probabilities must be positive, normalized, and tied to a specified readout map rather than to informal observer language.
- **Readout.** Born probabilities, detector records, post-measurement states, ensemble frequencies, or decision probabilities.
- **Check.** The interpretation is constrained by whether it changes the probability rule, the update rule, the detector model, or only the language used for them.

Topic Equations

Standard constructor skeleton: positive effects and generalized Born rule.

$$\begin{aligned} E_i &\geq 0, & \sum_i E_i &= I \\ p(i) &= \text{Tr}(\rho E_i) \\ E_i &= \sum_{\alpha} K_{i\alpha}^{\dagger} K_{i\alpha} \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.18, constraint closure=0.16

- [arXiv:1604.05385](#), score 0.574
- [arXiv:1801.03283](#), score 0.552
- [arXiv:1612.00682](#), score 0.546
- [arXiv:1506.05598](#), score 0.533
- [arXiv:quant-ph0205159](#), score 0.531

11.5 Wave function collapse

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.45. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a wave function collapse modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a wave function collapse acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Wave function collapse contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** State terms: Wave function, state vector, quantum state, or superposition. Context/domain terms: detector, measurement apparatus, or basis.
- **Operator or map.** Operator terms: observable. Protocol or update terms: collapse and Born rule.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenstate and eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels

- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.35, transport flow=0.25, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.535
- [arXiv:0912.2823](#), score 0.525
- [arXiv:1604.06537](#), score 0.524
- [arXiv:1612.00682](#), score 0.494
- [arXiv:1801.03283](#), score 0.493

11.6 Born rule

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.45. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Born rule is the probability-readout constructor: it maps a state and a spectral channel to an observed probability.

Mechanism Reading

The Born rule is the point where the constructor becomes predictive. It does not name an object; it connects state preparation and a legal question to frequencies over outcome channels. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Born rule contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** A state vector or density operator together with the measurement context in which outcome channels are defined.
- **Operator or map.** A projection-valued measure, POVM, update map, or instrument map connecting state to record.
- **Admissibility.** Outcome probabilities must be positive, normalized, and tied to a specified readout map rather than to informal observer language.
- **Readout.** Born probabilities, detector records, post-measurement states, ensemble frequencies, or decision probabilities.
- **Check.** The interpretation is constrained by whether it changes the probability rule, the update rule, the detector model, or only the language used for them.

Topic Equations

Standard constructor skeleton: probability assignment for projective and position readouts.

$$\begin{aligned}p(i|\rho, \{P_i\}) &= \text{Tr}(\rho P_i) \\p(x|\psi) &= |\psi(x)|^2 \\ \sum_i p(i) &= 1\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.27, constraint closure=0.20

- [arXiv:1604.05385](#), score 0.568
- [arXiv:1604.06537](#), score 0.539
- [arXiv:1801.03283](#), score 0.537
- [arXiv:1612.00682](#), score 0.530
- [arXiv:0912.2823](#), score 0.527

11.7 Measurement in quantum mechanics

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Measurement in quantum mechanics is the complete readout junction: it combines a state, a measurement model, probabilities, and sometimes an update rule.

Mechanism Reading

Measurement is not the root of quantum theory in this book. It is the junction where a prepared state and an observable or POVM are converted into probabilities and recorded outcomes. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Measurement in quantum mechanics contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** A state vector or density operator together with the measurement context in which outcome channels are defined.
- **Operator or map.** A projection-valued measure, POVM, update map, or instrument map connecting state to record.

- **Admissibility.** Outcome probabilities must be positive, normalized, and tied to a specified readout map rather than to informal observer language.
- **Readout.** Born probabilities, detector records, post-measurement states, ensemble frequencies, or decision probabilities.
- **Check.** The interpretation is constrained by whether it changes the probability rule, the update rule, the detector model, or only the language used for them.

Topic Equations

Standard constructor skeleton: generalized measurement probability and conditional update.

$$\begin{aligned}
 p(i) &= \text{Tr}(\rho E_i) \\
 \rho \mapsto \rho_i &= \frac{K_i \rho K_i^\dagger}{\text{Tr}(K_i \rho K_i^\dagger)} \\
 E_i &= K_i^\dagger K_i
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.23, constraint closure=0.16

- [arXiv:1604.05385](#), score 0.579
- [arXiv:1604.06537](#), score 0.551
- [arXiv:0912.2823](#), score 0.544
- [arXiv:1612.00682](#), score 0.535
- [arXiv:quant-ph0205159](#), score 0.535

11.8 Quantum jump

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.32. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum jump can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum jump is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, unitary operator or Hamiltonian, and energy level or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest

Quantum Mechanism Frame

- **Role.** Quantum jump contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: energy level. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.24, constraint closure=0.13

- [arXiv:1604.05385](#), score 0.511
- [arXiv:2108.07838](#), score 0.508
- [arXiv:1612.00682](#), score 0.497
- [arXiv:0805.4565](#), score 0.494
- [arXiv:quant-ph0205159](#), score 0.481

11.9 Measurement problem

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.28. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

the Measurement problem modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

the Measurement problem acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is state evolution, preparation, basis, or boundary context, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Measurement problem contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** State terms: wave function and superposition. Context/domain terms: measurement apparatus.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse and Born rule.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.

- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, boundary weak form=0.20, constraint closure=0.20

- [arXiv:1604.05385](#), score 0.470
- [arXiv:0912.2823](#), score 0.467

- [arXiv:1706.03846](#), score 0.459
- [arXiv:2501.07524](#), score 0.443
- [arXiv:1604.06537](#), score 0.438

11.10 Quantum eraser experiment

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.23. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum eraser experiment can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum eraser experiment is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through complementarity or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, preparation, basis, or boundary context, state evolution; the

Quantum Mechanism Frame

- **Role.** Quantum eraser experiment contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: detector and experimental setup.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: complementarity. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.

- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.32, boundary weak form=0.16, transport flow=0.14

- [arXiv:1604.05385](#), score 0.485
- [arXiv:0912.2823](#), score 0.484
- [arXiv:gr-qc0303063](#), score 0.467
- [arXiv:1604.06537](#), score 0.445
- [arXiv:2110.09771](#), score 0.436

11.11 Projection-valued measure

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.19. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Projection-valued measure is the sharp-readout constructor: mutually exclusive outcome projectors partition the identity.

Mechanism Reading

A projection-valued measure encodes an ideal sharp measurement. It defines outcome channels that are orthogonal and exhaustive. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Projection-valued measure contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** A state vector or density operator together with the measurement context in which outcome channels are defined.
- **Operator or map.** A projection-valued measure, POVM, update map, or instrument map connecting state to record.
- **Admissibility.** Outcome probabilities must be positive, normalized, and tied to a specified readout map rather than to informal observer language.
- **Readout.** Born probabilities, detector records, post-measurement states, ensemble frequencies, or decision probabilities.
- **Check.** The interpretation is constrained by whether it changes the probability rule, the update rule, the detector model, or only the language used for them.

Topic Equations

Standard constructor skeleton: sharp outcome channels, probability, and projective update.

$$\begin{aligned} P_i P_j &= \delta_{ij} P_i, & \sum_i P_i &= I \\ p(i) &= \text{Tr}(\rho P_i) \\ \rho &\mapsto \frac{P_i \rho P_i}{\text{Tr}(\rho P_i)} \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.18, constraint closure=0.16

- [arXiv:1604.05385](#), score 0.533
- [arXiv:1612.00682](#), score 0.519
- [arXiv:1801.03283](#), score 0.509
- [arXiv:1506.05598](#), score 0.496
- [arXiv:quant-ph0205159](#), score 0.495

11.12 Delayed-choice quantum eraser

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 1.06. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a delayed-choice quantum eraser modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a delayed-choice quantum eraser acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Delayed-choice quantum eraser contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** State terms: superposition. Context/domain terms: detector, experimental setup, or basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: complementarity and uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels

- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.29, transport flow=0.23, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.492
- [arXiv:0912.2823](#), score 0.492
- [arXiv:1604.06537](#), score 0.480
- [arXiv:quant-ph0205159](#), score 0.445
- [arXiv:1708.03640](#), score 0.444

11.13 Quantum metrology

Mechanism-tree role. Readout rule: how answers become probabilities; assignment score 0.88. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum metrology can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum metrology is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or unitary operator, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Quantum metrology contributes an instrument-mediated readout role to the quantum construction.
- **Placement.** This page is read first as a readout move: it connects the state and question to recorded outcomes.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian and unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.14

- [arXiv:1604.06537](#), score 0.556
- [arXiv:1604.05385](#), score 0.548
- [arXiv:0912.2823](#), score 0.547
- [arXiv:1612.00682](#), score 0.538
- [arXiv:1506.05598](#), score 0.536

Chapter 12

Compatibility limit: what cannot be jointly sharp

The non-classical part of the theory appears when two otherwise legal questions do not compose into one common sharp question. Commutators, uncertainty relations, contextuality, Bell tests, and entanglement live here.

12.1 Why This Branch Exists

The non-classical core appears as failure of joint sharpness. Entanglement, Bell phenomena, and uncertainty are different faces of this compatibility structure.

12.2 Page Map

Page	Dominant evidence signal	Score
Bell's theorem	observables and spectra	1.42
Quantum entanglement	state evolution	1.38
Commutator	observables and spectra	1.33
Uncertainty principle	observables and spectra	1.32
Einstein–Podolsky–Rosen paradox	state evolution	1.13
Quantum nonlocality	observables and spectra	1.10

12.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

12.4 Bell's theorem

Mechanism-tree role. Compatibility limit: what cannot be jointly sharp; assignment score 1.42. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Bell's theorem is a compatibility/locality stress test: quantum correlations violate bounds satisfied by local hidden-variable assignments.

Mechanism Reading

Bell's theorem is not a page about a mysterious object. It is a falsifier for a classical joint-assignment model of measurement outcomes. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, spectral profile.

Quantum Mechanism Frame

- **Role.** Bell's theorem contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a compatibility move: it asks which otherwise legal questions cannot share one sharp answer set.
- **Carrier or domain.** One state space or a multipartite state space on which several questions can be asked.
- **Operator or map.** Two or more observables, contexts, correlation operators, or hidden-variable assignments being compared.
- **Admissibility.** Commutators, uncertainty bounds, contextuality constraints, or Bell-type inequalities decide which joint assignments are possible.
- **Readout.** Joint spectra, correlations, inequality violations, uncertainty products, or incompatible outcome statistics.
- **Check.** The non-classical content appears only if the incompatible questions cannot be replaced by one common sharp classical assignment.

Topic Equations

Standard constructor skeleton: CHSH inequality and quantum violation bound.

$$|E(a, b) + E(a, b') + E(a', b) - E(a', b')| \leq 2$$

quantum prediction can reach $2\sqrt{2}$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.24, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.511
- [arXiv:1604.06537](#), score 0.472
- [arXiv:0912.2823](#), score 0.466
- [arXiv:1801.03283](#), score 0.445
- [arXiv:quant-ph0205159](#), score 0.443

12.5 Quantum entanglement

Mechanism-tree role. Compatibility limit: what cannot be jointly sharp; assignment score 1.38. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum entanglement is the non-factorization constructor: a composite state can carry correlations not reducible to independent subsystem states.

Mechanism Reading

Entanglement belongs to composition and compatibility. It appears when the tensor-product state cannot be written as a product or mixture of local states, producing correlations that stress classical separability. In the source-evidence profile for this page, the strongest construction signal is state evolution, operator-to-spectrum readout, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum entanglement contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a compatibility move: it asks which otherwise legal questions cannot share one sharp answer set.
- **Carrier or domain.** One state space or a multipartite state space on which several questions can be asked.
- **Operator or map.** Two or more observables, contexts, correlation operators, or hidden-variable assignments being compared.
- **Admissibility.** Commutators, uncertainty bounds, contextuality constraints, or Bell-type inequalities decide which joint assignments are possible.
- **Readout.** Joint spectra, correlations, inequality violations, uncertainty products, or incompatible outcome statistics.
- **Check.** The non-classical content appears only if the incompatible questions cannot be replaced by one common sharp classical assignment.

Topic Equations

Standard constructor skeleton: tensor composition, non-factorization, and reduced state.

$$\begin{aligned}\mathcal{H}_{AB} &= \mathcal{H}_A \otimes \mathcal{H}_B \\ |\Psi\rangle_{AB} &\neq |\psi\rangle_A \otimes |\phi\rangle_B \\ \rho_A &= \text{Tr}_B |\Psi\rangle\langle\Psi|\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, spectral operator=0.22, constraint closure=0.21

- [arXiv:1604.05385](#), score 0.477
- [arXiv:0912.2823](#), score 0.476
- [arXiv:1604.06537](#), score 0.443
- [arXiv:1801.03283](#), score 0.432
- [arXiv:quant-ph0205159](#), score 0.427

12.6 Commutator

Mechanism-tree role. Compatibility limit: what cannot be jointly sharp; assignment score 1.33. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Commutator is the incompatibility constructor: it measures the failure of two transformations or questions to be freely exchanged.

Mechanism Reading

The commutator is the algebraic source of many non-classical restrictions. If two observables do not commute, they generally cannot be resolved in one common sharp basis. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, non-commuting compatibility limits; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Commutator contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a compatibility move: it asks which otherwise legal questions cannot share one sharp answer set.
- **Carrier or domain.** A common state space on which two transformations, observables, or questions are both defined.
- **Operator or map.** The ordered products AB and BA , compared through the obstruction $[A,B]=AB-BA$.

- **Admissibility.** A nonzero commutator marks an order-dependence or compatibility limit; a zero commutator permits a common sharp refinement only when the remaining spectral conditions hold.
- **Readout.** Compatibility tests, uncertainty bounds, common eigenspaces, or canonical commutation relations.
- **Check.** The mechanism is supported only when changing operator order changes the algebraic or statistical prediction.

Topic Equations

Standard constructor skeleton: order obstruction and canonical commutation.

$$\begin{aligned}
 [A, B] &= AB - BA \\
 [A, B] = 0 &\Rightarrow \text{common spectral refinement may exist} \\
 [x, p] &= i\hbar
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.19, commutator incompatibility=0.13

- [arXiv:1604.06537](#), score 0.550
- [arXiv:2111.12617](#), score 0.543
- [arXiv:1612.00682](#), score 0.525
- [arXiv:2108.07838](#), score 0.519
- [arXiv:0805.4565](#), score 0.516

12.7 Uncertainty principle

Mechanism-tree role. Compatibility limit: what cannot be jointly sharp; assignment score 1.32. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Uncertainty principle is a compatibility-limit theorem: non-commuting observables impose lower bounds on joint sharpness.

Mechanism Reading

Uncertainty is not detector imperfection. It is a structural consequence of state variance and non-commuting observables. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Uncertainty principle contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a compatibility move: it asks which otherwise legal questions cannot share one sharp answer set.
- **Carrier or domain.** One state space or a multipartite state space on which several questions can be asked.
- **Operator or map.** Two or more observables, contexts, correlation operators, or hidden-variable assignments being compared.
- **Admissibility.** Commutators, uncertainty bounds, contextuality constraints, or Bell-type inequalities decide which joint assignments are possible.
- **Readout.** Joint spectra, correlations, inequality violations, uncertainty products, or incompatible outcome statistics.
- **Check.** The non-classical content appears only if the incompatible questions cannot be replaced by one common sharp classical assignment.

Topic Equations

Standard constructor skeleton: variance bound from commutator structure.

$$\Delta_{\rho}A \Delta_{\rho}B \geq \frac{1}{2} |\text{Tr}(\rho[A, B])|$$
$$\Delta x \Delta p \geq \frac{\hbar}{2}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.18, constraint closure=0.13

- [arXiv:1604.05385](#), score 0.539
- [arXiv:1604.06537](#), score 0.525
- [arXiv:2111.12617](#), score 0.504
- [arXiv:0912.2823](#), score 0.502
- [arXiv:1612.00682](#), score 0.499

12.8 Einstein–Podolsky–Rosen paradox

Mechanism-tree role. Compatibility limit: what cannot be jointly sharp; assignment score 1.13. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

the Einstein–Podolsky–Rosen paradox modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

the Einstein–Podolsky–Rosen paradox acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or admissibility, non-commuting compatibility limits; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Einstein–Podolsky–Rosen paradox contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a compatibility move: it asks which otherwise legal questions cannot share one sharp answer set.
- **Carrier or domain.** State terms: wave function, quantum state, or superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: incompatible and uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, constraint closure=0.21, commutator incompatibility=0.14

- [arXiv:1604.05385](#), score 0.449
- [arXiv:0912.2823](#), score 0.447
- [arXiv:2501.07524](#), score 0.440
- [arXiv:1706.03846](#), score 0.439
- [arXiv:1604.06537](#), score 0.437

12.9 Quantum nonlocality

Mechanism-tree role. Compatibility limit: what cannot be jointly sharp; assignment score 1.10. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a quantum nonlocality modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum nonlocality acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, controlled update protocol; the strongest carrier signal is local notation, information profile, spectral profile.

Quantum Mechanism Frame

- **Role.** Quantum nonlocality contributes a compatibility or joint-readout role to the quantum construction.
- **Placement.** This page is read first as a compatibility move: it asks which otherwise legal questions cannot share one sharp answer set.
- **Carrier or domain.** State terms: wavefunction, quantum state, or wave function. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.24, discrete protocol=0.20

- [arXiv:1604.05385](#), score 0.555
- [arXiv:1612.00682](#), score 0.485
- [arXiv:1801.03283](#), score 0.480
- [arXiv:1506.05598](#), score 0.477
- [arXiv:quant-ph0205159](#), score 0.475

Chapter 13

Boundary realization: how effects appear

Many named quantum effects are boundary realizations of the same construction. A potential, barrier, box, cavity, detector, or medium changes the allowed spectral channels without changing the basic prediction problem.

13.1 Why This Branch Exists

Named effects such as tunnelling and particle-in-a-box are boundary realizations of the state-operator-readout construction.

13.2 Page Map

Page	Dominant evidence signal	Score
Potential well	observables and spectra	1.44
Particle in a box	observables and spectra	1.42
Scattering	state evolution	1.39
Wave interference	observables and spectra	1.37
Quantum optics	observables and spectra	1.37
Spectral line	observables and spectra	0.92
S-matrix	observables and spectra	1.44
Quantum tunnelling	observables and spectra	1.42
Quantum imaging	preparation and boundary context	1.38
Quantum harmonic oscillator	observables and spectra	1.34
Macroscopic quantum phenomena	state evolution	1.08
Quantum metamaterial	state evolution	0.78

13.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

13.4 Potential well

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.44. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A potential well can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a potential well is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through wave function or state vector, Hamiltonian or observable operator, and spectrum or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest

Quantum Mechanism Frame

- **Role.** Potential well contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: wave function. Context/domain terms: Potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum, energy level, or mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.13

- [arXiv:1604.05385](#), score 0.579
- [arXiv:0912.2823](#), score 0.575
- [arXiv:1506.05598](#), score 0.570
- [arXiv:1612.00682](#), score 0.570
- [arXiv:2108.07838](#), score 0.570

13.5 Particle in a box

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.42. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Particle in a box is a boundary-spectrum constructor: a spatial domain and boundary condition discretize the allowed energy spectrum.

Mechanism Reading

The page shows how a boundary condition changes the domain of the Hamiltonian and therefore the allowed spectra. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Particle in a box contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.
- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Standard constructor skeleton: boundary condition and discrete spectrum.

$$\begin{aligned}\psi(0) &= \psi(L) = 0 \\ \psi_n(x) &= \sqrt{\frac{2}{L}} \sin \frac{n\pi x}{L} \\ E_n &= \frac{\hbar^2 \pi^2 n^2}{2mL^2}\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.16

- [arXiv:0912.2823](#), score 0.599
- [arXiv:1801.03283](#), score 0.592
- [arXiv:1612.00682](#), score 0.591
- [arXiv:1506.05598](#), score 0.590
- [arXiv:quant-ph0205159](#), score 0.583

13.6 Scattering

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.39.
Dominant evidence signal. state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A scattering can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a scattering is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Matrix or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or admissibility,

Quantum Mechanism Frame

- **Role.** Scattering contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: boundary condition and potential.
- **Operator or map.** Operator terms: Matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, constraint closure=0.21, spectral operator=0.20

- [arXiv:1604.05385](#), score 0.530
- [arXiv:0912.2823](#), score 0.527
- [arXiv:1604.06537](#), score 0.503
- [arXiv:1708.03640](#), score 0.495
- [arXiv:1801.03283](#), score 0.495

13.7 Wave interference

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.37. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A wave interference can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a wave interference is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through superposition or wave function, observable operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the

Quantum Mechanism Frame

- **Role.** Wave interference contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: observable. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.36, transport flow=0.25, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.561
- [arXiv:1506.05598](#), score 0.541
- [arXiv:1801.03283](#), score 0.538
- [arXiv:1612.00682](#), score 0.535
- [arXiv:0908.0752](#), score 0.532

13.8 Quantum optics

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.37. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum optics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum optics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wavefunction or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum optics contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: wavefunction. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.34, transport flow=0.22, constraint closure=0.13

- [arXiv:1604.05385](#), score 0.571
- [arXiv:0912.2823](#), score 0.570
- [arXiv:0805.4565](#), score 0.549
- [arXiv:1708.03640](#), score 0.548
- [arXiv:quant-ph0205159](#), score 0.548

13.9 Spectral line

Mechanism-tree role. Boundary realization: how effects appear; assignment score 0.92. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A spectral line can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a spectral line is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, preparation, basis, or boundary context, state evolution; the

Quantum Mechanism Frame

- **Role.** Spectral line contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.27, boundary weak form=0.21, transport flow=0.18

- [arXiv:0912.2823](#), score 0.552
- [arXiv:1604.06537](#), score 0.516
- [arXiv:1604.05385](#), score 0.516
- [arXiv:1801.03283](#), score 0.497
- [arXiv:quant-ph0205159](#), score 0.497

13.10 S-matrix

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.44. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The s-matrix can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the S-matrix is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through wave function or probability amplitude, matrix or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** S-matrix contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: wave function and probability amplitude. Context/-domain terms: potential, context, or basis.
- **Operator or map.** Operator terms: matrix, Hamiltonian, or unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.27, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.567
- [arXiv:1604.06537](#), score 0.550
- [arXiv:1801.03283](#), score 0.547
- [arXiv:1612.00682](#), score 0.539
- [arXiv:1506.05598](#), score 0.535

13.11 Quantum tunnelling

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.42. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum tunnelling is a boundary-realization constructor: a state has nonzero transmission through a classically forbidden region.

Mechanism Reading

Tunnelling shows that the realization layer matters. A potential barrier changes the admissible wave solutions and produces transmission even where classical kinetic energy would be negative. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum tunnelling contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** A Hilbert space with a selected domain, potential, interface, asymptotic channel, cavity, well, or boundary condition.
- **Operator or map.** A Hamiltonian, wave operator, transfer operator, or scattering map whose domain is changed by the boundary.
- **Admissibility.** Boundary conditions and matching conditions determine allowed states, resonances, transmission amplitudes, and spectra.
- **Readout.** Eigenvalues, resonances, tunnelling probabilities, phase shifts, reflection/transmission amplitudes, or scattering data.
- **Check.** Changing the boundary should change the spectrum or amplitude in the predicted way, while the free or asymptotic limit is recovered when the boundary is removed.

Topic Equations

Standard constructor skeleton: barrier-domain Schrödinger equation and WKB transmission.

$$T \sim \exp\left(-2 \int_{x_1}^{x_2} \sqrt{\frac{2m(V(x) - E)}{\hbar^2}} dx\right)$$
$$-\frac{\hbar^2}{2m}\psi''(x) + V(x)\psi(x) = E\psi(x)$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.24, constraint closure=0.18

- [arXiv:1604.05385](#), score 0.569
- [arXiv:0912.2823](#), score 0.568
- [arXiv:1612.00682](#), score 0.562
- [arXiv:quant-ph0205159](#), score 0.558
- [arXiv:1801.03283](#), score 0.555

13.12 Quantum imaging

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.38. **Dominant evidence signal.** preparation and boundary context. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum imaging can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum imaging is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the

Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is preparation, basis, or boundary context, operator-to-spectrum readout, state evolution; the

Quantum Mechanism Frame

- **Role.** Quantum imaging contributes an instrument-mediated readout role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential and detector.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

boundary weak form=0.25, spectral operator=0.25, transport flow=0.22

- [arXiv:0912.2823](#), score 0.523
- [arXiv:1604.05385](#), score 0.484
- [arXiv:1612.00682](#), score 0.452
- [arXiv:quant-ph0205159](#), score 0.449
- [arXiv:gr-qc0411110](#), score 0.447

13.13 Quantum harmonic oscillator

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.34. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum harmonic oscillator can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum harmonic oscillator is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through wave function or wavefunction, Hamiltonian or observable operator, and energy level

or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum harmonic oscillator contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: wave function and wavefunction. Context/domain terms: potential and basis.
- **Operator or map.** Operator terms: Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: energy level, eigenvalue, eigenstate, or spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.23, constraint closure=0.16

- [arXiv:1604.06537](#), score 0.603
- [arXiv:1801.03283](#), score 0.568
- [arXiv:1612.00682](#), score 0.560
- [arXiv:0912.2823](#), score 0.556
- [arXiv:1506.05598](#), score 0.555

13.14 Macroscopic quantum phenomena

Mechanism-tree role. Boundary realization: how effects appear; assignment score 1.08.

Dominant evidence signal. state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A macroscopic quantum phenomena can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a macroscopic quantum phenomena is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through wave function or state vector, Hamiltonian or observable operator, and eigenvalue or

energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Macroscopic quantum phenomena contributes an open-system transport and coherence role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: wave function. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.28, boundary weak form=0.23, constraint closure=0.18

- [arXiv:0912.2823](#), score 0.544
- [arXiv:1706.03846](#), score 0.539
- [arXiv:hep-lat9608080](#), score 0.518
- [arXiv:1708.03640](#), score 0.511
- [arXiv:astro-ph0604157](#), score 0.510

13.15 Quantum metamaterial

Mechanism-tree role. Boundary realization: how effects appear; assignment score 0.78. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum metamaterial can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum metamaterial is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and

eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Quantum metamaterial contributes a boundary-shaped spectrum role to the quantum construction.
- **Placement.** This page is read first as a realization move: it changes the domain, boundary, geometry, or interface in which the operator acts.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.30, spectral operator=0.13, constraint closure=0.07

- [arXiv:1706.03846](#), score 0.540
- [arXiv:1708.03640](#), score 0.516
- [arXiv:quant-ph0312187](#), score 0.513
- [arXiv:0908.0752](#), score 0.506
- [arXiv:0805.4565](#), score 0.505

Chapter 14

Many-mode extension: fields, particles, and scaling

Quantum field theory, gauge theory, renormalization, photons, fermions, and related topics extend the same state-operator-spectrum logic to variable particle number, local fields, and scale-dependent descriptions.

14.1 Why This Branch Exists

Field theory and gauge theory extend the same construction to many modes, local generators, and scale-dependent descriptions.

14.2 Page Map

Page	Dominant evidence signal	Score
Fermi–Dirac statistics	state evolution	1.50
Dirac equation	observables and spectra	1.50
Quantum electrodynamics	observables and spectra	1.50
Renormalization	observables and spectra	1.50
Gauge theory	observables and spectra	1.49
Photon	observables and spectra	1.48
Quantum field theory	observables and spectra	1.47
Fermion	state evolution	1.42
Boson	normalization and admissibility	1.40
Quantum chromodynamics	observables and spectra	1.30
AdS/CFT correspondence	state evolution	1.43
Klein–Gordon equation	observables and spectra	1.22
Quantum gravity	state evolution	1.20
Spin network	observables and spectra	1.11
Electron	observables and spectra	0.94
Loop quantum gravity	observables and spectra	0.92
Spin foam	observables and spectra	0.92
String theory	observables and spectra	0.90
Quantum geometry	observables and spectra	0.88
Fock space	observables and spectra	0.86

Page	Dominant evidence signal	Score
Quantum spacetime	incompatible questions	0.85

14.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

14.4 Fermi–Dirac statistics

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.50. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A fermi–dirac statistics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a fermi–Dirac statistics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** Fermi–Dirac statistics contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, boundary weak form=0.24, spectral operator=0.23

- [arXiv:0912.2823](#), score 0.491
- [arXiv:1604.05385](#), score 0.452
- [arXiv:1801.03283](#), score 0.421
- [arXiv:1612.00682](#), score 0.421
- [arXiv:quant-ph0205159](#), score 0.417

14.5 Dirac equation

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.50. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The dirac equation can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Dirac equation is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wavefunction or wave function, matrix or Hamiltonian, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Dirac equation contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: wavefunction and wave function. Context/domain terms: context.
- **Operator or map.** Operator terms: matrix and Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.26, constraint closure=0.21

- [arXiv:0912.2823](#), score 0.525
- [arXiv:1604.06537](#), score 0.523
- [arXiv:1801.03283](#), score 0.515
- [arXiv:quant-ph0205159](#), score 0.504
- [arXiv:1612.00682](#), score 0.503

14.6 Quantum electrodynamics

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.50. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum electrodynamics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum electrodynamics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through probability amplitude or wave function, observable operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through incompatible observables or commutator. In the source-evidence profile for this page, the strongest construction signal is

Quantum Mechanism Frame

- **Role.** Quantum electrodynamics contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: probability amplitude. Context/domain terms: experimental setup.
- **Operator or map.** Operator terms: observable. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.

- **Admissibility.** Compatibility or closure terms: incompatible. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.26, transport flow=0.22, constraint closure=0.21

- [arXiv:0912.2823](#), score 0.470
- [arXiv:1604.06537](#), score 0.452
- [arXiv:2111.12617](#), score 0.422
- [arXiv:1801.03283](#), score 0.416
- [arXiv:2501.07524](#), score 0.413

14.7 Renormalization

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.50. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Renormalization is the scale-flow constructor: the effective parameters of the theory change with resolution while predictions remain controlled.

Mechanism Reading

Renormalization explains why a mechanism can preserve its operator role while changing its apparent parameters across scales. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Renormalization contributes an unresolved constructor role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** A context-selected state space or effective carrier for prediction.
- **Operator or map.** The relevant Hamiltonian, observable, channel, constraint, or update map.
- **Admissibility.** Domain, normalization, positivity, compatibility, boundary, or gauge requirements state what is legal.
- **Readout.** The outcome probabilities, spectra, correlations, amplitudes, or records used to test the mechanism.
- **Check.** A complete account must specify state carrier, operator or map, admissibility condition, readout, and at least one possible falsifier.

Topic Equations

Standard constructor skeleton: beta flow and effective operator expansion.

$$\begin{aligned}\mu \frac{dg}{d\mu} &= \beta(g) \\ g &= g(\mu) \\ \mathcal{L}_{\text{eff}}(\mu) &= \sum_i c_i(\mu) \mathcal{O}_i\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.19, constraint closure=0.15

- [arXiv:0912.2823](#), score 0.498
- [arXiv:1604.06537](#), score 0.494
- [arXiv:1612.00682](#), score 0.465
- [arXiv:1801.03283](#), score 0.464
- [arXiv:2111.12617](#), score 0.461

14.8 Gauge theory

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.49. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Gauge theory is a redundancy-and-constraint constructor: different local presentations can represent the same physical state.

Mechanism Reading

Gauge theory belongs at the field/geometry frontier. It separates physical degrees of freedom from representational choices and imposes covariant transport through a connection. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Gauge theory contributes an unresolved constructor role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** A context-selected state space or effective carrier for prediction.
- **Operator or map.** The relevant Hamiltonian, observable, channel, constraint, or update map.
- **Admissibility.** Domain, normalization, positivity, compatibility, boundary, or gauge requirements state what is legal.
- **Readout.** The outcome probabilities, spectra, correlations, amplitudes, or records used to test the mechanism.
- **Check.** A complete account must specify state carrier, operator or map, admissibility condition, readout, and at least one possible falsifier.

Topic Equations

Standard constructor skeleton: covariant derivative, curvature, and local gauge transformation.

$$\begin{aligned}D_\mu &= \partial_\mu + igA_\mu \\F_{\mu\nu} &= \partial_\mu A_\nu - \partial_\nu A_\mu + ig[A_\mu, A_\nu] \\ \psi(x) &\mapsto U(x)\psi(x)\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels

- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.25, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.511
- [arXiv:1604.05385](#), score 0.478
- [arXiv:1612.00682](#), score 0.468
- [arXiv:1801.03283](#), score 0.463
- [arXiv:quant-ph0205159](#), score 0.459

14.9 Photon

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.48. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Photon is a field-mode constructor: a one-quantum excitation of the electromagnetic field, constrained by massless dispersion and transverse polarization.

Mechanism Reading

The native photon mechanism is field quantization. The electromagnetic field is decomposed into modes, creation and annihilation operators act on those modes, and a one-photon state is created from the vacuum. The relevant readouts are occupation number, energy, momentum, polarization, and detector clicks; the constraints are dispersion and gauge-compatible transversality. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Photon contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.
- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Topic-specific constructor: the equations express massless dispersion, one-mode occupation, number readout, and transverse polarization.

$$\begin{aligned} E &= \hbar\omega, & \mathbf{p} &= \hbar\mathbf{k}, & \omega &= c|\mathbf{k}| \\ |1_{\mathbf{k},\lambda}\rangle &= a_{\mathbf{k},\lambda}^\dagger |0\rangle \\ \hat{N}_{\mathbf{k},\lambda} &= a_{\mathbf{k},\lambda}^\dagger a_{\mathbf{k},\lambda} \\ \mathbf{k} \cdot \boldsymbol{\epsilon}_{\mathbf{k},\lambda} &= 0 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.25, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.536
- [arXiv:0912.2823](#), score 0.504
- [arXiv:1612.00682](#), score 0.495
- [arXiv:1801.03283](#), score 0.494
- [arXiv:quant-ph0205159](#), score 0.488

14.10 Quantum field theory

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.47. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum field theory is the many-mode local-field extension of the quantum constructor.

Mechanism Reading

Quantum field theory lifts the state-operator-spectrum construction to fields, local operators, creation and annihilation modes, and scattering amplitudes. Particles become stable excitation/readout roles of fields. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum field theory contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.

- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Standard constructor skeleton: field expansion, mode algebra, and correlation readout.

$$\begin{aligned}\Phi(x) &= \sum_k \left(a_k u_k(x) + a_k^\dagger u_k^*(x) \right) \\ [a_k, a_l^\dagger]_{\mp} &= \delta_{kl} \\ \langle 0 | T \{ \Phi(x) \Phi(y) \} | 0 \rangle\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.27, transport flow=0.26, constraint closure=0.19

- [arXiv:0912.2823](#), score 0.518
- [arXiv:1604.06537](#), score 0.508
- [arXiv:1708.03640](#), score 0.473
- [arXiv:quant-ph0205159](#), score 0.471
- [arXiv:1801.03283](#), score 0.468

14.11 Fermion

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.42. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Fermion is an exchange-symmetry constructor: identical fermions live in antisymmetric sectors and obey Pauli exclusion.

Mechanism Reading

The mechanism is an admissibility rule on many-body state space. Antisymmetry, anticommutation, and zero-or-one mode occupation define the portable role. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or admissibility, preparation, basis, or boundary context; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Fermion contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.
- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Topic-specific constructor: the equations express antisymmetric sectors, anticommutation, and occupation restriction.

$$\begin{aligned}\mathcal{F}_-(\mathcal{H}) &= \bigoplus_{n=0}^{\infty} \wedge^n \mathcal{H} \\ \{a_i, a_j^\dagger\} &= \delta_{ij}, \quad \{a_i, a_j\} = 0 \\ n_i &= a_i^\dagger a_i \in \{0, 1\}\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.28, constraint closure=0.26, boundary weak form=0.18

- [arXiv:1706.03846](#), score 0.487
- [arXiv:0912.2823](#), score 0.479
- [arXiv:1708.03640](#), score 0.465
- [arXiv:0908.0752](#), score 0.453
- [arXiv:astro-ph0604157](#), score 0.452

14.12 Boson

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.40. **Dominant evidence signal.** normalization and admissibility. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Boson is an exchange-symmetry constructor: identical bosons live in symmetric sectors and can share the same mode.

Mechanism Reading

The mechanism is symmetric exchange. Commuting creation and annihilation operators allow arbitrary nonnegative occupation of a mode, which supports field modes, coherent states, condensates, and photon-like readouts. In the source-evidence profile for this page, the strongest construction signal is normalization or admissibility, state evolution, operator-to-spectrum readout; the strongest carrier signal is local notation, information profile, spectral profile.

Quantum Mechanism Frame

- **Role.** Boson contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.
- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Topic-specific constructor: the equations express symmetric sectors, commutation, and unrestricted mode occupation.

$$\begin{aligned}\mathcal{F}_+(\mathcal{H}) &= \bigoplus_{n=0}^{\infty} \text{Sym}^n \mathcal{H} \\ [a_i, a_j^\dagger] &= \delta_{ij}, \quad [a_i, a_j] = 0 \\ n_i &= a_i^\dagger a_i \in \{0, 1, 2, \dots\}\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

constraint closure=0.26, transport flow=0.16, spectral operator=0.16

- [arXiv:1706.03846](#), score 0.448
- [arXiv:1708.03640](#), score 0.430
- [arXiv:1801.08949](#), score 0.424
- [arXiv:0908.0752](#), score 0.421
- [arXiv:1801.03283](#), score 0.413

14.13 Quantum chromodynamics

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.30. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum chromodynamics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum chromodynamics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum chromodynamics contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: context and basis.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.33, transport flow=0.27, constraint closure=0.21

- [arXiv:0912.2823](#), score 0.480
- [arXiv:1604.05385](#), score 0.480
- [arXiv:1604.06537](#), score 0.476
- [arXiv:1801.03283](#), score 0.462
- [arXiv:quant-ph0205159](#), score 0.455

14.14 AdS/CFT correspondence

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.43. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

AdS/CFT correspondence is a geometry-translation constructor: bulk gravitational data and boundary field data are treated as dual presentations of one operator structure.

Mechanism Reading

AdS/CFT belongs at the frontier where geometry becomes a realization rather than the invariant root. The practical content is a dictionary between bulk fields and boundary operators. In the source-evidence profile for this page, the strongest construction signal is state evolution, operator-to-spectrum readout, preparation, basis, or boundary context; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** AdS/CFT correspondence contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** A spacetime, boundary algebra, gauge orbit, spin network, bulk/boundary pair, or geometric representation of a quantum state space.
- **Operator or map.** Hamiltonian, action, constraint, boundary operator, correlation map, or dictionary between two representations.
- **Admissibility.** Gauge, boundary, metric, covariance, and constraint conditions decide which geometric descriptions represent the same physical content.
- **Readout.** Boundary correlators, spectra, entropies, scattering data, geometric invariants, or reconstructed bulk quantities.
- **Check.** A geometric reformulation is physical only to the extent that it preserves observables or correlation functions across the representation change.

Topic Equations

Standard constructor skeleton: boundary-source/bulk-field dictionary.

$$\begin{aligned} Z_{\text{grav}}[\phi_0] &\simeq Z_{\text{CFT}}[J = \phi_0] \\ \left\langle \exp \int J \mathcal{O} \right\rangle_{\text{CFT}} &= Z_{\text{bulk}}[\phi|_{\partial} = J] \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, spectral operator=0.26, boundary weak form=0.22

- [arXiv:0912.2823](#), score 0.576
- [arXiv:quant-ph0205159](#), score 0.514
- [arXiv:1801.03283](#), score 0.511
- [arXiv:1708.03640](#), score 0.511
- [arXiv:1506.05598](#), score 0.509

14.15 Klein–Gordon equation

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.22. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

The klein–gordon equation can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, the Klein–Gordon equation is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and spectrum or eigenvalue.

Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Klein–Gordon equation contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.27, transport flow=0.26, commutator incompatibility=0.21

- [arXiv:1604.06537](#), score 0.573
- [arXiv:0908.0752](#), score 0.537
- [arXiv:1506.05598](#), score 0.537
- [arXiv:1708.03640](#), score 0.536
- [arXiv:quant-ph0205159](#), score 0.536

14.16 Quantum gravity

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.20. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum gravity can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum gravity is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through superposition or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue.

Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or

Quantum Mechanism Frame

- **Role.** Quantum gravity contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: superposition. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, constraint closure=0.20, spectral operator=0.20

- [arXiv:1604.05385](#), score 0.535
- [arXiv:0912.2823](#), score 0.535
- [arXiv:1708.03640](#), score 0.500
- [arXiv:1801.03283](#), score 0.497
- [arXiv:quant-ph0205159](#), score 0.495

14.17 Spin network

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 1.11. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A spin network can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a spin network is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and spectrum or eigenstate. Probabilities

enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Spin network contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: context and basis.
- **Operator or map.** Operator terms: matrix and Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum, eigenstate, or eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.35, transport flow=0.23, constraint closure=0.17

- [arXiv:0912.2823](#), score 0.587
- [arXiv:1801.03283](#), score 0.545
- [arXiv:1612.00682](#), score 0.545
- [arXiv:quant-ph0205159](#), score 0.540
- [arXiv:1506.05598](#), score 0.539

14.18 Electron

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.94. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Electron is a charged spinor constructor: its identity is fixed by mass, charge, spin-1/2 representation, fermionic statistics, and electromagnetic coupling.

Mechanism Reading

The electron is not a generic object label in this tree. Its native mechanism combines a spinor state, a Schrödinger/Pauli/Dirac generator depending on regime, conserved charge, and fermionic anticommutation. The readouts are charge, spin, momentum, energy, and scattering response. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, preparation, basis, or boundary context, state evolution; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Electron contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.
- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Topic-specific constructor: the equations express relativistic spinor transport, electromagnetic coupling, and fermionic field statistics.

$$\begin{aligned}(i\hbar\gamma^\mu D_\mu - mc)\psi &= 0 \\ D_\mu &= \partial_\mu + \frac{ie}{\hbar c}A_\mu \\ \{\psi_\alpha(x), \psi_\beta^\dagger(y)\} &= \delta_{\alpha\beta}\delta(x-y)\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.26, boundary weak form=0.25, transport flow=0.22

- [arXiv:0912.2823](#), score 0.507
- [arXiv:1604.05385](#), score 0.467
- [arXiv:1612.00682](#), score 0.436
- [arXiv:quant-ph0205159](#), score 0.435
- [arXiv:gr-qc0411110](#), score 0.431

14.19 Loop quantum gravity

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.92. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A loop quantum gravity can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a loop quantum gravity is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Loop quantum gravity contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.35, transport flow=0.26, constraint closure=0.18

- [arXiv:0912.2823](#), score 0.577
- [arXiv:quant-ph0205159](#), score 0.543
- [arXiv:1612.00682](#), score 0.542
- [arXiv:1801.03283](#), score 0.542
- [arXiv:1506.05598](#), score 0.535

14.20 Spin foam

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.92. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A spin foam can be read as a quantum construction: the boundary conditions that define which histories are included fixes the admissible state space; the action functional assigning phases to histories defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a spin foam is described by boundary states or field configurations at the endpoints of a process. The physical question is represented by the action functional assigning phases to histories; the experimental or mathematical setting is the boundary conditions that define which histories are included. The observable content is obtained from transition amplitudes and probabilities obtained by summing over histories. In the local terminology of this topic, the same construction appears through superposition or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or

Quantum Mechanism Frame

- **Role.** Spin foam contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: path integral.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.44, transport flow=0.23, constraint closure=0.13

- [arXiv:0912.2823](#), score 0.487
- [arXiv:1604.06537](#), score 0.485
- [arXiv:1612.00682](#), score 0.470
- [arXiv:2108.07838](#), score 0.456
- [arXiv:quant-ph0205159](#), score 0.449

14.21 String theory

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.90. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

String theory can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, string theory is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest

Quantum Mechanism Frame

- **Role.** String theory contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.15

- [arXiv:0912.2823](#), score 0.599
- [arXiv:1612.00682](#), score 0.586
- [arXiv:quant-ph0205159](#), score 0.582
- [arXiv:1506.05598](#), score 0.578
- [arXiv:1801.03283](#), score 0.578

14.22 Quantum geometry

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.88. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum geometry is a geometry-realization page: geometric quantities are promoted to quantum observables rather than assumed as a smooth background.

Mechanism Reading

Quantum geometry uses a quantum state of geometry, often represented by graph or spin-network data. The operator-to-spectrum step asks for eigenvalues of geometric observables such as area or volume. This places the page near the geometry/boundary frontier rather than inside a generic many-mode field layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, normalization or admissibility, state evolution; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum geometry contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** A spacetime, boundary algebra, gauge orbit, spin network, bulk/boundary pair, or geometric representation of a quantum state space.

- **Operator or map.** Hamiltonian, action, constraint, boundary operator, correlation map, or dictionary between two representations.
- **Admissibility.** Gauge, boundary, metric, covariance, and constraint conditions decide which geometric descriptions represent the same physical content.
- **Readout.** Boundary correlators, spectra, entropies, scattering data, geometric invariants, or reconstructed bulk quantities.
- **Check.** A geometric reformulation is physical only to the extent that it preserves observables or correlation functions across the representation change.

Topic Equations

Topic-specific constructor: the equations express graph-based geometry states and spectral readout of geometric observables.

$$\begin{aligned}\mathcal{H}_\Gamma &= L^2(SU(2)^E/SU(2)^V), \quad |\Gamma, j_e, \iota_v\rangle \\ \hat{A}(S)|\Gamma, j, \iota\rangle &= 8\pi\gamma\ell_P^2 \sum_{e \in S} \sqrt{j_e(j_e+1)} |\Gamma, j, \iota\rangle \\ \text{geometry readout : } \hat{G} |g_i\rangle &= g_i |g_i\rangle\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.37, constraint closure=0.16, transport flow=0.14

- [arXiv:1708.03640](#), score 0.596
- [arXiv:quant-ph0205159](#), score 0.595
- [arXiv:1801.03283](#), score 0.594
- [arXiv:1612.00682](#), score 0.593
- [arXiv:1506.05598](#), score 0.592

14.23 Fock space

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.86. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Fock space is the occupation-number state space: the construction that replaces a fixed-particle Hilbert space by a direct sum over particle number.

Mechanism Reading

Fock space changes the carrier of the quantum state. Instead of describing one system in one Hilbert space, it builds sectors with zero, one, two, and more identical quanta, then imposes the bosonic or fermionic exchange rule. Creation and annihilation operators are the native coordinates of this page because they move the state between occupation sectors. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Fock space contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** Fock space, field configuration space, or a sector selected by charge, spin, momentum, statistics, or gauge data.
- **Operator or map.** Creation, annihilation, field, charge, spin, Hamiltonian, or scattering operators acting on the admissible sector.
- **Admissibility.** Statistics, gauge constraints, commutation or anticommutation rules, domain conditions, and sector labels decide which states are legal.
- **Readout.** Occupation number, charge, spin, momentum, energy, correlation function, cross-section, or scattering amplitude.
- **Check.** The field description must preserve the relevant observables under changes of representation and reduce to the expected particle or quasiparticle limit when that limit exists.

Topic Equations

Topic-specific constructor: the equations express variable particle number, exchange symmetry, and occupation-number readout.

$$\begin{aligned}\mathcal{F}_{\pm}(\mathcal{H}) &= \bigoplus_{n=0}^{\infty} \mathcal{S}_{\pm} \mathcal{H}^{\otimes n} \\ [a_i, a_j^{\dagger}]_{\mp} &= \delta_{ij}, \quad [a_i, a_j]_{\mp} = 0 \\ N &= \sum_i a_i^{\dagger} a_i, \quad N |n_1, n_2, \dots\rangle = \left(\sum_i n_i \right) |n_1, n_2, \dots\rangle\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.44, transport flow=0.09, constraint closure=0.06

- [arXiv:2308.15676](#), score 0.548
- [arXiv:2108.07838](#), score 0.539
- [arXiv:1612.00682](#), score 0.533
- [arXiv:0809.5271](#), score 0.528
- [arXiv:2105.11733](#), score 0.520

14.24 Quantum spacetime

Mechanism-tree role. Many-mode extension: fields, particles, and scaling; assignment score 0.85. **Dominant evidence signal.** incompatible questions. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum spacetime can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum spacetime is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is non-commuting compatibility limits, state

Quantum Mechanism Frame

- **Role.** Quantum spacetime contributes a geometry or holographic realization role to the quantum construction.
- **Placement.** This page is read first as a many-mode or field-realization move: it extends the state and operator construction beyond a single-particle carrier.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

commutator incompatibility=0.26, transport flow=0.26, constraint closure=0.21

- [arXiv:1604.06537](#), score 0.500
- [arXiv:1706.03846](#), score 0.489
- [arXiv:1604.05385](#), score 0.478
- [arXiv:2111.12617](#), score 0.476
- [arXiv:1708.03640](#), score 0.470

Chapter 15

Protocol layer: engineered transformations

Quantum computing, channels, circuits, algorithms, networks, sensors, and error correction turn the same formal machinery into controlled sequences of operations.

15.1 Why This Branch Exists

Quantum information is the engineering layer: the same state-operator-readout machinery becomes a controlled sequence of transformations.

15.2 Page Map

Page	Dominant evidence signal	Score
Quantum information science	observables and spectra	1.42
Quantum network	observables and spectra	1.42
Quantum algorithm	observables and spectra	1.42
Quantum error correction	observables and spectra	1.41
Quantum channel	observables and spectra	1.40
Quantum logic gate	observables and spectra	1.39
Quantum neural network	observables and spectra	1.39
Quantum circuit	observables and spectra	1.39
Quantum information	observables and spectra	1.38
Quantum computing	observables and spectra	1.20
Quantum key distribution	observables and spectra	0.81
Quantum sensor	observables and spectra	1.39
Quantum teleportation	observables and spectra	1.39
Quantum bus	observables and spectra	1.24
Quantum image processing	observables and spectra	1.23
Quantum finite automaton	state evolution	1.15
Quantum cryptography	observables and spectra	1.13
Quantum programming	observables and spectra	0.83

15.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

15.4 Quantum information science

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.42. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum information science can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum information science is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through wave function or superposition, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum information science contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: wave function and superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.

- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.35, transport flow=0.13, discrete protocol=0.10

- [arXiv:1604.05385](#), score 0.561
- [arXiv:2308.15676](#), score 0.535

- [arXiv:2105.11733](#), score 0.533
- [arXiv:0809.5271](#), score 0.531
- [arXiv:cond-mat0108470](#), score 0.530

15.5 Quantum network

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.42. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum network can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum network is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum network contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.

- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.32, transport flow=0.22, constraint closure=0.13

- [arXiv:1604.05385](#), score 0.577
- [arXiv:0912.2823](#), score 0.574
- [arXiv:0805.4565](#), score 0.555
- [arXiv:1708.03640](#), score 0.551
- [arXiv:quant-ph0205159](#), score 0.551

15.6 Quantum algorithm

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.42. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum algorithm can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum algorithm is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or superposition, Hamiltonian or matrix, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum algorithm contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state and superposition. Context/domain terms: domain.
- **Operator or map.** Operator terms: Hamiltonian, matrix, or unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.32, transport flow=0.22, constraint closure=0.16

- [arXiv:0912.2823](#), score 0.576
- [arXiv:1604.05385](#), score 0.576
- [arXiv:1708.03640](#), score 0.559
- [arXiv:quant-ph0205159](#), score 0.556
- [arXiv:1801.03283](#), score 0.556

15.7 Quantum error correction

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.41. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum error correction can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum error correction is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through superposition or wave function, Hamiltonian or unitary operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum error correction contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: superposition. Context/domain terms: preparation.
- **Operator or map.** Operator terms: Hamiltonian and unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.39, transport flow=0.22, constraint closure=0.15

- [arXiv:1604.05385](#), score 0.553
- [arXiv:1604.06537](#), score 0.525
- [arXiv:0912.2823](#), score 0.523
- [arXiv:1612.00682](#), score 0.516
- [arXiv:quant-ph0205159](#), score 0.513

15.8 Quantum channel

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.40. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum channel is the open-system protocol constructor: it maps input states to output states while preserving complete positivity and trace.

Mechanism Reading

A channel is the mechanism for noisy transformations, measurements with forgotten outcomes, and subsystem evolution. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum channel contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** Input and output density operators, possibly on different Hilbert spaces or subsystem carriers.
- **Operator or map.** A completely positive trace-preserving map, often represented by Kraus operators or by a Stinespring dilation.
- **Admissibility.** Complete positivity and trace preservation are the legal conditions; non-trace-preserving maps require an explicitly conditioned outcome.
- **Readout.** Output state, final POVM probabilities, fidelity, capacity, error rate, or recovered subsystem statistics.
- **Check.** The channel claim requires a map that stays positive under extension by an untouched reference system and preserves total probability.

Topic Equations

Standard constructor skeleton: completely positive trace-preserving map and readout.

$$\begin{aligned}\mathcal{E}(\rho) &= \sum_a K_a \rho K_a^\dagger \\ \sum_a K_a^\dagger K_a &= I \\ p(y) &= \text{Tr}(M_y \mathcal{E}(\rho))\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.21, constraint closure=0.14

- [arXiv:1604.05385](#), score 0.560
- [arXiv:1604.06537](#), score 0.555
- [arXiv:1506.05598](#), score 0.542
- [arXiv:0805.4565](#), score 0.542
- [arXiv:1708.03640](#), score 0.539

15.9 Quantum logic gate

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum logic gate can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum logic gate is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum logic gate contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: basis.
- **Operator or map.** Operator terms: matrix, Hamiltonian, or unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation

- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.43, transport flow=0.26, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.581
- [arXiv:1801.03283](#), score 0.552
- [arXiv:1612.00682](#), score 0.550
- [arXiv:quant-ph0205159](#), score 0.545
- [arXiv:1506.05598](#), score 0.539

15.10 Quantum neural network

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

a quantum neural network modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum neural network acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum neural network contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.22, constraint closure=0.11

- [arXiv:1604.05385](#), score 0.550
- [arXiv:2108.07838](#), score 0.545
- [arXiv:0805.4565](#), score 0.544
- [arXiv:1612.00682](#), score 0.537
- [arXiv:quant-ph0205159](#), score 0.530

15.11 Quantum circuit

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum circuit is the engineered-composition constructor: a finite sequence of admissible maps prepares, transforms, and measures a register.

Mechanism Reading

A circuit is the protocol layer of the same state-operator-readout machinery. Gates are controlled unitary or channel maps; measurement converts final states into output probabilities. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure.

Quantum Mechanism Frame

- **Role.** Quantum circuit contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** An input state, register, channel state, error syndrome, key, or controlled experimental configuration.
- **Operator or map.** An ordered sequence of gates, channels, measurements, corrections, encodings, or conditional maps.
- **Admissibility.** Each step must belong to the claimed map class: unitary, completely positive, trace-preserving, projective, conditional, or corrective.
- **Readout.** Output state, key, error rate, fidelity, channel capacity, algorithmic success probability, or sensor estimate.
- **Check.** Changing operation order, inserting classical controls, or replacing a quantum channel should identify which step carries the effect.

Topic Equations

Standard constructor skeleton: composed gates and final measurement.

$$\rho_{\text{out}} = U_m \cdots U_2 U_1 \rho_{\text{in}} U_1^\dagger U_2^\dagger \cdots U_m^\dagger$$
$$p(y) = \text{Tr}(M_y \rho_{\text{out}})$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.22, constraint closure=0.12

- [arXiv:1604.05385](#), score 0.552
- [arXiv:0912.2823](#), score 0.551
- [arXiv:1612.00682](#), score 0.537
- [arXiv:2108.07838](#), score 0.532
- [arXiv:quant-ph0205159](#), score 0.530

15.12 Quantum information

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.38. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum information can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum information is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or density matrix, observable operator or matrix, and eigenstate or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or non-commuting observables. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum information contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state and density matrix. Context/domain terms: preparation and basis.
- **Operator or map.** Operator terms: observable, matrix, or unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: uncertainty and non-commuting. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenstate and eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.26, constraint closure=0.21

- [arXiv:1604.06537](#), score 0.521
- [arXiv:0912.2823](#), score 0.501
- [arXiv:1604.05385](#), score 0.500
- [arXiv:1612.00682](#), score 0.493
- [arXiv:1801.03283](#), score 0.492

15.13 Quantum computing

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.20. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum computing can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum computing is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or superposition, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Quantum computing contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state and superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: Born rule.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.23, discrete protocol=0.20

- [arXiv:1604.05385](#), score 0.579
- [arXiv:1801.03283](#), score 0.510
- [arXiv:quant-ph0205159](#), score 0.507
- [arXiv:1708.03640](#), score 0.505
- [arXiv:1612.00682](#), score 0.503

15.14 Quantum key distribution

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 0.81. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum key distribution can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum key distribution is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and Mode or eigenstate. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Quantum key distribution contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: Mode and eigenstate. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.18, constraint closure=0.16

- [arXiv:1604.05385](#), score 0.583
- [arXiv:1612.00682](#), score 0.573
- [arXiv:quant-ph0205159](#), score 0.566
- [arXiv:1801.03283](#), score 0.563
- [arXiv:1506.05598](#), score 0.560

15.15 Quantum sensor

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum sensor can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum sensor is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through superposition or wave function, Hamiltonian or observable operator, and mode or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum sensor contributes an instrument-mediated readout role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: superposition. Context/domain terms: potential.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: projection.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: mode. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.19, constraint closure=0.12

- [arXiv:1604.05385](#), score 0.528
- [arXiv:0912.2823](#), score 0.528
- [arXiv:1604.06537](#), score 0.526
- [arXiv:quant-ph0205159](#), score 0.503
- [arXiv:1612.00682](#), score 0.501

15.16 Quantum teleportation

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum teleportation can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum teleportation is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or superposition, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum teleportation contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state and superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse and projection.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.38, transport flow=0.17, discrete protocol=0.11

- [arXiv:1604.05385](#), score 0.536
- [arXiv:0912.2823](#), score 0.497
- [arXiv:2308.15676](#), score 0.486
- [arXiv:1612.00682](#), score 0.484
- [arXiv:quant-ph0205159](#), score 0.474

15.17 Quantum bus

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.24. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum bus can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum bus is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through superposition or wave function, hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state

Quantum Mechanism Frame

- **Role.** Quantum bus contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.37, transport flow=0.21, boundary weak form=0.11

- [arXiv:0912.2823](#), score 0.574
- [arXiv:2108.07838](#), score 0.556
- [arXiv:0805.4565](#), score 0.546
- [arXiv:0908.0752](#), score 0.545
- [arXiv:1612.00682](#), score 0.545

15.18 Quantum image processing

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.23. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum image processing can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum image processing is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum image processing contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.15, commutator incompatibility=0.14

- [arXiv:2111.12617](#), score 0.559
- [arXiv:1604.06537](#), score 0.541
- [arXiv:2308.15676](#), score 0.509
- [arXiv:1506.05598](#), score 0.508
- [arXiv:1612.00682](#), score 0.502

15.19 Quantum finite automaton

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.15.
Dominant evidence signal. state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum finite automaton can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum finite automaton is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or state vector, matrix or unitary operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state evolution, normalization or

Quantum Mechanism Frame

- **Role.** Quantum finite automaton contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state and state vector. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: matrix and unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.28, constraint closure=0.24, spectral operator=0.19

- [arXiv:1604.05385](#), score 0.446
- [arXiv:1706.03846](#), score 0.442
- [arXiv:1708.03640](#), score 0.420
- [arXiv:0908.0752](#), score 0.410
- [arXiv:1801.03283](#), score 0.408

15.20 Quantum cryptography

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 1.13. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

A quantum cryptography can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum cryptography is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** Quantum cryptography contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state, wave function, or superposition. Context/domain terms: potential and basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.33, transport flow=0.26, constraint closure=0.20

- [arXiv:0912.2823](#), score 0.527
- [arXiv:1604.05385](#), score 0.526
- [arXiv:1801.03283](#), score 0.497
- [arXiv:quant-ph0205159](#), score 0.496
- [arXiv:1612.00682](#), score 0.495

15.21 Quantum programming

Mechanism-tree role. Protocol layer: engineered transformations; assignment score 0.83. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

Quantum programming can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, quantum programming is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, unitary operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Quantum programming contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an operation-sequence move: it specifies an ordered composition of allowed maps.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.40, transport flow=0.27, constraint closure=0.15

- [arXiv:1604.05385](#), score 0.480
- [arXiv:0805.4565](#), score 0.447
- [arXiv:1612.00682](#), score 0.442
- [arXiv:2108.07838](#), score 0.441
- [arXiv:quant-ph0205159](#), score 0.435

Chapter 16

Annotations: history, interpretations, and popular frames

Some pages help readers navigate the subject but do not form steps in the mechanism. They are kept as annotations so books, historical figures, interpretations, and popular frames do not distort the constructive tree.

16.1 Why This Branch Exists

These pages remain useful for orientation and are placed downstream of the construction steps. This prevents biographies, books, and interpretations from becoming false roots of the mechanism.

16.2 Page Map

Page	Dominant evidence signal	Score
Introduction to quantum mechanics	observables and spectra	0.72
Old quantum theory	observables and spectra	0.91
History of quantum mechanics (<i>annotation</i>)	observables and spectra	1.50
Introduction to Quantum Mechanics (book) (<i>annotation</i>)	state evolution	1.45
Quantum mind (<i>annotation</i>)	observables and spectra	1.44
History of quantum field theory (<i>annotation</i>)	state evolution	1.41
Erwin Schrödinger (<i>annotation</i>)	observables and spectra	1.39
Interpretations of quantum mechanics (<i>annotation</i>)	observables and spectra	1.38
Quantum mysticism (<i>annotation</i>)	incompatible questions	1.37
David Hilbert (<i>annotation</i>)	observables and spectra	1.36
Quantum Computing: A Gentle Introduction (<i>annotation</i>)	state evolution	1.30
QBism (<i>annotation</i>)	observables and spectra	1.43
Relational quantum mechanics (<i>annotation</i>)	observables and spectra	1.43

Page	Dominant evidence signal	Score
Modern Quantum Mechanics (<i>annotation</i>)	observables and spectra	1.23
The Quantum Universe (<i>annotation</i>)	state evolution	1.20
Quantum Theory: Concepts and Methods (<i>annotation</i>)	observables and spectra	1.10

16.3 Mechanism Pages

Each page below is read as a specialization of the compact quantum constructor. Topic-specific pages use explicit equations or overrides; core-derived pages use the branch-level equation form associated with their mechanism role.

16.4 Introduction to quantum mechanics

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 0.72. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

An introduction to quantum mechanics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, an introduction to quantum mechanics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or complementarity. In the source-evidence profile for this page, the strongest construction signal is

Quantum Mechanism Frame

- **Role.** Introduction to quantum mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.

- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: Born rule.
- **Admissibility.** Compatibility or closure terms: uncertainty and complementarity. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.29, transport flow=0.26, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.508
- [arXiv:1604.06537](#), score 0.496
- [arXiv:0912.2823](#), score 0.476
- [arXiv:1708.03640](#), score 0.462
- [arXiv:quant-ph0205159](#), score 0.462

16.5 Old quantum theory

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 0.91. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

Central Claim

An old quantum theory can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, an old quantum theory is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, matrix or Hamiltonian, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution,

Quantum Mechanism Frame

- **Role.** Old quantum theory contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.

- **Operator or map.** Operator terms: matrix and Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.34, transport flow=0.24, constraint closure=0.18

- [arXiv:1604.06537](#), score 0.564
- [arXiv:0912.2823](#), score 0.564
- [arXiv:1604.05385](#), score 0.563
- [arXiv:1708.03640](#), score 0.547
- [arXiv:1801.03283](#), score 0.546

16.6 History of quantum mechanics

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.50. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

A history of quantum mechanics can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a history of quantum mechanics is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum

Quantum Mechanism Frame

- **Role.** History of quantum mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.

- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.25, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.532
- [arXiv:1604.06537](#), score 0.531
- [arXiv:1612.00682](#), score 0.522
- [arXiv:quant-ph0205159](#), score 0.516
- [arXiv:1801.03283](#), score 0.514

16.7 Introduction to Quantum Mechanics (book)

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.45. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

An introduction to quantum mechanics (book) can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, an introduction to Quantum Mechanics (book) is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through Wave Function or state vector, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state

Quantum Mechanism Frame

- **Role.** Introduction to Quantum Mechanics (book) contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.

- **Carrier or domain.** State terms: Wave Function. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.28, constraint closure=0.24, spectral operator=0.18

- [arXiv:1706.03846](#), score 0.454
- [arXiv:1708.03640](#), score 0.427
- [arXiv:0806.4515](#), score 0.425
- [arXiv:0908.0752](#), score 0.418
- [arXiv:1801.03283](#), score 0.416

16.8 Quantum mind

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.44. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

a quantum mind modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum mind acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, controlled update protocol; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** Quantum mind contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: wave function and superposition. Context/domain terms: basis.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.31, transport flow=0.20, discrete protocol=0.17

- [arXiv:1604.05385](#), score 0.547
- [arXiv:0912.2823](#), score 0.510
- [arXiv:quant-ph0205159](#), score 0.483
- [arXiv:1612.00682](#), score 0.480
- [arXiv:0805.4565](#), score 0.480

16.9 History of quantum field theory

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.41. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

A history of quantum field theory can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a history of quantum field theory is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and spectrum or eigenvalue. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is state evolution,

Quantum Mechanism Frame

- **Role.** History of quantum field theory contributes a many-mode field or particle-realization role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.

- **Carrier or domain.** State terms: quantum state. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: Uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: spectrum. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
\rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
[O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.26, spectral operator=0.24, commutator incompatibility=0.22

- [arXiv:1604.06537](#), score 0.536
- [arXiv:1604.05385](#), score 0.517
- [arXiv:0912.2823](#), score 0.515
- [arXiv:1708.03640](#), score 0.511
- [arXiv:1801.03283](#), score 0.509

16.10 Erwin Schrödinger

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.39. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

an erwin Schrödinger modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

an erwin Schrödinger acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** Erwin Schrödinger contributes a lawful state-transport role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: wave function. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.

- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: Eigenvalue. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.32, transport flow=0.26, constraint closure=0.22

- [arXiv:1604.05385](#), score 0.519
- [arXiv:0912.2823](#), score 0.518
- [arXiv:1604.06537](#), score 0.516
- [arXiv:1801.03283](#), score 0.507
- [arXiv:1708.03640](#), score 0.500

16.11 Interpretations of quantum mechanics

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.38. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

an interpretations of quantum mechanics modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

an interpretations of quantum mechanics acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** Interpretations of quantum mechanics contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: wave function, wavefunction, state vector, quantum state, or superposition. Context/domain terms: domain.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: Born rule and collapse.

- **Admissibility.** Compatibility or closure terms: Complementarity. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenstate. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
\rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
[O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
\end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.26, transport flow=0.26, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.515
- [arXiv:0912.2823](#), score 0.514
- [arXiv:1604.06537](#), score 0.507
- [arXiv:1708.03640](#), score 0.479
- [arXiv:quant-ph0205159](#), score 0.475

16.12 Quantum mysticism

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.37. **Dominant evidence signal.** incompatible questions. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

a quantum mysticism modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum mysticism acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is non-commuting compatibility limits, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** Quantum mysticism contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.

- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

commutator incompatibility=0.27, transport flow=0.25, constraint closure=0.17

- [arXiv:2111.12617](#), score 0.373
- [arXiv:1604.06537](#), score 0.361
- [arXiv:1604.05385](#), score 0.349
- [arXiv:1706.03846](#), score 0.325
- [arXiv:2410.23449](#), score 0.319

16.13 David Hilbert

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.36. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

A david hilbert can be read as a quantum construction: the measurement basis and experimental arrangement fixes the admissible state space; the self-adjoint observable being asked of that state defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a david Hilbert is described by a prepared quantum state before the measurement. The physical question is represented by the self-adjoint observable being asked of that state; the experimental or mathematical setting is the measurement basis and experimental arrangement. The observable content is obtained from the observable's spectral projectors and the Born probabilities assigned to them. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest

Quantum Mechanism Frame

- **Role.** David Hilbert contributes a state-carrier role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: basis.

- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.42, transport flow=0.17, constraint closure=0.15

- [arXiv:0912.2823](#), score 0.528
- [arXiv:1604.06537](#), score 0.523
- [arXiv:1612.00682](#), score 0.518
- [arXiv:quant-ph0205159](#), score 0.505
- [arXiv:1801.03283](#), score 0.503

16.14 Quantum Computing: A Gentle Introduction

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.30. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

A quantum computing: a gentle introduction can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a quantum Computing: A Gentle Introduction is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through superposition or wave function, unitary operator or Hamiltonian, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is state

Quantum Mechanism Frame

- **Role.** Quantum Computing: A Gentle Introduction contributes an engineered operation-sequence role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.

- **Carrier or domain.** State terms: superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: unitary. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.27, constraint closure=0.25, spectral operator=0.13

- [arXiv:1706.03846](#), score 0.534
- [arXiv:1708.03640](#), score 0.510
- [arXiv:0908.0752](#), score 0.505
- [arXiv:1801.03283](#), score 0.490
- [arXiv:quant-ph0205159](#), score 0.487

16.15 QBism

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.43. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

QBism modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

QBism acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** QBism contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: wavefunction, quantum state, wave function, or superposition. Context/domain terms: measurement apparatus.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: Born rule and collapse.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.

- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, transport flow=0.26, constraint closure=0.19

- [arXiv:1604.05385](#), score 0.486
- [arXiv:1604.06537](#), score 0.471

- [arXiv:0912.2823](#), score 0.452
- [arXiv:quant-ph0205159](#), score 0.429
- [arXiv:1612.00682](#), score 0.428

16.16 Relational quantum mechanics

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.43. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

a relational quantum mechanics modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a relational quantum mechanics acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, controlled update protocol, state evolution; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** Relational quantum mechanics contributes a probability/readout role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: state vector, Wavefunction, or superposition. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: matrix. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenstate. These name the outcome labels, projectors, amplitudes, or records used for testing.

- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.30, discrete protocol=0.19, transport flow=0.17

- [arXiv:1604.05385](#), score 0.508
- [arXiv:0912.2823](#), score 0.453
- [arXiv:quant-ph0607206](#), score 0.435
- [arXiv:2306.13129](#), score 0.430
- [arXiv:1612.00682](#), score 0.424

16.17 Modern Quantum Mechanics

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.23. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

A modern quantum mechanics can be read as a quantum construction: the potential, domain, initial condition, or boundary condition fixes the admissible state space; the Hamiltonian, whose exponential gives unitary time evolution defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, a modern Quantum Mechanics is described by a wave function or density operator defined on the Hilbert space allowed by the system's domain. The physical question is represented by the Hamiltonian, whose exponential gives unitary time evolution; the experimental or mathematical setting is the potential, domain, initial condition, or boundary condition. The observable content is obtained from the eigenvalues and eigenfunctions of the relevant observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through commutator or uncertainty relation. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout,

Quantum Mechanism Frame

- **Role.** Modern Quantum Mechanics contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian. Protocol or update terms: unitary evolution, projection or measurement update, or path integral weighting.
- **Admissibility.** Compatibility or closure terms: commutator, uncertainty relation, or non-commuting transformations. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.

- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned}
 B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\
 \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\
 O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\
 [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis})
 \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.41, transport flow=0.27, constraint closure=0.21

- [arXiv:1612.00682](#), score 0.507
- [arXiv:1801.03283](#), score 0.505
- [arXiv:quant-ph0205159](#), score 0.504
- [arXiv:1708.03640](#), score 0.502
- [arXiv:1506.05598](#), score 0.498

16.18 The Quantum Universe

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.20. **Dominant evidence signal.** state evolution. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

The quantum universe can be read as a quantum construction: the chosen basis, pulse sequence, or measurement axis fixes the admissible state space; a Hamiltonian or unitary matrix rotating that state between preparation and measurement defines the transformation or question; and spectral projectors with the Born rule determine the recorded probability distribution.

Mechanism Reading

In quantum-mechanical terms, The Quantum Universe is described by a two-dimensional Hilbert space, usually written as a qubit state or a density matrix. The physical question is represented by a Hamiltonian or unitary matrix rotating that state between preparation and measurement; the experimental or mathematical setting is the chosen basis, pulse sequence, or measurement axis. The observable content is obtained from projectors onto the two eigenstates of the measured observable. In the local terminology of this topic, the same construction appears through quantum state or wave function, Hamiltonian or observable operator, and eigenvalue or energy level. Probabilities enter only after this spectral decomposition: the Born rule assigns weights to projectors, not to informal object names. When two observables have a non-zero commutator, no single basis diagonalizes both; the limitation is therefore a statement about jointly available spectra, not about detector imperfection. In this page the compatibility condition is expressed through uncertainty relation or commutator. In the source-evidence profile for this page, the strongest construction signal is state evolution, non-commuting

Quantum Mechanism Frame

- **Role.** The Quantum Universe contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: quantum state, wave function, or density operator. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: path integral.
- **Admissibility.** Compatibility or closure terms: uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

transport flow=0.25, commutator incompatibility=0.25, constraint closure=0.16

- [arXiv:1706.03846](#), score 0.533
- [arXiv:1604.06537](#), score 0.516
- [arXiv:2111.12617](#), score 0.515
- [arXiv:1708.03640](#), score 0.510
- [arXiv:0908.0752](#), score 0.501

16.19 Quantum Theory: Concepts and Methods

Mechanism-tree role. Annotations: history, interpretations, and popular frames; assignment score 1.10. **Dominant evidence signal.** observables and spectra. **Lagrangian construction road.** no Lagrangian road map available; road score 0.00.

This page is treated as historical, interpretive, or popular context rather than as a conceptual root.

Central Claim

a quantum Theory: Concepts and Methods modifies the interpretation of the probability/readout layer while preserving the formal quantum dynamics.

Mechanism Reading

a quantum Theory: Concepts and Methods acts on the readout layer of the quantum constructor. The formal ingredients remain the state assignment, the operator or measurement being applied, and the Born-rule map from projectors to probabilities. What changes is the status assigned to those ingredients: for this topic, the state or probability is treated through the agent, measurement context, or interpretive stance attached to the formalism. The page should therefore be read as a statement about the interpretation of state, probability, update, or recorded outcome while the Hamiltonian, spectral resolution, and commutator structure remain the formal reference layer. In the source-evidence profile for this page, the strongest construction signal is operator-to-spectrum readout, state evolution, normalization or admissibility; the strongest carrier signal is local notation, information profile, formula structure. Its constructive use is to identify which formal layer is being interpreted: state assignment, probability, update, readout, or ontology.

Quantum Mechanism Frame

- **Role.** Quantum Theory: Concepts and Methods contributes a topic-native constructor role to the quantum construction.
- **Placement.** This page is read first as an interpretive or historical move: it clarifies which formal layer is being discussed.
- **Carrier or domain.** State terms: wavefunction and quantum state. Context/domain terms: preparation condition, measurement setup, or potential or domain.
- **Operator or map.** Operator terms: Hamiltonian, observable operator, or generator. Protocol or update terms: collapse.
- **Admissibility.** Compatibility or closure terms: incompatible and uncertainty. These determine which questions, states, or updates are legal.
- **Readout.** Readout terms: eigenvalue, energy level, or measurement outcome. These name the outcome labels, projectors, amplitudes, or records used for testing.
- **Check.** A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.

Topic Equations

$$\begin{aligned} B &\longmapsto \rho_B \quad (\text{context specifies an admissible state}) \\ \rho_t &= U_t \rho_B U_t^\dagger \quad (\text{unitary evolution from preparation to readout}) \\ O &= \sum_i \lambda_i P_i, \quad p_i = \text{Tr}(P_i \rho_t) \quad (\text{spectral probability measure}) \\ [O_1, O_2] &\neq 0 \quad (\text{incompatible observables: no common sharp basis}) \end{aligned}$$

What Remains Stable

- the rule connecting prepared states, observables, and spectral probability measures across wave, matrix, path-integral, circuit, or field notation
- the operator-to-spectrum relation: admissible observations are represented through eigenvalues, projections, modes, or outcome channels
- the dependence of admissible readout on measurement context or boundary condition
- the non-commuting compatibility structure, which survives changes of representation

What Changes With Realization

- the name of the carrier: particle, wave, field, qubit, or excitation
- where time dependence is represented: on the state, on the operator, or in a path weight
- the coordinate system, basis, or geometric picture used to display the same relation
- the physical implementation of detector, boundary, preparation, or readout

Checks

- A concrete transfer target is a material, biological, or collective system with a state, a transformation, and a spectral or categorical readout, but without a tested incompatibility relation.
- The validation criterion is that varying the context changes the admissible readout while the transformation law remains identifiable; shuffled or erased contexts should weaken the effect.

Evidence Profile

spectral operator=0.36, transport flow=0.26, constraint closure=0.20

- [arXiv:1604.05385](#), score 0.485
- [arXiv:1604.06537](#), score 0.477
- [arXiv:1612.00682](#), score 0.452
- [arXiv:quant-ph0205159](#), score 0.452
- [arXiv:1801.03283](#), score 0.451

Source Boundary

The source pages are stored under `discoveries/morphwiki_quantum/pages`. The mechanism tree is stored as `discoveries/morphwiki_quantum/quantum_mechanism_tree.json`. ArXiv links are evidence pointers, not claims that any individual paper proves the whole branch.